The torque transfer coefficient in DNA under torsional stress

Alexey K. Mazur*

UPR9080 CNRS, Université Paris Diderot, Sorbonne Paris Cité, Institut de Biologie Physico-Chimique, 13, rue Pierre et Marie Curie, Paris,75005, France.

In recent years, significant progress in understanding the properties of supercoiled DNA has been obtained due to nanotechniques that made stretching and twisting of single molecules possible. Quantitative interpretation of such experiments requires accurate knowledge of torques inside manipulated DNA. This paper argues that it is not possible to transfer the entire magnitudes of external torques to the twisting stress of the double helix, and that a reducing torque transfer coefficient (TTC<1) should always be assumed. This assertion agrees with simple physical intuition and is supported by the results of all-atom molecular dynamics (MD) simulations. According to MD, the TTCs around 0.8 are observed in nearly optimal conditions. Reaching higher values requires special efforts and it should be difficult in practice. The TTC can be partially responsible for the persistent discrepancies between the twisting rigidity of DNA measured by different methods.

PACS numbers: 87.14.gk 87.15.H- 87.15.ap 87.15.ak

The twisting stiffness is a unique property of DNA that makes possible its supercoiling, which is essential for genome compaction and regulation [1]. At present, the mechanics of supercoiled DNA is frequently studied by using nanotechniques that offer a means to stretch and twist single molecules [2–4]. Due to its remarkable conceptual simplicity this method has gained broad recognition. The effects of external twisting depend upon the torsional rigidity of DNA and the values of applied torques. Accurate knowledge of these quantities is necessary for interpretation of experimental data. The recent methodological advances made twisting of single DNA with external torques of known magnitude [5, 6] possible. However, one question has never been asked: Is it evident that the twisting stress created inside DNA always corresponds to the external torque measured from outside?

A simple example shown in Fig. 1 demonstrates that the answer to the above question can be negative. Consider the left panel. An elastic rod is fixed on a solid base, with a constant torque τ applied to its top, which causes a deviation of the equilibrium twist angle $\Delta \Phi_{\tau} = \Phi_{\tau} - \Phi_0$. For simplicity, assume that the bending rigidity is very high and bending deformations can be neglected. If the torque axis is deviated by angle θ only the vertical torque projection works and the twisting stress inside the cylinder as well as $\Delta \Phi_{\tau}$ decrease as $\cos \theta$. The above effect persists with the addition of other degrees of freedom. Fig. 1 illustrates this for Brownian dynamics (BD) of a discrete wormlike rod (WLR) model where bending, twisting and translational motions of DNA are reproduced in agreement with experiments [8]. In the harmonic approximation the torsional part of the free energy is

$$\frac{U_{\tau}(\Phi)}{kT} = \frac{l_t}{2L} \left(\Phi - \Phi_{\tau}\right)^2 \tag{1}$$

where L is the DNA length, k is Boltzmann's constant,




FIG. 1: (Color online) (Left panel) An elastic rod is fixed on a solid base. A twisting torque is applied at the top with the torque axis inclined by angle θ form the vertical axis. (Right panel) A discrete WLR model of a DNA fragment of three base pairs. The external torques are applied to terminal bases under angle θ with respect to the chain axis. (Middle panel) Twisting caused by external torques in BD of a 14mer DNA fragment with elasticity corresponding to MD data [7]. The torques were applied with $\theta = 0^{\circ}$ (black dots), 30° (red squares), and 60° (blue circles). The black dashed line displays the theoretical dependence for the input value of l_t .

and T is the temperature. The twisting rigidity of DNA is characterized by parameter l_t called the torsional persistence length. In this case

$$\Delta \Phi_{\tau} = \frac{\tau L}{kT l_t} \tag{2}$$

and the probability distribution of torsional fluctuations is Gaussian

$$P_{\Phi} \sim \exp\left[-\frac{l_t}{2L} \left(\Phi - \Phi_{\tau}\right)^2\right].$$
 (3)

The right panel of Fig. 1 shows a coarse-grained WLR representation of a DNA trimer. Each base pair is modeled by a rigid composite bead of four particles O, X,

Y, and Z that define a local Cartesian frame. The OZaxis is constrained to pass through the center of the following bead in the chain. Other degrees of freedom are restrained by harmonic nearest-neighbor potentials adjusted to provide the desired macroscopic rigidity for bending and twisting [8]. The twisting torques are applied to terminal composite beads by using the earlier algorithm [9], with a small modification to deviate the torque vector by a fixed angle θ with respect to the local OZ axes. The external torque at one end is compensated by reactions at the opposite end so that the integral external force and torque on the molecule are always zero. This can be viewed as if a demon in the moving global frame twists one end of the molecule and holds the opposite end so that the overall translation and rotation are not perturbed. The results of BD simulations of a 14mer DNA fragment are shown in the middle panel. As expected, the effective torque decreases approximately as $\cos \theta$. The analytical linear dependence corresponding to the true l_t value is reproduced in BD only with $\theta=0$.




FIG. 2: (Color online) The normalized probability distributions of twisting fluctuations in DNA obtained by all-atom MD simulations [7]. The Φ angles are measured directly between the terminal base pair frames; therefore, $\Phi = 0$ for one helical turn. The CG dodecamer was considered in relaxed state ($\tau = 0$) and under steady torsional stress induced by external torques of $\tau = -80$ and +60 pN nm, respectively. The dashed lines show the corresponding Gaussian distributions [Eq. (3)] for the computed values of Φ_{τ} and l_t . The solid lines display the analytical distributions for the best-fit anharmonic torsional potential

It might seem that the above example is irrelevant because in experiments the external torques are always applied along the stretched DNA double helix. The problem is that local twisting axes in real DNA may be inclined with respect to the helix direction. This issue surfaced unexpectedly in the recent study of elastic properties of strained DNA by all-atom molecular dynamics (MD) [7, 10]. In these computations steady twisting torques were applied to two short DNA fragments that I denote here as TA and CG (because of the ATand GC-alternating sequences, respectively). The classical all-atom MD produce the probability distribution of twisting P_{Φ} . The three distributions shown in Fig. 2 were obtained for dodecamer CG (one helical turn) with contrasting values of the external torque. From these data, the l_t value can be extracted by using either Eq.

(2) or (3) and the resulting two values should be identical. However, according to Eq. (3) the variance of the distribution should be constant whereas in Fig. 2 it evidently changes with τ ; that is, the harmonic approximation breaks down. Therefore, a broader issue should be considered, namely, whether or not these MD results agree with statistical mechanics of an anharmonic elastic rod. To answer this question we look for an anharmonic potential that would consistently describe the MD data. Assuming that $\Phi_{\tau} = 0$ for zero load the free energy is

$$U_{\tau}(\Phi) = -\tau \Phi + U(\Phi),$$

where $U(\Phi)$ is the torsional potential of bare DNA, therefore,

$$\tau = \frac{dU}{d\Phi}\Big|_{\Phi_{\tau}}.$$
(4)

According to Fig. 2 all the probability distributions are nearly Gaussian; that is, in the vicinity of Φ_{τ} the harmonic approximation is valid and one can use the Taylor series to obtain

$$l_t(\tau) = \frac{L}{kT} U''(\Phi_\tau).$$
(5)

Let us look for a polynomial form of $U(\Phi)$. We choose U(0)=0; in this case the first term is quadratic. For $l_t(\tau)$ not to be constant the polynomial should be of the third degree or higher, for instance,

$$U(\Phi) = q_1 \frac{1}{2} \Phi^2 + q_2 \frac{1}{6} \Phi^3 + q_3 \frac{1}{24} \Phi^4.$$
 (6)

In this case Eq. (4) and (5) give

$$\tau = q_1 \Phi_\tau + q_2 \frac{1}{2} \Phi_\tau^2 + q_3 \frac{1}{6} \Phi_\tau^3 \tag{7}$$

$$l_t = \frac{L}{kT} \left(q_1 + q_2 \Phi_\tau + q_3 \frac{1}{2} \Phi_\tau^2 \right).$$
 (8)

MD provides Φ_{τ} and l_t measured for several values of τ . Coefficients q_i are found by the least squares minimization of the discrepancy between the left- and right-hand parts of Eq. (7) and (8). The results of this fitting are shown in Fig. 3 by dashed red lines. It is seen that they significantly deviate from the MD points. The deviations are clearly systematic and they go far beyond the error ranges. This situation does not change with increased degree of the polynomial. Note that in all plots the deviations grow with $|\tau|$. It turned out that the quality of fitting is radically improved if a torque transfer coefficient (TTC) q_0 is added to Eq. (7) as follows:

$$q_0\tau = q_1\Phi_\tau + q_2\frac{1}{2}\Phi_\tau^2 + q_3\frac{1}{6}\Phi_\tau^3,\tag{9}$$

with q_0 optimized together with other q_i . In this case, the agreement within the range of statistical errors is easily

reached for polynomials of the fourth degree and higher. Interestingly, the optimized TTC values are similar for CG and TA (0.81 and 0.79, respectively) and they vary by only a few percent when the degree of the polynomial is increased up to six. The quality of fitting is illustrated by the solid traces in Fig. 2 and 3. The agreement is quite good, notably, small deviations of P_{Φ} from Gaussians in Fig. 2 are well reproduced. These subtleties have low statistical weight and actually do not affect the values of Φ_{τ} and l_t involved in the fitting. One can conclude that the MD results qualitatively and quantitatively agree with statistical mechanics, but only if the occurrence of the TTC $q_0 < 1$ in Eq. (9) can be reasonably explained.




FIG. 3: (Color online) Fitting of analytical models to the results of all-atom MD simulations of two DNA fragments (TA and CG) [7]. Error bars show statistical errors. In the upper two panels the symbol size corresponds to maximal errors. The dashed red lines show the results of fitting for a polynomial torsional potential without a TTC applied. The solid black lines show best fit dependences for a fourth order polynomial with optimized TTC.

The difference between Eq. (7), and (9) indicates that good fitting requires that the q_i values in Eq. (7) and (8) differ by a constant factor. Therefore, the scaling might be transferred from Eq. (9) to Eq. (8), which would give equally good fitting. At first sight, this seems reasonable because the scaling can be interpreted as a correction of the uncertainty in the L/T value. Actually, however, this factor is used in computing l_t from MD data and a correcting scaling should be applied simultaneously to both sides of Eq. (8). Therefore, the occurrence of q_0 can be reasonably explained only in Eq. (9) by assuming that a fraction of external torque applied to DNA is somehow lost.

The possible origin of incomplete torque transfer was shown in Fig. 1 above. The DNA can respond to external torques by either twisting or bending or both. The torque fraction that causes twisting varies with angle θ whereas the remaining part is largely canceled by auxiliary reaction forces. Such forces are always necessary. In the left panel of Fig. 1 and in single molecule experiments the reaction forces maintain DNA straight and fixed in place. In computer simulations, they are used to zero the integral external force and torque that otherwise would accelerate the overall translation and rotation [9]. All these reactions do not affect torsional fluctuations, but effectively damp all other deformations, which is where a part of the external torque goes.

The orientation of the twisting axis in real DNA is not known. The external torques in MD are applied to terminal bases, with the torque axes orthogonal to the base plane [9]. On closer inspection, however, one notices that this choice may not be optimal. There are other distinguished directions, for instance, the helical axis or the perpendicular to the base-pair plane. However, these directions are well defined and coincident only in the ideal B-DNA model. In reality, bases are not perpendicular to the helix direction and paired bases are not co-planar, therefore, the definition of the base pair plane as well as the helical axis is a matter of convention. Moreover, local twisting occurs due to many rotatable covalent bonds and it is possible that the transfer of the torsional stress to DNA depends upon specific orientations of these bonds near the point where the torque is applied.

To get an idea of the situation in real DNA, I tried to vary the θ angle in MD similarly to the BD simulations presented in Fig. 1. The protocols of MD simulations were the same as before [7, 10]. The external torque is applied by using a Cartesian frame rigidly attached to the base [9]. The axes of this frame approximately correspond to the standard convention (see Fig. 4), and by default the torque is directed along the Z axis. Unlike the WLR model, the all-atom DNA does not have cylindrical symmetry, therefore with $\theta \neq 0$, we should also check different torque azimuths. The natural qualitatively distinct directions of deviation correspond to XZ and YZplanes. With $\theta \neq 0$ in the XZ plane, the torque vector is inclined toward the major $(\theta < 0)$ or minor $(\theta > 0)$ groove. In the YZ plane, the torque can be inclined toward the backbone of the same ($\theta < 0$) or the opposite $(\theta > 0)$ base. The magnitude of the external torque was fixed at ± 40 pN nm. A relatively large value is required to obtain variations beyond the range of statistical errors. Earlier studies showed that, in MD, the properties of short DNA fragments change smoothly in this range of torques [7, 10].

The results shown in Fig. 4 confirm that the effect is also significant in the all-atom MD. Computed points are compared with theoretical dependences of the form $\delta \cos(\theta - \theta_s)$, with δ and θ_s adjusted manually. Angle θ_s specifies the softest local twisting axis. Parameter δ affects simultaneously the vertical displacement and the




FIG. 4: (Color online) The upper panel displays the orientation of the principal local axes according to the standard convention [11]. The Z axis is directed along the double helix. The X axis looks towards the major groove. The Y axis is directed along the base pair so that the three axes form a right-hand triple. The torque vector is inclined by a variable angle θ with respect to the Z axis. The lower panel displays the results of all-atom MD simulations of DNA subjected to external torques $\tau = \pm 40$ pN nm applied to terminal bases. Red open circles show the results for the torque vector inclined in the XZ plane (toward the grooves), with $\theta > 0$ corresponding to deviations toward the minor groove. Similar data for the YZ plane (deviations toward the backbones) are shown by blue open squares. In this case, the positive θ values correspond to deviations toward the opposite base. The solid lines display the plots of $\delta \cos(\theta - \theta_s)$, with δ and θ_s fit manually to match the computed points.

inclination of the curves, i.e., the fitting is constrained. The agreement is relatively good for twisting and less so for untwisting. In both planes the softest axes are found close to, but not exactly perpendicular to the base. For deviations in the YZ plane the optimal θ_s value is about -5° for both signs of the torque. In the XZ plane the softest twisting direction is observed at $\theta_s = +10^\circ$ and $+20^{\circ}$ for twisting and untwisting, respectively. Therefore, the all-azimuth softest axis may be inclined by 20- 25° toward the minor groove, which would correspond to a TTC value around 0.9. Taking into account the details of the original algorithm the TTC can be reduced to 0.87 [12]. The remaining discrepancy with the best fit value $q_0 \approx 0.8$ is not negligible. Moreover, when the twisting is measured by summing the local twist angle along the chain [7] the optimized TTC values decrease to 0.72 and 0.76 for TA and CG fragments, respectively. This residual discrepancy can be attributed to conformational fluctuations. In the ideal WLR model and BD simulations shown in Fig. 1, the fixed angle θ is measured between the torque and the axis of local twisting.

In contrast, in Fig. 4 and MD simulations, the external torque is applied with θ fixed with respect to the base plane whereas the local twisting axis may fluctuate with the DNA structure. As a result, even in an optimal orientation, the torque is not strictly collinear with the twisting axis, and this should further reduce the TTC.

The foregoing results prove that it is difficult, if ever possible, to transfer the desired value of external torque to real double helical DNA. This conclusion should also be valid for nanomanipulations with isolated DNA molecules. The samples used in experiments commonly involve a few kilobase pairs of nearly random DNA. Earlier studies indicate that the WLR model is valid for any integral number of helical turns starting from one [8, 13]; therefore, one helical turn used in MD already involves all essential properties and the conclusions obtained should be invariant to length scaling. In experiments the double helix is attached to macroscopic beads and extended by a stretching force. The torques are applied to the beads in different ways and transferred to DNA through a number of point contacts. However, any multi-point interaction can be decomposed to individual components, each characterized by a specific TTC. The quasi-randomness of natural DNA sequences and the possible sequence dependence cannot smooth out the effect because the TTC never exceeds 1. Therefore, the induced twisting is always downgraded.

The TTC can explain some discrepancies in the recent experimental measurements of the torsional rigidity of DNA. The value of l_t (or $C = kTl_t$) has been measured since the 1970s [14] because it plays a fundamental role in all manifestations of DNA supercoiling. Before the recent single-molecule studies, $C \approx 310 \text{ pN nm}^2$ was considered as a consensus value for random sequence DNA. This value reasonably agrees with some indirect estimates that used single-molecule data [15, 16], but it is at least 20-30% lower than in direct measurements using forced DNA twisting by calibrated external torques [5, 6]. Several pitfalls possibly responsible for such a difference have been discussed in the literature [5, 17]. Assuming the TTC values around 0.8, which may slightly vary between different experimental installations, all these data can be accounted for without additional assumptions. Only one observation cannot be explained, namely, the very high apparent torsional rigidity ($C = 440 \text{ pN nm}^2$) reported for equilibrium thermal torsional fluctuations of single stretched DNA [5]. I believe that some other factors could affect this result and it requires further consideration.

In summary, the present paper shows that because of the structural details of the double helix, it is not possible to transfer the entire magnitudes of external torques to the internal twisting stress of DNA, and that a reducing torque transfer coefficient should always be assumed. This assertion agrees with simple physical intuition and it is supported by the results of all-atom MD simulations. Theoretical estimates indicate that the TTCs around 0.8 should be observed for torques nearly parallel to the double helix. Reaching higher values requires special efforts that should take into account the details of the DNA structure and it should be very difficult for experiments with long molecules. The TTC explains some controversies in the experimental values of DNA twisting rigidity measured by different methods.

The author is grateful to Alex Vologodskii for useful discussions and comments to the paper. I also thank Zev Bryant for correspondence and valuable clarifications concerning the rotor bead tracking method.

* Electronic address: alexey@ibpc.fr

- A. V. Vologodskii and N. R. Cozzarelli, Annu. Rev. Biophys. Biomol. Struct. 23, 609 (1994).
- [2] S. B. Smith, L. Finzi, and C. Bustamante, Science. 258, 1122 (1992).
- [3] T. R. Strick, J. F. Allemand, D. Bensimon, A. Bensimon, and V. Croquette, Science 271, 1835 (1996).
- [4] M. D. Wang, H. Yin, R. Landick, J. Gelles, and S. M. Block, Biophys. J. 72, 1335 (1997).
- [5] Z. Bryant, M. D. Stone, J. Gore, S. B. Smith, N. R.

Cozzarelli, and C. Bustamante, Nature 424, 338 (2003).

- [6] S. Forth, C. Deufel, M. Y. Sheinin, B. Daniels, J. P. Sethna, and M. D. Wang, Phys. Rev. Lett. **100**, 148301 (2008).
- [7] A. K. Mazur, Phys. Rev. E 84, 021903 (2011).
- [8] A. K. Mazur, J. Phys. Chem. B **113**, 2077 (2009).
- [9] A. K. Mazur, J. Chem. Theory Comput. 5, 2149 (2009).
- [10] A. K. Mazur, Phys. Rev. Lett. 105, 018102 (2010).
- [11] W. K. Olson, M. Bansal, S. K. Burley, R. E. Dickerson, M. Gerstein, S. C. Harvey, U. Heinemann, X.-J. Lu, S. Neidle, Z. Shakked, et al., J. Mol. Biol. **313**, 229 (2001).
- [12] In the previous work [7, 10], the input torque vector was projected on the approximate DNA axis to avoid spurious bending deformations [9]. Analysis of resulting trajectories indicates that this gave a supplementary reduction factor of 0.96. The new data displayed in Fig. 1 and 4 were obtained without such reduction.
- [13] A. K. Mazur, Phys. Rev. Lett. 98, 218102 (2007).
- [14] P. J. Hagerman, Annu. Rev. Biophys. Biophys. Chem. 17, 265 (1988).
- [15] A. V. Vologodskii and J. F. Marko, Biophys. J. 73, 123 (1997).
- [16] C. Bouchiat and M. Mezard, Phys. Rev. Lett. 80, 1556 (1998).
- [17] B. S. Fujimoto, G. P. Brewood, and J. M. Schurr, Biophys. J. 91, 4166 (2006).