Bound state in continuum like solutions in one dimensional hetero-structures

O. PANELLA¹ and P. ROY^2

¹Istituto Nazionale di Fisica Nucleare, Sezione di Perugia, Via A. Pascoli, I-06123 Perugia, Italy^{*} ²Physics and Applied Mathematics Unit Indian Statistical Institute Kolkata, India (Dated: September 28, 2018)

Abstract

We examine the one dimensional Dirac equation with modulated or position dependent velocity. In particular, it is shown that using suitable velocity profiles it is possible to create bound state in continuum (BIC) like, as well as, discrete energy bound state solutions.

PACS numbers:

^{*}Electronic address: orlando.panella@pg.infn.it (**Corresponding Author**). ; Permanent Address: INFN Sezione di Perugia, Via A Pascoli, I-06123, Perugia, Italy. Phone: +39 075 5852762; Fax: +39 075 44666.

I. INTRODUCTION

Since the rise of graphene [1, 2] there have been a lot of studies on the application of Dirac equation in condensed matter systems. In such systems the effective low energy model of the quasi particles is described by the Dirac equation with a Fermi velocity v_F which is 1/300 times the velocity of light. The peculiar property of charge carriers in graphene is that they behave as relativistic massless fermions. Subsequently, there have been numerous studies on various theoretical as well as experimental aspects of graphene. In particular, confinement of the quasi particles using various types of magnetic and electrostatic fields have been studied [3]. Other types of physically interesting states like the quasi bound states [4] have also been investigated. Here our interest lies in finding bound states in the continuum (BIC states) as well as discrete energy states in one dimensional heterostructures governed by the Dirac equation. The BIC are bound states embedded in the continuum [5]. Such states have a long history [6] and some years back the possibility of detecting BIC in semiconductor heterostructures has also been pointed out [7]. Later, BIC has been detected in photonic systems [8, 9]. The possibility of creating BIC in graphene quantum dot structures has also been studied [10].

On the other hand, there have been a number of studies on observable effects of modulated velocity or position dependent velocity in one dimensional heterostructures as well as graphene [11–14]. In such a scenario, the Fermi velocity is different in parts of the material. Backscattering in such systems described by one dimensional Dirac equation has recently been investigated thoroughly in ref [11]. Here our objective is to examine the(1 + 1) dimensional Dirac equation with position dependent velocity. To be more specific, we shall investigate the existence of BIC like states as well as discrete energy bound states in the presence of velocity profiles which continuously depends on the position.

II. THE MODEL

The Hamiltonian of a quasi-particle is of the form

$$H = v_F \sigma_x p_x \tag{1}$$

where σ_x denotes the Pauli matrix. Now in the case of a heterostructure, it is quite possible that velocity may be different in two regions of the material. In other words, the velocity becomes space dependent. However, this can not be achieved by simply replacing v_F by $v_F(x)$ in Eq.(1) since it would render the Hamiltonian non Hermitian. To keep the Hamiltonian Hermitian if the velocity v_F becomes position dependent it is necessary to write it in the following way [11]

$$H = \sqrt{v_F(x)}\sigma_x p_x \sqrt{v_F(x)} \tag{2}$$

It is easy to see that in case the velocity is a constant, the Hamiltonian (2) reduces to the form (1). The Hamiltonian in Eq. (2) being hermitian we can also easily derive the associated continuity equation which expresses the conservation of probability:

$$\frac{\partial \rho}{\partial t} + \frac{\partial j_x}{\partial x} = 0 \tag{3}$$

where the current associated to the probability density $\rho = \psi^{\dagger}\psi = \psi_1^*\psi_1 + \psi_2^*\psi_2$ is given by:

$$j_x = \sqrt{v_F} \psi_1^* \sqrt{v_F} \psi_2 + \text{c.c.}$$
(4)

The Hamiltonian in Eq. (2) operates on two component spinors $\psi(x) = (\psi_1(x), \psi_2(x))^T$ and the coupled equations for the components can now be written as

$$\sqrt{v_F(x)} p_x \left[\sqrt{v_F(x)} \psi_2(x) \right] = E \psi_1(x)$$
(5)

$$\sqrt{v_F(x)} p_x \left[\sqrt{v_F(x)} \psi_1(x) \right] = E \psi_2(x) \tag{6}$$

Now introducing an auxiliary spinor $u_{1,2}(x) = \sqrt{v_F(x)} \psi_{1,2}(x)$, the above pair of equations may be written as

$$v_F p_x u_1 = E u_2 \tag{7}$$

$$v_F p_x u_2 = E u_1 \tag{8}$$

Let us now, for example, obtain the equation for the component u_1 by eliminating u_2 from Eqs.(7) and (8) :

$$v_F^2 \frac{d^2 u_1}{dx^2} + v_F v_F' \frac{du_1}{dx} = \epsilon^2 u_1, \quad \epsilon^2 = E^2/\hbar^2$$
(9)

where the prime indicates differentiation w.r.t x. Let us now apply a further transformation involving the independent variable as

$$z = \int^x \frac{1}{v_F(t)} dt \tag{10}$$

and obtain from Eq.(9)

$$\frac{d^2 u_1}{dz^2} + \epsilon^2 u_1 = 0 \tag{11}$$

Eq.(11) looks like a free particle equation. However, whether or not it is a free particle equation depends on the range of z which in turn depends on $v_F(x)$. We shall now examine the scenario with a couple of different choices of $v_F(x)$.

III. BIC LIKE SOLUTIONS

Here we shall examine the existence of BIC like solutions of Eq.(11) using different choices of the velocity $v_F(x)$.

Example 1.

Let us first consider

$$v_F(x) = v_0 \cosh^2(\alpha x), \quad v_0 \text{ is a constant} > 0$$
 (12)

It follows that for $\alpha \to 0$ we recover the constant velocity scenario. Then from (10) we find $-\frac{1}{\alpha v_0} \leq z = \frac{1}{\alpha v_0} \tanh \alpha x \leq \frac{1}{\alpha v_0}$ and consequently Eq.(11) does not represent a free particle. However, we choose a formal or unphysical solution [15] of Eq.(11), namely,

$$u_1 = \sin\left(\epsilon z\right) \tag{13}$$

Then the component $\psi_1(x)$ of the original problem is given by

$$\psi_1(x) = \operatorname{sech}(\alpha x) \, \sin\left(\frac{\epsilon}{\alpha v_0} \, \tanh \alpha x\right)$$
(14)

Clearly, the second part of $\psi_1(x)$ is an oscillatory function bounded between ± 1 while the first part is a function decreasing at both ends. Consequently, $\psi_1(x)$ as given by Eq.(14) is a normalizable solution for any value of the energy E and we conclude that Eq.(14) represents a BIC like solution. Now using the relation (7) the second component of the spinor can be found to be

$$\psi_2(x) = -i \operatorname{sech}(\alpha x) \cos\left[\frac{\epsilon}{\alpha v_0} \tanh(\alpha x)\right]$$
 (15)

Clearly both components $\psi_{1,2}$ are square integrable functions. One can indeed exactly evaluate the normalization constant N for the complete Dirac spinor. Choosing to normalize to 1 the total probability density $\rho = \psi^{\dagger} \psi$ we have:

$$\int_{-\infty}^{+\infty} dx \rho(x) = \int_{-\infty}^{+\infty} dx \left[|\psi_1(x)|^2 + |\psi_2(x)|^2 \right] \\ = \frac{2}{\alpha} N^2$$
(16)



FIG. 1: (Left panel) Plot of the component $\psi_1(x)$, see Eq. 14 for the following set of values for the parameters: $v_0 = 1, \epsilon = 10, \alpha = 0.5$ (solid curve), $\alpha = 1$ (dashed curve), $\alpha = 2$ (dot-dashed curve) (arbitrary units). (Right panel) Plot of the (normalized) probability density $\rho(x)$ for the same values of the parameters as in the left panel. We clearly see that increasing the parameter α increases the localization of the BIC state.

and the normalization constant is given by $N = \sqrt{\alpha/2}$. We note that the two components of our spinor solution contribute differently, in general, to the total probability density. In any case, both components $\psi_{1,2}(x)$ and consequently the Dirac spinor $\psi = (\psi_1, \psi_2)^T$ represent a BIC like solution. In Fig. 1 we have drawn a plot of $\psi_1(x)$ (left panel) and the normalized probability density $\rho(x) = \psi^{\dagger}\psi = \psi_1^*\psi_1 + \psi_2^*\psi_2$ (right panel) for a particular set of parameters in arbitrary units (see figure caption). It can be seen that the wavefunction oscillates with decreasing amplitude and eventually goes to zero. Also, the plot of the probability density (Fig. 1 right panel) shows that increasing the parameter α increases the localization of the BIC state.

We can also easily check that the states described by Eqs. (14,15) have a vanishing probability current density as it is expected for a bound state:

$$j_x = N\sqrt{v_0} \left[(i-i) \sin\left(\frac{\epsilon}{\alpha v_0} \tanh \alpha x\right) \cos\left(\frac{\epsilon}{\alpha v_0} \tanh \alpha x\right) \right] = 0.$$
(17)

Example 2.

Here we choose

$$v_F(x) = v_0 (1 + \alpha x^2)^2, \quad \alpha > 0$$
 (18)

Then it follows that

$$-\frac{\pi}{2\sqrt{\alpha}v_0} \le z = \frac{1}{2v_0} \left[\frac{x}{(1+\alpha x^2)} + \frac{1}{\sqrt{\alpha}} \arctan\sqrt{\alpha}x \right] \le \frac{\pi}{2\sqrt{\alpha}v_0}$$
(19)

Now proceeding as before the solutions $\psi_{1,2}(x)$ are found to be

$$\psi_1(x) = \frac{1}{1 + \alpha x^2} \sin\left[\frac{\epsilon}{2v_0} \left(\frac{x}{1 + \alpha x^2} + \frac{1}{\sqrt{\alpha}} \arctan\sqrt{\alpha}x\right)\right]$$
(20)

$$\psi_2(x) = -i\frac{1}{1+\alpha x^2} \cos\left[\frac{\epsilon}{2v_0} \left(\frac{x}{1+\alpha x^2} + \frac{1}{\sqrt{\alpha}} \arctan\sqrt{\alpha x}\right)\right]$$
(21)

As in the previous example both the components are square integrable functions and they contribute differently to the total probability density $\rho = \psi^{\dagger}\psi$. Normalizing to 1 the total probability density we find:

$$\int_{-\infty}^{+\infty} dx \rho(x) = \int_{-\infty}^{+\infty} dx \left[|\psi_1(x)|^2 + |\psi_2(x)|^2 \right] \\ = \frac{\pi}{\sqrt{\alpha}} N^2$$
(22)

where $N = \sqrt{\sqrt{\alpha}/\pi}$.

It is readily seen that the solutions (20) and (21) are valid for any value of ϵ , and hence for any value of the energy eigenvalue E. In Fig. 2 we have plotted the component wave function ψ_1 of Eq. (20) (left panel) and the normalized probability density (right panel) for the same set of parameters (in arbitrary units) as in Example 1. As in the first example, it is also seen here that the wave function oscillates with non constant amplitude before dying out. The probability density shown in the right panel again shows that with larger values of the parameter α there is stronger localization.

From these two examples it is clear that one may choose many different profiles for $v_F(x)$ which would produce a BIC like solution.

A. Bound state solutions

In both the examples above we have considered unphysical solutions of Eq.(11). It now remains to be seen what happens when a physical solution i.e., a solution which vanishes



FIG. 2: (Left panel) Plot of the component $\psi_1(x)$, see Eq. 20 relative to Example 2, for the following set of values for the parameters: $v_0 = 1, \epsilon = 10, \alpha = 0.5$ (solid curve), $\alpha = 1$ (dashed curve), $\alpha = 2$ (dot-dashed curve) (arbitrary units). (Right panel) Plot of the (normalized) probability density $\rho(x)$ for the same values of the parameters as in the left panel. We clearly see also in this second example that increasing the parameter α increases the localization of the BIC state.

at the boundaries is chosen. Here we shall only examine Example 1 in detail. In this case the problem is clearly that of a particle trapped in an infinite square well potential. The solutions of this problem are well known and are given by

$$u_1(z) = \sin\left[\frac{\pi\alpha v_0 n}{2}(z + \frac{1}{\alpha v_0})\right], \quad \epsilon_n^2 = \frac{\alpha^2 v_0^2 \pi^2 n^2}{4}, \quad n = 1, 2, 3, \dots$$
(23)

Hence the solutions of the original problem are given by

$$\psi_{1,n}(x) = \operatorname{sech}(\alpha x) \, \sin\left[\frac{\pi n}{2}(\tanh \alpha x + 1)\right], \quad E_{1,n} = \frac{\hbar \alpha v_0 \pi n}{2} \tag{24}$$

One may easily check that the solutions (24) are normalizable. Now again using relation (7) we obtain

$$\psi_{2,n}(x) = -i \operatorname{sech}(\alpha x) \cos\left[\frac{\pi n}{2}(\tanh \alpha x + 1)\right], \quad E_{2,n} = \frac{\hbar \alpha v_0 \pi n}{2}$$
(25)

It may be observed that the solution $\psi_{2,n}(x)$ represents the same energy level as $\psi_{1,n}(x)$. Therefore the complete solution is given by

$$\psi_n(x) = \sqrt{\frac{\alpha}{2}} \operatorname{sech}\alpha x \left(\begin{array}{c} \sin\left[\frac{\pi n}{2}(\tanh\alpha x + 1)\right] \\ -i \ \cos\left[\frac{\pi n}{2}(\tanh\alpha x + 1)\right] \end{array} \right), \quad E_n = \frac{\hbar\alpha v_0 \pi n}{2}$$
(26)

So, in conclusion, whether or not the solutions of the original problem would represent a BIC or bound state depends on whether we choose an unphysical or a physical solution of the transformed equation given by Eq.(11).

IV. CONCLUSION

Here we have examined the possibility of obtaining BIC as well as bound state solutions within the framework of one dimensional Dirac equation with a velocity which depends continuously on position. In particular, it has been shown that such solutions do exist for suitable velocity profiles. We feel it would be of interest to investigate BIC in the presence of a position dependent mass or a scalar potential as well looking for similar solutions in two dimensions.

Acknowledgments

One of us (P. R.) wishes to thank INFN Sezione di Perugia for supporting a visit during which this work was carried out. The Physics Department of the University of Perugia is also acknowledged for the kind hospitality.

- K. S. Novoselov, A. K. Geim, S. V. Morozov, D. Jiang, Y. Zhang, S. V. Dubonos, I. V. Grigorieva, and A. A. Firsov, Science **306**, 666 (2004).
- K. S. Novoselov, A. K. Geim, S. V. Morozov, D. Jiang, M. I. Katsnelson, I. V. Grigorieva, S. V. Dubonos, and A. A. Firsov, Nature 438, 197 (2005).
- [3] A. De Martino, L. Dell'Anna, and R. Egger, Phys. Rev. Lett 98, 066802 (2007)
 A. De Martino, Solid State Comm. 144, 547 (2007)
 L. Dell'Anna and A. De Martino, Phys. Rev. B79, 045420 (2009).
 - J. M. Pereira, F. M. Peeters and P. Vasilopoulos, Phys. Rev. B75, 125433 (2007)
 - T. K. Ghosh, J. Phys: Condens. Matter 21, 045505 (2009).
 - S. Kuru, J. M. Negro and L. M. Nieto, J. Phys: Condens.Matter 21, 455305 (2009).
- [4] P.G. Silvestrov, Phys.Rev.Lett 98, 016802 (2007)
 A. Matulis and F.M. Peeters, Phys.Rev B77, 115423 (2008)

M. R. Masir, A. Matulis, and F. M. Peeters, Phys. Rev. B79, 155451 (2009)

H. Chau Nguyen, M. Tien Hoang, and V. Lien Nguyen, Phys.Rev B79, 035411 (2009).

- [5] J. von Neumann and E. Wigner, Phys. Z. **30**, 465 (1929).
- [6] J. Pappademos, U. Sukhatme and A. Pagnamenta, Phys.Rev A48, 3525 (1993) and references therein.
- [7] F. Capasso, C. Sirtori, J. Faist, D.L. Sivco, S.G. Chu and A.Y. Cho, Nature 358, 5656 (1992).
- [8] D.C. Marinica and A.G. Borisov, Phys. Rev. Lett. 100, 183902 (2008).
- [9] E. N. Bulgakov and A. F. Sadreev, Phys. Rev. B 78, 075105 (2008).
- [10] J.W. Gonzalez, M. Pacheko, L. Rosales and P.A. Orellana, Euro.Phys.Lett 91, 66001 (2010).
- [11] N.M.R. Peres, J.Phys.Condens.Matter **21**, 095501 (2009).
- [12] P.M. Krăstajić and P Vasilopoulos, J.Phys.Condens.Matter 23, 135302 (2010).
- [13] A. Raoux, M. Polini, R. Asgari, A. R. Hamilton, R. Fazio, and A. H. MacDonald, Phys. Rev. B81, 073407 (2010).
- [14] A. Concha and Z. Tesanovic, Phys.Rev **B82**, 033413 (2010).
- [15] This solution does not satisfy the boundary condition at $\pm \frac{1}{\alpha v_0}$