

MULTIPLICITY OF THE LOWEST EIGENVALUE OF NON-COMMUTATIVE HARMONIC OSCILLATORS

FUMIO HIROSHIMA AND ITARU SASAKI

ABSTRACT. The multiplicity of the lowest eigenvalue E of the so-called non-commutative harmonic oscillator $Q(\alpha, \beta)$ is studied. It is shown that E is simple for α and β in some region.

1. DEFINITION AND MAIN RESULTS

Recently a special attention is paid to studying the spectrum of self-adjoint operators with *non-commutative* coefficients. It is considered not only as mathematics but also physics experiments. A historically important model is the Dirac operator, and the Rabi model and the Jaynes-Cumming model are prevalent in cavity QED. See [HH12] and references therein. The non-commutative harmonic oscillator is a quantum system defined by the Hamiltonian:

$$Q = Q(\alpha, \beta) = A \otimes \left(-\frac{1}{2} \frac{d^2}{dx^2} + \frac{1}{2} x^2 \right) + J \otimes \left(x \frac{d}{dx} + \frac{1}{2} \right), \quad (1.1)$$

where $A = \begin{pmatrix} \alpha & 0 \\ 0 & \beta \end{pmatrix}$, $J = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$, and $\alpha, \beta > 0$ parameters with $\alpha\beta > 1$. Operator Q defines a positive self-adjoint operator acting in the Hilbert space $\mathcal{H} = \mathbb{C}^2 \otimes L^2(\mathbb{R})$.

The non-commutative harmonic oscillator Q has been introduced by Parmeggiani and Wakayama [PW01, PW02a, PW02b, PW03], and the spectral property of Q is considered in [Par04, Par06, Par08a] from the pseudo-differential-calculus point of view. It can be seen that Q has purely discrete spectrum $\lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_n \leq \dots \uparrow \infty$, where the eigenvalues are counted with multiplicity. One can define the so-called spectral zeta function associated with Q as

$$\zeta_Q(s) = \sum_{n=1}^{\infty} \frac{1}{\lambda_n^s}.$$

When $\alpha = \beta$, Q is unitarily equivalent to the direct sum of harmonic oscillators, and $\lambda_{2m-1} = \lambda_{2m} = \sqrt{\alpha^2 - 1}(m + \frac{1}{2})$, and thus ζ_Q with $\alpha \neq \beta$ can be regarded as a q -deform of the Riemann zeta-function. Analytic properties of the spectral zeta-function

Date: May 20, 2018.

2000 Mathematics Subject Classification. 35P05, 35P15.

Key words and phrases. non-commutative harmonic oscillator, multiplicity, lowest eigenvalue.

is studied in [IW05a, IW05b, IW07, KW06, KW07, KY09]. Furthermore it is also known that the set of *odd* eigenvectors of non-commutative harmonic oscillator is deeply related to the set of some solutions of the Heun differential equation [IW05b, Och01, Och04]:

$$\frac{\partial^2}{\partial w^2} f + \left(\frac{1-n}{w} + \frac{-n}{w-1} + \frac{n+3/2}{w-a} \right) \frac{\partial}{\partial w} f + \frac{-(3/2)nw - q}{w(w-1)(w-a)} f = 0,$$

where $n \in \mathbb{N} \cup \{0\}$, $a \in \mathbb{C}$ with $|a| < 1$ and $q \in \mathbb{C}$.

In this paper we concentrate on the study of the lowest eigenvalue λ_1 of Q . We set

$$E = \lambda_1 \tag{1.2}$$

and $p = -id/dx$. In particular we are interested in determining the dimension of $\text{Ker}(Q - E)$. The eigenvector associated with the lowest eigenvalue is called the ground state. In the case of $\alpha = \beta$, as is mentioned above, Q can be diagonalized as $Q \cong \begin{pmatrix} h & 0 \\ 0 & h \end{pmatrix}$ with

$$h = \frac{1}{2}p^2 + \frac{\alpha^2 - 1}{2}x^2, \tag{1.3}$$

where \cong denotes the unitary equivalence. Then all the eigenvalues of $Q(\alpha, \alpha)$ are two-fold degenerate. In particular, its lowest eigenvalue

$$E_0 = \frac{1}{2}\sqrt{\alpha^2 - 1} \tag{1.4}$$

is two-fold degenerate. In the general case, $\alpha \neq \beta$, the so-called Ichinose-Wakayama bound is established in [IW07]:

$$\left(j - \frac{1}{2} \right) \min\{\alpha, \beta\} \sqrt{\frac{\alpha\beta - 1}{\alpha\beta}} \leq \lambda_{2j-1} \leq \lambda_{2j} \leq \left(j - \frac{1}{2} \right) \max\{\alpha, \beta\} \sqrt{\frac{\alpha\beta - 1}{\alpha\beta}} \tag{1.5}$$

for $j \in \mathbb{N}$. By this inequality we see that the multiplicity of E is at most two if $\beta < 3\alpha$ or $\alpha < 3\beta$.

Furthermore beyond above results one may expect that E is simple for $\alpha \neq \beta$. In [NNW02] it can be numerically shown that E is simple for $\alpha \neq \beta$ and, in [Par04] the simplicity is proven but only for sufficiently large $\alpha\beta$. It is then mentioned in [Par08b, 8.3 Notes] that the determining the multiplicity of the lowest eigenvalue should be explored.

In this paper we show that

- (a): E is at most twofold degenerate for $(\alpha, \beta) \in (2, \infty) \times (2, \infty)$,
- (b): E is simple for some region of α and β .

In order to prove (a), we apply the method in [Hir05], where the so-called pull-through formula [GJ68] is a key. The second result (b) consists of two estimates. The first is for large $|\beta - \alpha|$ and the second for small $|\beta - \alpha|$ but $\alpha \neq \beta$. The first case is proven in

a similar manner to (a) and the second by the regular perturbation theory of discrete spectrum.

Let $\mathcal{G} = \text{Ker}(Q - E)$ be the set of ground states. Let $L_+ \subset L^2(\mathbb{R})$ (resp. L_-) be the set of even functions (resp. odd functions). We define $\mathcal{H}_\pm = \mathbb{C}^2 \otimes L_\pm$. Since Q conserves the parity, Q is reduced by \mathcal{H}_\pm . Set $Q|_{\mathcal{H}_\pm} = Q_\pm$ and then $Q = Q_+ \oplus Q_-$.

Theorem 1.1. *Suppose that $(\alpha, \beta) \in (2, \infty) \times (2, \infty)$. Then $\dim \mathcal{G} \leq 2$ and $\mathcal{G} \subset \mathcal{H}_+$. I.e., the multiplicity of E is at most two and ground states are even functions.*

We can furthermore show that E is simple.

Theorem 1.2. *Suppose $\beta > \alpha > 2$ and*

$$\frac{1}{2} > \left(\frac{1}{2}\beta - E \right)^{-2} \frac{1}{\alpha^2 - 1} + \frac{1}{\alpha^2 - 1}. \quad (1.6)$$

Then E is simple.

Condition (1.6) includes the implicit value E . Let

$$E_{\text{upper}} = \frac{\sqrt{\alpha\beta}\sqrt{\alpha\beta-1}}{\alpha + \beta + |\alpha - \beta| \frac{(\alpha\beta-1)^{1/4}}{\sqrt{\alpha\beta}} \text{Re} \rho},$$

where $\rho^2 = \sqrt{\alpha\beta-1} - i$ with $\text{Re} \omega > 0$, i.e., $\text{Re} \rho = \sqrt{\frac{\sqrt{\alpha\beta}(\sqrt{\alpha\beta-1}+1)}{2}}$. Bound $E < E_{\text{upper}}$ holds. See [Par08b, Theorem 8.2.1]. Combining this with (1.6) we have the corollary:

Corollary 1.3. *Suppose $\beta > \alpha > 2$ and*

$$\frac{1}{2} > \left(\frac{1}{2}\beta - E_{\text{upper}} \right)^{-2} \frac{1}{\alpha^2 - 1} + \frac{1}{\alpha^2 - 1}. \quad (1.7)$$

Then E is simple.

Theorem 1.2 does not valid for (α, β) in a neighborhood of the diagonal line on $\alpha - \beta$ plane. See Figure 1. We can also however show that E is simple for α and β in a




FIGURE 1. The region satisfying (1.7)
(The case of $\alpha > \beta$ is also drawn.)

neighborhood of the diagonal line. We define g_1, \dots, g_4 by

$$g_1 = (\alpha - 1)^{-1} \left(3 + \frac{\sqrt{3}}{\sqrt{\alpha^2 - 1}} \right), \quad (1.8)$$

$$g_2 = \frac{\sqrt{\alpha^2 - 1}}{2|\sqrt{\alpha^2 - 1} - \lambda_2|} g_1^2 \quad (1.9)$$

$$g_3 = \frac{\alpha}{2\sqrt{\alpha^2 - 1}} \quad (1.10)$$

$$g_4 = \frac{(\sqrt{\alpha^2 - 1})^{3/2}}{4\alpha^{3/2}}. \quad (1.11)$$

Theorem 1.4. *Let $\varepsilon = \beta - \alpha$. Assume that $\beta > \alpha > 1$, $\sqrt{\beta^2 - 1} \leq 3\sqrt{\alpha^2 - 1}$ and $\varepsilon^2 g_2 < 1/2$. Let*

$$\kappa(\varepsilon) = E_0 g_1^2 + \varepsilon g_2 (E_0 g_1 + g_3 + g_4) + \varepsilon^2 2E_0 g_1^2 g_2 + \varepsilon^3 2E_0 g_1 g_2^2, \quad (1.12)$$

$$\ell(\varepsilon) = (1 - \varepsilon^2 g_2) \sqrt{1 - 2\varepsilon^2 g_2^2}. \quad (1.13)$$

Then

$$|\lambda_1 - \lambda_2| \geq \frac{2\varepsilon}{\ell(\varepsilon)} (g_4 - \varepsilon \kappa(\varepsilon)) \quad (1.14)$$




FIGURE 2. A wedge-shaped region included in the region satisfying $\varepsilon\kappa(\varepsilon) < g_4$

In particular when $\varepsilon\kappa(\varepsilon) < g_4$, E is simple.

Note that we know the bound $\lambda_2 \leq \frac{\beta}{2} \frac{\sqrt{\alpha\beta-1}}{\sqrt{\alpha\beta}}$ by the Ichinose-Wakayama bound. Then, the region of α, β satisfying $\varepsilon\kappa(\varepsilon) < g_4$, includes a wedge-shaped region illustrated in Figure 1, where we also drew the case of $\alpha > \beta$.

2. PROOFS OF THEOREMS

2.1. Proof of Theorem 1.1. In the following we omit the symbol \otimes for the notational simplicity, and we can suppose that $\alpha < \beta$ without loss of generality. Let

$$N = \frac{1}{2} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} (p^2 + x^2 - 1).$$

The spectrum of N is $\sigma(N) = \{0\} \cup \mathbb{N}$ and the multiplicity of each eigenvalues. Let $a = \frac{1}{\sqrt{2}}(x + ip)$ and $a^* = \frac{1}{\sqrt{2}}(x - ip)$. They satisfy canonical commutation relations $[a, a^*] = 1$ and $[a, a] = [a^*, a^*] = 0$, and we have $a^*a = N$

Proof of Theorem 1.1: In terms of a and a^* , the operator Q can be realized as

$$Q = a^*Aa + \frac{1}{2}A + \frac{1}{2}(aJa - a^*Ja). \quad (2.1)$$

Let $\varphi_g \in \mathcal{G}$. We have $(Q - E)a\varphi_g = [Q, a]\varphi_g$. Since $[Q, a] = -Aa + Ja^*$ by canonical commutation relation, we have $(Q - E)a\varphi_g = (-Aa + Ja^*)\varphi_g$ and then $(Q - E + A)a\varphi_g =$

$Ja^*\varphi_g$. Notice that $Q - E + A \geq \alpha > 0$. We have

$$a\varphi_g = (Q - E + A)^{-1}Ja^*\varphi_g. \quad (2.2)$$

Taking the norm on both sides above, we have

$$\|a\varphi_g\|^2 \leq \frac{1}{\alpha^2}\|a^*\varphi_g\|^2. \quad (2.3)$$

Since $\|a\varphi_g\|^2 = \langle \varphi_g, N\varphi_g \rangle$ and $\|a^*\Omega\| = \langle \varphi_g, N\varphi_g \rangle + \|\varphi_g\|^2$, we see that

$$\langle \varphi_g, N\varphi_g \rangle \leq \frac{1}{\alpha^2 - 1}\|\varphi_g\|^2. \quad (2.4)$$

Let P_Ω be the projection onto $\ker N = \ker a$. Note that $N + P_\Omega \geq 1$. Let $\mathcal{G} = \mathcal{G}_+ \oplus \mathcal{G}_-$, where $\mathcal{G}_\pm = \mathcal{G} \cap \mathcal{H}_\pm$. Let P^\pm be the projection onto \mathcal{G}_\pm . Then, by (2.4), we have

$$P^+P_\Omega P^+ \geq P^+(1 - N)P^+ \geq \left(1 - \frac{1}{\alpha^2 - 1}\right)P^+. \quad (2.5)$$

Taking the trace of both sides, we have

$$2 \geq \text{Tr}(P^+P_\Omega P^+) \geq \frac{\alpha^2 - 2}{\alpha^2 - 1}\text{Tr}P^+ \quad (2.6)$$

Thus we have the bound

$$\dim \ker P^+ \leq 2\frac{\alpha^2 - 1}{\alpha^2 - 2}. \quad (2.7)$$

Then the right hand side above is less than three for $\alpha > 2$. Then $\dim \mathcal{G}^+ \leq 2$. Similarly but replacing P^+ with P^- , we can also see that

$$P^-P_\Omega P^- \geq P^-(1 - N)P^- \geq \left(1 - \frac{1}{\alpha^2 - 1}\right)P^-. \quad (2.8)$$

Note that $P^-P_\Omega P^- = 0$, since P_Ω is the projection to the set of even functions. Then we have

$$0 = \text{Tr}(P^-P_\Omega P^-) \geq \frac{\alpha^2 - 2}{\alpha^2 - 1}\text{Tr}P^-. \quad (2.9)$$

In particular for $\alpha > \sqrt{2}$, the dimension of \mathcal{G}_- equal to zero. Then the theorem follows. \square

2.2. Proof of Theorem 1.2. In this subsection we show that the lowest eigenvalue is simple. The strategy is parallel with that of previous subsection but P^+ is replaced by a projection R with the dimension of $\text{Ran}R = 1$. Let $\sigma_1, \sigma_2, \sigma_3$ be the 2×2 Pauli matrices given by

$$\sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \quad (2.10)$$

Proof of Theorem 1.2: The Hamiltonian Q can be written in the form:

$$Q = \frac{1}{2}A(p^2 + x^2) + \frac{1}{2}\sigma_2(px + xp). \quad (2.11)$$

We set $M = \frac{1}{2}(1 + \sigma_3) = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$, $M^\perp = \frac{1}{2}(1 - \sigma_3) = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$ and $R = M^\perp P_\Omega$. Then we have

$$(Q - E)R\varphi_g = M^\perp[Q, P_\Omega]\varphi_g + [Q, M^\perp]P_\Omega\varphi_g. \quad (2.12)$$

The commutator $[Q, M^\perp]$ can be computed as

$$[Q, M^\perp] = -\frac{1}{2}(aa - a^*a^*)\sigma_1. \quad (2.13)$$

Thus we have

$$[Q, R] = M^\perp \frac{i}{2}\sigma_2[aa - a^*a^*, P_\Omega] + \frac{1}{2}a^*a^*\sigma_1P_\Omega, \quad (2.14)$$

where we used the fact that $aP_\Omega = 0$. Hence we have

$$\begin{aligned} \langle R\varphi_g, (Q - E)R\varphi_g \rangle &= \left\langle \varphi_g, \frac{i}{2}RM^\perp\sigma_2[aa - a^*a^*, P_\Omega]\varphi_g \right\rangle + \frac{1}{2}\langle \varphi_g, Ra^*a^*\sigma_1P_\Omega\varphi_g \rangle \\ &= \frac{i}{2}\langle \varphi_g, R\sigma_2a^2\varphi_g \rangle. \end{aligned} \quad (2.15)$$

On the other hand $R(Q - E)R = (\frac{1}{2}\beta - E)R^2$. Then (2.4) and $\|a^*R\varphi_g\| = \|R\varphi_g\|$ yield that

$$\left(\frac{1}{2}\beta - E\right) \|R\varphi_g\|^2 \leq \frac{1}{2}\|a^*R\varphi_g\| \|\sigma_2a\varphi_g\| \leq \frac{1}{\sqrt{\alpha^2 - 1}}\|R\varphi_g\| \|\varphi_g\|.$$

Therefore

$$\|R\varphi_g\|^2 \leq \left(\frac{1}{2}\beta - E\right)^{-2} \frac{1}{\alpha^2 - 1} \|\varphi_g\|^2.$$

Since $M + M^\perp = 1$, it holds that $P_\Omega M + R + N \geq 1$. Then, by using (2.4) we have

$$P(P_\Omega M)P \geq P(1 - P_\Omega M^\perp - N)P = \left(1 - \left(\frac{1}{2}\beta - E\right)^{-2} \frac{1}{\alpha^2 - 1} - \frac{1}{\alpha^2 - 1}\right) P$$

where $P = P^+ + P^-$ is the orthogonal projection onto \mathcal{G} . Taking the trace of both sides above, we have

$$1 \geq \left(1 - \left(\frac{1}{2}\beta - E\right)^{-2} \frac{1}{\alpha^2 - 1} - \frac{1}{\alpha^2 - 1}\right) \text{Tr}P,$$

and the theorem follows. \square

2.3. Proof of Theorem 1.4. Recall that $\varepsilon = \beta - \alpha$. In this section, we fix an arbitrary $\alpha > 1$ and set

$$Q(\alpha, \beta) = Q = Q(\varepsilon) = Q_0 + \varepsilon V, \quad (2.16)$$

where $Q_0 = Q(\alpha, \alpha)$ and $V = \frac{1}{2} \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} (p^2 + x^2)$.

Lemma 2.1. *For all $\Phi \in D(Q_0)$, it follows that*

$$\|V\Phi\| \leq (\alpha - 1)^{-1} \|Q_0\Phi\| + \frac{\sqrt{3}}{2} (\alpha - 1)^{-1} \|\Phi\|. \quad (2.17)$$

Proof. One can show that $\|(px + xp)u\|^2 \leq \|(p^2 + x^2)u\|^2 + 3\|u\|^2$. Since $\sigma_2^2 = 1$, we have

$$\begin{aligned} \|Q_0\Phi\| &\geq \alpha \left\| \frac{1}{2}(p^2 + x^2)\Phi \right\| - \left\| \frac{1}{2}(px + xp)\Phi \right\| \\ &\geq \alpha \left\| \frac{1}{2}(p^2 + x^2)\Phi \right\| - \frac{1}{2} \left(\|(p^2 + x^2)\Phi\|^2 + 3\|\Phi\|^2 \right)^{1/2}, \end{aligned}$$

and hence

$$\|Q_0\Phi\| \geq \frac{\alpha - 1}{2} \|(p^2 + x^2)\Phi\| - \frac{\sqrt{3}}{2} \|\Phi\|. \quad (2.18)$$

Noticing $\|V\Phi\| \leq \frac{1}{2} \|(p^2 + x^2)\Phi\|$ we have the bound (2.17). \square

By (1.5) or the sandwich estimate $Q(\alpha, \alpha) \leq Q(\alpha, \beta) \leq Q(\beta, \beta)$ we see bounds: $\frac{\sqrt{\alpha^2 - 1}}{2} \leq \lambda_1 \leq \lambda_2 \leq \frac{\sqrt{\beta^2 - 1}}{2}$ and $\frac{3}{2}\sqrt{\alpha^2 - 1} \leq \lambda_3$. When $\sqrt{\beta^2 - 1} \leq 3\sqrt{\alpha^2 - 1}$, Q has exactly two eigenvalues in the interval $\left[\frac{1}{2}\sqrt{\alpha^2 - 1}, \frac{1}{2}\sqrt{\beta^2 - 1} \right]$. Let C be the closed disk centered at $\frac{\sqrt{\alpha^2 - 1}}{2}$ with the radius $\frac{\sqrt{\alpha^2 - 1}}{2}$ in the complex plane:

$$C = \left\{ \frac{\sqrt{\alpha^2 - 1}}{2} + re^{i\theta} \in \mathbb{C} \mid 0 \leq r \leq \frac{\sqrt{\alpha^2 - 1}}{2}, 0 \leq \theta \leq 2\pi \right\}. \quad (2.19)$$

Thus Q has exactly two eigenvalues, λ_1 and λ_2 , inside of C .

Lemma 2.2. *It follows that $\|V(Q_0 - z)^{-1}\| \leq g_1$ for all $z \in \partial C$.*

Proof. By Lemma 2.1, we have

$$\|V(Q_0 - z)^{-1}\| \leq (\alpha - 1)^{-1} \|Q_0(Q_0 - z)^{-1}\| + \frac{\sqrt{3}}{2} (\alpha - 1)^{-1} \|(Q_0 - z)^{-1}\|. \quad (2.20)$$

Since the eigenvalues of Q_0 are $\left\{ \left(\frac{1}{2} + n \right) \sqrt{\alpha^2 - 1} \right\}_{n=0}^{\infty}$ we have $\sup_{z \in \partial C} \|Q_0(Q_0 - z)^{-1}\| = 3$ and $\sup_{z \in \partial C} \|(Q_0 - z)^{-1}\| = \frac{2}{\sqrt{\alpha^2 - 1}}$. Then the lemma follows. \square

The two-dimensional subspace spanned by eigenvectors associated with eigenvalues λ_1 and λ_2 is denoted by \mathcal{F} . The orthogonal projection onto \mathcal{F} is then given by

$$P = P(\varepsilon) = -\frac{1}{2\pi i} \oint_{\partial C} (Q - z)^{-1} dz. \quad (2.21)$$

We expand $P(\varepsilon)$ with respect to ε up to the second order:

$$P = P_0 + \varepsilon P_1 + \varepsilon^2 R, \quad (2.22)$$

where P_0 is the orthogonal projection onto the ground states of Q_0 and

$$P_1 = -\frac{1}{2\pi i} \oint_{\partial C} (Q_0 - z)^{-1} V (Q_0 - z)^{-1} dz, \quad (2.23)$$

$$R = R(\varepsilon) = -\frac{1}{2\pi i} \oint_{\partial C} (Q_0 - z)^{-1} V (Q_0 - z)^{-1} V (Q - z)^{-1} dz. \quad (2.24)$$

Lemma 2.3. *We have $\|R\| \leq g_2$ and $\|VP_1\| \leq E_0 g_1^2$.*

Proof. By Lemma 2.2, we have

$$\|R\| \leq \frac{|C|}{2\pi} \sup_{z \in \partial C} \|V(Q_0 - z)^{-1}\|^2 \|(Q - z)^{-1}\| \leq \frac{\sqrt{\alpha^2 - 1}}{2} g_1^2 \|(Q - \sqrt{\alpha^2 - 1})^{-1}\|. \quad (2.25)$$

Since $\lambda_3 \geq \frac{3}{2}\sqrt{\alpha^2 - 1}$, we have $\|(Q - \sqrt{\alpha^2 - 1})^{-1}\| = |\lambda_2 - \sqrt{\alpha^2 - 1}|^{-1}$. Hence $\|R\| \leq g_2$ holds. Similarly one can prove the second bound. \square

Let $v_0 \in L^2(\mathbb{R})$ be the normalized ground state of $h = \frac{1}{2}p^2 + \frac{\alpha^2 - 1}{2}x^2$. Namely

$$v_0(x) = \left(\frac{\sqrt{\alpha^2 - 1}}{\pi} \right)^{1/4} e^{-\sqrt{\alpha^2 - 1}x^2/2}. \quad (2.26)$$

Let S_a be the dilation defined by $S_a f(x) = \frac{1}{\sqrt{a}} f(ax)$ for $a > 0$. We define the unitary operator U on \mathcal{H} by

$$U = \frac{1}{\sqrt{2}} S_{\sqrt{\alpha}} \begin{pmatrix} e^{ix^2/(2\alpha)} & 0 \\ 0 & e^{-ix^2/(2\alpha)} \end{pmatrix} \begin{pmatrix} 1 & -i \\ 1 & i \end{pmatrix}. \quad (2.27)$$

Then $UQ_0U^* = \begin{pmatrix} h & 0 \\ 0 & h \end{pmatrix}$ and vectors $u_1 = U^* \begin{pmatrix} v_0 \\ 0 \end{pmatrix}$ and $u_2 = U^* \begin{pmatrix} 0 \\ v_0 \end{pmatrix}$ are two fold ground states of $Q(\alpha, \alpha)$. Since P is a projection onto \mathcal{F} , each of the ground state and the first excited state can be expressed as a linear combination of Pu_1 and Pu_2 as long as both Pu_1 and Pu_2 are linearly independent, which is proven in the lemma below:

Lemma 2.4. *Assume that $\varepsilon^2 g_2 < 1$. Then Pu_1 and Pu_2 are non-zero vectors. Moreover, if $\varepsilon^2 g_2 < 1/2$, then Pu_1 and Pu_2 are linearly independent.*

Proof. By (2.22) we have

$$\|Pu_1\|^2 = 1 + \varepsilon \langle u_1, P_1u_1 \rangle + \varepsilon^2 \langle u_1, Ru_1 \rangle. \quad (2.28)$$

The second term on the right hand side above is zero, since

$$\langle u_1, P_1u_1 \rangle = \frac{-1}{2\pi i} \oint_{\partial C} \frac{1}{(E_0 - z)^2} dz \langle u_1, Vu_1 \rangle = 0. \quad (2.29)$$

Hence $\|Pu_1\|^2 \geq 1 - \varepsilon^2 g_2^2 > 0$ holds by Lemma 2.3 and the assumption. Thus Pu_i ($i = 1, 2$) are non-zero vectors. Next we assume that $\varepsilon^2 g_2 < 1/2$. Then we have

$$\begin{aligned} \|Pu_1\|^2 \|Pu_2\|^2 - |\langle Pu_1, Pu_2 \rangle|^2 &\geq (1 - \varepsilon^2 g_2)(1 - \varepsilon^2 g_2) - \varepsilon^4 g_2^2 \\ &= 1 - 2\varepsilon^2 g_2 > 0 \end{aligned} \quad (2.30)$$

which implies that Pu_1 and Pu_2 are linearly independent. \square

Lemma 2.5. *Let g_3 and g_4 be given in (1.10) and (1.11), respectively. Then it follows that where we used the bound*

$$\|Vu_i\| \leq g_1 E_0, \quad i = 1, 2, \quad (2.31)$$

$$|\langle u_1, Vu_1 \rangle| = g_3, \quad (2.32)$$

$$|\langle u_1, Vu_2 \rangle| \leq g_4. \quad (2.33)$$

Proof. (2.31) follows from

$$\|Vu_i\| \leq \|V(Q_0 - \sqrt{\alpha^2 - 1})^{-1}\| \|(Q_0 - \sqrt{\alpha^2 - 1})u_i\| \leq g_1 E_0. \quad (2.34)$$

The proofs of (2.32) and (2.33) are given in Appendix. \square

Proof of Theorem 1.4: Suppose that $\varepsilon^2 g_2 < 1/2$. We define

$$\Phi_1 = \frac{Pu_1}{\|Pu_1\|}, \quad \Phi_2 = \frac{Pu_2 - \langle \Phi_1, u_2 \rangle \Phi_1}{\|Pu_2 - \langle \Phi_1, u_2 \rangle \Phi_1\|} \quad (2.35)$$

Then, by Lemma 2.4, Φ_1 and Φ_2 are orthogonal vectors in \mathcal{F} . Let $\mathcal{V} = L.H.\{\Phi_1, \Phi_2\}$ be the two dimensional vector space and $Q : \mathcal{V} \rightarrow \mathcal{V}$ can be regarded as a linear operator and its matrix representation is given by

$$m = \begin{pmatrix} \langle \Phi_1, Q\Phi_1 \rangle & \langle \Phi_1, Q\Phi_2 \rangle \\ \langle \Phi_2, Q\Phi_1 \rangle & \langle \Phi_2, Q\Phi_2 \rangle \end{pmatrix}. \quad (2.36)$$

Thus the eigenvalues λ_1 and λ_2 of Q are also the eigenvalue of m . Therefore, the difference of λ_1 and λ_2 can be computed by

$$(\lambda_2 - \lambda_1)^2 = (\langle \Phi_1, Q\Phi_1 \rangle - \langle \Phi_2, Q\Phi_2 \rangle)^2 + 4|\langle \Phi_1, Q\Phi_2 \rangle|^2, \quad (2.37)$$

which implies that

$$|\lambda_2 - \lambda_1| \geq 2|\langle \Phi_1, Q\Phi_2 \rangle|. \quad (2.38)$$

We estimate $|\langle \Phi_1, Q\Phi_2 \rangle|$ from below. Inserting the definition of Φ_j into $\langle \Phi_i, Q\Phi_j \rangle$ we have

$$\langle \Phi_1, Q\Phi_2 \rangle = \frac{\langle Pu_1, Q(\|Pu_1\|^2 Pu_2 - (Pu_1, Pu_2)Pu_1) \rangle}{\|Pu_1\|^3 \|Pu_2 - (Pu_1, Pu_2)Pu_1 / \|Pu_1\|^2\|}. \quad (2.39)$$

Notice that

$$\langle Pu_i, QPu_j \rangle = (Pu_i, Qu_j) = E_0 \delta_{ij} + \varepsilon \langle V \rangle_{ij} + \varepsilon^2 (E_0 \langle R \rangle_{ij} + \langle P_1 V \rangle_{ij}) + \varepsilon^3 \langle RV \rangle_{ij},$$

where $\langle K \rangle_{ij} = (u_i, Ku_j)$. We see that the denominator of (2.39) is expanded as

$$\begin{aligned} & \|Pu_1\| \| \|Pu_1\|^2 Pu_2 - (Pu_1, Pu_2)Pu_1 \| \\ &= (1 + \varepsilon^2 \langle R \rangle_{11}) \sqrt{(1 + \varepsilon^2 \langle R \rangle_{11})(1 + \varepsilon^2 \langle R \rangle_{22}) - \varepsilon^4 |\langle R \rangle_{12}|^2}. \end{aligned}$$

By the bound $\|R\| \leq g_2$ we have the lower bound

$$\|Pu_1\| \| \|Pu_1\|^2 Pu_2 - (Pu_1, Pu_2)Pu_1 \| \geq (1 - \varepsilon^2 g_2) \sqrt{1 - 2\varepsilon^2 g_2^2}. \quad (2.40)$$

The numerator of (2.39) can be also expanded as

$$\begin{aligned} & (Pu_1, Q(\|Pu_1\|^2 Pu_2 - (Pu_1, Pu_2)Pu_1)) \\ &= \varepsilon \langle V \rangle_{12} + \varepsilon^2 \langle P_1 V \rangle_{12} + \varepsilon^3 \left(\langle RV \rangle_{12} + \langle V \rangle_{12} \langle R \rangle_{11} - \langle R \rangle_{12} \langle V \rangle_{11} \right) \\ &+ \varepsilon^4 \left(\langle P_1 V \rangle_{12} \langle R \rangle_{11} - \langle R \rangle_{12} \langle P_1 V \rangle_{11} \right) + \varepsilon^5 \left(\langle R \rangle_{11} \langle RV \rangle_{12} - \langle R \rangle_{12} \langle RV \rangle_{12} \right). \end{aligned}$$

By using Lemmas 2.3 and 2.5, each term can be evaluated as

$$\begin{aligned} \varepsilon^2 |\langle P_1 V \rangle_{12}| &\leq \varepsilon^2 E_0 g_1^2 \\ \varepsilon^3 |\langle RV \rangle_{12} + \langle V \rangle_{12} \langle R \rangle_{11} - \langle R \rangle_{12} \langle V \rangle_{11}| &\leq \varepsilon^3 (E_0 g_1 g_2 + g_2 g_4 + g_2 g_3) \\ \varepsilon^4 |\langle P_1 V \rangle_{12} \langle R \rangle_{11} - \langle R \rangle_{12} \langle P_1 V \rangle_{11}| &\leq \varepsilon^4 2E_0 g_1^2 g_2 \\ \varepsilon^5 |\langle R \rangle_{11} \langle RV \rangle_{12} - \langle R \rangle_{12} \langle RV \rangle_{12}| &\leq \varepsilon^5 2E_0 g_1 g_2^2. \end{aligned}$$

By combining all the estimates stated above, we have

$$|\lambda_1 - \lambda_2| \geq \frac{2\varepsilon}{\ell(\varepsilon)} (g_4 - \varepsilon \kappa(\varepsilon)), \quad (2.41)$$

where $\ell(\varepsilon) = (1 - \varepsilon^2 g_2) \sqrt{1 - 2\varepsilon^2 g_2^2}$. Hence the theorem follows. \square

APPENDIX A. COMPUTATION OF $\langle V \rangle_{ij}$

We recall that $v_0 = (\omega/\pi)^{1/4} e^{-\omega x^2/2}$ with $\omega = \sqrt{\alpha^2 - 1}$, $V = \frac{1}{2} \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} (p^2 + x^2)$ and U is given by (2.27). We directly see that

$$\begin{aligned} & UVU^{-1} \\ &= \frac{1}{4} S_{\sqrt{\alpha}} \begin{pmatrix} e^{ix^2/2\alpha} & 0 \\ 0 & e^{-ix^2/2\alpha} \end{pmatrix} \begin{pmatrix} 1 & -i \\ 1 & i \end{pmatrix} \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ i & -i \end{pmatrix} (p^2 + x^2) \begin{pmatrix} e^{-ix^2/2\alpha} & 0 \\ 0 & e^{ix^2/2\alpha} \end{pmatrix} S_{1/\sqrt{\alpha}} \\ &= \frac{1}{4} \begin{pmatrix} e^{ix^2/2\alpha} \left(\frac{p^2}{\alpha} + \alpha x^2 \right) e^{-ix^2/2\alpha} & e^{-ix^2/2\alpha} \left(\frac{p^2}{\alpha} + \alpha x^2 \right) e^{ix^2/2\alpha} \\ -e^{-ix^2/2\alpha} \left(\frac{p^2}{\alpha} + \alpha x^2 \right) e^{-ix^2/2\alpha} & e^{ix^2/2\alpha} \left(\frac{p^2}{\alpha} + \alpha x^2 \right) e^{ix^2/2\alpha} \end{pmatrix}. \end{aligned}$$

Then we have

$$\left\langle \begin{pmatrix} 1 \\ 0 \end{pmatrix}, UVU^{-1} \begin{pmatrix} 1 \\ 0 \end{pmatrix} \right\rangle_{\mathbb{C}^2} = \frac{1}{4} e^{ix^2/2\alpha} \left(\frac{p^2}{\alpha} + \alpha x^2 \right) e^{-ix^2/2\alpha} \quad (1.1)$$

$$\left\langle \begin{pmatrix} 1 \\ 0 \end{pmatrix}, UVU^{-1} \begin{pmatrix} 0 \\ 1 \end{pmatrix} \right\rangle_{\mathbb{C}^2} = -\frac{1}{4} e^{ix^2/2\alpha} \left(\frac{p^2}{\alpha} + \alpha x^2 \right) e^{ix^2/2\alpha}. \quad (1.2)$$

Lemma A.1. *It follows that $\langle u_1, Vu_1 \rangle = \frac{\alpha}{4\omega}$, and then $|\langle u_1, Vu_1 \rangle| = g_3$.*

Proof. By (1.1) we have

$$\langle u_1, Vu_1 \rangle = \frac{1}{4} \left\langle v_0, e^{ix^2/2} \left(\frac{p^2}{\alpha} + \alpha x^2 \right) e^{-ix^2/2} v_0 \right\rangle = \frac{1}{4} \left\langle v_0, \left(\frac{(p-x)^2}{\alpha} + \alpha x^2 \right) v_0 \right\rangle.$$

Since $\langle v_0, (px + xp)v_0 \rangle = 0$, we obtain that

$$\langle u_1, Vu_1 \rangle = \frac{1}{4} \left\langle v_0, \left(\frac{p^2 + x^2}{\alpha} + \alpha x^2 \right) v_0 \right\rangle.$$

By $\langle v_0, x^2 v_0 \rangle = 1/(2\omega)$ and $\langle v_0, p^2 v_0 \rangle = \omega/2$ we have $\langle u_1, Vu_1 \rangle = \alpha/(4\omega)$. \square

Lemma A.2. *It follows that $\langle u_1, Vu_2 \rangle = -\frac{\omega^{3/2}}{4\alpha} (\omega - i)^{-1/2}$. In particular $|\langle u_1, Vu_2 \rangle| \leq \frac{\omega^{3/2}}{4\alpha^{3/2}}$.*

Proof. By (1.2) we have

$$\begin{aligned} \langle u_1, Vu_2 \rangle &= -\frac{1}{4} \left\langle v_0, e^{ix^2/2} \left(\frac{p^2}{\alpha} + \alpha x^2 \right) e^{ix^2/2} v_0 \right\rangle \\ &= -\frac{1}{4\alpha} \left\langle (-i - \omega) x e^{-ix^2/2} v_0, (i - \omega) x e^{ix^2/2} v_0 \right\rangle - \frac{\alpha}{4} \left\langle v_0, x^2 e^{ix^2} v_0 \right\rangle \\ &= -\frac{1}{4\alpha} ((\omega - i)^2 + \alpha^2) \left\langle v_0, x^2 e^{ix^2} v_0 \right\rangle. \end{aligned}$$

Since $\langle v_0, x^2 e^{ix^2} v_0 \rangle = (\omega/\pi)^{1/2} \int_{\mathbb{R}} x^2 e^{-(\omega-i)x^2} dx = \frac{1}{2} \omega^{1/2} (\omega-i)^{-3/2}$, we have the lemma. \square

Acknowledgment

We thank Takashi Ichinose and Masato Wakayama for sending [IW07] to our attention and giving helpful comments. I. S.'s work was partially supported by Y22740087 from JSPS, and was performed through the Program for Dissemination of Tenure-Track System funded by the Ministry of Education and Science, Japan. F.H. thanks for the financial support by Grant-in-Aid for Science Research (B) 20340032 from JSPS.

REFERENCES

- [GJ68] J. Glimm and A. Jaffe, The $\lambda(\phi^4)_2$ quantum field theory without cutoffs. I. *Phys. Rev.* **176** (1968), 1945–1951.
- [HH12] M. Hirokawa and F. Hiroshima, Absence of energy level crossing for the ground state energy of the Rabi model, preprint 2012.
- [Hir05] F. Hiroshima, Multiplicity of ground states in quantum field models: applications of asymptotic fields, *J. Funct. Anal.* **224** (2005), 431–470.
- [IW05a] T. Ichinose and M. Wakayama, Zeta functions for the spectrum of the non-commutative harmonic oscillators, *Commun. Math. Phys.* **258** (2005), 697–739.
- [IW05b] T. Ichinose and M. Wakayama, Special values of the spectral zeta function of the non-commutative harmonic oscillator and confluent Heun equations, *Kyushu. J. Math.* **59** (2005), 39–100.
- [IW07] T. Ichinose and M. Wakayama, On the spectral zeta function for the non-commutative harmonic oscillator and, *Rep. Math. Phys.* **59** (2007), 421–432.
- [KW06] K. Kimoto and M. Wakayama, Apéry-like number arising from special values of spectral zeta functions for non-commutative harmonic oscillators, *Kyushu. J. Math.* **60** (2006), 383–404.
- [KW07] K. Kimoto and M. Wakayama, Elliptic curves arising from the spectral zeta function for non-commutative harmonic oscillators and $\Gamma_0(4)$ -modular forms, *The Conference on L-Functions*, World Sci. Publ., Hackensack, NJ, 2007, 201–218.
- [KY09] K. Kimoto and Y. Yamasaki, A variation of multiple L-values arising from the spectral zeta function of the non-commutative harmonic oscillator, *Proc. Amer. Math. Soc.* **137** (2009), 2503–2515.
- [NNW02] K. Nagatou, M. T. Nakao and M. Wakayama, Verified numerical computations for eigenvalues of non-commutative harmonic oscillators, *Num. Funct. Anal. and Opt.* **23** (2002), 633–650.
- [Och01] H. Ochiai Non-commutative harmonic oscillators and Fuchsian ordinary differential operators, *Commun. Math. Phys.* **217** (2001), 357–373.
- [Och04] H. Ochiai Non-commutative harmonic oscillators and the connection problem for the Heun differential equation, *Lett. Math. Phys.* **70** (2004), 133–139.
- [PW01] A. Parmeggiani and M. Wakayama, Oscillator representations and systems of ordinary differential equations, *Proceedings of the National Academy of Sciences U.S.A* **98** (2001), 26–30.
- [PW02a] A. Parmeggiani and M. Wakayama, Non-commutative harmonic oscillator I, *Form Math.* **14** (2002), 539–604.
- [PW02b] A. Parmeggiani and M. Wakayama, Non-commutative harmonic oscillator II, *Form Math.* **14** (2002), 669–690.
- [PW03] A. Parmeggiani and M. Wakayama, Corrigenda and remarks to: "Non-commutative harmonic oscillators. I." [Forum Math. 14 (2002), 539–604] *Forum Math.* **15** (2003), 955–963.

- [Par04] A. Parmeggiani, On the spectrum and the lowest eigenvalue of certain non-commutative harmonic oscillators, *Kyushu J. Math.* **58** (2004), 277–322.
- [Par06] A. Parmeggiani, On the spectrum of certain noncommutative harmonic oscillators, *Ann. Univ. Ferrara Sez. VII Sci. Mat.* **52** (2006), 431–456.
- [Par08a] A. Parmeggiani, On the spectrum of certain non-commutative harmonic oscillators and semi-classical analysis, *Comm. Math. Phys.* **279** (2008), 285–308.
- [Par08b] A. Parmeggiani, *Introduction to the spectral theory of non-commutative harmonic oscillators*, COE Lecture Note **8**, Kyushu University, 2008.
- [Par10] A. Parmeggiani, *Spectral theory of non-commutative harmonic oscillators: an introduction*, Lecture Notes in Mathematics, **1992**, Springer-Verlag, Berlin, 2010.

FACULTY OF MATHEMATICS, KYUSHU UNIVERSITY, FUKUOKA, 819-0395, JAPAN
E-mail address: hiroshima@math.kyushu-u.ac.jp.

FIBER-NANOTECH YOUNG RESEARCHER EMPOWERMENT CENTER, SHINSHU UNIVERSITY, MATSUMOTO 390-8621, JAPAN
E-mail address: isasaki@shinshu-u.ac.jp