

Designing FPT algorithms for cut problems using randomized contractions

Rajesh Chitnis ^{*} Marek Cygan [†] MohammadTaghi Hajiaghayi [‡]
Marcin Pilipczuk [§] Michał Pilipczuk [¶]

Abstract

We introduce a new technique for designing fixed-parameter algorithms for cut problems, namely *randomized contractions*. We apply our framework to obtain the first FPT algorithm for the UNIQUE LABEL COVER problem and new FPT algorithms with exponential speed up for the STEINER CUT and NODE MULTIWAY CUT-UNCUT problems. More precisely, we show the following:

- We prove that the parameterized version of the UNIQUE LABEL COVER problem, which is the base of the UNIQUE GAMES CONJECTURE, can be solved in $2^{O(k^2 \log |\Sigma|)} n^4 \log n$ deterministic time (even in the stronger, vertex-deletion variant) where k is the number of unsatisfied edges and $|\Sigma|$ is the size of the alphabet. As a consequence, we show that one can in polynomial time solve instances of UNIQUE GAMES where the number of edges allowed not to be satisfied is upper bounded by $O(\sqrt{\log n})$ to optimality, which improves over the trivial $O(1)$ upper bound.
- We prove that the STEINER CUT problem can be solved in $2^{O(k^2 \log k)} n^4 \log n$ deterministic time and $\tilde{O}(2^{O(k^2 \log k)} n^2)$ randomized time where k is the size of the cutset. This result improves the double exponential running time of the recent work of Kawarabayashi and Thorup (FOCS'11).
- We show how to combine considering ‘cut’ and ‘uncut’ constraints at the same time. More precisely, we define a robust problem NODE MULTIWAY CUT-UNCUT that can serve as an abstraction of introducing uncut constraints, and show that it admits an algorithm running in $2^{O(k^2 \log k)} n^4 \log n$ deterministic time where k is the size of the cutset. To the best of our knowledge, the only known way of tackling uncut constraints was via the approach of Marx, O’Sullivan and Razgon (STACS’10), which yields algorithms with double exponential running time.

An interesting aspect of our algorithms is that they can handle real weights; to the best of our knowledge, the technique of important separators does not work in the weighted version.

^{*}Department of Computer Science, University of Maryland at College Park, USA, rchitnis@cs.umd.edu, Supported in part by NSF CAREER award 1053605, ONR YIP award N000141110662, DARPA/AFRL award FA8650-11-1-7162 and a University of Maryland Research and Scholarship Award (RASA).

[†]Institute of Informatics, University of Warsaw, Poland, cygan@mimuw.edu.pl, Supported in part by NSF CAREER award 1053605, ONR YIP award N000141110662, DARPA/AFRL award FA8650-11-1-7162, University of Maryland Research and Scholarship Award (RASA) NCN grant N206567140, Foundation for Polish Science.

[‡]Department of Computer Science, University of Maryland at College Park, USA, hajiagha@cs.umd.edu, Supported in part by NSF CAREER award 1053605, ONR YIP award N000141110662, DARPA/AFRL award FA8650-11-1-7162 and a University of Maryland Research and Scholarship Award (RASA).

[§]Institute of Informatics, University of Warsaw, Poland, malcin@mimuw.edu.pl, Partially supported by NCN grant N206567140 and Foundation for Polish Science.

[¶]Department of Informatics, University of Bergen, Norway, michal.pilipczuk@ii.uib.no, Partially supported by European Research Council (ERC) grant “Rigorous Theory of Preprocessing”, reference 267959.

1 Introduction

Graph cut problems is a class of problems where, given a graph, one is asked to find a *cutset* of minimum size whose removal makes the graph satisfy a global separation property. The motivation of studying graph cut problems stems from the fundamental minimum cut problem, where the goal is to separate two terminals from each other by removing the least possible number of vertices or edges, depending on the variant. Even though the minimum cut problem can be solved in polynomial time, many of its natural generalizations become NP-hard. Moreover, many problems, whose classical definitions do not resemble cut formulations, after choosing an appropriate combinatorial viewpoint show deep links with finding minimum separators; the most important examples are FEEDBACK VERTEX SET and ODD CYCLE TRANSVERSAL.

Therefore, circumventing NP-hardness of fundamental graph cut problems, like MULTIWAY CUT (given a graph with a set of terminals, separate the terminals from each other using minimum size cutset) or MULTICUT (given a graph with a set of terminal pairs, separate terminals in the pairs using minimum size cutset), became an important algorithmic challenge. It is then no surprise that graph cut problems were studied intensively from the point of view of approximation; (cf. [1, 4, 8, 15, 16, 20, 21, 26, 36, 38, 41]).

In this paper we address a different paradigm of tackling NP-hard problems, that is, *fixed-parameter tractability* (FPT). Recall that in the parameterized complexity setting the instance of the problem comes with an additional integer k , called the *parameter*, which intuitively measures the hardness of the instance. The goal is to devise an algorithm solving the problem with running time of form $f(k)n^c$, where f is some computable function and c is a fixed constant. In other words, for every fixed parameter the algorithm has to work in polynomial time, where the degree of the polynomial is independent of the parameter. Algorithms with such running time guarantee are called *fixed-parameter* algorithms, and if a problem admits one, then we say that it is *fixed-parameter tractable*. For a more detailed introduction to fixed-parameter tractability we address an interested reader to the books of Downey and Fellows [14] or Flum and Grohe [18].

Graph separation problems in the context of parameterized complexity were probably first considered in the seminal work of Marx [32]. Marx established fixed-parameterized complexity of MULTIWAY CUT parameterized by the size of the cutset and MULTICUT parameterized by the size of the cutset plus the number of terminal pairs. Perhaps the most fruitful consequence of his work was the introduction of the concept of an *important separator*. Important separators proved to be a robust tool that enable us to capture the bounded-in-parameter character of the family of sensible cutsets. The technique has found a number of applications [9, 10, 11, 13, 23, 31, 32, 34, 40], including proving fixed-parameter tractability of MULTICUT parameterized by the cutsize only, which was resolved by Marx and Razgon [34] and, independently, by Bousquet et al. [6], after resisting attacks as a long-standing open problem. The latest advances, by Chitnis et al. [11] and Kratsch et al. [30], try to find a proper understanding of this notion in directed graphs.

The important separators technique is based on greedy arguments, which unfortunately makes this approach difficult to extend. Consider, for instance, adding constraints of type 'uncut', i.e., we would like to find a cutset that separates some pairs of terminals, but is required *not* to separate some other pairs. Any greedy choice of the farthest possible cutset, which is precisely the idea behind the notion of an important separator, can spoil the delicate requirements of existence of some paths. Similar obstacles arise when considering problems with weights or in directed setting.

Another approach has been recently presented by Marx, O'Sullivan and Razgon [33]. Intuitively, the authors prove that one can find a subset of vertices in which all the minimal cutsets of size at most k are contained, and which induces (for an appropriate meaning of 'induce') a graph

of treewidth bounded by an exponential function of k . Thus, having expressed the problem in MSO_2 logic, we can use Courcelle’s lemma on the obtained tree decomposition. The technique can tackle more general constraints than important separators, such as ‘uncut’ constraints, and works elegantly for easy cut problems with strong parameterizations. A drawback is that algorithms constructed in this manner almost always have double exponential dependence on k of the running time, as we run a dynamic program on a tree decomposition of width exponential in k .

1.1 Our techniques

We introduce a new technique, called *randomized contractions*, of constructing fixed-parameter algorithms for graph cut problems. The technique is based on a WIN/WIN approach: either we find a well-balanced separation of small order, whose one side can be simplified by a recurrential call, or the graph admits a highly-connected structure, which can be used to identify the solution. First we test, whether the graph admits a well-balanced separation. If this is the case, we run the algorithm recurrentially on one of the sides for all possible behaviors on the boundary, for each marking some optimal solution. Then we argue that unmarked parts can be conveniently reduced. However, if no separation is present, we find that after removing the solution the graph can split only into a bounded number of connected components of bounded size, and at most one connected component that can be arbitrarily large. Then, we randomly contract parts of the graph; the highly-connected structure ensures us that with high probability the optimal solution will be ‘highlighted’ by the contraction step, so that we can easily extract it. Intuitively, the event we aim for is that all the components of bounded size and a sufficiently large neighborhood of the solution are distinguished.

Finding a small separator and recurrentially reducing one of the sides is the core idea of many algorithms for parameterized problems. For example, a novel concept of finding and reducing *protrusions*, large subgraphs of constant treewidth and constant boundary, led to construction of linear kernels for a large number of problems on classes of graphs with topological constraints [5, 19]. Perhaps a better example is the recent fixed-parameter algorithm for the k -WAY CUT problem of Kawarabayashi and Thorup [27], which is in fact the original motivation of our technique. The authors observe that if they find a small, well-balanced separation in the graph, they may reduce one of the sides up to bounded size, in a very similar manner as we do. Basing on this, they analyze the graph in a minor-style manner and whenever they encounter a feasible separation, they can apply the reduction.

One of the tools we use in our algorithms is the color coding technique introduced by Alon et al. [2] to solve some special cases of the SUBGRAPH ISOMORPHISM problem. The main idea is to color the graph at random and ensure that with high probability the solution gets sufficiently highlighted to be recognizable quickly. It has now become a classical tool in the parameterized complexity toolbox. Our technique is both similar and different to color coding. The similarity lies in the first application of color coding in our approach, i.e., finding a feasible separation to make the reduction step. However, when no good separation can be found, we use different ideas not only to highlight the solution, but also to expose the highly connected structure of the graph. Although the intuition behind color coding is of probabilistic nature, the algorithms obtained using this approach can be derandomized using the technique of *splitters* of Naor et al. [37]. In fact, we find it more convenient to present our algorithms already in the derandomized version, so in spite of the name of the technique there will be no randomization at all; instead we use the following abstraction:

Lemma 1.1 (\spadesuit). *Given a set U of size n together with integers $0 \leq a, b \leq n$, one can in $O(2^{O(\min(a,b) \log(a+b))} n \log n)$ time construct a family \mathcal{F} of at most $O(2^{O(\min(a,b) \log(a+b))} \log n)$*

subsets of U , such that the following holds: for any sets $A, B \subseteq U$, $A \cap B = \emptyset$, $|A| \leq a$, $|B| \leq b$, there exists a set $S \in \mathcal{F}$ with $A \subseteq S$ and $B \cap S = \emptyset$.

Our approach is most natural for edge-deletion problems; however, we can also extend it to node-deletion variants. For the node deletion problems however, the situation is more complicated and we need to define two kinds of separations. Only when the graph does not have both kinds of separations do we get enough structure to perform the recursive step. Additionally, one needs to be much more careful in the case when no feasible separation can be found, as we obtain much weaker structural properties of the graph.

1.2 Our results

We use the technique to provide the first fixed-parameter algorithm solving an important problem in parameterized complexity, and moreover we show how our approach can be applied to reduce the time complexity of the best known algorithms from double exponential to single exponential for some problems already known to be FPT.

Unique Label Cover. In the UNIQUE LABEL COVER problem we are given an undirected graph G , where each edge $uv = e \in E(G)$ is associated with a permutation $\psi_{e,u}$ of a constant size alphabet Σ . The goal is to construct a labeling $\Psi : V(G) \rightarrow \Sigma$ maximizing the number of satisfied edge constraints, that is, edges for which $(\Psi(u), \Psi(v)) \in \psi_{uv,u}$ holds. At the first glance UNIQUE LABEL COVER does not seem related to the previously mentioned cut problems, however it is not hard to show that the node deletion version of UNIQUE LABEL COVER is a generalization of GROUP FEEDBACK VERTEX SET problem [24], and hence ODD CYCLE TRANSVERSAL and FEEDBACK VERTEX SET, as well as MULTIWAY CUT.

The optimization version of UNIQUE LABEL COVER is the subject of the very extensively studied UNIQUE GAMES CONJECTURE proposed by Khot [28] in 2002, which is used as a hardness assumption for showing several tight inapproximability results. The UNIQUE GAMES CONJECTURE states that for every $\epsilon, \delta > 0$, there exists an alphabet size $|\Sigma|(\epsilon, \delta)$, such that given an instance $(G, \Sigma, (\psi_{e,v})_{e \in E(G), v \in e})$ it is NP-hard to distinguish between the cases $|OPT| \leq \delta |E(G)|$ and $|OPT| \geq (1 - \epsilon) |E(G)|$. In 2010 Arora et al. [3] presented a breakthrough subexponential time algorithm, which in $2^{O(|\Sigma|n^\epsilon)}$ running time satisfies $(1 - \epsilon) |E(G)|$ edge constraints, assuming the given instance satisfies $|OPT| \geq (1 - \epsilon^c) |E(G)|$. We refer the reader to a recent survey of Khot [29] for more detailed discussion on the UNIQUE GAMES CONJECTURE.

Since all the edge constraints are permutations, fixing a label for one vertex gives only one possibility for each of its neighbors, assuming we want to satisfy all edges. For this reason we can verify in polynomial time, whether $OPT = |E(G)|$. In this paper we show that we can efficiently solve the UNIQUE LABEL COVER problem, assuming almost all the edges are to be satisfied. In particular we design an fixed parameter algorithm for NODE UNIQUE LABEL COVER, which is a generalization of EDGE UNIQUE LABEL COVER.

NODE UNIQUE LABEL COVER

Parameter: $k + s$

Input: An undirected graph G , a finite alphabet Σ of size s , an integer k , and for each edge $e \in E(G)$ and each of its endpoints v a permutation $\psi_{e,v}$ of Σ , such that if $e = uv$ then $\psi_{e,u} = \psi_{e,v}^{-1}$.

Question: Does there exist a set $X \subseteq V(G)$ of size at most k and a function $\Psi : V(G) \setminus X \rightarrow \Sigma$ such that for any $uv \in E(G \setminus X)$ we have $(\Psi(u), \Psi(v)) \in \psi_{uv,u}$?

A brief sketch of the proof of the following theorem is given in Section 4; the details are in Appendix D.

Theorem 1.2. *There is an $O(2^{O(k^2 \log s)} n^4 \log n)$ time algorithm solving NODE UNIQUE LABEL COVER, and consequently EDGE UNIQUE LABEL COVER.*

To justify our parameterization, we would like to note that there is a long line of polynomial time approximation algorithms designed for instances of UNIQUE LABEL COVER, with currently best by Charikar et al. [7], working under the assumption $|OPT| \geq (1 - \epsilon)|E(G)|$, and where the alphabet is of constant size. Therefore, it is reasonable to assume that only a small number of constraints is not going to be satisfied. Our results imply that one can in polynomial time verify whether it is possible to satisfy $|E(G)| - O(\sqrt{\log n})$ constraints; consequently, we extend the range of instances that can be solved optimally in polynomial time.

Finally, we show that the dependence on the alphabet size in Theorem 1.2 is probably necessary, since the problem parameterized by the cutsize only is $W[1]$ -hard. Hence, existence of an algorithm parameterized by the cutsize only would cause $FPT = W[1]$, which is considered implausible. For a more detailed introduction to the hierarchy of parameterized problems and consequences of its collapse, we refer to the books of Downey and Fellows [14] or Flum and Grohe [18]. We consider this result an interesting counterposition of the parameterized status of GROUP FEEDBACK VERTEX SET [12], which is FPT even when the group size is not a parameter.

Theorem 1.3. *The EDGE UNIQUE LABEL COVER problem, and consequently NODE UNIQUE LABEL COVER, is $W[1]$ -hard when parameterized by k only.*

Steiner Cut. Next, we address a robust generalization of both k -WAY CUT and MULTIWAY CUT problems, namely the STEINER CUT problem.

STEINER CUT

Parameter: k

Input: A graph G , a set of terminals $T \subseteq V(G)$, and integers s and k .

Question: Does there exist a set X of at most k edges of G , such that in $G \setminus X$ at least s connected components contain at least one terminal?

Using our technique we present an FPT algorithm working in $O(2^{O(k^2 \log k)} n^4 \log n)$, where the polynomial factor can be improved to $\tilde{O}(n^2)$ at the cost of our algorithm being randomized. These results improve the double exponential time complexity of the recent algorithm of Kawarabayashi and Thorup [27]¹. A brief sketch of the proof of the following theorem is given in Section 3 and the details are in Appendix E.

Theorem 1.4. *There is a deterministic $O(2^{O(k^2 \log k)} n^4 \log n)$ and randomized $\tilde{O}(2^{O(k^2 \log s)} n^2)$ running time algorithm solving STEINER CUT.*

Connectivity constraints. We define the following problem as an abstraction of introducing 'cut' and 'uncut' constraints at the same time.

NODE MULTIWAY CUT-UNCUT (N-MWCU)

Parameter: k

Input: A graph G together with a set of terminals $T \subseteq V(G)$, an equivalence relation \mathcal{R} on the set T , and an integer k .

Question: Does there exist a set $X \subseteq V(G) \setminus T$ of at most k nonterminals such that for any $u, v \in T$, the vertices u and v belong to the same connected component of $G \setminus X$ if and only if $(u, v) \in \mathcal{R}$?

Fixed-parameter tractability of this problem can be derived from the framework of Marx, Razgon and O'Sullivan [33], complemented with a reduction of the number of equivalence classes of \mathcal{R} in flavour of the reduction for MULTIWAY CUT of Razgon [39] (we formalize this reduction in Appendix F). However, the dependence on k of the running time is double exponential. Using

¹In [27] the authors solve the k -WAY CUT problem, however a straightforward generalization of their algorithm solves the STEINER CUT problem as well.

our framework we provide an algorithm working in $O(2^{O(k^2 \log k)} n^4 \log n)$ time, which can be constructed by combining (i) ideas from the edge-deletion variant, (ii) the general approach to node-deletion problems, (iii) the aforementioned reduction of the number of equivalence classes of \mathcal{R} and (iv) simple reductions of terminals sharing large parts of neighborhoods. The full proof of the following theorem is in Appendix F, whereas in Section 2 we present a sketch of it for the edge-deletion variant.

Theorem 1.5. *There is an $O(2^{O(k^2 \log k)} n^4 \log n)$ time algorithm solving NODE MULTIWAY CUT-UNCUT.*

Organization of the paper. We firstly describe the general idea of the technique in Section 2, illustrating them on the edge-deletion version of the N-MWCU problem. Then, in Section 3 we discuss how this general framework can be put to work for the STEINER CUT problem. In Section 4 we sketch the algorithm for the NODE UNIQUE LABEL COVER problem; we find this problem best-suited to present the challenges that arise when considering node-deletion variants, and means of overcoming them. Section 5 is devoted to brief concluding remarks. Although the intuition behind our algorithms is very natural, their formal description needs paying attention to a plethora of technical details. Therefore, in the main body of the paper we present only sketches, and the full and formal description of the proofs of Theorems 1.2, 1.4, 1.5 can be found in Appendices D, E, F, respectively. Theorem 1.3 is proved in Appendix G. Appendix A contains notation and preliminaries used in the formal proofs, Appendix B contains proofs of lemmas omitted in the main body of the paper (marked with ♠), Appendix C contains discussion on separations in node-deletion problems, and Appendix H contains a short description of how weights can be incorporated to our framework.

2 Illustration

In this section we present the outline of the technique, illustrating it with a running example of the EDGE MULTIWAY CUT-UNCUT problem, the edge-deletion variant of NODE MULTIWAY CUT-UNCUT.

<p>EDGE MULTIWAY CUT-UNCUT (E-MWCU)</p> <p>Input: A graph G together with a set of terminals $T \subseteq V(G)$, an equivalence relation \mathcal{R} on the set T, and an integer k.</p> <p>Question: Does there exist a set $X \subseteq E(G)$ of at most k edges such that for any $u, v \in T$, the vertices u and v belong to the same connected component of $G \setminus X$ if and only if $(u, v) \in \mathcal{R}$?</p>	<p>Parameter: k</p>
--	---

As the edge-deletion variant can be easily reduced to the node-deletion variant (see Appendix F for a formal reduction), the fixed-parameter tractability of E-MWCU follows from Theorem 1.5. However, we find this particular problem best-suited to serve as an illustration of our technique, to complement the description of abstract and often intuitive concepts with a ‘real-life’ example. Throughout this section we sketch an algorithm with running time $2^{O(k^2 \log k)} n^4 \log n$, resolving the E-MWCU problem. As we consider the edge-deletion version, we use edge cuts throughout this section. However, as our general framework can be also applied to node-deletion problems, we comment along the description where additional argumentation is needed.

We assume that the graph given in the input is connected, as it is easy to reduce the problem to considering each connected component separately. This is true for all the considered problems. Connectivity of the graph will be maintained during the whole course of the algorithm. Note that this means that the graph after excluding X can have at most $k+1$ connected components. Hence, we can assume that \mathcal{R} has at most $k+1$ equivalence classes, as otherwise we may safely return a negative answer.

2.1 High-level intuition

The starting point of our approach is an observation that if two vertices of the graph v, w can be connected via $k + 1$ edge-disjoint paths, then after removing at most k edges they remain in the same connected component. Therefore, in this situation we may apply a reduction rule that simplifies the graph basing on the knowledge that v and w cannot be separated by a small separator. For example, in the EDGE MULTIWAY CUT-UNCUT problem it is safe to simply identify v and w . Note that in this way we may obtain a multigraph.

Of course, this simple reduction cannot solve the problem completely in general, as the input graph can have, for instance, degrees bounded by a constant. However, we still would like to pursue the intuition that making the input graph well-connected leads to better understanding of its structure. To this end, we introduce the notion of *good separations*.

Definition 2.1. Let G be a connected graph. A partition (V_1, V_2) of $V(G)$ is called a (q, k) -good edge separation, if (i) $|V_1|, |V_2| > q$; (ii) $|\delta(V_1, V_2)| \leq k$; (iii) $G[V_1]$ and $G[V_2]$ are connected.

In the first phase, named *recursive understanding*, we iteratively find a good edge separation and reduce one of its sides up to the size bounded by a function of the parameter. We use the lower bound on the number of vertices of either side to ensure that we indeed make some simplification. The applied reduction step needs introducing a more general problem, in which, intuitively, we have to prepare for every possible behavior on a bounded number of distinguished vertices of the graph, called *border terminals*.

When no good edge separation can be found, by Menger's theorem we know that between every two disjoint connected subgraphs of size larger than q we can find $k + 1$ edge-disjoint paths. Then we proceed to the second phase, named *high connectivity phase*, where we exploit this highly connected structure to identify the solution.

2.2 Recursive understanding

First, we show how to find a good edge separation in the graph. Then we show how a good edge separation can be used to reduce the instance.

Lemma 2.2 (♠). *There exists a deterministic algorithm that, given an undirected, connected graph G on n vertices along with integers q and k , in time $O(2^{O(\min(q,k) \log(q+k))} n^3 \log n)$ either finds a (q, k) -good edge separation, or correctly concludes that no such separation exists.*

In Appendix B we provide an even faster algorithm for the same problem, that is randomized but runs in $\tilde{O}(2^{O(\min(q,k) \log(q+k))} (|V(G)| + |E(G)|))$ time. The approach is based on the classical Karger's algorithm for minimum cut [25].

Having found a good edge separation we can proceed to simplification of one of the sides. To this end, we consider a more general problem, where the input graph is equipped with a set of *border terminals* T_b , whose number is bounded by a function of the budget for edge deletions. Intuitively, each considered instance of the border problem corresponds to solving some small part of the graph, which can be adjacent to the remaining part only via a small boundary — the border terminals. Our goal in the border version is, for every fixed behavior on the border terminals, to find some minimum size solution or conclude that the size of the minimum solution exceeds the given budget. Of course, the definition of behavior is problem-dependent; therefore, we present this concept on the example of the E-MWCU problem.

The behavior on the border terminals, whose number will be bounded by $2k$, is defined by a pair of equivalence relations $\mathcal{P} = (\mathcal{R}_b, E_b)$. \mathcal{R}_b is defined on $T \cup T_b$, but we require that $\mathcal{R}_b|_T = \mathcal{R}$, i.e., \mathcal{R}_b extends \mathcal{R} . Informally, \mathcal{R}_b expresses which border terminals are required to be in the same connected with which terminals after removing the solution. The second relation E_b is

defined on T_b and we require it to be a subset of \mathcal{R}_b . Informally, E_b expresses, which pairs of border terminals are assumed to be reachable via paths outside the considered subgraph. The reader may view E_b as a *torso* operation, commonly used in the context of graph minors; thus, in the border problem we prepare ourselves for *all* the possible torso operations. For an instance of the border problem \mathcal{J}_b by $\mathbb{P}(\mathcal{J}_b)$ we denote the set of possible pairs $\mathcal{P} = (\mathcal{R}_b, E_b)$; note that $|\mathbb{P}(\mathcal{J}_b)| \leq f(k) = 2^{O(k \log k)}$, as the number of equivalence classes of \mathcal{R} is bounded by $k + 1$. Formally, we say that a set $X \subseteq E(G)$ is a *solution* to $(\mathcal{J}_b, \mathcal{P})$ if $|X| \leq k$ and in the graph $G_{\mathcal{P}} := (V(G), E(G) \cup E_b)$ after removing X every two vertices $u, v \in T \cup T_b$ are in the same connected component if and only if they are equivalent with respect to \mathcal{R}_b . Note that the graph $G_{\mathcal{P}}$ is exactly G with torso operation performed on equivalence classes of E_b . We can now formally define the border problem:

BORDER E-MWCU

Input: An E-MWCU instance $\mathcal{J} = (G, T, \mathcal{R}, k)$ with G being connected, and a set $T_b \subseteq V(G)$ of size at most $2k$; denote $\mathcal{J}_b = (G, T, \mathcal{R}, k, T_b)$.

Output: For each $\mathcal{P} \in \mathbb{P}(\mathcal{J}_b)$ output a solution $\text{sol}_{\mathcal{P}} = X_{\mathcal{P}}$ to $(\mathcal{J}_b, \mathcal{P})$ with $|X_{\mathcal{P}}|$ minimum possible, or output $\text{sol}_{\mathcal{P}} = \perp$ if such a solution does not exist.

BORDER E-MWCU generalizes E-MWCU: we ask for $T_b = \emptyset$ and take the output for the pair (\mathcal{R}, \emptyset) .

Now assume that (V_1, V_2) is a (q, k) -good separation of the graph G , for the input instance $\mathcal{J}_b = (G, T, \mathcal{R}, k, T_b)$ of BORDER E-MWCU, where $q = k \cdot f(k) + 1$ is the maximum number of edges that can appear on the output plus 1. As $|T_b| \leq 2k$, at least one of the sides contains at most k border terminals. Without loss of generality we assume that $|V_1 \cap T_b| \leq k$. Now consider an instance $\mathcal{J}_b^* = (G[V_1], T \cap V_1, \mathcal{R}|_{T \cap V_1}, k, (T_b \cap V_1) \cup (V_1 \cap V(\delta(V_1, V_2))))$ of BORDER E-MWCU. In other words, we trim the instance to the part V_1 and incorporate all the vertices incident to V_2 to the set of border terminals. Note that \mathcal{J}_b^* is a correct instance, as $|(T_b \cap V_1) \cup (V_1 \cap V(\delta(V_1, V_2)))| \leq k + k = 2k$. We now recursively solve the instance \mathcal{J}_b^* , obtaining a set of at most $q - 1$ edges that appear in any solution for any behavior on border terminals. Consider the remaining edges, i.e., those that do not appear in any solution to \mathcal{J}_b^* for any behavior on the terminals. It is not hard to see that all these edges can be in fact contracted, as it is useless to incorporate them in the solution to \mathcal{J}_b for any behavior on the border terminals. Intuitively, for every solution that uses some of these edges we can replace the part contained in $G[V_1]$ with the optimal solution for \mathcal{J}_b^* that imposes the same behavior on border terminals of \mathcal{J}_b^* , computed by the recursive call. We omit here the formal proof. After applying the contractions, part $G[V_1]$ shrinks to size at most q , as it is still connected and contains at most $q - 1$ edges. The sets of terminals, border terminals and relations are defined naturally.

We remark that the operation applied to reduce parts of the graph determined to be useless is problem-dependent. In E-MWCU we use edge contraction; however, more complex problems or node-deletion versions require more careful simplification rules.

We are left with estimating the running time of the algorithm. By Lemma 2.2 the time required to find a (q, k) -good edge separation is $O(2^{O(k \log f(k))} n^3 \log n) = O(2^{O(k^2 \log k)} n^3 \log n)$; hence, the recurrence is

$$T(n) \leq \max_{q+1 \leq n' \leq n-q-1} \left(O(2^{O(k^2 \log k)} n^3 \log n) + T(n') + T(n - n' + q) \right). \quad (1)$$

It is not difficult to prove by induction that $T(n) \leq O(2^{O(k^2 \log k)} n^4 \log n)$. We note that if f , the bound on the number of behaviors on the border terminals, is only a function of k , then this recurrence always yields running time of the form $T(n) \leq O(g(k) n^4 \log n)$ for some function g .

2.3 High connectivity phase

We are left with the more involved part of our approach, namely, what to do when no (q, k) -good edge separation is present in the graph. Note that we can assume that the graph has more than $q(k+1)$ vertices, as otherwise a brute-force search, which checks all the subsets of edges of size at most k , runs within the claimed time complexity bound.

The following simple lemma formalizes the structural properties of the graph after removing the solution. Note that this structure is precisely the gain of the first phase of the algorithm. We remark that we have this structure only in graph G , not $G_{\mathcal{P}}$. Fortunately, adding the edges of E_b will not be a problem.

Lemma 2.3 (\spadesuit). *Let G be a connected graph that admits no (q, k) -good edge separation. Let F be a set of edges of size at most k , such that $G \setminus F$ has connected components C_0, C_1, \dots, C_ℓ . Then (i) $\ell \leq k$, and (ii) all the components C_i except at most one contain at most q vertices.*

From now on, we always assume that $|V(C_i)| \leq q$ for $i = 1, 2, \dots, \ell$, while C_0 can have unbounded size. In fact, as $|V(G)| > q(k+1)$ we find that $|V(C_0)| > q$. We refer to components C_i for $i = 1, 2, \dots, \ell$ as to *small components*, while C_0 is called the *big component*. We would like to remark that if we apply the framework directly to the node-deletion problems, we do not have any bound on ℓ , i.e., the number of small components — in the node-deletion setting we need additional tools here.

Fix some behavior on the border terminals $\mathcal{P} = (\mathcal{R}_b, E_b) \in \mathbb{P}(\mathcal{J}_b)$; we iterate through all of them, which gives $2^{O(k \log k)}$ overhead to the running time. Assume that there exists a solution $X \subseteq E(G)$ for this particular choice. Without loss of generality let X be minimum. Let C_0, \dots, C_ℓ be components of $G \setminus X$, as in Lemma 2.3, where $|V(C_i)| \leq q$ for $i = 1, 2, \dots, \ell$. For every component C_i , choose an arbitrary its spanning tree T_i . Let $A_1 = \bigcup_{i=1}^{\ell} E(T_i)$ be the set of edges of the spanning trees of small components. As $\ell \leq k$, we have that $|A_1| \leq (q-1)k$. For every vertex $u \in V(X) \cap V(C_0)$ construct an arbitrary subtree T_0^u of T_0 such that $u \in V(T_0^u)$ and $|V(T_0^u)| = q+1$, and let $A_2 = \bigcup_{u \in V(X) \cap V(C_0)} E(T_0^u)$. As X is minimum, we have that $|V(X) \cap V(C_0)| \leq k$ and hence $|A_2| \leq qk$.

Now, we would like to apply edge contractions once more, in order to condensate the high-connectivity between large connected subgraphs of G to single vertices. Let \mathcal{F} be the family obtained from Lemma 1.1 for the universe $E(G)$ and constants $a = (2q-1)k$ and $b = k$. By Lemma 1.1, there exists $S_0 \in \mathcal{F}$ such that all the edges from $A_1 \cup A_2$ belong to S_0 , whereas all the edges from X are not in S_0 . We branch into $|\mathcal{F}|$ subcases labeled by the sets $S \in \mathcal{F}$. In each branch we either find a candidate for an optimal solution among edges that do not belong to S , or conclude that no such exists. We argue that if a (minimum) solution X exists, then some optimal solution will be found in the branch where S_0 is chosen.

For the sake of analysis, we present the routine performed in every branch assuming that we consider S_0 . First, we can contract all the edges of S_0 , as we seek a solution that is disjoint with S_0 . Let H_0 be the graph obtained in this operation. Observe that each small component C_i , $i = 1, 2, \dots, \ell$, forms exactly the subgraph contracted to a single vertex c_i . Moreover, all the vertices c_i can be incident only to edges from X , which are preserved due to the properties of S_0 and X being minimum. The set of terminals, border terminals and relations in H_0 are defined naturally as projections of the corresponding objects in G ; if we encounter any mismatch, e.g., two terminals that are non-equivalent with respect to \mathcal{R} are contracted onto the same vertex, we can safely terminate the branch. Observe that the branch when S_0 is considered is not terminated. Moreover, X survives the contractions and remains a feasible solution.

For a vertex $u \in V(H_0)$ we define its *weight* to be the number of vertices of G that were contracted onto it. A vertex $u \in V(H_0)$ is called *heavy* if it has weight at least $q+1$. Observe that all the vertices of $V(X) \cap V(C_0)$, that is, endpoints of edges from the solution that lie in C_0 ,

are contracted onto heavy vertices. Now we may make use of the high-connectivity structure of the graph. As G does not admit any (q, k) -good edge separation, by Menger's theorem between all the big vertices in H_0 we can find $k + 1$ edge-disjoint paths. This means that after removing the solution, all the big vertices are still in the same connected component. Hence, it is safe to identify them into one vertex b , which will be denoted the *core vertex*. Let H be the graph obtained after identification. The set of terminals and border terminals are defined naturally as before; again we can provide a negative answer if we encounter any mismatch during identifications. Moreover, all the edges from X are still present in H , as they do not connect big vertices, and X remains a feasible solution.

We remark that we can assume that the optimal deletion set is nonempty, as this particular possibility can be checked in polynomial time. Hence, there exists at least one heavy vertex or otherwise we may terminate the branch. Therefore, the core vertex b is well-defined.

Let B'_1, B'_2, \dots, B'_p be the components of $H \setminus \{b\}$ and let $B_i = H[V(B'_i) \cup \{b\}]$ for $i = 1, 2, \dots, p$. Observe that B'_i are connected, edge-disjoint and b separates them. We claim that for every component B_i , the solution X contains either all edges of $E(B_i)$ or none of them. This follows directly from the fact that all the small components are contracted into single vertices, while all endpoints of edges from X that are contained in C_0 are contracted onto b .




Figure 1: Obtaining the graph H from the graph G . Red edges are the edges of X that are preserved, while blue edges belong to $A_1 \cup A_2$ and are contracted. Note that trees T_0^u inside T_0 are not necessarily disjoint, but to make the presentation clearer they are disjoint in the figure. Moreover, on the left side of the picture we have drawn only edges belonging to X and trees T_i . However, there may be many more edges that, in particular, can influence the shape of the graph H after contraction and identification.

We conclude the algorithm by showing how to find an optimal solution inside the graph H in $O(kn^2)$ time. First, if b is not a border terminal nor a terminal, we iterate through all the possible equivalence class of \mathcal{R}_b , with which the vertex b is in the same connected component (plus one possibility, with none of them). Observe that we can assume that the number of equivalence classes of \mathcal{R}_b is bounded by $k + 1$, so we have at most $k + 2$ possibilities. Let D be the guessed subset of vertices from $T \cup T_b$ that are reachable from b after removing the solution. Observe that now we know exactly how the solution needs to look like. It simply needs to contain the entire edge sets of components that contain terminals or border terminals that do not belong to D , while all the other components may be left disjoint with the solution. The last

claim is asserted by $E_b \subseteq \mathcal{R}_b$: the additional edges inserted in $G_{\mathcal{P}}$ cannot force us to include into the solution also an edge set of a component, that contains only terminals and border terminals from D . Having constructed a candidate for the solution, we can check if it satisfies all the constraints in $G_{\mathcal{P}}$ in $O(n^2)$ time. It follows from the presented construction that when S_0 is considered, some solution of optimal size is found.

We remark that the last part of the algorithm, i.e., solving the problem inside the graph H , is highly problem-dependent. In our simple example of EDGE MULTIWAY CUT-UNCUT we were able to derive a simple routine with running time polynomial in k and n , but for more involved problems we can still need time exponential in k . All the other parts of the approach are to some extent generic and can be applied to various problems. We also would like to note that contractions of big trees on the side of C_0 were not necessary in this particular problem: one could iterate through a smaller family \mathcal{F} assuming only contractions of small components into single vertices. However, we chose to present the algorithm in this way, as the idea of condensating the big component into the core vertex significantly strengthens the obtained structure of the graph, and is an essential ingredient in all the other algorithms presented in this paper.

3 FPT algorithm for STEINER CUT

In this section we sketch the fixed-parameter algorithm for STEINER CUT. The algorithm will be similar to the one sketched in the illustration, so we concentrate on differences. Full description of the algorithm can be found in Appendix E. Again, we can assume that the graph given in the input is connected.

3.1 Recursive understanding

This phase is even simpler than the corresponding one in the illustration. In the border problem the graph is equipped with a set T_b of at most $2k$ border terminals. The behavior on border terminals is expressed by a triple $\mathcal{P} = (\mathcal{R}_b, Y_b, s_b)$. \mathcal{R}_b is an equivalence relation on T_b expressing, which border terminals are required to be in the same connected components after removing the solution. Y_b is a subset of T_b such that if $(u, v) \in \mathcal{R}_b$, then $u \in Y_b$ if and only if $v \in Y_b$. Y_b expresses which border terminals are required to be in a connected component that contains a terminal. Finally, s_b denotes how many connected components after removing the solution need to contain a terminal. If \mathcal{J}_b is an instance of the border problem, we say that $X \subseteq E(G)$ is a solution to $(\mathcal{J}_b, \mathcal{P})$ if $|X| \leq k$ and in $G \setminus X$ all these constraints expressed by \mathcal{P} are satisfied. The border problem is defined in the same manner as in the illustration: for each behavior on the border terminals we have to output a solution with minimum $|X|$, or \perp if no solution exists. Note that we do not need to remember also the connectivity outside, i.e., E_b . Intuitively, we can avoid it because the support of \mathcal{R}_b is bounded by $2k$, so after trimming the instance to one side of the separation we do not risk uncontrolled partitioning of equivalence classes of \mathcal{R}_b .

The recurrent application and simplification steps are very similar, so we omit the details in this sketch. If the recursive understanding cannot be applied further, we know that the graph does not admit a (q, k) -good edge separation for $q = 2^{O(k \log k)}$.

3.2 High connectivity phase

The first part of the high connectivity phase can be done in exactly the same manner as in the illustration. We guess a member S of an appropriately chosen family \mathcal{F} given by Lemma 1.1, assuming that S is disjoint with the solution but contains spanning trees of all small connected components after removing the solution as well as, for every vertex of the big component incident

to an edge of the solution, a tree of size $q+1$ attached to it and contained in C_0 . Having guessed S correctly, it is safe to contract the edges of S and identify all the vertices onto which more than q vertices were contracted. Thus, we obtain the same structure as in the illustration: a core vertex b and a number of components B_i attached to it, whose edge sets have to be either disjoint with the solution or entirely contained in it.

As previously, if b is not a border terminal, then we guess the equivalence class (or its lack) of \mathcal{R}_b , whose vertices will be in the same connected component as b . Having fixed the alignment of b , we can immediately resolve all the components B_i that contain border terminals. Those that contain border terminals guessed not to be in the same component with b have to be included in the solution, whereas those containing terminals required to be reachable from b need to be disjoint with the solution. We are left with a number of components B_i , and using the least possible number of additional edges we need to cut out a given number of components containing terminals. Each component B_i can be described by two numbers: a_i , the number of edges of B_i , and b_i , the number of terminals in $V(B_i) \setminus \{b\}$. Hence, we need to minimize the sum of a_i in a subset of components subject to obtaining a given sum of b_i . This can be done in polynomial time via standard dynamic programming. We remark that we have to be careful about one technical detail: the equivalence class guessed to be in the same connected component as b might require a terminal, so it may happen that we need to make sure that at least one component B_i that contains a terminal is disjoint with the solution. Fortunately, this can be solved by adding an additional binary dimension to the dynamic program, which keeps track whether some component B_i containing a terminal has been already chosen to be disjoint with the solution. Details can be found in Appendix E.

4 FPT algorithm for NODE UNIQUE LABEL COVER

In this section we present a brief sketch of the algorithm for NODE UNIQUE LABEL COVER. The full version with all the formal proofs can be found in Appendix D. In the following description we point out the difficulties that arise due to considering a node-deletion variant, as well as means that we use to overcome them. In fact, the edge-deletion variant can be reduced to the node-deletion variant via a simple polynomial parameterized transformation; see Appendix D for an appropriate reduction.

NODE UNIQUE LABEL COVER **Parameter:** $k + s$
Input: An undirected graph G , a finite alphabet Σ of size s , an integer k , and for each edge $e \in E(G)$ and each its endpoint v a permutation $\psi_{e,v}$ of Σ , such that if $e = uv$ then $\psi_{e,u} = \psi_{e,v}^{-1}$.
Question: Does there exist a set $X \subseteq V(G)$ of size at most k and a function $\Psi : V(G) \setminus X \rightarrow \Sigma$ such that for any $uv \in E(G \setminus X)$ we have $(\Psi(u), \Psi(v)) \in \psi_{uv,u}$?

Relations $\psi_{e,u}$ are called *edge constraints*, function Ψ is called a *labeling* and set X is the *deletion set*. We can naturally extend the notion of labelings to subsets: a labeling of a subset $S \subseteq V(G)$ is a function from S to the alphabet that respects all the constraints in $G[S]$. Given a set S one can check in $O(s^2n^2)$ time whether S admits a consistent labeling via a simple breadth-first search on each connected component.

In case of the edge-deletion problems we could simply contract edges determined to be useless. Since NODE UNIQUE LABEL COVER is a vertex-deletion type of problem, when we decide that a vertex is not going to be a part of a solution, then we *bypass* this vertex. When bypassing a vertex v , we remove it from the graph and for each pair of neighbors $u_1, u_2 \in N_G(v)$ the constraint $\psi_{u_1u_2}$ needs to be restricted to simultaneously satisfy also $\psi_{vu_2,v} \circ \psi_{vu_1,u_1}$. For this reason, in Appendix D we consider a slightly more general version of the NODE UNIQUE

LABEL COVER problem, where edge constraints are partial permutations and each vertex has a list of available labels $\phi_v \subseteq \Sigma$. The additional *vertex constraints*, as we call lists ϕ_v , are due to the fact that a solution might remove all but one vertex of $N(v)$, and then even the restricted edge constraints between vertices of $N(v)$ are not enough. Details can be found in Appendix D.

4.1 Good cuts in the node-deletion variant

We need to adjust the notion of good separations also to the node-deletion problems. We gather formal definitions and proofs on good separations in node-deletion setting in Appendix C; the definitions there are slightly more general, as they are adjusted to work also for the technical details of the algorithm for the N-MWCU problem. Unfortunately, due to technical difficulties we need two different notions of separations. The first one is a natural extension of the corresponding definition in the edge-deletion variant.

Definition 4.1. Let G be a connected graph. A triple (Z, V_1, V_2) of subsets of $V(G)$ is called a (q, k) -good node separation, if $|Z| \leq k$, V_1 and V_2 are vertex sets of two different connected components of $G \setminus Z$ and $|V_1|, |V_2| > q$.

Similarly to edge-deletion variant, one can find in time $O(2^{O(\min(q,k)\log(q+k))}n^3 \log n)$ a (q, k) -good node separation in the graph, or conclude that no such exists. The second notion, called (q, k) -flower separation, tries to capture the situation, where a large number of small components is attached to a common, small interface (for a formal definition, see Appendix C). Again, there exists an $O(2^{O(\min(q,k)\log(q+k))}n^3 \log n)$ algorithm that either finds a (q, k) -flower separation, or concludes that no such exists. If a graph does not admit a (q, k) -good node separation or a (q, k) -flower separation, we can prove a similar result to Lemma 2.3 that encapsulates the high-connectivity behavior in G (a more general result is proven as Lemma C.5 in Appendix C).

Lemma 4.2. *If a connected graph G with terminals $T_b \subseteq V(G)$ does not contain a (q, k) -good node separation or a (q, k) -flower separation w.r.t. T_b then, for any $Z \subseteq V(G)$ of size at most k , the graph $G \setminus Z$ contains at most $(2q + 2)(2^k - 1) + |T_b| + 1$ connected components, out of which at most one has more than q vertices.*

Note that the number of connected components could not be bounded if we only used the first type of separations. This problem is the sole purpose of introducing flower separations as well. Fortunately, they admit enough structure to make the recursive understanding step still work.

4.2 Border problem and recursive understanding

The definition of the border problem is very natural. In the border problem the graph is also equipped with a set T_b of at most $4k$ border terminals. The behavior on the border terminals is expressed by a function $\mathcal{P} : T_b \rightarrow \Sigma \cup \{\emptyset\}$, which simply fixes the values of the labeling. We say that a solution (X, Ψ) is *consistent with \mathcal{P}* if $X \cap T_b = \mathcal{P}^{-1}(\emptyset)$ and $\Psi|_{T_b \setminus X} = \mathcal{P}|_{T_b \setminus X}$.

BORDER N-ULC

Input: A NODE UNIQUE LABEL COVER instance $\mathcal{J} = (G, \Sigma, k, (\phi_v)_{v \in V(G)}, (\psi_{e,v})_{e \in E(G), v \in e})$ with G being connected, and a set $T_b \subseteq V(G)$ of size at most $4k$.

Output: For each $\mathcal{P} \in \mathbb{P}(\mathcal{J})$, output a solution $\text{sol}_{\mathcal{P}} = (X_{\mathcal{P}}, \Psi_{\mathcal{P}})$ to the instance \mathcal{J} that is consistent with \mathcal{P} and $|X_{\mathcal{P}}|$ is minimum possible, or outputs $\text{sol}_{\mathcal{P}} = \perp$ if such a pair does not exist.

This definition allows a standard recursive understanding routine. Take $q = k(s + 1)^{4k} + 2k$ and suppose we are given a $(q, 2k)$ -good node separation (Z, V_1, V_2) of G . At least one set V_1, V_2

has at most $2k$ terminals, let it be V_1 . We recursively apply the algorithm to the instance trimmed to the graph induced by the set $N[V_1] \subseteq V_1 \cup Z$, where we also incorporate $N(V_1) \subseteq Z$ to the border terminals. Let us mark all the border terminals and all the vertices that appeared in any output solution. One can easily show that all the unmarked vertices can be safely bypassed, as their usage can be always replaced with using some marked ones. As we chose q to be large enough, we bypass at least one vertex and we obtain a recurrential equation that gives the claimed complexity bound. Given a (q, k) -flower separation with respect to T_b we can make a recursive understanding step in the same manner. Once neither a $(q, 2k)$ -good node separation nor a (q, k) -flower separation with respect to T_b is present, the graph admits the structure expressed by Lemma 4.2 and we may proceed to the high connectivity phase. Note that we also excluded good node separations with larger cutsets, as it will be useful in the future.

4.3 High connectivity phase

Recall that we are to compute the optimal solution for every behavior \mathcal{P} on border terminals. We iterate through all possible \mathcal{P} and perform computation for each separately; note that this gives $2^{O(k \log s)}$ overhead to the running time. From now on we can assume that \mathcal{P} is fixed.

Let us examine the structure of the instance after removing an optimal deletion set X , given by Lemma 4.2. We have at most $d = (q + 1)(2^k - 1) + 4k$ small components, each containing at most q vertices, and one component of unbounded size. Note that we can assume that this big component, denote its vertex set by $\mathbf{big}(X)$, contains more than q vertices, as otherwise the whole graph has size at most $q(d + 1)$ and a brute-force search runs within the claimed complexity bound. As feasibility of an empty deletion set can be checked in polynomial time, we assume that X is nonempty. Let Ψ be a labeling witnessing that X is a feasible deletion set.

We now would like to use a similar approach to the one presented in the illustration, that is, to condensate the high-connectivity behavior in G . Let us take A_1 to be the union of vertex sets of small connected components in $G \setminus X$; note that $|A_1| \leq qd$. For every $u \in X \cap N(\mathbf{big}(X))$ we construct a set $A^u \subseteq \mathbf{big}(X)$ such that $|A^u| = q + 1$, $G[A^u]$ is connected and A^u contains a neighbor of u ; note that this is possible due to $|\mathbf{big}(X)| > q$. Let us now take $A_2 = \bigcup_{u \in X \cap N(\mathbf{big}(X))} A^u$; note that $|A_2| \leq k(q + 1)$. Therefore, if using Lemma 1.1 we generate the random family \mathcal{F} for universe $U = V(G)$ and constants $a = qd + k(q + 1)$, $b = k$, then we know that there is a set $S_0 \in \mathcal{F}$ such that $A_1 \cup A_2 \subseteq S_0$ and $X \cap S_0 = \emptyset$. We guess the right S_0 by branching into $|\mathcal{F}|$ subcases, labeled by $S \in \mathcal{F}$. Formally, each branch produces a candidate for an optimal solution and the algorithm picks the smallest possible; we argue that the branch with S_0 guessed correctly produces an optimal solution. From now on, we have fixed a set S and we seek a deletion set X that is disjoint with S , but S contains all the vertex sets of small components after removing X , as well as, for every $u \in X$ adjacent to the big component, a vertex set inducing a connected subgraph of size at least $q + 1$ adjacent to u .

We refer to vertex sets of connected components of $G[S]$ as *stains*; note that these sets will induce connected subgraphs even after removing the solution. A stain is *big* if it is of size at least $q + 1$, otherwise it is *small*. Note that each big stain has to be entirely contained in $\mathbf{big}(X)$. Moreover, for each stain there must exist a consistent labeling of it, or otherwise we may immediately terminate the branch. We would like to mimic the identification step from the illustration for big stains. Consider two big stains C_1, C_2 . As the graph does not admit a $(q, 2k)$ -good node separation, by Menger's theorem we can distinguish $2k + 1$ paths P^0, P^1, \dots, P^{2k} from C_1 to C_2 that are internally vertex-disjoint. Having fixed $\Psi|_{C_1}$ (recall that we assume that all the vertices of S do not belong to X), for every path P^i we can find a labeling Ψ^i of C_2 that is Ψ propagated via the path P^i to C_2 assuming that P^i is not hit by the deletion set, or conclude that P^i must be hit. As at most k paths P^i are hit, majority of the paths are not hit.

It follows that for these paths $\Psi^i = \Psi|_{C_2}$. Therefore, having fixed a labeling on one big stain, we can also fix the labeling on all the big stains, even if we do not know the deletion set: for each big stain we just compute all the labelings Ψ^i and take the one that appears most of the time. Let S^{big} be the union of all big stains. We branch into at most s branches, in each fixing a labeling of one vertex from S^{big} , thus fixing the labeling of the whole S^{big} .

Denote $D^{\text{big}} = N(S^{\text{big}})$; note that $D^{\text{big}} \cap S = \emptyset$. By our assumptions on S , we know that $X \cap N(\text{big}(X)) \subseteq D^{\text{big}}$. Intuitively, S^{big} together with D^{big} play the role of the core vertex b , while the components of $G \setminus (S^{\text{big}} \cup D^{\text{big}})$ can be solved more or less independently. In fact, it still holds that for each such component, let C be its vertex set, either C is disjoint with X and entirely contained in $\text{big}(X)$, or all the small stains in C are separated from each other and from $\text{big}(X)$ by X , while the deletion set X contains $C \setminus S$ as well as the whole $N(C)$. This corresponds to the observation in the edge-deletion variant, that no component of the graph after removing the core has a nontrivial intersection with the deletion set.

We now construct an optimal deletion set by iteratively considering components of $G \setminus (S^{\text{big}} \cup D^{\text{big}})$ and for each trying to consistently label it. Recall that we begin with S^{big} already labeled. Let C be the vertex set of the next component. If the whole $N(C)$ is already included in the deletion set, small stains contained in C are already separated from $\text{big}(X)$ so we can simply include the whole $C \setminus S$ to the deletion set and label all the remaining small stains arbitrarily. Otherwise, let $C^* = G[S^{\text{big}} \cup N[C]]$ be the graph induced by C , S^{big} and the subset of D^{big} that is adjacent to both of them. There are three possibilities:

- The labeling produced so far can be consistently extended on the whole $V(C^*)$, excluding vertices already chosen to the deletion set. In this case it is not hard to prove that it is safe to simply label C with this labeling and proceed to the next component.
- Even $G[C]$ cannot be consistently labeled. We infer that C is disjoint with $\text{big}(X)$ and $C \setminus S$ as well as $N(C)$ has to be entirely in the deletion set, so we include $C \setminus S$ and $N(C)$ to the constructed deletion set, label the remaining small stains in $G[C \cap S]$ arbitrarily and proceed. Note that in this manner we strictly increase the size of the constructed deletion set, as $N(C)$ was not entirely contained in the so far constructed deletion set.
- There exist some consistent labelings of $G[C]$, but none of them can be extended to vertices of $N(C) \subseteq D^{\text{big}}$ that are not yet included into the deletion set, so that it is consistent with already labeled vertices. We branch into several subcases:
 - Either C is disjoint with $\text{big}(X)$; in this branch we proceed as in the previous point.
 - Otherwise, we create a branch for every labeling of $G[C]$ (note that there are at most s of them). In the corresponding branch we label C accordingly. Moreover, as the labeling cannot be consistently extended to vertices of $N(C)$ that are not yet included in the deletion set, it follows that one can find just at most two vertices that are creating the problem. More precisely, in polynomial time one can find a set $B \subseteq N(C)$ of at most two vertices that are not yet included in the deletion set, such that the current labeling cannot be extended even to B . At least one vertex of B has to be included in the deletion set, so we branch further into at most two subcases, in each including one of them.

Once we have dealt with all the components of $G \setminus (S^{\text{big}} \cup D^{\text{big}})$, the set of vertices unlabeled so far is a subset of D^{big} , which is dominated by S^{big} . Again, we can use the same observation that if the labeling cannot be extended consistently on the whole remaining graph, there are just two vertices causing a problem, and the algorithm can branch which of them to delete.

Note that in the whole procedure we produce at most $2s + 1$ branches, and in each the size of the constructed deletion set increases. Thus, the branching routine works in $O((2s + 1)^k (ns)^{O(1)})$ time. For detailed analysis, see Appendix D.

5 Conclusions

In this paper we presented a new technique of designing parameterized algorithms for cut problems. The natural next step is to try to find further applications of this framework. On the other hand, the algorithms obtained using our framework run in time complexity $2^{O(k^2 \log k)} \cdot \text{poly}(n)$. Is the dependence on k optimal (for example, under Exponential Time Hypothesis)? Or is it possible to design algorithms with running time $2^{O(k \log k)} \cdot \text{poly}(n)$ or even $2^{O(k)} \cdot \text{poly}(n)$ for at least some of the considered problems?

A Preliminaries

A.1 Notation

We use standard graph notation. As the definitions vary among the algorithms, we introduce problem-specific notation at the beginning of each corresponding section, describing whether we work on graphs, multigraphs or some other structures. Generally, by a graph we denote the pair $G = (V, E)$ consisting vertex set V and edge set E . By $V(G)$ we denote the vertex set of G and by $E(G)$ the edge set. For $F \subseteq E(G)$ by $V(F)$ we denote the set of endpoints of F . For $V_1, V_2 \subseteq V(G)$, by $\delta(V_1, V_2)$ we denote the set of edges with one endpoint in V_1 and second in V_2 . For $W \subseteq V(G)$, by $G[W]$ we denote the graph induced by W . For $u \in V(G)$, by $N(u)$ we denote the neighborhood of u , i.e., $N(u) = \{v \mid uv \in E(G)\}$, and the closed neighborhood is defined by $N[u] = N(u) \cup \{u\}$. We extend this notion to subsets in the following manner: for $W \subseteq V(G)$, $N[W] = \bigcup_{u \in W} N[u]$, and $N(W) = N[W] \setminus W$. If X is a set of vertices or edges, by $G \setminus X$ we denote the graph G with edges or vertices of X removed.

A.2 Contractions

In this section we gather the definitions and simple facts connected to the notion of an *edge contraction*. Our definition works in multigraphs.

Definition A.1. Given a multigraph G and an edge $uv \in E(G)$, *contraction of uv* is the operation that yields a new multigraph G' with following properties:

- $V(G') = V(G) \setminus \{u, v\} \cup \{w_{uv}\}$, where $w_{uv} \notin V(G)$ is a new vertex;
- $E(G')$ is first constructed from $E(G)$ by deleting all edges uv , and then substituting all occurrences of u or v by w_{uv} in all the other edges.

In other words, we preserve multiple edges but delete loops. With contraction of an edge uv we can associate a mapping $\iota_{uv} : V(G) \rightarrow V(G')$ by setting $\iota_{uv}(u) = \iota_{uv}(v) = w_{uv}$ and $\iota_{uv}(t) = t$ for all $t \in V(G) \setminus \{u, v\}$. For $w \in V(G)$, we say that vertex w is *contracted onto* w_{uv} . By somewhat abusing the notation we identify all the edges of $E(G')$ with the edges from $E(G)$ in which they originated. By contracting the edge set $S \subseteq E(G)$ we mean consecutively contracting edges of S in an arbitrary order. Note that if some edge already disappeared from the graph because of becoming a loop, we omit this contraction. We usually use ι to denote the composition of all the mappings ι_{uv} corresponding to the performed contractions. The following lemma, which can be considered a folklore, implies that the order of performing the contractions does not matter.

Lemma A.2. Let G be a multigraph, $D \subseteq E(G)$ be a set of edges and G' be the graph obtained by contracting D in an arbitrary order. Then the following holds:

- $\iota(u) = \iota(v)$ if and only if u and v can be connected via a path consisting of edges from D , for $u, v \in V(G)$.
- $\iota^{-1}(v)$ induces a connected subgraph of G , for $v \in V(G')$;
- $E(G') \subseteq E(G)$;
- an edge $vw \in E(G)$ is contained also in $E(G')$ if and only if $\iota(v) \neq \iota(w)$;
- if $X \subseteq V(G')$, then $G'[X]$ is a maximal connected component if and only if $G[\iota^{-1}(X)]$ is;
- in particular, G is connected if and only if G' is;

- for every set F such that $D \cap F = \emptyset$, $G' \setminus F$ can be obtained by contracting D in $G \setminus F$.

From Lemma A.2 it follows, that given a graph $G = (V, E)$ and the set $D \subseteq E$, in time $O(|V| + |E|)$ we can construct the graph G' obtained by contracting edges of D . We simply find connected components of the graph (V, D) , construct a new vertex for each of them, and for every edge of E check whether it should be introduced in G' , and where.

B Proofs of Lemmas omitted in the main body of the paper

Proof of Lemma 1.1. For $a = 0$ or $b = 0$ the lemma is trivial; assume then $a, b \geq 1$.

We use the standard technique of splitters. A (n, r, r^2) -*splitter* is a family of functions from $\{1, 2, \dots, n\}$ to $\{1, 2, \dots, r^2\}$, such that for any subset $X \subseteq \{1, 2, \dots, n\}$ of size r , one of the functions in the family is injective on X . Naor et al. [37] gave an explicit construction of an (n, r, r^2) -splitter of size $O(r^6 \log r \log n)$ using $O(\text{poly}(r) \cdot n \log n)$ time.

Without loss of generality, assume that $a \leq b$ and that $U = \{1, 2, \dots, n\}$. Let $c = \min(a + b, n)$. We construct a (n, c, c^2) -splitter using the algorithm of Naor et al. and, for each function f in the splitter and for each subset $S' \subseteq \{1, 2, \dots, c^2\}$ of size a , we put into the family \mathcal{F} the set $f^{-1}(S') \subseteq U$. Assume now that we have $A, B \subseteq U$ such that $|A| \leq a$ and $|B| \leq b$. Obtain A' and B' by adding arbitrary elements of $U \setminus (A \cup B)$ to A and B so that $|A'| + |B'| = c$. By definition of the splitter, there exists some f in the splitter that is injective on $A' \cup B'$. To finish the proof one needs to observe that if we take $S = f^{-1}(f(A'))$, then $A \subseteq S$ and $B \cap S = \emptyset$.

The time bound and the size of the constructed family \mathcal{F} follow from the bound on the size of the splitter and the fact that there are at most $\binom{a+b}{a} = 2^{O(a \log(a+b))}$ choices for the set S' ; note that for fixed f and S' , the set $f^{-1}(S')$ can be computed in linear time. \square

A well-known result by Nagamochi and Ibaraki [35] states that the graph can be efficiently sparsified while preserving all the essential connectivity.

Lemma B.1 ([35]). *Given an undirected graph $G = (V, E)$ and an integer k , in $O(k(|V| + |E|))$ time we can obtain a set of edges $E_0 \subseteq E$ of size at most $(k+1)(|V| - 1)$, such that for any edge $uv \in E \setminus E_0$ in the graph (V, E_0) there are at least $k+1$ edge-disjoint paths between u and v .*

Proof. The algorithm performs exactly $k+1$ iterations. In each iteration it finds a spanning forest F of the graph G , adds all the edges of F to E_0 and removes all the edges of F from the graph G .

Observe that for any edge uv remaining in the graph G , the vertices u and v are in the same connected components in each of the forests found. Hence in each of those forests we can find a path between u and v ; thus, we obtain $k+1$ edge-disjoint paths between u and v . \square

We are ready to present proofs of lemmas regarding algorithms finding good edge separations.

Proof of Lemma 2.2. The algorithm iterates through all the sets from the family \mathcal{F} , obtained from Lemma 1.1 for universe $U = E(G)$ and constants $a = 2q$ and $b = k$. For a set $S \in \mathcal{F}$, we obtain a new graph H by contracting all the edges of S . Let $\iota : V(G) \rightarrow V(H)$ be the mapping that maps every vertex of G to the vertex it is contracted onto. We say that a vertex $u \in V(H)$ is *big* if $|\iota^{-1}(u)| > q$. Now, for every pair of big vertices $u_1, u_2 \in V(H)$ we compute some minimum edge cut between u_1 and u_2 if it is of size at most k , or find that it has to have larger size. This can be done in $O(k^2 n^3)$ time, since first we can sparsify the graph by removing all the edges outside of the set E_0 returned by Lemma B.1, and next for each of the $O(n^2)$ pairs of big vertices using the classical algorithm by Ford and Fulkerson in $O(k^2 n)$ time find a cut of size at most k if it exists. Assume that for some pair of big vertices u_1, u_2 we have found a

minimum edge cut F_{u_1, u_2} , of size at most k . We claim that F_{u_1, u_2} induces a (q, k) -good edge separation of G , which can be returned as the output of the algorithm.

Let $v_1 \in \iota^{-1}(u_1)$ and $v_2 \in \iota^{-1}(u_2)$ be arbitrary vertices. Let V_1, V_2 be the sets of vertices reachable from v_1, v_2 in $G \setminus F_{u_1, u_2}$, respectively. We claim that (V_1, V_2) is a (q, k) -good edge separation of G . Firstly, observe that V_1 and V_2 are disjoint. Otherwise there would be a path from v_1 to v_2 in G that avoids F_{u_1, u_2} , which after applying the contractions would become a path from u_1 to u_2 in H that avoids F_{u_1, u_2} . Secondly, observe that $V_1 \cup V_2 = V(G)$. It follows from the well-known properties of minimum cuts that in $H \setminus F_{u_1, u_2}$ every vertex is reachable either from u_1 or from u_2 . As graphs $G[\iota^{-1}(u)]$ are connected for $u \in H$, we find that in G every vertex is reachable either from v_1 or from v_2 . Thirdly, observe that $|V_1|, |V_2| \geq q$, as $\iota^{-1}(u_1) \subseteq V_1$ and $\iota^{-1}(u_2) \subseteq V_2$.

We are left with proving that if the graph admits a (q, k) -good edge separation, then for at least one set $S_0 \in \mathcal{F}$ we obtain two big vertices that can be separated by an edge cut of size at most k . This ensures that if no solution has been found for any $S \in \mathcal{F}$, then the algorithm can safely provide a negative answer. Fix some (q, k) -good edge separation (V_1, V_2) and let T_1, T_2 be arbitrary subtrees of $G[V_1]$ and $G[V_2]$, respectively, each having exactly $q + 1$ vertices. By the choice of family \mathcal{F} , there exists $S_0 \in \mathcal{F}$ that contains all the edges of T_1 and T_2 , but is disjoint with $\delta(V_1, V_2)$. In the step when S_0 is considered, after applying contractions all the vertices of T_1 are contracted onto one vertex u_1 , all the vertices of T_2 are contracted onto one vertex u_2 , but edges from $\delta(V_1, V_2)$ are not being contracted. Hence, we obtain big vertices u_1, u_2 that can be separated by an edge cut of size at most k . \square

Proof of Lemma 2.3. Claim (i) follows directly from the fact, that removing an edge from the graph can increase the number of connected components by at most one. For Claim (ii), observe that if two components had at least q vertices, then F could serve as an edge cut between their vertex sets of size at most k . It follows that the minimum edge cut between their vertex sets would also have size bounded by k , hence it would induce a (q, k) -good edge separation in G . \square

Lemma B.2. *There exists a randomized algorithm that, given an undirected, connected graph $G = (V, E)$ along with integers q and k , in time $\tilde{O}(2^{O(\min(q, k) \log(q+k))}(|V| + |E|))$ either finds a (q, k) -good edge separation, or correctly concludes that no such separation exists with probability at least $(1 - 1/|V|^2)$.*

Proof. Let (V_1, V_2) be a (q, k) -good edge separation. Intuitively, we want to have an edge-contraction process such that no edge of $\delta(V_1, V_2)$ is contracted and each vertex which remains is big, because then any cut of size at most k gives a (q, k) -good edge separation, which we can find by using Karger's algorithm.

The algorithm iterates through all the sets from the family \mathcal{F} , obtained from Lemma 1.1 for universe $U = E(G)$ and constants $a = 2q$ and $b = k$. For a set $S \in \mathcal{F}$, we obtain a new graph H' by contracting all the edges of S . Let $\iota' : V(G) \rightarrow V(H')$ be the mapping that maps every vertex of G to the vertex it is contracted onto. We say that a vertex $u' \in V(H')$ is *big* if $|\iota'^{-1}(u')| > q$ and *small* otherwise. Let $S' \subseteq E(H')$ be the set of edges of H' having at least one small endpoint. We construct a graph H , by contracting all the edges of S' in H' . Let $\iota : V(G) \rightarrow V(H)$ be the mapping from the graph G to the graph H . Note that after contracting all the edges of S' all the vertices are big in the graph H with respect to ι . By using Karger's algorithm [25], in $\tilde{O}(\min(q, k) \log(q + k)(|V| + |E|))$ time we find the minimum cut in the graph H with probability at least $\left(1 - \frac{1}{2^{c(\min(q, k) \log(q+k))|V|^2 \log |V|}}\right)$, for some constant c . If the minimum cut found is of size at most k , it immediately gives a (q, k) -good edge separation in the graph G , since all the vertices of H are big.

We are left with proving that if G admits a (q, k) -good edge separation (V_1, V_2) , then for at least one set $S_0 \in \mathcal{F}$ the graph H contains a cut of size at most k , providing some (possibly

different) (q, k) -good edge separation. This ensures that if no solution has been found for any $S \in \mathcal{F}$, then the algorithm can safely provide a negative answer. For each vertex $u \in N(V_2) \subseteq V_1$ let T^u be an arbitrary subtree of $G[V_1]$ containing the vertex u , having exactly $q + 1$ vertices. Similarly, for each vertex $u \in N(V_1)$ let T^u be an arbitrary subtree of $G[V_2]$ containing u , having exactly $q + 1$ vertices. By the choice of the family \mathcal{F} , there exists $S_0 \in \mathcal{F}$ that contains all the edges of T^u for each $u \in V(\delta(V_1, V_2))$, but at the same time S_0 is disjoint with $\delta(V_1, V_2)$. In the step when S_0 is considered, after applying contractions, for each $u \in V(\delta(V_1, V_2))$ all the vertices of T_u are contracted onto one vertex u' , which is big. However, the edges from $\delta(V_1, V_2)$ are not being contracted. Observe, that in the graph H' no edge of $\delta(V_1, V_2)$ has a small endpoint, and consequently all of the edges of $\delta(V_1, V_2)$ are present in the graph H , and they induce a cut of size at most k .

Note that, the algorithm of Karger is used $O(2^{O(\min(q,k)\log(q+k))} \log |V|)$ times, and therefore, by the union bound, if our algorithm does not find a (q, k) -good edge separation, with probability at least $(1 - 1/|V|^2)$ it does not exist. \square

C Good separations in node-deletion problems

As we consider node-deletion problems in the next two sections, we need to define an appropriate variant of good separations. In the edge-deletion variant, we might have assumed that we only consider cuts that separate the graph into exactly two connected components; this is no longer a case in the node-deletion variant. In the node-deletion problems we use two types of separations. In the first one, we require that, after removal of the separator, at least two connected components are large.

Definition C.1. Let G be a connected graph and $V^\infty \subseteq V(G)$ a set of undeletable vertices. A triple (Z, V_1, V_2) of subsets of $V(G)$ is called a (q, k) -good node separation, if $|Z| \leq k$, $Z \cap V^\infty = \emptyset$, V_1 and V_2 are vertex sets of two different connected components of $G \setminus Z$ and $|V_1 \setminus V^\infty|, |V_2 \setminus V^\infty| > q$.

In the second one we require a bunch of connected components with the same neighbourhood.

Definition C.2. Let G be a connected graph, $V^\infty \subseteq V(G)$ a set of undeletable vertices, and $T_b \subseteq V(G)$ a set of border terminals in G . A pair $(Z, (V_i)_{i=1}^\ell)$ is called a (q, k) -flower separation in G (with regard to border terminals T_b), if the following holds:

- $1 \leq |Z| \leq k$ and $Z \cap V^\infty = \emptyset$; the set Z is the *core* of the flower separation $(Z, (V_i)_{i=1}^\ell)$;
- V_i are vertex sets of pairwise different connected components of $G \setminus Z$, each set V_i is a *petal* of the flower separation $(Z, (V_i)_{i=1}^\ell)$;
- $V(G) \setminus (Z \cup \bigcup_{i=1}^\ell V_i)$, called a *stalk*, contains more than q vertices of $V \setminus V^\infty$;
- for each petal V_i we have $V_i \cap T_b = \emptyset$, $|V_i \setminus V^\infty| \leq q$ and $N_G(V_i) = Z$;
- $|\bigcup_{i=1}^\ell V_i \setminus V^\infty| > q$.

We now show how to detect the aforementioned separations using Lemma 1.1, similarly as it is done in the case of good edge separations.

Lemma C.3. *Given a connected graph G with undeletable vertices $V^\infty \subseteq V(G)$ and integers q and k , one may find in $O(2^{O(\min(q,k)\log(q+k))} n^3 \log n)$ time a (q, k) -good node separation of G , or correctly conclude that no such separation exists.*

Proof. The algorithm iterates through all the sets from the family \mathcal{F} , obtained from Lemma 1.1 for universe $U = V(G) \setminus V^\infty$ and constants $a = 2q + 2$ and $b = k$. For a set $S \in \mathcal{F}$, we obtain a new graph H by contracting all the edges between vertices of $S \cup V^\infty$ in G . Let $\iota : V(G) \rightarrow V(H)$ be the mapping that maps every vertex of G to the vertex it is contracted onto, and let $S' = \iota(S \cup V^\infty)$. We say that a vertex $u \in S'$ is *big* if $|\iota^{-1}(u) \setminus V^\infty| > q$.

In the graph H , we assign weight ∞ to all vertices of S' and weight 1 to all vertices of $V(H) \setminus S'$. In this weighted graph, for every pair of big vertices u_1 and u_2 , we compute a minimum node cut between u_1 and u_2 if it is of size at most k , or find that it has to have larger size. This can be done in $O(kn^3)$ time using the Gomory-Hu tree extended to node weighted separations by Granot and Hassin [22]. That is we can use $|V(H)| - 1$ applications of the classic Ford-Fulkerson algorithm, each of which consuming $O(kn^2)$ time, since after finding $k + 1$ vertex disjoint paths we may stop the algorithm. Assume that for some pair of big vertices u_1, u_2 we have found a minimum node cut F_{u_1, u_2} , of size at most k . We claim that F_{u_1, u_2} induces a (q, k) -good node separation of G , which can be returned as the output of the algorithm.

Let $v_1 \in \iota^{-1}(u_1) \setminus V^\infty$ and $v_2 \in \iota^{-1}(u_2) \setminus V^\infty$ be arbitrary vertices. Note that $F_{u_1, u_2} \subseteq V(G) \setminus V^\infty$, as only vertices of $V(H) \setminus S' = V(G) \setminus (S \cup V^\infty)$ have finite weights. Let V_1, V_2 be the sets of vertices reachable from v_1, v_2 in $G \setminus F_{u_1, u_2}$, respectively. We claim that (F_{u_1, u_2}, V_1, V_2) is a (q, k) -good node separation of G . Indeed: V_1 and V_2 are defined as vertex sets of two connected components of $G \setminus F_{u_1, u_2}$; moreover, $V_1 \neq V_2$ as F_{u_1, u_2} separates u_1 from u_2 in H , and therefore v_1 from v_2 in G . Finally, observe that $|V_1 \setminus V^\infty|, |V_2 \setminus V^\infty| > q$, as $\iota^{-1}(u_1) \subseteq V_1$ and $\iota^{-1}(u_2) \subseteq V_2$.

We are left with proving that if the graph admits a (q, k) -good node separation, then for at least one set $S_0 \in \mathcal{F}$ we obtain two big vertices that can be separated by a node cut of size at most k . This ensures that if no solution has been found for any $S \in \mathcal{F}$, then the algorithm can safely provide a negative answer. Fix some (q, k) -good node separation (Z, V_1, V_2) and let T_1, T_2 be arbitrary subtrees of $G[V_1]$ and $G[V_2]$, respectively, each having exactly $q + 1$ vertices that are not in V^∞ . As $|Z| \leq k$, by the choice of family \mathcal{F} , there exists $S_0 \in \mathcal{F}$ that contains $(V(T_1) \cup V(T_2)) \setminus V^\infty$, but is disjoint with Z . In the step when S_0 is considered, after applying contractions all the vertices of T_1 are contracted onto one vertex u_1 , all the vertices of T_2 are contracted onto one vertex u_2 , but vertices of Z get weight 1. Hence, we obtain big vertices u_1, u_2 that can be separated by a node cut of size at most k (note that the algorithm does not necessarily find precisely the cut Z in this step). \square

Lemma C.4. *Given a connected graph G with undeletable vertices $V^\infty \subseteq V(G)$ and border terminals $T_b \subseteq V(G)$ and integers q and k , one may find in $O(2^{O(\min(q, k) \log(q+k))} n^3 \log n)$ time a (q, k) -flower separation in G w.r.t. T_b , or correctly conclude that no such flower separation exists.*

Proof. We first note that, given a set $Z \subseteq V(G)$ of size at most k , we can in $O(n^2)$ time verify whether there exists a (q, k) -flower separation with Z as the core, that is, $(Z, (V_i)_{i=1}^\ell)$ for some choice of the family of petals $(V_i)_{i=1}^\ell$. Indeed, we may simply iterate over connected components of $G \setminus Z$ using a simple dynamic program. For each prefix of the sequence of connected components and for each $n' \leq n$ we compute, whether some of the components can be chosen to be petals so that the total number of vertices of $V(G) \setminus V^\infty$ in the petals is equal to n' . When we consider the next connected component, if it does not satisfy requirements for a petal then we cannot take it as a petal (and we take the value of the cell computed in the last iteration for the same value of n'). However, if it does satisfy these requirements, then we either not take it to be a petal (and do the same as previously) or take it (and we take the value of the cell computed in the last iteration for the value n' decremented by the number of vertices from $V \setminus V^\infty$ in the considered component). There exists a flower separation with Z as the centre if and only if some of the values for $q + 1 \leq n' \leq |V(G) \setminus (V^\infty \cup Z)| - q - 1$ is true

in the last iteration. It is trivial to augment the dynamic program with backlinks, so that the flower separation can be retrieved.

To prove the lemma, we iterate through all the sets from the family \mathcal{F} , obtained from Lemma 1.1 for universe $U = V(G) \setminus V^\infty$ and constants $a = q$ and $b = k$. For a set $S \in \mathcal{F}$, we obtain a new graph H by contracting all the edges between vertices of $S \cup V^\infty$ in G . Let $\iota : V(G) \rightarrow V(H)$ be the mapping that maps every vertex of G to the vertex it is contracted onto, and let $S' = \iota(S \cup V^\infty)$. We say that a vertex $u \in S'$ is *interesting* if $|\iota^{-1}(u) \setminus V^\infty| \leq q$. For each interesting vertex u with $|N_H(u)| \leq k$ we verify whether there exists a (q, k) -flower separation $(Z, (V_i)_{i=1}^\ell)$ in G w.r.t. T_b with the core $Z = N_H(u)$; note that $N_H(u) \subseteq V(G)$. We output such a flower separation if we find one. If no flower separation is found for any choice of S and u , we conclude that no (q, k) -flower separation exists in G w.r.t. T_b . The time bound follows from the fact that for each vertex u , we can verify whether u is interesting and compute $N_H(u)$ in $O(n^2)$ time, and then within the same complexity check if $N_H(u)$ is the core of some (q, k) -flower separation. To finish the proof of the lemma we need to show that if the algorithm concludes that there is no appropriate flower separation in the graph, then this conclusion is correct.

To this end, assume that there exists a (q, k) -flower separation $(Z, (V_i)_{i=1}^\ell)$ in G w.r.t. T_b . Note that $|V_1 \setminus V^\infty| \leq q$ ($\ell \geq 1$ since $|\bigcup_{i=1}^\ell V_i \setminus V^\infty| > q$) and $|Z| \leq k$, so by the properties of the family \mathcal{F} there exists a set $S_0 \in \mathcal{F}$ with $(V_1 \setminus V^\infty) \subseteq S_0$ and $Z \cap S_0 = \emptyset$. Recall that $N_G(V_1) = Z$; thus, in the graph H constructed for the set S_0 there is a vertex $u \in V(H)$ with $\iota^{-1}(u) = V_1$. Note that u is an interesting vertex (as $|V_1 \setminus V^\infty| \leq q$) and $N_H(u) = Z$ (as $N_G(V_1) = Z$ by the definition of the flower separation). Therefore the algorithm considers $Z = N_H(u)$ and finds a (q, k) -flower separation in G w.r.t. T_b . \square

We conclude this section with a lemma that shows that if we do not have any good node or flower separations, then any k -cut not only cannot split the graph into two large components, but also cannot split the graph into too many small ones.

Lemma C.5. *If a connected graph G with undeletable vertices $V^\infty \subseteq V(G)$ and border terminals $T_b \subseteq V(G)$ does not contain a (q, k) -good node separation or a (q, k) -flower separation w.r.t. T_b then, for any $Z \subseteq V(G) \setminus V^\infty$ of size at most k , the graph $G \setminus Z$ contains at most $(2q + 2)(2^k - 1) + |T_b| + 1$ connected components containing a vertex of $V \setminus V^\infty$, out of which at most one has more than q vertices not in V^∞ .*

Proof. Let $Z \subseteq V(G) \setminus V^\infty$, $|Z| \leq k$. First, if there are at least two connected components of $G \setminus Z$ with more than q vertices outside V^∞ , then a minimal subset of Z separating these two components would induce a (q, k) -good node separation in G . Thus, in $G \setminus Z$ we have at most one connected component with more than q vertices outside V^∞ and at most $|T_b|$ connected components that contain a vertex from T_b . We denote the remaining connected components containing at least one vertex of $V \setminus V^\infty$ as *nice* ones; they have at most q vertices outside V^∞ each. Let us partition them with respect to their neighbourhood (which is a subset of Z). Note that, if there exists a set $Z' \subseteq Z$, such that more than $2q$ nice connected components of $G \setminus Z$ that are adjacent to exactly Z' , then there exists a (q, k) -flower separation in G w.r.t. T_b with core Z' and petals being $q + 1$ of aforementioned nice connected components of $G \setminus Z$. As there are at most $2^k - 1$ nonempty subsets of Z , the lemma follows. \square

D Full details of the algorithm for NODE UNIQUE LABEL COVER

This section is devoted to fixed-parameter tractability of the NODE UNIQUE LABEL COVER problem, parameterized by both the size of the cutset and the size of the alphabet. We solve a bit more general version of the problem; the new definition makes the presentation easier.

For an alphabet Σ , a binary relation $\psi \subseteq \Sigma \times \Sigma$ is called a *partial permutation* if for any $\alpha \in \Sigma$ both $(\{\alpha\} \times \Sigma) \cap \psi$ and $(\Sigma \times \{\alpha\}) \cap \psi$ are of size at most one. For a partial permutation ψ , its reverse is defined as $\psi^{-1} = \{(\beta, \alpha) : (\alpha, \beta) \in \psi\}$. For any two partial permutations ψ_1, ψ_2 their composition is defined as

$$\psi_2 \circ \psi_1 = \{(\alpha, \gamma) : \exists \beta \in \Sigma (\alpha, \beta) \in \psi_1 \wedge (\beta, \gamma) \in \psi_2\}.$$

Note that a composition of two partial permutations is a partial permutation itself.

We are now ready to define the considered problem.

<p>NODE UNIQUE LABEL COVER</p> <p>Input: An undirected graph G, a finite alphabet Σ of size s, an integer k, for each vertex $v \in V(G)$ a set $\phi_v \subseteq \Sigma$ and for each edge $e \in E(G)$ and each its endpoint v a partial permutation $\psi_{e,v}$ of Σ, such that if $e = uv$ then $\psi_{e,u} = \psi_{e,v}^{-1}$.</p> <p>Question: Does there exist a set $X \subseteq V(G)$ of size at most k and a function $\Psi : V(G) \setminus X \rightarrow \Sigma$ such that for any $v \in V(G) \setminus X$ we have $\Psi(v) \in \phi_v$ and for any $uv \in E(G \setminus X)$ we have $(\Psi(u), \Psi(v)) \in \psi_{uv,u}$?</p>	<p>Parameter: $k + s$</p>
---	---

The relations $\psi_{e,u}$ are called *edge constraints*, the sets ϕ_v are called *vertex constraints*, the function Ψ is called a *labeling* and the set X is the *deletion set*.

We note that often in literature in the standard NODE UNIQUE LABEL COVER definition the constraints are required to be (non-partial) permutations and vertex constraints are omitted (i.e., are equal to Σ); we here relax these requirements to ease the presentation.

Before we start, we note that the edge-deletion variant (where we look for a deletion set being a subset of edges; we are to label all vertices, but we do not need to satisfy the constraints on the deleted edges) reduces to the defined above node-deletion variant.

Indeed, first observe that in the NODE UNIQUE LABEL COVER problem we can assume that additionally we are given in the input a set of undeletable vertices $V^\infty \subseteq V(G)$ and we are to find a deletion set X disjoint with V^∞ : we can reduce this variant to the original one by replacing each undeletable vertex with a clique on $k + 1$ vertices, with constraints on the edges of the clique being identities. Second, given an EDGE UNIQUE LABEL COVER instance $(G, \Sigma, k, (\phi_v)_{v \in V(G)}, (\psi_{e,v})_{e \in E(G), v \in e})$, we can first make all vertices of G undeletable, and then subdivide each edge, so that the edge constraints on the two halves of the edge $e = uv$ of G compose to the constraint $\psi_{e,u}$; the new vertices introduced in this operation are kept deletable.

Note that we may assume that the input graph G in the NODE UNIQUE LABEL COVER problem is connected; otherwise, we may solve the problem on each connected component, for all budgets between 0 and k , separately. During the course of the algorithm, we maintain the connectivity of G . We denote by n the number of vertices of the graph of the currently considered NODE UNIQUE LABEL COVER instance.

We also assume that we are working in the random access memory model, and that the elements of Σ can be compared in constant time.

The description of the algorithm consists of a sequence of *steps*. Each step is accompanied with some lemmas and a discussion that justifies its correctness and verifies complexity bounds.

D.1 Labelings

We first extend the notion of labeling to arbitrary subsets of $V(G)$.

Definition D.1. Given an instance $\mathcal{J} = (G, \Sigma, k, (\phi_v)_{v \in V(G)}, (\psi_{e,v})_{e \in E(G), v \in e})$ and a set $S \subseteq V(G)$, a function $\Psi^S : S \rightarrow \Sigma$ is called a *labeling* if it satisfies all constraints on $G[S]$ in \mathcal{J} , that is: for each $v \in S$ we have $\Psi^S(v) \in \phi_v$ and for each $uv \in E(G[S])$ we have $(\Psi^S(u), \Psi^S(v)) \in \psi_{uv,u}$.

The following lemma is a straightforward corollary of the fact that the edge constraints are partial permutations.

Lemma D.2. *Let $\mathcal{J} = (G, \Sigma, k, (\phi_v)_{v \in V(G)}, (\psi_{e,v})_{e \in E(G), v \in e})$ be a NODE UNIQUE LABEL COVER instance and let $A \subseteq V(G)$ be an arbitrary subset of the vertex set that induces a connected subgraph of G . Then, for any $v \in A$ and $\alpha \in \Sigma$ there exists at most one labeling $\Psi^A : A \rightarrow \Sigma$ such that $\Psi^A(v) = \alpha$, and in $O(s|A|^2)$ time one can find such a labeling or correctly conclude that it does not exist. Consequently, for each set $A \subseteq V(G)$ such that $G[A]$ is connected, there are at most s labelings of A and those can be enumerated in $O(s^2|A|^2)$ time.*

Proof. Note that for any $uw \in E(G[A])$, if $\Psi^A(u)$ is fixed, then there exists at most one value $\Psi^A(w)$ such that $(\Psi^A(u), \Psi^A(w)) \in \psi_{uw,u}$, and such a value $\Psi^A(w)$ can be found in $O(s)$ time. The first claim of the lemma follows from the assumption that $G[A]$ is connected: the labeling Ψ^A can be found using a breadth-first search, and then verified to satisfy all the constraints in $O(s|A|^2)$ time. For the second claim, we simply iterate over all possible values $\Psi^A(v)$ for one fixed vertex $v \in A$. \square

D.2 Operations on the input graph

In this section we define two basic operations the algorithm repetitively applies on the graph and show their key properties.

Definition D.3. Let $\mathcal{J} = (G, \Sigma, k, (\phi_v)_{v \in V(G)}, (\psi_{e,v})_{e \in E(G), v \in e})$ be a NODE UNIQUE LABEL COVER instance, let $u, v \in V(G)$ and ψ be a partial permutation of Σ . By *updating an edge uv with a constraint ψ* we mean the following operation: if $uv \notin E(G)$, then we add an edge uv to the graph G with constraints $\psi_{uv,u} = \psi$, $\psi_{uv,v} = \psi^{-1}$; otherwise, we modify the constraints on the edge uv in G by replacing $\psi_{uv,u}$ with $\psi_{uv,u} \cap \psi$ and $\psi_{uv,v}$ with $\psi_{uv,v} \cap \psi^{-1}$.

Informally speaking, updating an edge uv with ψ is equivalent to adding a new edge between u and v with this constraint; however, we use the definition above to avoid multiple edges in G . Note that obviously updating an edge cannot spoil the assumption of connectivity of G . The following lemma is immediate.

Lemma D.4. *Let \mathcal{J}' be a NODE UNIQUE LABEL COVER instance obtained from \mathcal{J} by updating an edge uv with a constraint ψ . Then a pair (X, Ψ) is a solution to \mathcal{J}' if and only if it is a solution to \mathcal{J} that satisfies the following additional property: either $u \in X$ or $v \in X$ or $(\Psi(u), \Psi(v)) \in \psi$.*

The second operation allows us to remove a vertex that, for some reason, may be omitted by X .

Definition D.5. Let $\mathcal{J} = (G, \Sigma, k, (\phi_v)_{v \in V(G)}, (\psi_{e,v})_{e \in E(G), v \in e})$ be a NODE UNIQUE LABEL COVER instance and $v \in V(G)$. By *bypassing the vertex v* we mean the following operation:

1. remove the vertex v with its incident edges from the graph G ;
2. for each $u \in N_G(v)$ we replace ϕ_u with $\phi_u \cap \{\beta : \exists \alpha \in \phi_v, (\alpha, \beta) \in \psi_{uv,v}\}$;
3. for each $u_1, u_2 \in N_G(v)$, $u_1 \neq u_2$, we update an edge u_1u_2 with a constraint $\psi_{vu_2,v} \circ \psi_{vu_1,u_1}$.

Lemma D.6. *Let \mathcal{J}' be a NODE UNIQUE LABEL COVER instance obtained from an instance*

$$\mathcal{J} = (G, \Sigma, k, (\phi_v)_{v \in V(G)}, (\psi_{e,v})_{e \in E(G), v \in e})$$

by bypassing a vertex v with $\phi_v \neq \emptyset$. Then the following holds:

- if (X, Ψ) is a solution to \mathcal{J}' , then there exists $\alpha \in \Sigma$ such that $(X, \Psi \cup \{(v, \alpha)\})$ is a solution to \mathcal{J} ;
- if (X, Ψ) is a solution to \mathcal{J} that satisfies $v \notin X$, then $(X, \Psi|_{V(G) \setminus \{v\}})$ is a solution to \mathcal{J}' .

Proof. For the first claim, pick α as follows: if $N_G(v) \subseteq X$, pick $\alpha \in \phi_v$ arbitrarily, and otherwise take arbitrary $w \in N_G(v) \setminus X$ and set α such that $(\Psi(w), \alpha) \in \psi_{vw,w}$ and $\alpha \in \phi_v$. Note that such α exists as, by the definition of the bypassing operation, in \mathcal{J}' the vertex constraint for w are contained in $\{\beta : \exists \alpha' \in \phi_v(\beta, \alpha') \in \psi_{vw,w}\}$. We claim that $(X, \Psi \cup \{(v, \alpha)\})$ is a solution to \mathcal{J} . Clearly, $\Psi \cup \{(v, \alpha)\}$ satisfies all vertex constraints of $V(G) \setminus (X \cup \{v\})$ as well as all edge constraints on edges not incident to v , as those constraints in \mathcal{J} are supersets of the corresponding constraints in \mathcal{J}' . Moreover, the choice of α ensures that $\alpha \in \phi_v$. We are left with verifying edge constraints $\psi_{uv,v}$ for $u \in N_G(v)$. If $u = w$, clearly $(\alpha, \Psi(w)) \in \psi_{vw,v}$ by the choice of α . Otherwise, by the definition of the bypassing operation, $(\Psi(w), \Psi(u)) \in \psi_{uw,v} \circ \psi_{vw,w}$. Since $(\Psi(w), \alpha) \in \psi_{vw,w}$, we infer that $(\alpha, \Psi(u)) \in \psi_{uw,v}$ and the claim is proven.

For the second claim, denote $\alpha = \Psi(v)$. To prove the claim we need to verify that $\Psi|_{V(G) \setminus \{v\}}$ satisfies vertex constraints on $N_G(v)$ (that may shrink during the bypassing operation) and edge constraints on edges between vertices in $N_G(v)$ (that are updated during the bypassing operation). First consider a vertex $u \in N_G(v) \setminus X$. Since (X, Ψ) is a solution to \mathcal{J} and $v \notin X$, we have $\Psi(u) \in \phi_u$, $\alpha \in \phi_v$ and $(\Psi(u), \alpha) \in \psi_{uv,u}$. Thus $\Psi(u) \in \{\beta : \exists \alpha' \in \phi_v(\alpha', \beta) \in \psi_{uv,v}\}$ and $\Psi(u)$ satisfies the vertex constraint at u in \mathcal{J}' . Second, consider two vertices $u_1, u_2 \in N_G(v) \setminus X$, $u_1 \neq u_2$. Since (X, Ψ) is a solution to \mathcal{J} and $v \notin X$, we have $(\Psi(u_1), \alpha) \in \psi_{vu_1,u_1}$ and $(\alpha, \Psi(u_2)) \in \psi_{vu_2,v}$. Therefore $(\Psi(u_1), \Psi(u_2)) \in \psi_{vu_2,v} \circ \psi_{vu_1,u_1}$ and the claim is proven. \square

During the course of the algorithm we perform bypassing operations multiple times, which can drastically increase the number of edges, even if the graph was sparse in the beginning. Therefore, we measure the complexity of our algorithm only in n , the number of vertices, and always use only the trivial quadratic bound on the number of edges.

D.3 Borders and recursive understanding

In the case of the NODE UNIQUE LABEL COVER problem, the definition of the border variant is much more natural than in the previously considered problems: informally speaking, for each vertex on the border, we need to know whether it is deleted and if not, what label is assigned to it. More formally, given a NODE UNIQUE LABEL COVER instance $\mathcal{J} = (G, \Sigma, k, (\phi_v)_{v \in V(G)}, (\psi_{e,v})_{e \in E(G), v \in e})$ and a set of border terminals $T_b \subseteq V(G)$, for any function $\mathcal{P} : T_b \rightarrow \Sigma \cup \{\text{⊗}\}$, we say that a solution (X, Ψ) to \mathcal{J} is consistent with \mathcal{P} , if $X \cap T_b = \mathcal{P}^{-1}(\text{⊗})$ and $\Psi|_{T_b \setminus X} = \mathcal{P}|_{T_b \setminus X}$. Let $\mathbb{P}(\mathcal{J})$ be the set of all functions from T_b to $\Sigma \cup \{\text{⊗}\}$. We define the border problem as follows.

BORDER N-ULC

Input: A NODE UNIQUE LABEL COVER instance $\mathcal{J} = (G, \Sigma, k, (\phi_v)_{v \in V(G)}, (\psi_{e,v})_{e \in E(G), v \in e})$ with G being connected, and a set $T_b \subseteq V(G)$ of size at most $4k$.

Output: For each $\mathcal{P} \in \mathbb{P}(\mathcal{J})$, output a solution $\text{sol}_{\mathcal{P}} = (X_{\mathcal{P}}, \Psi_{\mathcal{P}})$ to the instance \mathcal{J} that is consistent with \mathcal{P} and $|X_{\mathcal{P}}|$ is minimum possible, or outputs $\text{sol}_{\mathcal{P}} = \perp$ if such a pair does not exist.

Note that $|\mathbb{P}(\mathcal{J})| \leq (s+1)^{4k}$ and all output sets $X_{\mathcal{P}}$ contain at most $k(s+1)^{4k}$ vertices in total. Let $q = k(s+1)^{4k} + 2k$; if $|V(G)| > q + 2k$, then there are at least $|V(G)| - q - 2k$ vertices in G that are not in T_b nor in any of the output sets $X_{\mathcal{P}}$.

The following lemma prepares the ground for a recursive step in our BORDER N-ULC algorithm.

Lemma D.7. *Assume we are given an instance $\mathcal{J}_b = (G, \Sigma, k, (\phi_v)_{v \in V(G)}, (\psi_{e,v})_{e \in E(G), v \in e}, T_b)$ of BORDER N-ULC, and two disjoint sets of vertices $Z, V^* \subseteq V(G)$, such that $|Z| \leq 2k$, $N_G(V^*) \subseteq Z$, $|V^* \cap T_b| \leq 2k$ and the subgraph of G induced by $W := V^* \cup Z_W$ is connected, where $Z_W := N_G(V^*)$. Denote $T_b^* = (T_b \cup Z_W) \cap W$ and $\mathcal{J}_b^* = (G[W], \Sigma, k, (\phi_v)_{v \in W}, (\psi_{e,v})_{e \in E(G[W]), v \in e}, T_b^*)$. Then \mathcal{J}_b^* is a proper BORDER N-ULC instance. Moreover, if we denote by $(\mathbf{sol}_{\mathcal{P}}^*)_{\mathcal{P} \in \mathbb{P}(\mathcal{J}_b^*)}$ an arbitrary output to the BORDER N-ULC instance \mathcal{J}_b^* and*

$$U(\mathcal{J}_b^*) = T_b^* \cup \bigcup \{X_{\mathcal{P}}^* : \mathcal{P} \in \mathbb{P}(\mathcal{J}_b^*), \mathbf{sol}_{\mathcal{P}}^* = (X_{\mathcal{P}}^*, \Psi_{\mathcal{P}}^*) \neq \perp\},$$

then there exists a correct output $(\mathbf{sol}_{\mathcal{P}})_{\mathcal{P} \in \mathbb{P}(\mathcal{J}_b)}$ to the BORDER N-ULC instance \mathcal{J}_b such that whenever $\mathbf{sol}_{\mathcal{P}} = (X_{\mathcal{P}}, \Psi_{\mathcal{P}}) \neq \perp$ then $X_{\mathcal{P}} \cap V^ \subseteq U(\mathcal{J}_b^*)$.*

Proof. The claim that \mathcal{J}_b^* is a proper BORDER N-ULC instance follows directly from the assumptions that $G[W]$ is connected, $|Z_W| \leq |Z| \leq 2k$ and $|V^* \cap T_b| \leq 2k$. In the rest of the proof we justify the second claim of the lemma.

Fix $\mathcal{P} \in \mathbb{P}(\mathcal{J}_b)$. Assume that there exists a solution to the instance \mathcal{J}_b that is consistent with \mathcal{P} ; let $(X_{\mathcal{P}}, \Psi_{\mathcal{P}})$ be such a solution with minimum possible $|X_{\mathcal{P}}|$. To prove the lemma we need to show a second solution $(X'_{\mathcal{P}}, \Psi'_{\mathcal{P}})$ to \mathcal{J}_b that is consistent with \mathcal{P} , $|X'_{\mathcal{P}}| \leq |X_{\mathcal{P}}|$ and $X'_{\mathcal{P}} \cap V^* \subseteq U(\mathcal{J}_b^*)$.

Define $\mathcal{P}^* : T_b^* \rightarrow \Sigma \cup \{\bullet\}$ as follows: $\mathcal{P}^*(v) = \mathcal{P}(v)$ if $v \in T_b$, $\mathcal{P}^*(v) = \bullet$ if $v \in Z_W \cap X_{\mathcal{P}}$ and $\mathcal{P}^*(v) = \Psi_{\mathcal{P}}(v)$ if $v \in Z_W \setminus X_{\mathcal{P}}$. Note that, as $(X_{\mathcal{P}}, \Psi_{\mathcal{P}})$ is consistent with \mathcal{P} , this definition is proper even for vertices in $Z \cap T_b$.

Moreover, for the same reason $(X_{\mathcal{P}} \cap W, \Psi_{\mathcal{P}}|_{W \setminus X_{\mathcal{P}}})$ is a solution to \mathcal{J}_b^* that is consistent with \mathcal{P}^* . Therefore the output $\mathbf{sol}_{\mathcal{P}^*}$ to \mathcal{J}_b^* is different than \perp ; let $\mathbf{sol}_{\mathcal{P}^*} = (X_{\mathcal{P}^*}^*, \Psi_{\mathcal{P}^*}^*)$. By definition, $|X_{\mathcal{P}^*}^*| \leq |(X_{\mathcal{P}} \cap W)|$.

Let us define $X'_{\mathcal{P}} = (X_{\mathcal{P}} \setminus W) \cup X_{\mathcal{P}^*}^*$ and $\Psi'_{\mathcal{P}} : V(G) \setminus X'_{\mathcal{P}} \rightarrow \Sigma \cup \{\bullet\}$ as $\Psi'_{\mathcal{P}}(v) = \Psi_{\mathcal{P}^*}^*(v)$ if $v \in W$ and $\Psi'_{\mathcal{P}}(v) = \Psi_{\mathcal{P}}(v)$ otherwise. Note that this definition is consistent as $X'_{\mathcal{P}} \cap W = X_{\mathcal{P}^*}^*$ and $X'_{\mathcal{P}} \setminus W = X_{\mathcal{P}} \setminus W$. Clearly, $|X'_{\mathcal{P}}| \leq |X_{\mathcal{P}}|$. To finish the proof we need to show two things: first, that $\Psi'_{\mathcal{P}}$ satisfies the properties so that $(X'_{\mathcal{P}}, \Psi'_{\mathcal{P}})$ is a solution to \mathcal{J}_b , and, second, that $(X'_{\mathcal{P}}, \Psi'_{\mathcal{P}})$ is consistent with \mathcal{P} .

For the first claim, recall that $Z_W = W \cap Z = N(V^*)$ and $Z_W \subseteq T_b^*$. By the definition of T_b^* and \mathcal{P}^* , we have that $X_{\mathcal{P}^*}^* \cap Z_W = X'_{\mathcal{P}} \cap Z_W = X_{\mathcal{P}} \cap Z_W$ and $\Psi_{\mathcal{P}^*}^*|_{Z_W} = \Psi'_{\mathcal{P}}|_{Z_W} = \Psi_{\mathcal{P}}|_{Z_W}$. First consider a vertex $v \notin X'_{\mathcal{P}}$. If $v \in W$ then $v \notin X_{\mathcal{P}^*}^*$ and $\Psi'_{\mathcal{P}}(v) = \Psi_{\mathcal{P}^*}^*(v) \in \phi_v$. Otherwise, $v \notin X_{\mathcal{P}}$ and $\Psi'_{\mathcal{P}}(v) = \Psi_{\mathcal{P}}(v) \in \phi_v$. Second, consider an edge $e = uv$ of $G \setminus X'_{\mathcal{P}}$. If $e \in E(G[W])$, then $u, v \in W \setminus X_{\mathcal{P}^*}^*$ and

$$(\Psi'_{\mathcal{P}}(u), \Psi'_{\mathcal{P}}(v)) = (\Psi_{\mathcal{P}^*}^*(u), \Psi_{\mathcal{P}^*}^*(v)) \in \psi_{e,u}.$$

Similarly, in the other case we have $e \in E(G \setminus V^*)$, $u, v \in V(G) \setminus (V^* \cup X_{\mathcal{P}})$ and

$$(\Psi'_{\mathcal{P}}(u), \Psi'_{\mathcal{P}}(v)) = (\Psi_{\mathcal{P}}(u), \Psi_{\mathcal{P}}(v)) \in \psi_{e,u}.$$

We conclude that $(X'_{\mathcal{P}}, \Psi'_{\mathcal{P}})$ is a solution to the BORDER N-ULC instance \mathcal{J}_b .

For the second claim, let $v \in T_b$ and consider two cases: either $v \in W$ or $v \notin W$. If $v \in W$, then $v \in T_b^*$ and $\mathcal{P}(v) = \mathcal{P}^*(v)$. We have $v \in X'_{\mathcal{P}}$ if and only if $v \in X_{\mathcal{P}^*}^*$ which happens if and only if $\mathcal{P}^*(v) = \bullet$; moreover, if $v \notin X'_{\mathcal{P}}$, we have $\Psi'_{\mathcal{P}}(v) = \Psi_{\mathcal{P}^*}^*(v) = \mathcal{P}^*(v) = \mathcal{P}(v)$. In the case $v \notin W$, then $v \in X'_{\mathcal{P}}$ if and only if $v \in X_{\mathcal{P}}$ which happens if and only if $\mathcal{P}(v) = \bullet$; moreover, if $v \notin X'_{\mathcal{P}}$ then $\Psi'_{\mathcal{P}}(v) = \Psi_{\mathcal{P}}(v) = \mathcal{P}(v)$. This finishes the proof that $(X'_{\mathcal{P}}, \Psi'_{\mathcal{P}})$ is consistent with \mathcal{P} and concludes the proof of the lemma. \square

Note that in Lemma D.7 we have $|U(\mathcal{J}_b^*) \cap V^*| \leq q$. We are now ready to present the recursive steps of the algorithm.

Step D.1. Assume we are given an instance $\mathcal{J}_b = (G, \Sigma, k, (\phi_v)_{v \in V(G)}, (\psi_{e,v})_{e \in E(G), v \in e}, T_b)$ of BORDER N-ULC. Invoke first the algorithm of Lemma C.3 in a search for a $(q, 2k)$ -good node separation (with $V^\infty = \emptyset$). If it returns a good node separation (Z, V_1, V_2) , let $j \in \{1, 2\}$ be such that $|V_j \cap T_b| \leq 2k$ and denote $Z^* = Z$, $V^* = V_j$. Otherwise, if it returns that no such good node separation exists in G , invoke the algorithm of Lemma C.4 in a search for a (q, k) -flower separation w.r.t. T_b (with $V^\infty = \emptyset$ again). If it returns that no such flower separation exists in G , pass the instance \mathcal{J}_b to the next step. Otherwise, if it returns a flower separation $(Z, (V_i)_{i=1}^\ell)$, denote $Z^* = Z$ and $V^* = \bigcup_{i=1}^\ell V_i$.

In the case we have obtained Z^* and V^* (either from Lemma C.3 or Lemma C.4), invoke the algorithm recursively for the BORDER N-ULC instance \mathcal{J}_b^* defined as in the statement of Lemma D.7 for separator Z^* and set V^* , obtaining an output $(\mathbf{sol}_\mathcal{P}^*)_{\mathcal{P} \in \mathbb{P}(\mathcal{J}_b^*)}$. Compute the set $U(\mathcal{J}_b^*)$. Bypass (in an arbitrary order) all vertices of $V^* \setminus U(\mathcal{J}_b^*)$ to obtain a new instance \mathcal{J}'_b (observe that for each bypassed vertex v we have $\phi_v \neq \emptyset$, which is a necessary condition for bypassing). Recall that $T_b^* \subseteq U(\mathcal{J}_b^*)$, so no border terminal get bypassed. Restart the algorithm on the new instance \mathcal{J}'_b and obtain a family of solutions $(\mathbf{sol}'_\mathcal{P})_{\mathcal{P} \in \mathbb{P}(\mathcal{J}'_b)}$. For every $\mathcal{P} \in \mathbb{P}(\mathcal{J}_b)$, if $\mathbf{sol}'_\mathcal{P} = \perp$ then output $\mathbf{sol}_\mathcal{P} = \perp$ as well, while if $\mathbf{sol}_\mathcal{P} = (X_\mathcal{P}, \Psi'_\mathcal{P})$ then obtain $\Psi_\mathcal{P}$ by extending $\Psi'_\mathcal{P}$ on $U(\mathcal{J}_b^*)$ using Lemma D.2 (we justify that such an extension exists in Lemma D.8) and output $\mathbf{sol}_\mathcal{P} = (X_\mathcal{P}, \Psi_\mathcal{P})$.

Let us first verify that the application of Lemma D.7 is justified. Indeed, by the definitions of the good node separation and the flower separation, as well as the choice of V^* , we have in both cases $|V^* \cap T_b| \leq 2k$ and that $G[V^* \cup N_G(V^*)]$ is connected. Moreover, note that the recursive call is applied to the graph with strictly smaller number of vertices than G : in the case of a good node separation, V_2 is removed from the graph, and in the case of a flower separation, recall that the definition of the flower separation requires $Z \cup \bigcup_{i=1}^\ell V_i$ to be a proper subset of $V(G)$. Finally, in both cases $|V^*| > q$, and $|V^* \setminus U(\mathcal{J}_b^*)| \geq |V^*| - q \geq 1$ vertices are bypassed in Step D.1.

The following lemma verifies the correctness of Step D.1.

Lemma D.8. *Assume we are given an instance $\mathcal{J}_b = (G, \Sigma, k, (\phi_v)_{v \in V(G)}, (\psi_{e,v})_{e \in E(G), v \in e}, T_b)$ of BORDER N-ULC, on which Step D.1 is applied, and let \mathcal{J}'_b be an instance after all bypassing operations of Step D.1 are applied. Let $(\mathbf{sol}'_\mathcal{P})_{\mathcal{P} \in \mathbb{P}(\mathcal{J}'_b)}$ be a correct output to \mathcal{J}'_b . Then there exists a correct output $(\mathbf{sol}_\mathcal{P})_{\mathcal{P} \in \mathbb{P}(\mathcal{J}_b)}$ to \mathcal{J}_b , such that:*

- $\mathbf{sol}_\mathcal{P} = \perp$ if $\mathbf{sol}'_\mathcal{P} = \perp$;
- if $\mathbf{sol}'_\mathcal{P} = (X'_\mathcal{P}, \Psi'_\mathcal{P})$ then $\Psi'_\mathcal{P}$ can be consistently extended to $V(G)$ and for every such extension $\Psi_\mathcal{P}$ the pair $(X'_\mathcal{P}, \Psi_\mathcal{P})$ is a correct output for \mathcal{P} in \mathcal{J}_b ;

Proof. The lemma is a straightforward corollary of Lemma D.7 and the properties of the bypassing operation described in Lemma D.6. Lemma D.7 ensures us that each vertex of $V^* \setminus U(\mathcal{J}_b^*)$ is omitted by some optimal solution for every $\mathcal{P} \in \mathbb{P}(\mathcal{J}_b)$, which enables us to use Lemma D.6. Note that existence of the extension is asserted by the second claim of Lemma D.6. \square

We are left with an analysis of the time complexity of Step D.1. The applications of Lemmas C.3 and C.4 use $O(2^{O(\min(q, 2k) \log(q+2k))} n^3 \log n) = O(2^{O(k^2 \log s)} n^3 \log n)$ time. Let $n' = |V^*|$; the recursive step is applied to the graph with at most $n' + 2k$ vertices and, after bypassing, there are at most $n - n' + q$ vertices left. Moreover, each bypassing operation takes $O(sn^2)$ time, the computation of $U(\mathcal{J}_b^*)$ takes $O((s+1)^{4k} n)$ time, and extending the labelings from the trimmed instance takes $O((s+1)^{4k} sn^2)$ time. The values of $s = |\Sigma|$ and k do not change in this step. Therefore, we have the following recursive formula for time complexity as a function

of the number of vertices of G :

$$T(n) \leq \max_{q+1 \leq n' \leq n-2k-1} \left(2^{O(k^2 \log s)} n^3 \log n + T(n' + 2k) + T(n - n' + q) \right). \quad (2)$$

Note that the function $p(t) = t^4 \log t$ is convex, so the maximum of the expression is attained at one of the ends. A straightforward inductive check of both of the ends proves that we have indeed the claimed bound on the complexity, i.e., $T(n) = O(2^{O(k^2 \log s)} n^4 \log n)$.

We conclude this section with a note that Lemma C.5 asserts that, if Step D.1 is not applicable, then for any set $Z \subseteq V(G)$ of size at most k , the graph $G \setminus Z$ contains at most $t := (2q + 2)(2^k - 1) + 4k + 1$ connected components, out of which at most one has more than q vertices.

D.4 Brute force approach

If the graph is reduced by Step D.1 is small, the algorithm may apply a straightforward brute-force approach to the BORDER N-ULC problem. In this section we describe this method formally.

Lemma D.9. *Assume we are given an instance $\mathcal{J}_b = (G, \Sigma, k, (\phi_v)_{v \in V(G)}, (\psi_{e,v})_{e \in E(G), v \in e}, T_b)$ of BORDER N-ULC. Let $X \subseteq V(G)$ be a set of size at most k and let $\mathcal{P} \in \mathbb{P}(\mathcal{J}_b)$. Then, in time $O(s^2 n^2)$, one can compute a function $\Psi : V(G) \setminus X \rightarrow \Sigma$ such that (X, Ψ) is a solution to \mathcal{J}_b consistent with \mathcal{P} , or correctly conclude that no such function exists.*

Proof. We apply Lemma D.2 to the vertex set of every connected component of the graph induced by $A = V(G) \setminus X$. For each output labeling, we verify if it is consistent with \mathcal{P} . \square

Lemma D.10. *A correct output to an instance $\mathcal{J}_b = (G, \Sigma, k, (\phi_v)_{v \in V(G)}, (\psi_{e,v})_{e \in E(G), v \in e}, T_b)$ of BORDER N-ULC can be computed in $O((s + 1)^{4k} s^2 k n^{k+2})$ time.*

Proof. We apply Lemma D.9 for each $\mathcal{P} \in \mathbb{P}(\mathcal{J}_b)$ (there are at most $(s + 1)^{4k}$ choices) and for each deletion set $X \subseteq V(G)$ with $|X| \leq k$ (at most $(k + 1)n^k$ choices). \square

Step D.2. If $|V(G)| \leq qt + k$, apply Lemma D.10 to find a correct output to a BORDER N-ULC instance $\mathcal{J}_b = (G, \Sigma, k, (\phi_v)_{v \in V(G)}, (\psi_{e,v})_{e \in E(G), v \in e}, T_b)$.

Recall that $q = (s + 1)^{2k} k = 2^{O(k \log s)}$ and $t = (2q + 2)(2^k - 1) + 2k + 1 = 2^{O(k \log s)}$. Thus, if Step D.2 is applicable, its running time is $O(2^{O(k^2 \log s)})$.

D.5 High connectivity phase

Assume we have a BORDER N-ULC instance $\mathcal{J}_b = (G, \Sigma, k, (\phi_v)_{v \in V(G)}, (\psi_{e,v})_{e \in E(G), v \in e}, T_b)$ where Steps D.1 and D.2 are not applicable. In this section we show how to exploit high connectivity of the graph implied by Lemma C.5 to compute a correct output to \mathcal{J}_b . To this end, fix $\mathcal{P} \in \mathbb{P}(\mathcal{J}_b)$; we focus on finding the solution $\text{sol}_{\mathcal{P}}$. First, let us solve some simple cases.

Step D.3. For each $\mathcal{P} \in \mathbb{P}(\mathcal{J}_b)$, verify using Lemma D.9 whether there exists solution $\text{sol}_{\mathcal{P}} = (X_{\mathcal{P}}, \Psi_{\mathcal{P}})$ with $X_{\mathcal{P}} = \emptyset$. If yes, output such a solution.

Note that, if $|V(G)|$ is too large for Step D.2 to be applicable, for any set $Z \subseteq V(G)$ of size at most k , the bound on the number of connected components from Lemma C.5 implies that there exists exactly one connected component of $G \setminus Z$ with more than q vertices; denote its vertex set by $\text{big}(Z)$.

The set $\text{big}(Z)$ is an analogue of the component C_0 in the algorithm for the STEINER CUT problem. From now on the notation differs, as we try to avoid in the subsequent description

usage of the bypassing operation, which is an analogue of edge contraction. We remark that we could proceed similarly to the algorithm from Section E; however, we choose not to, as we find this way overcomplicated and harder to digest for this particular problem.

D.5.1 Interrogating sets

We now use Lemma 1.1 to get some more structure of the graph G .

Definition D.11. Let $Z \subseteq V(G)$ be a set of size at most k and let $S \subseteq V(G)$. We say that S *interrogates* Z if the following holds:

1. $S \cap Z = \emptyset$;
2. for any connected component C of $G \setminus Z$ with at most q vertices, all vertices of C belong to S ;
3. for any $v \in Z$, such that $N_G(v) \cap \mathbf{big}(Z) \neq \emptyset$, there exists a connected component of $G[S]$ that has more than q vertices and contains at least one neighbour of v .

Note that in the third point, the considered component has to be entirely contained in $\mathbf{big}(Z)$ due to its size.

Lemma D.12. *Let \mathcal{F} be a family obtained by the algorithm of Lemma 1.1 for universe $U = V(G)$ and constants $a = qt + (q + 1)k$ and $b = k$. Then, for any $Z \subseteq V(G)$ with $1 \leq |Z| \leq k$, there exists a set $S \in \mathcal{F}$ that interrogates Z .*

Proof. Fix $Z \subseteq V(G)$ with $|Z| \leq k$. Let A_1 be the union of vertex sets of all connected components of $G \setminus Z$ that have at most q vertices; by Lemma C.5, $|A_1| \leq qt$. For each $v \in Z$ such that $N_G(v) \cap \mathbf{big}(Z) \neq \emptyset$, fix $w_v \in N_G(v) \cap \mathbf{big}(Z)$ and a tree T_v with exactly $q + 1$ vertices that contains w_v and is contained in $\mathbf{big}(Z)$; note that this is possible due to $|\mathbf{big}(Z)| > q$. Let A_2 be the union of vertex sets of all trees T_v for $v \in Z$; clearly $|A_2| \leq (q + 1)k$. By Lemma 1.1, as $|A_1 \cup A_2| \leq qt + (q + 1)k$ and $|Z| \leq k$, there exists a set $S \in \mathcal{F}$ that contains $A_1 \cup A_2$ and is disjoint with Z . By the construction of the sets A_1 and A_2 , the set S interrogates Z and the lemma is proven. \square

Note that, as $q, t = 2^{O(k \log s)}$, the family \mathcal{F} of Lemma D.12 is of size $2^{O(k^2 \log s)} \log n$ and can be computed in $O(2^{O(k^2 \log s)} n \log n)$ time. Therefore we may branch, guessing a set S that interrogates a deletion set $X_{\mathcal{P}}$ for the solution $\mathbf{sol}_{\mathcal{P}} = (X_{\mathcal{P}}, \Psi_{\mathcal{P}})$ we are looking for.

Step D.4. Compute the family \mathcal{F} from Lemma D.12 and branch into $|\mathcal{F}|$ subcases, indexed by sets $S \in \mathcal{F}$. In a branch S we seek for a pair $(X_{\mathcal{P}}, \Psi_{\mathcal{P}})$ with minimum possible $|X_{\mathcal{P}}|$ that not only is a solution to J_b consistent with \mathcal{P} , but also satisfies that $X_{\mathcal{P}}$ is interrogated by S .

Lemma D.12 verifies the correctness of the branching of Step D.4; as discussed, the step is applied in $O(2^{O(k^2 \log s)} n \log n)$ time and leads to $O(2^{O(k^2 \log s)} \log n)$ subcases.

After choosing a set S , we may now slightly modify the set S to make it more regular.

Definition D.13. A vertex $v \in V(G)$ is said to be *forsaken*, if

- $v \in T_b$ and $\mathcal{P}(v) = \mathfrak{B}$; or
- $\phi_v = \emptyset$.

The forsaken vertices are those to be necessarily included in any solution $(X_{\mathcal{P}}, \Psi_{\mathcal{P}})$ consistent with \mathcal{P} .

Step D.5. As long as there exists a vertex $v \in V(G)$ that is not forsaken and $N_G[v] \cap S = \emptyset$, add v to S .

Step D.5 can clearly be applied in $O(sn^2)$ time (for all vertices it is applied to; note that Step D.5 is applied to one vertex v at a time and, by its application to the vertex v , it may become not applicable to the neighbours of v). We now discuss its correctness. Let $(X_{\mathcal{P}}, \Psi_{\mathcal{P}})$ be a solution to \mathcal{J}_b that is interrogated by S and consistent with \mathcal{P} . Then $X_{\mathcal{P}}$ is interrogated by $S \cup \{v\}$ unless $v \in X_{\mathcal{P}}$; assume then $v \in X_{\mathcal{P}}$. As $N_G[v] \cap S = \emptyset$, by the last property of an interrogating set v is not adjacent to any vertex of $\text{big}(X_{\mathcal{P}})$. Moreover, by the second property of an interrogating set, v is not adjacent to any vertex of connected component of $G \setminus X_{\mathcal{P}}$ of size at most q . We infer that $N_G[v] \subseteq X_{\mathcal{P}}$. Since v is not a forsaken vertex, there exists $\alpha \in \phi_v$ such that $(X_{\mathcal{P}} \setminus \{v\}, \Psi_{\mathcal{P}} \cup \{(v, \alpha)\})$ is a solution to \mathcal{J}_b that is consistent with \mathcal{P} (but not necessarily interrogated by S). Therefore $(X_{\mathcal{P}}, \Psi_{\mathcal{P}})$ is not a solution to \mathcal{J}_b consistent with \mathcal{P} with minimum possible $|X_{\mathcal{P}}|$, and we may omit it from consideration.

Step D.5 gives us the following property of the set S .

Lemma D.14. *After Step D.5 is applied, any vertex v that is not forsaken is contained in $N_G[S]$.*

Proof. If v is not forsaken and $v \notin N_G[S]$, then $N_G[v] \cap S = \emptyset$ and Step D.5 is applicable to v . \square

D.5.2 Labelings of big stains

Let us now focus on a fixed branch $S \in \mathcal{F}$.

Definition D.15. Each connected component of $G[S]$ is called a *stain*. A stain is *big* if it has more than q vertices, and *small* otherwise.

Let $S^{\text{big}} \subseteq S$ be the union of all vertex sets of big stains of $G[S]$. We now establish a crucial observation that the fact that G admits no $(q, 2k)$ -good node separations implies that there are only very few reasonable labelings for S^{big} .

Lemma D.16. *One can in $O(ksn^3 + ks^2n^2)$ time compute a family **PSI** of at most s labelings of S^{big} , such that for any solution (X, Ψ) to \mathcal{J}_b such that S interrogates X , there exists $\Psi^{\text{big}} \in \mathbf{PSI}$ with $\Psi|_{S^{\text{big}}} = \Psi^{\text{big}}$.*

Proof. If $S^{\text{big}} = \emptyset$ the lemma is trivial; assume then $S^{\text{big}} \neq \emptyset$. For any big stain with vertex set C in $G[S]$, by Lemma D.2 there are at most s labelings of C and all these labelings can be computed in $O(s^2|C|^2)$ time. Moreover, we know that any such a labeling is induced by fixing a label of one vertex of C ; in other words, for any two different labelings Ψ', Ψ'' of $G[C]$ and any $v \in C$, we have $\Psi'(v) \neq \Psi''(v)$.

Let C_1 and C_2 be vertex sets of two different big stains in $G[S]$. As G admits no $(q, 2k)$ -good node separations, by Menger's theorem there exists a sequence P^0, P^1, \dots, P^{2k} of $2k + 1$ paths, where each path starts in C_1 , ends in C_2 and the sets of internal vertices of those paths are pairwise disjoint. Moreover, such a sequence of paths P^0, P^1, \dots, P^{2k} can be found in $O(kn^2)$ time by the classic Ford-Fulkerson algorithm.

Let (X, Ψ) be a solution to \mathcal{J}_b that is interrogated by S . As $|X| \leq k$, for at least $k + 1$ indices $0 \leq i \leq 2k$, the path P^i does not contain a vertex from X (note that the endpoints of P^i are in $C_1 \cup C_2 \subseteq S$, and thus not in X). Denote the endpoints of P^i as $v_1^i \in C_1$ and $v_2^i \in C_2$. If P^i does not contain a vertex from X , the composition of all edge constraints on P^i (denote it by ψ^i) is a partial permutation such that $(\Psi(v_1^i), \Psi(v_2^i)) \in \psi^i$. We infer that for any $0 \leq i \leq 2k$ and any labeling Ψ' of $G[C_1]$ there exists at most one labeling $\Psi'_{C_2, i}$ of $G[C_2]$ such

that $(\Psi'(v_1^i), \Psi'_{C_2,i}(v_2^i)) \in \psi^i$. Moreover, given Ψ' , all labelings $\Psi'_{C_2,i}$ for all $0 \leq i \leq 2k$ can be computed in $O(ks(n + s|C_2|^2))$ time using Lemma D.2.

Let $\Psi' = \Psi|_{C_1}$. As at least $k+1$ paths P^i do not contain any vertices of X , for a majority of indices $0 \leq i \leq 2k$, the labelings Ψ'_{i,C_2} are the same labelings. For a fixed big stain C_1 and for each labeling Ψ' of $G[C_1]$, we can compute this majority labeling Ψ'_{maj,C_2} of $G[C_2]$ for any big stain $C_2 \neq C_1$ in time $O(kn^2 + ks(n + s|C_2|^2))$ (including the time needed to compute paths P^i). As there are at most n big stains, and s labelings of a fixed big stain C_1 , the lemma follows. \square

Note that for any $Z \subseteq V(G)$, $1 \leq |Z| \leq k$, there exists the component $\mathbf{big}(Z)$ and, if S interrogates Z , then there exists at least one big stain in $G[S]$ (note that we require here that Z is nonempty; the solutions with empty deletion sets are found by Step D.3). This observation, together with Lemma D.16, justifies the following step.

Step D.6. For each $\mathcal{P} \in \mathcal{J}$, in a branch with index S , if $G[S]$ contains no big stains, terminate this branch, and otherwise invoke Lemma D.16 to obtain a family \mathbf{PSI} and branch into at most s subcases, indexed by labelings $\Psi^{\mathbf{big}} \in \mathbf{PSI}$. For fixed \mathcal{P} , in a branch with indices S and $\Psi^{\mathbf{big}}$, we seek for a pair $(X_{\mathcal{P}}, \Psi_{\mathcal{P}})$ with minimum possible $|X_{\mathcal{P}}|$ such that $(X_{\mathcal{P}}, \Psi_{\mathcal{P}})$ is a solution to \mathcal{J}_b consistent with \mathcal{P} , $X_{\mathcal{P}}$ is interrogated by S and $\Psi|_{S^{\mathbf{big}}} = \Psi^{\mathbf{big}}$.

Each application of Step D.6 takes $O(ksn^3 + ks^2n^2)$ time and leads to $O(2^{O(k^2 \log s)} \log n)$ subcases in total.

D.5.3 Final bounded search tree algorithm

In this section we show how to finish the search for an appropriate output $\text{sol}_{\mathcal{P}}$ for $\mathcal{P} \in \mathbb{P}(\mathcal{J}_b)$, in a fixed branch with indices S and $\Psi^{\mathbf{big}}$. This is done in a standard framework of a bounded search tree algorithm. Formally speaking, we maintain a tuple $(X_0, Y, Y^{\Psi}, \Psi^Y)$, where $X_0 \subseteq V(G)$, $Y^{\Psi} \subseteq Y \subseteq V(G) \setminus X_0$ and $\Psi^Y : Y^{\Psi} \rightarrow \Sigma$ is a labeling of $G[Y^{\Psi}]$, and we say that a pair (X, Ψ) is a solution to the tuple $(X_0, Y, Y^{\Psi}, \Psi^Y)$ if the following holds:

- (X, Ψ) is a solution to \mathcal{J}_b ;
- $X_0 \subseteq X$;
- S interrogates X ;
- $\Psi|_{Y^{\Psi}} = \Psi^Y$;

Moreover, we maintain the invariants that $S^{\mathbf{big}} \subseteq Y^{\Psi}$, $S \subseteq Y$ and that all forsaken vertices are in X_0 .

Given a tuple $(X_0, Y, Y^{\Psi}, \Psi^Y)$, the algorithm looks for a solution (X, Ψ) with minimum possible $|X|$. In the course of the algorithm, we only add new elements to the elements of the tuple (sometimes while branching into a few subcases).

The intuition on the tuple $(X_0, Y, Y^{\Psi}, \Psi^Y)$ is as follows: X_0 are the vertices that the algorithm decided to include in a deletion set, and Y the vertices the algorithm decided not to include in a deletion set. Moreover, for some vertices $Y^{\Psi} \subseteq Y$ the algorithm already decided on their labeling. By Lemma D.2, if we fix a label on a vertex of a connected component of $G[Y]$, it propagates to the whole connected component. Thus, we may easily maintain an additional invariant that Y^{Ψ} contains all vertices of some connected components of $G[Y]$.

Note that in the above definition of a solution to a tuple $(X_0, Y, Y^{\Psi}, \Psi^Y)$ we do not use the function \mathcal{P} ; instead, we incorporate this function at the beginning into the other parts of the tuple. Formally, we construct the initial tuple via the following step.

Step D.7. Construct the initial tuple (X_0, Y, Y^Ψ, Ψ^Y) as follows. First start with X_0 being the set of forsaken vertices, $Y = S$, $Y^\Psi = S^{\text{big}}$ and $\Psi^Y = \Psi^{\text{big}}$. Second, for each $v \in T_b$:

1. if $\mathcal{P}(v) = \text{skull}$ and $v \in Y$, then terminate the current branch;
2. if $\mathcal{P}(v) = \text{skull}$ and $v \notin Y$, then do nothing (v is forsaken and already in X_0);
3. if $\mathcal{P}(v) \in \Sigma$ and $v \in Y^\Psi$, then if $\mathcal{P}(v) = \Psi^Y(v)$, do nothing, and otherwise terminate this branch;
4. if $\mathcal{P}(v) \in \Sigma$ and $v \notin Y^\Psi$, then first include v into Y in the case $v \notin Y$, and afterwards apply Lemma D.2 to the connected component C of $G[Y]$ that contains v for the label $\mathcal{P}(v)$ for v ; if the algorithm of Lemma D.2 returns a labeling Ψ^C , put C into Y^Ψ and extend Ψ^Y with Ψ^C , and otherwise terminate this branch.

The correctness of this step is straightforward. The step can be performed in $O(ks^2n^2)$ time (as $|T_b| \leq 2k$). Moreover, by Step D.7, either we terminate the branch or for each $v \in T_b$, $v \in X_0$ if $\mathcal{P}(v) = \text{skull}$ or otherwise $v \in Y^\Psi$ and $\Psi^Y(v) = \mathcal{P}(v)$.

The branching algorithm is described as a set of three *rules*. At each moment, we apply the first applicable rule.

First, let us define the stopping condition for the branching algorithm.

Rule D.1 (Finishing Rule). If $|X_0| > k$, terminate the current branch. If $G \setminus X_0$ admits a labeling that equals Ψ^Y on Y^Ψ (that can be found by Lemma D.2), output X_0 and this labeling as a solution in the current branch.

Note that recognizing whether Finishing Rule can be applied takes $O(s^2n^2)$ time. We now show a simple branching rule, justified by the following lemma.

Lemma D.17. *Let $A \subseteq V(G)$ and $\Psi^A : A \rightarrow \Sigma$ be a labeling of $G[A]$. If there does not exist a labeling of $G[N_{G \setminus X_0}[A]]$ that equals Ψ^A on the vertices of A , then there exists a set $B \subseteq N_{G \setminus X_0}(A)$ of size at most two, such that there does not exist a labeling of $G[A \cup B]$ that equals Ψ^A on A . Moreover, such a set B can be found on $O(sn^2)$ time.*

Proof. For each $v \in N_{G \setminus X_0}(A)$ pick an arbitrary vertex $w(v) \in A \cap N_G(v)$ and define $\Psi'(v) \in \phi_v$ such that $(\Psi'(v), \Psi^A(w(v))) \in \psi_{vw(v),v}$. If such $\Psi'(v)$ does not exist, then $B = \{v\}$ satisfies the conditions of the lemma. From the assumptions of the lemma $\Psi' \cup \Psi^A$ is not a valid labeling of $G[N_{G \setminus X_0}[A]]$. By the definition, $\Psi' \cup \Psi^A$ satisfies all vertex constraints, thus let uv be an edge with a constraint not satisfied by $\Psi' \cup \Psi^A$. As Ψ^A is a labeling of A , either u or v is not in A , assume then $v \notin A$. If $u \in A$, then $B = \{v\}$ satisfies the conditions of the lemma: for any function $\Psi^A \cup \{(v, \alpha)\}$, either $\alpha = \Psi'(v)$ and the constraint $\psi_{uv,u}$ is not satisfied, or $\alpha \neq \Psi'(v)$ and the constraint $\psi_{vw(v),v}$ is not satisfied. If $u \notin A$, then $B = \{u, v\}$ satisfies the conditions of the lemma: for any function $\Psi^A \cup \{(v, \alpha), (u, \beta)\}$, either $\alpha \neq \Psi'(v)$ and the constraint $\psi_{vw(v),v}$ is not satisfied, or $\beta \neq \Psi'(u)$ and the constraint $\psi_{uw(u),u}$ is not satisfied, or $\alpha = \Psi'(v)$ and $\beta = \Psi'(u)$ and the constraint $\psi_{uv,u}$ is not satisfied.

We note that we can compute Ψ' and find a constraint not satisfied by $\Psi' \cup \Psi^A$ in $O(sn^2)$ time. \square

Rule D.2 (Neighborhood Branching Rule). Apply Lemma D.2 to find in $O(s^2n^2)$ time a labeling of $N_{G \setminus X_0}[Y^\Psi]$ that equals Ψ^Y on Y^Ψ . If no such labeling exists, apply Lemma D.17 for $A = Y^\Psi$ to find the set B and branch into at most 2 subcases, including one of the vertices of B to X_0 .

Unfortunately, the neighborhood branching rule is not always applicable. Our goal in the remainder of the section is to show one additional rule that either extends Y^Ψ (without any branching), or performs some limited branching that always leads to at least one application of the Neighbourhood Branching Rule.

Let $D^{\text{big}} = N_G(S^{\text{big}}) \subseteq V(G) \setminus S$, $D^{\text{small}} = V(G) \setminus (S \cup D^{\text{big}})$ and $D = D^{\text{big}} \cup D^{\text{small}} = V(G) \setminus S$. Moreover, denote $S^{\text{small}} = S \setminus S^{\text{big}}$. The following lemma shows that on $G \setminus N_G[S^{\text{big}}]$, we have only very restricted options, and is an analogue of Lemma E.11.

Lemma D.18. *Let C be a vertex set of an arbitrary connected component of $G \setminus (X_0 \cup N_G[S^{\text{big}}])$. Let (X, Ψ) be a solution to the tuple (X_0, Y, Y^Ψ, Ψ^Y) . Then either:*

- $C \subseteq \text{big}(X)$, in particular $C \cap X = \emptyset$, or
- $C \setminus X \subseteq S^{\text{small}}$, that is, C contains vertices of the deletion set X and some small stains of S .

Moreover, let $C^* = S^{\text{big}} \cup (N_G[C] \setminus X_0)$ and assume additionally that there exists a labeling Ψ^* of $G[C^*]$ such that $\Psi^*(v) = \Psi^Y(v)$ whenever $v \in C^* \cap Y^\Psi$. Define $\Psi' : V(G) \setminus X \rightarrow \Sigma$ as $\Psi'(v) = \Psi^*(v)$ if $v \in C^*$ and $\Psi'(v) = \Psi(v)$ otherwise. Then (X, Ψ') is a solution to (X_0, Y, Y^Ψ, Ψ^Y) as well.

Proof. For the first part of the lemma, assume there exists $v \in C \cap \text{big}(X)$. From the connectivity of $G[C]$, we have that either $C \subseteq \text{big}(X)$ or there exists $uw \in E(G[C])$ with $w \in X$ and $u \in \text{big}(X)$. Recall that S interrogates X ; from the last property of the interrogating set we infer that w is adjacent to a big stain of S , and $w \in D^{\text{big}}$, a contradiction with the definition of C . Therefore either $C \subseteq \text{big}(X)$ or $C \cap \text{big}(X) = \emptyset$.

In the latter case, pick any $v \in C \setminus X$. As $C \cap \text{big}(X) = \emptyset$, v is contained in a connected component $C(v)$ of $G \setminus X$ with at most q vertices. By the second property of the interrogating set S , $C(v)$ is a small stain of S and we have $C(v) \subseteq C$. This finishes the proof of the first part of the lemma.

For the second part of the lemma, note that Ψ' satisfies all vertex constraints of J_b , as Ψ^* and Ψ are both labelings of $G[C^*]$ and $G \setminus X$, respectively. Moreover, $\Psi'|_{Y^\Psi} = \Psi^Y$, by the definition of Ψ^* and the fact that (X, Ψ) is a solution to (X_0, Y, Y^Ψ, Ψ^Y) . The deletion set X remains in the pair (X, Ψ') ; to finish the proof of the lemma, we need to show that Ψ' satisfies all edge constraints of $G \setminus X$.

As Ψ^* and Ψ are both labelings of $G[C^*]$ and $G \setminus X$, respectively, this claim is not obvious only for edges $uv \in E(G \setminus X)$ where $u \in C^*$ but $v \notin C^*$. As $X_0 \subseteq X$ and C^* contains $N_G[C] \setminus X_0$, we infer that $u \notin C$. Thus either $u \in S^{\text{big}}$ or $u \in N_G(C)$. In the first case, $u \in Y^\Psi$, $\Psi(u) = \Psi^*(u)$, and $(\Psi'(u), \Psi'(v)) \in \psi_{uv,u}$ since Ψ is a labeling of $G \setminus X$. Assume then $u \in N_G(C) \setminus S^{\text{big}}$.

As $u \notin C$, but u is adjacent to C , we infer that $u \in D^{\text{big}}$ and there exists $w \in N_G(u) \cap S^{\text{big}}$. As $S^{\text{big}} \subseteq Y^\Psi$, we have $\Psi(w) = \Psi^*(w) = \Psi^Y(w)$ and, since $\psi_{uw,u}$ is a partial permutation satisfied by both Ψ and Ψ^* , $\Psi(u) = \Psi^*(u)$. Thus $\Psi'(u) = \Psi(u)$ and $\Psi'(v) = \Psi(v)$ (as $v \notin C^*$), and the lemma is proven. \square

We use Lemma D.18 in the following rule.

Rule D.3 (Small Stains Rule). Assume there exists a connected component of $G \setminus (N_G[S^{\text{big}}] \cup X_0)$ that contains a vertex of $Y \setminus Y^\Psi$. Let C be its vertex set. Perform the following operations.

1. Let $C^* = S^{\text{big}} \cup (N_G[C] \setminus X_0)$. Using Lemma D.2, find if there exists a labeling Ψ^* of $G[C^*]$ that equals Ψ^Y on $Y^\Psi \cap C^*$. If this is the case, add $C \cap (Y \setminus Y^\Psi)$ to Y^Ψ and add $\Psi^*|_{C \cap (Y \setminus Y^\Psi)}$ to Ψ^Y .

2. Otherwise, first generate the following branch that corresponds to the second option of Lemma D.18, i.e., $C \cap \mathbf{big}(X) = \emptyset$. In the branch add $C \setminus S$ to X_0 and, for each stain C' of $G[C \cap S]$ that is not contained in Y^Ψ , find a labeling of C' using Lemma D.2. If no such labeling is found, terminate the current branch, and otherwise add C' to Y^Ψ and update Ψ^Y with any of the labelings found for C' .
3. If $N_G(C) \subseteq X_0$, finish branching with the single case from the previous point, as for any solution (X, Ψ) to (X_0, Y, Y^Ψ, Ψ^Y) , we have $C \cap \mathbf{big}(X) = \emptyset$.
4. Otherwise, generate additionally the following branches that correspond to the option $C \subseteq \mathbf{big}(X)$. Apply Lemma D.2 to find a family \mathbf{PSI} of at most s labelings of $G[C]$. Branch into $|\mathbf{PSI}|$ subcases, indexed by elements $\Psi^C \in \mathbf{PSI}$. In branch Ψ^C , we first verify if $\Psi^C(v) = \Psi^Y(v)$ whenever $v \in C \cap Y^\Psi$. If this is not the case, we abandon the branch Ψ^C . Otherwise, we add C to Y and Y^Ψ , and add Ψ^C to Ψ^Y .

First, note that the Small Stains Rule is applicable in $O(s^2n^2)$ time. The Small Stains Rule either leads to an increase in the size of X_0 or Y^Ψ , or reaches the last point and branches into at most $s + 1$ subcases. In the first subcase — output in the second point — we add new vertices to X_0 , as $N_G(C) \not\subseteq X_0$. In the remaining s cases we only increase Y^Ψ . However, if the Neighbourhood Branching Rule is not applicable before the Small Stains Rule is applied, then in each of these s subcases the Neighbourhood Branching Rule becomes applicable: there is no valid labeling of $G[C^*]$ (as Small Stains Rule did not finish at the first point), $C^* \subseteq S^{\mathbf{big}} \cup (N_G[C] \setminus X_0)$ and C is added to Y^Ψ .

Finally, note that if the Small Stains Rule is not applicable, then $Y = Y^\Psi$, as $S^{\mathbf{big}} \subseteq Y^\Psi$. In this case, by Lemma D.14 and since $S \subseteq Y$ and all forsaken vertices are in X_0 , we have that $N_G[Y^\Psi] \cup X_0 = V(G)$. Therefore either the Neighbourhood Branching Rule is applicable, or $G \setminus X_0$ admits a labeling that equals Ψ^Y on Y^Ψ and the Finishing Rule is applicable.

We infer that always at least one rule is applicable and the search tree has at most $(2s + 1)^k$ leaves. As each rule is applicable in $O(s^2n^2)$ time and the number of rule applications before reaching a leaf is bounded by n , the whole branching algorithm (described by these three rules and Step D.7) works in $O(2^{O(k \log s)} n^3)$ time. This finishes the description of the fixed-parameter algorithm for BORDER N-ULC.

E Full details of the algorithm for STEINER CUT

This section is devoted to fixed-parameter tractability of the STEINER CUT problem, parameterized by the size of the cutset.

<p>STEINER CUT</p> <p>Input: A graph G, a set of terminals $T \subseteq V(G)$, and integers s and k.</p> <p>Question: Does there exist a set X of at most k edges of G, such that in $G \setminus X$ at least s connected components contain at least one terminal?</p>	<p>Parameter: k</p>
--	---

For a STEINER CUT instance (G, T, s, k) , a set $X \subseteq E(G)$ is called a *solution* if $|X| \leq k$ and $G \setminus X$ contains at least s connected components that contain at least one terminal.

First, observe that by Lemma B.1 in $O(kn^2)$ time we can ensure that the graph G has $O(kn)$ edges, by removing the edges outside of the set E_0 . Correctness of this step follows from the fact that in this operation all cuts of size at most k are preserved and moreover no new cut of size at most k appears, since for each of the removed edges at least $k + 1$ edge disjoint paths between its endpoints remain. We are going to use the assumption that there are $O(kn)$ edges in the graph during the course of our algorithm. To this end, we use Lemma B.1 after each reduction, thus always ensuring that the graph has at most $O(kn)$ edges, where n is the current number

of vertices. We note that the only reason of using Lemma B.1 is caring about the polynomial factor.

Second, we observe that in the STEINER CUT problem we may assume that the graph G is connected. Indeed, otherwise we may add to G a clique on $k + 2$ vertices (so that the clique cannot be split with removal of k edges), make all vertices of the clique adjacent to exactly one vertex of each connected component of G , and decrease s by the number of connected components of G containing a terminal minus one.

If G is connected, removal of k edges may lead to at most $k + 1$ connected components. Thus, we may assume that $s \leq k + 1$, as otherwise the answer is trivially negative.

Moreover, in the course of the algorithm we repetitively contract some edges of G . In the process of contraction, we remove loops, but we keep multiple edges. Thus we allow G to be a multigraph with multiple edges, but without loops. Note that, if an edge $uv \in E(G)$ has multiplicity more than k , u and v cannot be separated by an edge cut of size k . Thus, for such an edge uv , we may reduce the multiplicity of uv to $k + 1$.

The algorithm performs a number of steps. Description of each step is accompanied by discussion of correctness and analysis of running time.

E.1 Operations on the input graph

The basic operation the algorithm performs on the graph is an edge contraction. As mentioned in the last section, we assume that after performing a serie of contractions we reduce all multiplicities of the multiedges that exceed $k + 1$ down to $k + 1$. Moreover, if in G a set of terminals $T \subseteq V(G)$ is given, if $T \cap \{u, v\} \neq \emptyset$, we replace T with $(T \setminus \{u, v\}) \cup \{w_{uv}\}$, i.e., we put w_{uv} into T if and only if u or v belongs to T .

The following straightforward corollary of Lemma A.2 shows when we may contract an edge of G in the STEINER CUT case.

Lemma E.1. *Let $\mathcal{J} = (G, T, s, k)$ be a STEINER CUT instance, let $D \subseteq E(G)$ and let $\mathcal{J}' = (G', T', s, k)$ be the instance \mathcal{J} with the edges D contracted in an arbitrary order. Then:*

1. *Any solution X to \mathcal{J}' is a solution to \mathcal{J} as well (recall that we treat $E(G')$ as a subset of $E(G)$).*
2. *For any solution X to \mathcal{J} that is disjoint with D , the set*

$$X' = \{\iota(u)\iota(v) : uv \in X, \iota(u) \neq \iota(v)\} \subseteq E(G')$$

is a solution to \mathcal{J}' .

We also use the notion of *identifying* two vertices.

Definition E.2. Given a multigraph G and two vertices u, v , *identification* of u and v is the operation of adding an edge uv and contracting it.

As identification is modelled by edge addition and contraction, we apply the same terminology also to this notion.

E.2 Borders and recursive understanding

In the border problem the graph is additionally equipped with at most $2k$ border terminals T_b . For a border STEINER CUT instance $\mathcal{J}_b = (G, T, k, T_b)$, we need to remember an equivalence relation \mathcal{R}_b on T_b , that corresponds to how the border terminals are to be distributed among connected components, a set $Y_b \subseteq T_b$, that carries information which border terminal is in

connected component with some terminal of T , and an integer s_b , which means that in \mathcal{J}_b we are to obtain s_b connected components that contain a terminal. Formally speaking, we define $\mathbb{P}(\mathcal{J}_b)$ as the set of all triples $\mathcal{P} = (\mathcal{R}_b, Y_b, s_b)$, where \mathcal{R}_b is an equivalence relation on T_b , $Y_b \subseteq T_b$ and $0 \leq s_b \leq k + 1$ is an integer. Moreover, we require that if $(u, v) \in \mathcal{R}_b$, then $u \in Y_b$ if and only if $v \in Y_b$.

We say that a set $X \subseteq E(G)$ is a *solution* to $(\mathcal{J}_b, \mathcal{P})$ for a triple $\mathcal{P} = (\mathcal{R}_b, Y_b, s_b) \in \mathbb{P}(\mathcal{J}_b)$ if

- two border terminals $u, v \in T_b$ are in the same connected component of $G \setminus X$ if and only if $(u, v) \in \mathcal{R}_b$;
- for any border terminal $u \in T_b$, the connected component of $G \setminus X$ which contains u , contains a vertex of T if and only if $u \in Y_b$;
- $G \setminus X$ contains exactly s_b connected components that contain a vertex of T ;
- $|X| \leq k$.

We formally define the border problem as follows.

BORDER STEINER CUT

Input: A STEINER CUT instance $\mathcal{J} = (G, T, s, k)$ with G being connected, and a set $T_b \subseteq V(G)$ of size at most $2k$; denote $\mathcal{J}_b = (G, T, k, T_b)$.

Output: For each $\mathcal{P} \in \mathbb{P}(\mathcal{J}_b)$ output a solution $\text{sol}_{\mathcal{P}} = X_{\mathcal{P}}$ to $(\mathcal{J}_b, \mathcal{P})$ with minimum possible $|X_{\mathcal{P}}|$, or $\text{sol}_{\mathcal{P}} = \perp$ if such a solution does not exist.

Observe that BORDER STEINER CUT generalizes STEINER CUT as we may ask for $T_b = \emptyset$ and check the value of a solution consistent with $(\emptyset, \emptyset, s)$, as we can assume that after removing the minimum size solution there are exactly s connected components containing a terminal.

Note that $|\mathbb{P}(\mathcal{J}_b)| \leq (2k)^{2k} \cdot 2^{2k} \cdot (k+2)$, as there are at most $|T_b|^{|T_b|}$ choices for an equivalence relation \mathcal{R}_b , $2^{|T_b|}$ choices for Y_b and $k+2$ choices for the value of s_b . Denote

$$q = k(2k)^{2k} 2^{2k} (k+2) + 1 = 2^{O(k \log k)}.$$

Let $\mathcal{J}_b = (G, T, k, T_b)$ be the given instance of BORDER STEINER CUT. Assume that G admits a (q, k) -good edge separation (V_1, V_2) .

As V_1 and V_2 are disjoint, at least one of them contains at most k border terminals from T_b . Without loss of generality assume that $|T_b \cap V_1| \leq k$. Let $G^* = G[V_1]$, $T^* = T \cap V_1$ and $T_b^* = (T_b \cap V_1) \cup (V_1 \cap V(\delta(V_1, V_2)))$. Consider an instance $\mathcal{J}_b^* = (G^*, T^*, k, T_b^*)$. Note that \mathcal{J}_b^* is a correct instance of BORDER STEINER CUT, as $|(T_b \cap V_1) \cup (V_1 \cap V(\delta(V_1, V_2)))| \leq 2k$. Apply the algorithm recursively to the instance \mathcal{J}_b^* (note that it is strictly smaller instance as the vertex set V_2 is removed) and let $U(\mathcal{J}_b^*)$ be the set of edges that are contained in any output solution for any behaviour on the border terminals of \mathcal{J}_b^* . Observe that $|U(\mathcal{J}_b^*)| \leq q - 1$. Let $R = E(G^*) \setminus U(\mathcal{J}_b^*)$ be the set of remaining edges in $G^* = G[V_1]$. Contract the edges of R in G to obtain the new graph G' with terminals T' and border terminals T_b' . Let V_1' be the set of vertices of G' onto which vertices of V_1 were contracted. Observe that $G'[V_1']$ is still connected as a contraction of a connected graph, and has at most $q - 1$ edges, as $|U(\mathcal{J}_b^*)| \leq q - 1$. Therefore, $|V_1'| \leq q$. The contraction induces a mapping $\iota : V(G) \rightarrow V(G')$ that maps every vertex of G to the vertex of G' onto which it is contracted.

The following lemma is useful in arguing safeness of the described operation.

Lemma E.3. *Let $\mathcal{P} \in \mathbb{P}(\mathcal{J}_b)$ and let $X_{\mathcal{P}}$ be a solution to $(\mathcal{J}_b, \mathcal{P})$. Then there exists a second solution $X'_{\mathcal{P}}$ to $(\mathcal{J}_b, \mathcal{P}')$, such that $|X'_{\mathcal{P}}| \leq |X_{\mathcal{P}}|$ and additionally $X'_{\mathcal{P}} \cap R = \emptyset$.*

Proof. Consider the graph $G^* = G[V_1]$ and the set $X_{\mathcal{P}} \cap E(G^*)$. Define:

- \mathcal{R}_b^* to be an equivalence relation on T_b^* such that for any $u, v \in T_b^*$ we have $(u, v) \in \mathcal{R}_b^*$ if and only if u and v are in the same connected component of $G^* \setminus X_{\mathcal{P}}$.
- Y_b^* to be a set of those vertices $v \in T_b^*$, such that the connected component of $G^* \setminus X_{\mathcal{P}}$ that contains v contains a terminal from $T^* = T \cap V_1$ as well.
- s_b^* to be the number of connected components of $G^* \setminus X_{\mathcal{P}}$ that contain a vertex of T^* .

Let $\mathcal{P}^* = (\mathcal{R}_b^*, Y_b^*, s_b^*)$. Clearly, $\mathcal{P}^* \in \mathbb{P}(\mathcal{J}_b^*)$ and $X_{\mathcal{P}} \cap E(G^*)$ is a solution to $(\mathcal{J}_b^*, \mathcal{P}^*)$ (note that $s_b^* \leq |X_{\mathcal{P}} \cap E(G^*)| + 1 \leq k + 2$, as G^* is connected). Therefore $\text{sol}_{\mathcal{P}^*} = X_{\mathcal{P}^*}^* \neq \perp$, that is, there exists a solution $X_{\mathcal{P}^*}^*$ to $(\mathcal{J}_b^*, \mathcal{P}^*)$, such that $|X_{\mathcal{P}^*}^*| \leq |X_{\mathcal{P}} \cap E(G^*)|$ and $X_{\mathcal{P}^*}^* \cap R = \emptyset$.

Define $X'_{\mathcal{P}} = (X_{\mathcal{P}} \setminus E(G^*)) \cup X_{\mathcal{P}^*}^*$. Clearly $|X'_{\mathcal{P}}| \leq |X_{\mathcal{P}}|$. To finish the proof of the lemma we need to show that $X'_{\mathcal{P}}$ is a solution to $(\mathcal{J}_b, \mathcal{P})$.

First, we show the following claim: for any $u, v \in T_b \cup T_b^*$, u and v are in the same connected component of $G \setminus X_{\mathcal{P}}$ if and only if u and v are in the same connected component of $G \setminus X'_{\mathcal{P}}$. We show only a proof in one direction, as proofs in both directions are totally symmetric and use only the facts that $X_{\mathcal{P}} \setminus E(G^*) = X'_{\mathcal{P}} \setminus E(G^*)$ and that both $X_{\mathcal{P}} \cap E(G^*)$ and $X'_{\mathcal{P}} \cap E(G^*)$ are solutions to $(\mathcal{J}_b^*, \mathcal{P}^*)$.

Let $u, v \in T_b \cup T_b^*$ be two vertices that are connected by a path P in $G \setminus X_{\mathcal{P}}$. Let $u = v_0, v_1, \dots, v_r = v$ be the sequence of all vertices on P that belong to $T_b \cup T_b^*$, in the order they appear on P . To prove the claim we need to show that for any $0 \leq i < r$, the vertices v_i and v_{i+1} belong to the same connected component of $G \setminus X'_{\mathcal{P}}$. By definition, as $V_1 \cap V(\delta(V_1, V_2)) \subseteq T_b^*$, the subpath P_i of P between v_i and v_{i+1} lies entirely in G^* or entirely in $G \setminus E(G^*)$. In the first case, we infer that $v_i, v_{i+1} \in T_b^*$, $(v_i, v_{i+1}) \in \mathcal{R}_b^*$ and v_i and v_{i+1} are in the same connected component of $G^* \setminus X'_{\mathcal{P}}$ as $X'_{\mathcal{P}} \cap E(G^*)$ is a solution to $(\mathcal{J}_b, \mathcal{P}^*)$. In the second case, we infer that v_i and v_{i+1} are in the same connected component of $(G \setminus E(G^*)) \setminus X'_{\mathcal{P}}$, as $X_{\mathcal{P}} \setminus E(G^*) = X'_{\mathcal{P}} \setminus E(G^*)$. This finishes the proof of the claim.

As a straightforward corollary of the aforementioned claim, we infer that for any $u, v \in T_b$, we have $(u, v) \in \mathcal{R}_b$ if and only if u and v are in the same connected component of $G \setminus X'_{\mathcal{P}}$. We now show that for any $v \in T_b \cup T_b^*$, its connected component of $G \setminus X_{\mathcal{P}}$ contains a vertex of T if and only if its connected component of $G \setminus X'_{\mathcal{P}}$ contains a vertex of T . The proofs in both directions are again totally symmetric, thus we present only the forward implication.

Let P be a path that connects the vertex v with a terminal $w \in T$ in $G \setminus X_{\mathcal{P}}$. Let u be the last (closest to w) vertex on P that belongs to $T_b \cup T_b^*$ (as $v \in T_b \cup T_b^*$, such a vertex u exists). From the claim we infer that u and v are in the same connected component of $G \setminus X'_{\mathcal{P}}$. Let P_u be the subpath of P from u to w . We have two cases: either P_u is contained in G^* , or in $G \setminus E(G^*)$. In the first case, we infer that $u \in T_b^*$, $u \in Y_b^*$ (as $X_{\mathcal{P}} \cap E(G^*)$ is a solution to $(\mathcal{J}_b^*, \mathcal{P}^*)$) and that the connected component of $G^* \setminus X'_{\mathcal{P}}$ that contains u contains a vertex from $T^* = T \cap V_1$ (not necessarily the vertex w). In the second case, we infer that the path P_u is present in $(G \setminus E(G^*)) \setminus X'_{\mathcal{P}}$. This finishes the proof that $v \in Y_b$ if and only if there exists a terminal in the connected component of $G \setminus X'_{\mathcal{P}}$ that contains v .

To finish the proof of the lemma we need to show that the number of connected components of $G \setminus X'_{\mathcal{P}}$ that contain a terminal equals s_b . To this end, we partition the connected components of $G \setminus X_{\mathcal{P}}$ and $G \setminus X'_{\mathcal{P}}$ containing terminals into three types:

1. those that contain a vertex from $T_b \cup T_b^*$;
2. those that do not contain such a vertex, but are contained in G^* ;
3. and the rest — those that do not contain a vertex from $T_b \cup T_b^*$, and are contained in $G \setminus V_1$.

Our goal is to prove that for each type, the numbers of connected components containing terminals of corresponding types in $G \setminus X_{\mathcal{P}}$ and $G \setminus X'_{\mathcal{P}}$ are equal.

For the first type, the claim is a straightforward corollary of already proven facts that (i) any two vertices $u, v \in T_b \cup T_b^*$ are in the same connected component of $G \setminus X_{\mathcal{P}}$ if and only if they are in the same connected component of $G \setminus X'_{\mathcal{P}}$, and (ii) for every vertex $u \in T_b \cup T_b^*$, the connected component of $G \setminus X_{\mathcal{P}}$ containing u contains a terminal if and only if the connected component of $G \setminus X'_{\mathcal{P}}$ containing u contains a terminal.

For the second type, note that we are to count the number of connected components of $G^* \setminus X_{\mathcal{P}}$ and $G^* \setminus X'_{\mathcal{P}}$ that do not contain a vertex of $T_b \cup T_b^*$, or, equivalently, T_b^* , but contain a terminal from T^* . As both $X_{\mathcal{P}} \cap E(G^*)$ and $X'_{\mathcal{P}} \cap E(G^*)$ are solutions to $(\mathcal{J}_b^*, \mathcal{P}^*)$, this number is equal to s_b^* minus the number of equivalence classes of \mathcal{R}_b^* that contain vertices from Y_b .

For the third type, recall that $X_{\mathcal{P}} \setminus E(G^*) = X'_{\mathcal{P}} \setminus E(G^*)$, so the sets of connected components of the third type in $G \setminus X_{\mathcal{P}}$ and $G \setminus X'_{\mathcal{P}}$ are equal. This finishes the proof of the lemma. \square

We now show that the output for the new instance $\mathcal{J}'_b = (G', T', k, T'_b)$ can be easily transformed to the output for the original instance \mathcal{J}_b .

Lemma E.4. *Let $(\text{sol}'_{\mathcal{P}})_{\mathcal{P} \in \mathbb{P}(\mathcal{J}'_b)}$ be a correct output for the instance \mathcal{J}'_b . For any $\mathcal{P} = (\mathcal{R}_b, Y_b, s_b) \in \mathbb{P}(\mathcal{J}_b)$ define $\text{sol}_{\mathcal{P}}$ as follows.*

- If ι maps two border terminals $u, v \in T_b$ with $(u, v) \notin \mathcal{R}_b$ to the same vertex of T'_b , then we take $\text{sol}_{\mathcal{P}} = \perp$.
- Otherwise, we define $\mathcal{P}' = (\mathcal{R}'_b, Y'_b, s_b)$ as follows: $(u', v') \in \mathcal{R}'_b$ if $\iota^{-1}(u') \cap T_b$ are contained in the same equivalence class as $\iota^{-1}(v') \cap T_b$, and $v' \in Y'_b$ if $\iota^{-1}(v') \cap T_b \subseteq Y_b$; and take $\text{sol}_{\mathcal{P}} = \text{sol}'_{\mathcal{P}'}$.

Then the sequence $(\text{sol}_{\mathcal{P}})_{\mathcal{P} \in \mathbb{P}(\mathcal{J}_b)}$ is a correct output to the instance \mathcal{J}_b .

Proof. The lemma is a straightforward corollary of the contraction properties of Lemmas A.2 and E.1, as well as Lemma E.3. \square

We can now formally define the first step of the algorithm.

Step E.1. Using Lemma 2.2 we check, whether G admits a (q, k) -good edge separation. If this is not the case, we proceed to the second phase, i.e., high connectivity phase. Otherwise let (V_1, V_2) be this separation and without loss of generality assume that $|T_b \cap V_1| \leq k$. Construct the instance $\mathcal{J}_b^* = (G[V_1], T \cap V_1, k, (T_b \cap V_1) \cup (V_1 \cap V(\delta(V_1, V_2))))$, apply Lemma B.1 to it, solve it recursively and compute $U(\mathcal{J}_b^*)$, the set of edges that appear in any solution given in the output. Contract all the remaining edges of $G[V_1]$ in G to obtain new graph G' with terminals T' and border terminals T'_b . Define $\mathcal{J}'_b = (G', T', k, T'_b)$; recall that a vertex belongs to (border) terminals if and only if some (border) terminal was contracted onto it. Apply Lemma B.1 to \mathcal{J}'_b , recursively solve the instance \mathcal{J}'_b and transform the output according to Lemma E.4.

Let us now estimate the running time. First, we spend $O(2^{O(k^2 \log k)} n^3 \log n)$ time to check, whether there exists a (q, k) -good edge separation. We apply the algorithm recursively to the instance \mathcal{J}_b^* , which has n' vertices for some $q+1 \leq n' \leq n-q-1$. Construction of the instance \mathcal{J}_b^* takes $O(kn)$ time, construction of $U(\mathcal{J}_b^*)$ takes $O(2^{O(k \log k)} n)$ time, and construction of the instance \mathcal{J}'_b takes $O(kn)$ time. Then, we apply the algorithm recursively to the instance \mathcal{J}'_b that has at most $n - n' + q$ vertices. Therefore, we can derive the following recurrent inequality:

$$T(n) \leq \max_{q+1 \leq n' \leq n-q-1} \left(O(2^{O(k^2 \log k)} n^3 \log n) + T(n') + T(n - n' + q) \right), \quad (3)$$

Note that the function $p(t) = t^3 \log t$ is convex, so the maximum of the expression is attained at one of the ends. A straightforward inductive check of both of the ends proves that we have indeed the claimed bound on the complexity, i.e., $O(2^{O(k^2 \log k)} n^4 \log n)$.

Observe, that if we use the randomized algorithm for finding good edge separations from Lemma B.2, we obtain $T(v) \leq \tilde{O}(2^{O(k^2 \log k)} n^2)$ time complexity with success probability at least $(1 - 1/n)$, since the graph is partitioned using good edge separations less than n times.

E.3 Brute force approach

If the graph returned by Step E.1 turns out to be small, we apply a brute-force approach. In this section we describe this step formally.

Lemma E.5. *A correct output to a BORDER STEINER CUT instance $\mathcal{J}_b = (G, T, k, T_b)$ can be computed in $O(2^{O(k \log k)} n^{2k+2})$ time.*

Proof. For every $\mathcal{P} \in \mathbb{P}(\mathcal{J}_b)$, and for every set $X \subseteq E(G)$ of at most k edges that, for all $u, v \in V(G)$, takes either all or zero edges uv , in $O(n^2)$ time we verify whether X is a solution to $(\mathcal{J}_b, \mathcal{P})$. The time bound follows from the fact that $|\mathbb{P}(\mathcal{J}_b)| \leq 2^{O(k \log k)}$ and there are at most $(k+1)n^{2k}$ choices of the set X . \square

We are ready to provide the step of the algorithm that finishes resolving the problem, providing that the graph is sufficiently small.

Step E.2. If $|V(G)| \leq (k+1)q$, then apply Lemma E.5 to resolve the given BORDER STEINER CUT instance $\mathcal{J}_b = (G, T, k, T_b)$.

The correctness of this step is obvious, while from Lemma E.5 we find that the running time is $O(2^{O(k^2 \log k)})$ as $q = 2^{O(k \log k)}$. Therefore, from now on we can assume that $|V(G)| > (k+1)q$.

E.4 High connectivity phase

We now show how to solve BORDER STEINER CUT in $\tilde{O}(2^{O(k^2 \log k)} n)$ time for the remaining case, when the graph G does not admit a (q, k) -good edge separation, yet is still too big to apply brute-force. We need to output answers for all the possible triples $\mathcal{P} \in \mathbb{P}(\mathcal{J}_b)$. We iterate through all such \mathcal{P} ; note that this gives only $2^{O(k \log k)}$ overhead in the running time. Therefore, from now on we may assume that $\mathcal{P} = (\mathcal{R}_b, Y_b, s_b)$ is fixed.

Firstly, we make a quick check whether an empty deletion set is sufficient for our needs. We formally need this step in order to be able to use nontriviality of the solution in some technical reasonings.

Step E.3. Given BORDER STEINER CUT instance $\mathcal{J}_b = (G, T, k, T_b)$ and $\mathcal{P} \in \mathbb{P}(\mathcal{J}_b)$, verify in $O(kn)$ time whether \emptyset is a solution to $(\mathcal{J}_b, \mathcal{P})$. If this is the case, output $\text{sol}_{\mathcal{P}} = \emptyset$.

The described step requires $O(kn)$ time and its correctness is obvious. From now on we may assume that the minimum deletion set is nonempty.

E.4.1 Interrogating sets

We now prepare ourselves to use Lemma 1.1 to extract more structure of the graph G .

Definition E.6. Let $X \subseteq E(G)$, $1 \leq |X| \leq k$, and let C_0, C_1, \dots, C_ℓ be connected components of $G \setminus X$, where, due to Lemma 2.3, $\ell \leq k$ and $|V(C_i)| \leq q$ for $i \geq 1$. We say that a set of edges $S \subseteq E(G)$ *interrogates* X if the following properties are satisfied:

- $X \cap S = \emptyset$;

- for every component C_i , $i \geq 1$, S contains a spanning tree of C_i ;
- for every vertex $u \in V(C_0) \cap V(X)$, u is contained in a connected component of $(V(G), S)$ of size at least $q + 1$.

Note that the first property together with $|V(C_i)| \leq q$ for $i \geq 1$ imply that the connected component considered in the third property has to be entirely contained in C_0 . We now prove that a sufficiently large family given by Lemma 1.1 contains a set interrogating a solution.

Lemma E.7. *Let \mathcal{F} be a family obtained by an application of the algorithm of Lemma 1.1 for universe $U = E(G)$ and constants $a = 3qk$ and $b = k$, Then, for any $X \subseteq E(G)$ with $1 \leq |X| \leq k$, there exists a set $S \in \mathcal{F}$ that interrogates X .*

Proof. Let C_0, C_1, \dots, C_ℓ be connected components of $G \setminus X$, where, due to Lemma 2.3, $\ell \leq k$ and $|V(C_i)| \leq q$ for $i \geq 1$. As the algorithm did not finish when performing Step E.2, we have that $|V(G)| > (k + 1)q$, so $|V(C_0)| \geq q + 1$. Fix a spanning tree T_i of each component C_i . Let $A_1 = \bigcup_{i=1}^{\ell} E(T_i)$, note that $|A_1| \leq qk$. For every vertex $u \in V(C_0) \cap V(X)$ fix an arbitrary subtree T_0^u of T_0 that contains exactly $q + 1$ vertices, and define $A_2 = \bigcup_{u \in V(C_0) \cap V(X)} E(T_0^u)$; this is possible due to $|V(C_0)| \geq q + 1$. By Lemma 2.3, there exists a set $S \in \mathcal{F}$ such that $A_1 \cup A_2 \subseteq S$ and $S \cap X = \emptyset$. It follows from the construction that S interrogates X . \square

This gives rise to the following branching step.

Step E.4. Using Lemma 1.1 generate family \mathcal{F} for universe $U = E(G)$, and constants $a = 3qk$ and $b = k$. Branch into $|\mathcal{F}|$ subcases, labeled with $S \in \mathcal{F}$. In branch S we seek a solution X to $(\mathcal{J}_b, \mathcal{P})$ such that S interrogates X and, moreover, $|X|$ is minimum among these.

Lemma E.7 asserts that the deletion set of an optimal solution is interrogated by some set from family \mathcal{F} . Therefore, in order to find a solution with minimum possible size of deletion set it suffices to take minimum over solutions given by the branches. Note that in this manner we introduce $O(2^{O(\min(q,k) \log(q+k))} \log n) = O(2^{O(k^2 \log k)} \log n)$ branches. Moreover, as the family \mathcal{F} can be computed in $O(2^{O(k^2 \log k)} n \log n)$ time and the construction of every branch takes quadratic time, the whole branching procedure takes $O(2^{O(k^2 \log k)} n \log n)$ time. We now describe the subroutine performed in each branch, let it be labeled by $S \in \mathcal{F}$. To simplify the presentation we assume that S interrogates X_0 for some minimum solution X_0 and examine what happens with X_0 during the operations performed on the graph.

Let us contract all the edges from S to obtain a new graph H_0 . Let ι_0 be the mapping from $V(G)$ to $V(H_0)$ corresponding to these contractions. Then, we obtain the new graph H by identifying all the vertices $u \in H_0$ for which $|\iota_0^{-1}(u)| > q$ into a single vertex; such vertices u are called *heavy*. If there are no heavy vertices, we can safely terminate the branch, as a set that interrogates a nonempty solution must induce at least one connected component that has at least $q + 1$ vertices. Otherwise, denote b the vertex resulting in their identification; we will further refer to it as to the *core* vertex. Let ι_1 be the mapping from $V(H_0)$ to $V(H)$ corresponding to these identifications. Moreover, let $\iota = \iota_1 \circ \iota_0$ be the mapping from $V(G)$ to $V(H)$ corresponding to the composition of these operations.

We claim that the feasible deletion set X_0 'survives' both steps.

Lemma E.8. *Let $v, w \in V(G)$ such that $\iota(v) = \iota(w)$. Then v and w are in the same connected component of $G \setminus X$ for any set $X \subseteq E(G)$ interrogated by S .*

Proof. Assume otherwise, that is, that we have $\iota(v) = \iota(w)$ but $v \in V(C_i)$ and $w \in V(C_j)$ for $i \neq j$. Without loss of generality assume that $i \neq 0$.

Assume first that $\iota_0(v) = \iota_0(w)$. As v and w are contracted onto the same vertex, there exists a path from v to w in G that consists of edges of S . As $S \cap X = \emptyset$, this means that v and w are in the same connected component of $G \setminus X$, which is a contradiction.

Assume now that $\iota_0(v) \neq \iota_0(w)$. This means that $\iota_0(v)$ and $\iota_0(w)$ have to be identified while constructing graph H from H_0 . It follows that $|\iota_0^{-1}(v)| \geq q + 1$. Therefore, there are at least q vertices of G that are reachable from v via paths contained in S , hence disjoint with X . However, $i \neq 0$ so $|V(C_i)| \leq q$, which is a contradiction. \square

From Lemma E.8 we infer that all the edges of X_0 are still present in H , as, from the minimality of X_0 we may assume that the edges of X_0 connect different connected components of $G \setminus X_0$. Let us define the sets of terminals T' and border terminals T'_b in H by setting $u \in T'$ if and only if $\iota^{-1}(u) \cap T \neq \emptyset$ and $u \in T'_b$ if and only if $\iota^{-1}(u) \cap T_b \neq \emptyset$.

Moreover, due to Lemma E.8 and the fact that X is a solution to $(\mathcal{J}_b, \mathcal{P})$, we infer that for any $v, w \in T_b$ with $\iota(v) = \iota(w)$, we have $(v, w) \in \mathcal{R}_b$ and $v \in Y_b$ iff $w \in Y_b$. Thus we can project \mathcal{R}_b and Y_b on $T'_b \subseteq V(H)$, by defining a relation \mathcal{R}'_b by taking $(\iota(v), \iota(w)) \in \mathcal{R}'_b$ iff $(v, w) \in \mathcal{R}_b$ and a set Y'_b by taking $\iota(v) \in Y'_b$ iff $v \in Y_b$.

Define $\mathcal{J}'_b = (H, T', k, T'_b)$ and $\mathcal{P}' = (\mathcal{R}'_b, Y'_b, s_b)$; note that $\mathcal{P}' \in \mathbb{P}(\mathcal{J}'_b)$. The next lemma can be proven by a straightforward check of the definition of the solution, as Lemma E.8 asserts connected components of $G \setminus X_0$ correspond in a one-to-one manner to connected components of $H \setminus X_0$.

Lemma E.9. *The set X_0 is a solution to $(\mathcal{J}'_b, \mathcal{P}')$.*

In the same manner we can also obtain a converse implication.

Lemma E.10. *If a set $X' \subseteq E(H)$ is a solution to $(\mathcal{J}'_b, \mathcal{P}')$ of minimum possible size, then X' is a solution to $(\mathcal{J}_b, \mathcal{P})$ as well.*

Lemmas E.9 and E.10 justify correctness of the following step, which we now state formally.

Step E.5. Contract the edges of S to obtain the graph H_0 . If there are no heavy vertices in H_0 , terminate the branch as S cannot interrogate any feasible deletion set. Otherwise, identify all heavy vertices into one core vertex b and denote by H the resulting graph. Define the instance $\mathcal{J}'_b = (H, T', k, T'_b)$ and \mathcal{P}' in a natural manner, described in this section. Run the remaining part of the algorithm on the instance \mathcal{J}'_b and triple \mathcal{P}' to obtain a solution sol , which then output as $\text{sol}_{\mathcal{P}}$.

Note that Step E.5 can be performed in $O(kn)$ time.

E.4.2 Connected components of $H \setminus \{b\}$ and dynamic programming

We now establish some structural properties of the behaviour of X_0 in H . The goal is to limit the class of possible solutions in \mathcal{J}'_b we need to search through. Note that in the constructed graph H the vertex b plays a special role, as we know that $V(X_0) \cap V(C_0) \subseteq \iota^{-1}(b)$.

Let B'_1, B'_2, \dots, B'_p be the components of $H \setminus \{b\}$ and let $B_i = H[V(B'_i) \cup \{b\}]$ for $i = 1, 2, \dots, p$. Observe that B'_i are connected, edge-disjoint and b separates them. Moreover, we can compute them in $O(kn)$ time. We now claim that for each component B_i , the solution either takes $E(B_i)$ entirely, or is disjoint with it.

Lemma E.11. *For each $i = 1, 2, \dots, p$, either $E(B_i) \cap X_0 = \emptyset$ or $E(B_i) \subseteq X_0$.*

Proof. We consider two cases. Assume first that no vertex in $V(B'_i)$ is in the same connected component of $H \setminus X_0$ as the core b . In particular, this implies that edges connecting b with $V(B'_i)$ belong to X_0 . Moreover, in G the set $\iota^{-1}(V(B'_i))$ is a union of vertex sets of some components

C_i for $i \geq 1$. As S interrogates X_0 , each component C_i for $i \geq 1$ is projected by ι onto a single vertex in H . Consider an edge $e \in E(B'_i)$. We infer that e connects two different connected components of $G \setminus X_0$, thus $e \in X_0$. Therefore $E(B_i) \subseteq X_0$ in this case.

Assume now that there exists a vertex $v \in V(B'_i)$ that is in the same connected component of $H \setminus X_0$ as b , i.e., $\iota^{-1}(v) \subseteq V(C_0)$. Let $w \in N_{B'_i}(v)$. As $v \neq b$, we have $|\iota^{-1}(v)| \leq q$ and $|\iota^{-1}(w)| \leq q$, so we have $vw \notin X_0$, as otherwise S does not interrogate X_0 . Since the choice of v and its neighbour w was arbitrary, and since B'_i is connected, we infer that all vertices of B'_i belong to the same connected component of $H \setminus X_0$. As $\iota^{-1}(v) \subseteq V(C_0)$, we find that $\iota^{-1}(V(B'_i)) \subseteq V(C_0)$. From the minimality of X_0 we infer that $E(B_i) \cap X_0 = \emptyset$. \square

Lemma E.11 ensures us that we may seek the optimal solution among the ones that do not intersect edge sets of components B_i nontrivially. We now show how to construct the optimal solution in the remaining instance in $O(k^2n)$ time. First, we resolve components B'_i that contain border terminals.

Step E.6. We define $D \subseteq T'_b$ as follows. If $b \in T'_b$, we define D to be the equivalence class of \mathcal{R}'_b that contains b . Otherwise, we branch into at most $1 + |T'_b| \leq 1 + 2k$ subcases, taking D to be an empty set or one of the equivalence classes of \mathcal{R}'_b . Given D , we seek for a solution X where the set of border terminals being in the in the same connected component of $H \setminus X$ as b equals \mathcal{R}'_b .

For a fixed choice of D , we may immediately resolve the connected components B'_i that contain a border terminal of T'_b . Initiate a counter $s_0 = 0$. For each component B'_i with $V(B'_i) \cap T'_b \neq \emptyset$ perform the following.

1. If there exists a border terminal in $V(B'_i) \cap D$, as well as a border terminal in $(V(B'_i) \cap T'_b) \setminus D$, terminate this branch, as for any of the two cases given by Lemma E.11, we cannot satisfy the conditions implied by \mathcal{R}'_b and the set D .
2. If all border terminals in $V(B'_i) \cap T'_b$ belong to D , contract all edges of B_i (we have $E(B_i) \cap X = \emptyset$ in this case).
3. If all border terminals in $V(B'_i) \cap T'_b$ do not belong to D , include $E(B_i)$ into the constructed solution: decrease k by $|E(B_i)|$ and increase s_0 by $|V(B'_i) \cap T'|$ (by including $E(B_i)$ into a solution, we delete $|E(B_i)|$ edges and create $|V(B'_i) \cap T'|$ new connected components that contain a terminal).

Note that Step E.6 can be performed in $O(k^2n)$ time and results in at most $2k + 1$ branches. Its correctness is asserted by Lemma E.11. After Step E.6 is applied, all terminals of T'_b are either contracted onto b (if they belong to D), or became isolated vertices after the removal of edges included to the constructed solution. If some equivalence class of \mathcal{R}'_b different than D is larger than a single vertex of H , or we do not satisfy the conditions implied by the set Y'_b for some vertex of T'_b , we may immediately reject the current branch. Otherwise, we may forget the relation \mathcal{R}'_b (as all conditions imposed by it are already satisfied). Moreover we may also forget almost all information carried by the set Y'_b , except for the fact whether $D \subseteq Y'_b$. This is done in the following step.

Step E.7. Terminate the current branch if one of the following conditions is satisfied:

1. There exists a equivalence class of \mathcal{R}'_b that is different than D and contains at least two border terminals.
2. There is a vertex $v \in T'_b \setminus D$, such that v is exactly in one of the sets T' and Y'_b .
3. $D \neq \emptyset$, $D \cap Y'_b = \emptyset$ and $b \in T'$ after Step E.6 is applied.

Otherwise, denote $\alpha = \perp$ if $D \neq \emptyset$ and $D \cap Y'_b = \emptyset$, and $\alpha = \top$ otherwise.

We note that, from the minimality of X and the connectivity of G , we have that any connected component of $G \setminus X$ that does not contain a border terminal, contains a terminal from T . Indeed, otherwise, if $G \setminus X$ contains a connected component C that does not contain a terminal nor a border terminal, one edge incident to C may be removed from X and still X would be a solution to $(\mathcal{J}_b, \mathcal{P})$, a contradiction. Therefore, if $D = \emptyset$, we may assume that the connected component of $G \setminus X$ that contains $\iota^{-1}(b)$, contains at least one terminal.

From now on we know that all the remaining components B'_i do not contain border terminals. Without loss of generality, let $B'_1, B'_2, \dots, B'_{p'}$ be the remaining components. For every remaining component B_i we have two numbers: $a_i = |E(B_i)|$, the cost of incorporating it to the solution, and $b_i = |V(B'_i) \cap T'|$, the number of separated terminals. Computation of a_i, b_i can be done in $O(kn)$ time. We would like to know what is the optimal number of edges needed to separate exactly $s_1 = s_b - s_0$ connected components with terminals, with the additional constraint that the connected component containing b contains a terminal if and only if $\alpha = \top$.

This can be solved in time $O(s_b p')$ via a standard dynamic programming routine. We create a 3-dimensional table $T[j, \ell, \mathfrak{t}]$ for $\ell = 0, 1, \dots, s_1$, $j = 0, 1, \dots, p'$, $\mathfrak{t} = \{\perp, \top\}$ with the following meaning: $T[j, \ell, \mathfrak{t}]$ is the minimum cost of a solution contained in the prefix B_1, B_2, \dots, B_j that separates exactly ℓ isolated vertices being terminals and \mathfrak{t} denotes whether the remaining connected component with b contains a terminal different than b (or $+\infty$ if such a solution does not exist). Formally,

$$T[j, \ell, \mathfrak{t}] = \min \left\{ \sum_{\gamma \in \Gamma} a_\gamma \mid \Gamma \subseteq \{1, 2, \dots, j\} \wedge \sum_{\gamma \in \Gamma} b_\gamma = \ell \wedge (\mathfrak{t} = \top \Leftrightarrow \sum_{\gamma \in \{1, 2, \dots, j\} \setminus \Gamma} b_\gamma > 0) \right\}.$$

Observe that T admits the following recurrential formula (by somewhat abusing notation, we assume that cells of T with negative coordinates contain $+\infty$):

$$T[j, \ell, \perp] = \begin{cases} +\infty & \text{if } j = 0 \text{ and } \ell > 0, \\ 0 & \text{if } j = \ell = 0, \\ \min(T[j-1, \ell, \perp], a_j + T[j-1, \ell - b_j, \perp]) & \text{if } j > 0 \text{ and } b_j = 0, \\ a_j + T[j-1, \ell - b_j, \perp] & \text{otherwise.} \end{cases}$$

$$T[j, \ell, \top] = \begin{cases} +\infty & \text{if } j = 0, \\ \min(T[j-1, \ell, \top], a_j + T[j-1, \ell - b_j, \top]) & \text{if } j > 0, \text{ and } b_j = 0, \\ \min(T[j-1, \ell, \perp], T[j-1, \ell, \top], a_j + T[j-1, \ell - b_j, \top]) & \text{otherwise.} \end{cases}$$

Hence, we can fill the table T in time $O(s_1 p') = O(kn)$; the optimal value can be deduced from the cells $T[p', s_1, \perp]$ and $T[p', s_1, \top]$. Although we presented here only the algorithm for computing the optimal value, it is straightforward to implement the dynamic program so that it also maintains backlinks via which one can retrieve the corresponding set Γ from the definition of T . Thus, we can formally present the final step of our algorithm.

Step E.8. Compute numbers a_i and b_i and fill table T in $O(kn)$ time. Let $\mathfrak{t} \in \{\perp, \top\}$ be defined as: $\mathfrak{t} = \perp$ if $\alpha = \perp$, $\mathfrak{t} = \top$ if $\alpha = \top$ and $b \notin T'$, and otherwise pick $\mathfrak{t} \in \{\perp, \top\}$ to minimize the value $T[p', s_1, \mathfrak{t}]$. Let $\Gamma \subseteq \{1, 2, \dots, p'\}$ be the set from the definition of the value $T[p', s_1, \mathfrak{t}]$, computed in $O(n)$ time by following backlinks in the table T . If the value $T[p', s_1, \mathfrak{t}]$ exceeds the remaining budget k , terminate the branch. Otherwise, incorporate the set $\bigcup_{\gamma \in \Gamma} E(B_\gamma)$ to the constructed solution.

Step E.8 can be performed in $O(kn)$ time and its correctness follows from the definition of the table T and the previous steps of the algorithm.

This finishes the description of the fixed-parameter algorithm for STEINER CUT.

F Full details of the algorithm for NODE MULTIWAY CUT-UNCUT

In this section we show an FPT algorithm for the following generalization of the well-known MULTIWAY CUT problem.

<p>NODE MULTIWAY CUT-UNCUT (N-MWCU)</p> <p>Input: A graph G together with a set of terminals $T \subseteq V(G)$, an equivalence relation \mathcal{R} on the set T, and an integer k.</p> <p>Question: Does there exist a set $X \subseteq V(G) \setminus T$ of at most k nonterminals such that for any $u, v \in T$, the vertices u and v belong to the same connected component of $G \setminus X$ if and only if $(u, v) \in \mathcal{R}$?</p>	<p>Parameter: k</p>
---	---

In other words, we are to delete at most k vertices from the graph, so that the terminals are split between connected components exactly as it is given by the equivalence relation \mathcal{R} . Given a N-MWCU instance $\mathcal{J} = (G, T, \mathcal{R}, k)$, a set of vertices X is called a *solution* to \mathcal{J} , if $|X| \leq k$ and for any $u, v \in T$, the vertices u and v belong to the same connected component of $G \setminus X$ if and only if $(u, v) \in \mathcal{R}$.

We proceed similarly as in the case of EDGE MULTIWAY CUT-UNCUT in the illustration, but using node separations of Section C. In particular, the highly connected phase is similarly simple as it is in the E-MWCU case. However, there is one issue that does not appear in the edge-deletion variant: a priori, in N-MWCU the equivalence relation \mathcal{R} may have arbitrarily large number of equivalence classes, whereas in the edge-deletion variant the number of connected components of $G \setminus X$ was clearly bounded by a function of k . We resolve this issue using a reduction similar to the one used by Razgon (Theorem 5 in [39]), that shows that we can limit ourselves to relations \mathcal{R} with $O(k^2)$ equivalence classes.

Our algorithm not only resolves N-MWCU instance, but, in the case of a YES answer, it returns a solution X with minimum possible $|X|$. This property will be used in the course of the algorithm.

F.1 Reduction of the number of equivalence classes

We now show how to reduce the number of equivalence classes of \mathcal{R} in an N-MWCU instance $\mathcal{J} = (G, T, \mathcal{R}, k)$.

Lemma F.1. *Let $\mathcal{J} = (G, T, \mathcal{R}, k)$ be an N-MWCU instance and let $v \in V(G) \setminus T$. Assume that there exist $k + 2$ paths P_1, P_2, \dots, P_{k+2} in G , such that:*

- for each $1 \leq i \leq k + 2$, the path P_i is a simple path that starts at v and ends at $v_i \in T$;
- the paths P_i have pairwise disjoint sets of vertices, except for the vertex v ;
- for any $i \neq j$, $(v_i, v_j) \notin \mathcal{R}$.

Then for any solution X in \mathcal{J} we have $v \in X$.

Proof. Let $X \subseteq V(G) \setminus T$ with $v \notin X$ and $|X| \leq k$. As the paths P_i are disjoint (except for v), there exist two indices $1 \leq i < j \leq k + 2$, such that P_i and P_j does not contain any vertex from X . A concatenation of P_i and P_j is a path from v_i to v_j that avoids X . As $(v_i, v_j) \notin \mathcal{R}$, we infer that X is not a solution to \mathcal{J} . \square

Lemma F.2. *Let $\mathcal{J} = (G, T, \mathcal{R}, k)$ be an N-MWCU instance. For any $v \in V(G)$, we can verify if v satisfies the conditions of Lemma F.1 in $O(kn^2)$ time.*

Proof. Consider the following auxiliary graph H . For each equivalence class $A \subseteq T$ of \mathcal{R} , we attach a new vertex t_A that is adjacent to all vertices of A . We make v an infinite-capacity source and each vertex t_A a unit-capacity sink; each other vertex of H has unit capacity. Clearly, v satisfies the conditions of Lemma F.1 iff there exists a flow of size at least $k + 2$ in H . As vertices in H have unit capacities (except for v), this can be done in $O(n^3)$ time by the classic Ford-Fulkerson algorithm. However, in our case we may stop the algorithm once it finds $k + 2$ augmenting paths; this observation reduces the running time to $O(kn^2)$. \square

Lemma F.1 justifies the following step.

Step F.1. For each $v \in V(G)$, if v satisfies the conditions of Lemma F.1, delete v from the graph and decrease k by one; if k becomes negative by this operation, return NO. Afterwards, restart the algorithm.

By Lemma F.2, each application of Step F.1 takes $O(kn^3)$ time. As we cannot apply Step F.1 more than k times, all applications of this step take $O(k^2n^3)$ time.

Let us now show that Step F.1 leads to a bound on the number of equivalence classes of \mathcal{R} .

Lemma F.3. *Let $\mathcal{J} = (G, T, \mathcal{R}, k)$ be a N-MWCU instance where Step F.1 is not applicable. If there exists a connected component of G that contains terminals of more than $k^2 + k$ equivalence classes of \mathcal{R} , then \mathcal{J} is a NO-instance to N-MWCU.*

Proof. Let C be the vertex set of any connected component of G , and let X be a solution to \mathcal{J} . Fix arbitrary $v \in X$. We say that v sees an equivalence class A of \mathcal{R} if there exists a connected component C_A of $G[C \setminus X]$ that contains a terminal of A and such that $N_G(v) \cap C_A \neq \emptyset$. Note that if v sees $k + 2$ equivalence classes of \mathcal{R} , then in each component C_A for each equivalence class A seen by v we can find a path from v to a terminal of A . Thus v satisfies the assumptions of Lemma F.1, as $C_A \neq C_B$ for $A \neq B$ (recall that X is a solution to \mathcal{J}). Therefore, each vertex $v \in X$ sees at most $k + 1$ equivalence classes of \mathcal{R} . From connectivity of $G[C]$ we infer that each component C_A must be seen by some element of X , so $C \setminus X$ may contain vertices of at most $k(k + 1)$ equivalence classes of \mathcal{R} . \square

Step F.2. If there exist $u, v \in T$ such that u and v lie in different connected components of G , but $(u, v) \in \mathcal{R}$, or there exists a connected component of G with terminals of more than $k^2 + k$ equivalence classes of \mathcal{R} , return NO.

Clearly, Step F.2 can be applied in $O(n^2)$ time.

We now ensure connectivity of G , by considering separately all connected components. Recall that we are developing an algorithm that not only resolves the given N-MWCU instance, but also, in case of the positive answer, returns a solution of minimum possible size.

Step F.3. For each connected component of G with vertex set C , pass the instance $(G, T \cap C, \mathcal{R}|_{T \cap C}, k)$ to the next step. If any of the subinstances returns NO, or if the union of the solutions to the subcases is larger than k , return NO. Otherwise, return YES and the union of the solutions for the connected components as the solution to the given instance.

The correctness of Step F.3 is straightforward (note that Step F.2 refutes instances where one equivalence class of \mathcal{R} is scattered among more than one connected component of G) and splitting G into connected components takes linear time in the size of G . Thus, from this point we may assume that G is connected and that the number of equivalence classes of \mathcal{R} is bounded by $\ell := k^2 + k$.

F.2 Operations on the input graph

In this section we show basic operations the algorithm repetitively applies to the graph.

Definition F.4. Let $\mathcal{J} = (G, T, \mathcal{R}, k)$ be an N-MWCU instance and let $v \in V(G) \setminus T$. By *bypassing* a vertex v we mean the following operation: we delete the vertex v from the graph and, for any $u_1, u_2 \in N_G(v)$, we add an edge u_1u_2 if it is not already present in G .

We now state the properties of the bypassing operation.

Lemma F.5. *Let $\mathcal{J} = (G, T, \mathcal{R}, k)$ be an N-MWCU instance, let $v \in V(G) \setminus T$ and let $\mathcal{J}' = (G', T, \mathcal{R}, k)$ be the instance \mathcal{J} with v bypassed. Then:*

- *if X is a solution to \mathcal{J}' , then X is a solution to \mathcal{J} as well;*
- *if X is a solution to \mathcal{J} and $v \notin X$ then X is a solution to \mathcal{J}' as well.*

Proof. The claim follows from the following correspondence of the paths in G and G' : any path P' in G' has a corresponding walk P in G , where each occurrence of an edge of $E(G') \setminus E(G)$ is replaced with a length-2 subpath via v . Moreover, any path P in G that does not start nor end in v has a corresponding path P' in G' , where a possible occurrence of v is circumvented by an edge in G' between two neighbours of v . \square

Apart from the bypassing operation, we need to show a way to reduce the number of terminals.

Definition F.6. Let $\mathcal{J} = (G, T, \mathcal{R}, k)$ be an N-MWCU instance and let $u, v \in T$ be two terminals with $u \neq v$, $(u, v) \in \mathcal{R}$. By *identifying* u and v we mean the following operation: we replace vertices u and v with a new vertex w_{uv} that is adjacent to all vertices of $N_G(u) \cup N_G(v)$. Moreover, we update \mathcal{R} by substituting u and v with w in the equivalence class they belong to.

Lemma F.7. *Let $\mathcal{J} = (G, T, \mathcal{R}, k)$ be an N-MWCU instance and let $u, v \in T$ be two different terminals with $(u, v) \in \mathcal{R}$, such that $uv \in E(G)$ or $|N_G(u) \cap N_G(v)| > k$. Let \mathcal{J}' be instance \mathcal{J} with terminals u and v identified. Then the set of solutions to \mathcal{J}' and \mathcal{J} are equal.*

Proof. The lemma follows from the fact that, for any $X \subseteq V(G) \setminus T$ of size at most k , in $G \setminus X$ the vertices u and v lie in the same connected component. \square

Lemma F.8. *Let $\mathcal{J} = (G, T, \mathcal{R}, k)$ be a N-MWCU instance and let $u_1, u_2, u_3 \in T$ be three different terminals of the same equivalence class of \mathcal{R} , pairwise nonadjacent and such that $N_G(u_1) = N_G(u_2) = N_G(u_3) \subseteq V(G) \setminus T$. Let \mathcal{J}' be obtained from \mathcal{J} by deleting the terminal u_3 (and all pairs that contain u_3 in \mathcal{R}). Then the set of solutions to \mathcal{J}' and \mathcal{J} are equal.*

Proof. Let $X \subseteq V(G) \setminus T$. We claim that for any $u, v \in V(G) \setminus \{u_3\}$, u and v are in the same connected component of $G \setminus X$ if and only if they are in the same connected component of $G' \setminus X$. Indeed, the backward implication is trivial, whereas for the forward implication observe that any path from u to v in $G \setminus X$ that visits u_3 can be redirected via u_1 or u_2 .

The proven equivalence already shows that any solution to \mathcal{J} is a solution to \mathcal{J}' as well. For the other direction, we need to additionally verify that for any $v \in T$, we have $(u_3, v) \in \mathcal{R}$ if and only if v and u_3 are in the same connected component of $G \setminus X$, assuming that X is a solution to \mathcal{J}' . As X is a solution to \mathcal{J}' and $(u_1, u_2) \in \mathcal{R}$, there exists $w \in N_G(u_1) \setminus X$. Therefore u_1, u_2 and u_3 are in the same connected component of $G \setminus X$. As $(u_1, v) \in \mathcal{R}$ iff $(u_3, v) \in \mathcal{R}$, the claim follows. \square

F.3 Borders and recursive understanding

For the recursive understanding phase, we need to define the bordered problem. Let $\mathcal{J} = (G, T, \mathcal{R}, k)$ be an N-MWCU instance and let $T_b \subseteq V(G) \setminus T$ be a set of border terminals; we assume $|T_b| \leq 2k$. Define $\mathcal{J}_b = (G, T, \mathcal{R}, k, T_b)$ to be an instance of the bordered problem. By $\mathbb{P}(\mathcal{J}_b)$ we define the set of all triples $\mathcal{P} = (X_b, E_b, \mathcal{R}_b)$, such that $X_b \subseteq T_b$, E_b is an equivalence relation on $T_b \setminus X_b$ and \mathcal{R}_b is an equivalence relation on $T \cup (T_b \setminus X_b)$ such that $E_b \subseteq \mathcal{R}_b$ and $\mathcal{R}_b|_T = \mathcal{R}$. For a triple $\mathcal{P} = (X_b, E_b, \mathcal{R}_b)$, by $G_{\mathcal{P}}$ we denote the graph $G \cup E_b$, that is, the graph G with additional edges E_b .

We say that a set $X \subseteq V(G) \setminus T$ is a solution to $(\mathcal{J}_b, \mathcal{P})$ if $|X| \leq k$, $X \cap T_b = X_b$ and for any $u, v \in T \cup (T_b \setminus X_b)$, the vertices u and v are in the same connected component of the graph $G_{\mathcal{P}} \setminus X$ (i.e., we delete vertices X and add edges E_b) if and only if $(u, v) \in \mathcal{R}_b$. Note that, if X is a solution to $(\mathcal{J}_b, \mathcal{P})$, the set X is not necessarily a solution to \mathcal{J}_b ; however, X is a solution to the N-MWCU instance $(G_{\mathcal{P}}, T, \mathcal{R}, k)$.

We also say that X is a solution to $\mathcal{J}_b = (G, T, \mathcal{R}, k, T_b)$ whenever X is a solution to $\mathcal{J} = (G, T, \mathcal{R}, k)$.

We define the bordered problem as follows.

BORDER N-MWCU

Input: An N-MWCU instance $\mathcal{J} = (G, T, \mathcal{R}, k)$ with G being connected and a set $T_b \subseteq V(G) \setminus T$ of size at most $2k$; denote $\mathcal{J}_b = (G, T, \mathcal{R}, k, T_b)$

Output: For each $\mathcal{P} = (X_b, E_b, \mathcal{R}_b) \in \mathbb{P}(\mathcal{J}_b)$, output a $\text{sol}_{\mathcal{P}} = X_{\mathcal{P}}$ being a solution to $(\mathcal{J}_b, \mathcal{P})$ with minimum possible $|X_{\mathcal{P}}|$, or $\text{sol}_{\mathcal{P}} = \perp$ if such a solution does not exist.

Clearly, N-MWCU reduces to BORDER N-MWCU, as we may ask for an instance with $T_b = \emptyset$. Moreover, in this case the single answer to BORDER N-MWCU for $\mathcal{P} = (\emptyset, \emptyset, \mathcal{R})$ returns a solution of minimum possible size.

We note that

$$|\mathbb{P}(\mathcal{J}_b)| \leq (1 + |T_b|(|T_b| + \ell))^{|T_b|} \leq (2k^3 + 6k^2 + 1)^{2k} = 2^{O(k \log k)},$$

as \mathcal{R}_b has at most $\ell + |T_b|$ equivalence classes, E_b has at most $|T_b|$ equivalence classes, and each $v \in T_b$ can go either to X_b or choose an equivalence class in \mathcal{R}_b and E_b . Let $q = k(2k^3 + 6k^2 + 1)^{2k} + k$; all output solutions to a BORDER N-MWCU instance \mathcal{J}_b contain at most $q - k$ vertices in total.

The following lemma prepares the ground for a recursive step in our BORDER N-MWCU algorithm.

Lemma F.9. *Assume we are given a BORDER N-MWCU instance $\mathcal{J}_b = (G, T, \mathcal{R}, k, T_b)$ and two disjoint sets of vertices $Z, V^* \subseteq V(G)$, such that $|Z| \leq k$, $Z \cap T = \emptyset$, $Z_W := N_G(V^*) \subseteq Z$, $|V^* \cap T_b| \leq k$ and the subgraph of G induced by $W := V^* \cup Z_W$ is connected. Denote $G^* = G[W]$, $T_b^* = (T_b \cup Z_W) \cap W$, $T^* = T \cap W$, $\mathcal{R}^* = \mathcal{R}|_{T \cap W}$ and $\mathcal{J}_b^* = (G^*, T^*, \mathcal{R}^*, k, T_b^*)$. Then \mathcal{J}_b^* is a proper BORDER N-MWCU instance. Moreover, if we denote by $(\text{sol}_{\mathcal{P}^*}^*)_{\mathcal{P}^* \in \mathbb{P}(\mathcal{J}_b^*)}$ an arbitrary output to the BORDER N-MWCU instance \mathcal{J}_b^* and*

$$U(\mathcal{J}_b^*) = T_b^* \cup \bigcup \{X_{\mathcal{P}^*}^* : \mathcal{P}^* \in \mathbb{P}(\mathcal{J}_b^*), \text{sol}_{\mathcal{P}^*}^* = X_{\mathcal{P}^*}^* \neq \perp\},$$

then there exists a correct output $(\text{sol}_{\mathcal{P}})_{\mathcal{P} \in \mathbb{P}(\mathcal{J}_b)}$ to the BORDER N-MWCU instance \mathcal{J}_b such that whenever $\text{sol}_{\mathcal{P}} = X_{\mathcal{P}} \neq \perp$ then $X_{\mathcal{P}} \cap V^ \subseteq U(\mathcal{J}_b^*)$.*

Proof. The claim that \mathcal{J}_b^* is a proper BORDER N-MWCU instance follows directly from the assumptions that $G^* = G[W]$ is connected, $|Z_W| \leq |Z| \leq k$ and $|V^* \cap T_b| \leq k$. In the rest of the proof we justify the second claim of the lemma.

Fix $\mathcal{P} = (X_b, E_b, \mathcal{R}_b) \in \mathbb{P}(\mathcal{J}_b)$ and recall that $G_{\mathcal{P}} = G \cup E_b$. Assume that there exists a solution to the instance $(\mathcal{J}_b, \mathcal{P})$; let $X_{\mathcal{P}}$ be such a solution with minimum possible $|X_{\mathcal{P}}|$. To prove the lemma we need to show a second solution $X'_{\mathcal{P}}$ to $(\mathcal{J}_b, \mathcal{P})$, $|X'_{\mathcal{P}}| \leq |X_{\mathcal{P}}|$ and $X'_{\mathcal{P}} \cap V^* \subseteq U(\mathcal{J}_b^*)$.

Define a triple $\mathcal{P}^* = (X_b^*, E_b^*, \mathcal{R}_b^*)$ as follows.

- Let $X_b^* = X_{\mathcal{P}} \cap T_b^*$; note that $X_b \cap W \subseteq X_b^*$ since $X_{\mathcal{P}}$ is a solution to $(\mathcal{J}_b, \mathcal{P})$.
- We define E_b^* to be the following relation on $T_b^* \setminus X_b^*$: $(u, v) \in E_b^*$ iff u and v are in the same connected component of $G_{\mathcal{P}} \setminus ((V^* \setminus T_b) \cup X_{\mathcal{P}})$ (in particular, if $(u, v) \in E_b$).
- Finally, we define \mathcal{R}_b^* to be the following relation on $T^* \cup (T_b^* \setminus X_b^*)$: $(u, v) \in \mathcal{R}_b^*$ iff u and v are in the same connected component of $G_{\mathcal{P}} \setminus X_{\mathcal{P}}$. As $X_{\mathcal{P}}$ is a solution to $(\mathcal{J}_b, \mathcal{P})$, $\mathcal{R}_b^*|_{T^*} = \mathcal{R}^*$.

Moreover, note that $E_b^* \subseteq \mathcal{R}_b^*$, as both E_b^* and \mathcal{R}_b^* corresponds to the relation of being in the same connected component, but in E_b^* we consider a smaller graph than in \mathcal{R}_b^* .

Clearly, $\mathcal{P}^* \in \mathbb{P}(\mathcal{J}_b^*)$. Moreover, we claim that $X_{\mathcal{P}} \cap W$ is a solution to $(\mathcal{J}_b^*, \mathcal{P}^*)$. Clearly, $|X_{\mathcal{P}} \cap W| \leq k$. By the definition of X_b^* , we have $X_{\mathcal{P}} \cap W \cap T_b^* = X_b^*$. Consider two vertices $u, v \in T^* \cup (T_b^* \setminus X_b^*)$. We have $(u, v) \in \mathcal{R}_b^*$ iff there exists a path P between u and v in $G_{\mathcal{P}} \setminus X_{\mathcal{P}}$. By the definition of E_b^* , such a path P exists iff there exists a path P^* connecting u and v in $G_{\mathcal{P}^*}^* \setminus X_{\mathcal{P}}$: each subpath of P with internal vertices in $V(G) \setminus W$ corresponds to an edge in E_b^* and vice versa. Thus, u and v are in the same connected component of $G_{\mathcal{P}^*}^* \setminus X_{\mathcal{P}}$ if and only if $(u, v) \in \mathcal{R}_b^*$ and the claim is proven.

We infer that $\text{sol}_{\mathcal{P}^*}^* = X_{\mathcal{P}^*}^* \neq \perp$ and $|X_{\mathcal{P}^*}^*| \leq |X_{\mathcal{P}} \cap W|$. Let $X'_{\mathcal{P}} = (X_{\mathcal{P}} \setminus W) \cup X_{\mathcal{P}^*}^*$. We claim that $X'_{\mathcal{P}}$ is a solution to $(\mathcal{J}_b, \mathcal{P})$ as well; as $|X'_{\mathcal{P}}| \leq |X_{\mathcal{P}}|$, this would finish the proof of the lemma.

As X_b^* is defined as $X_{\mathcal{P}} \cap T_b^*$ and $X_{\mathcal{P}}$ is a solution to $(\mathcal{J}_b, \mathcal{P})$, we have $X'_{\mathcal{P}} \cap T_b = X_b$.

Let $u, v \in T \cup (T_b \setminus X_b)$. Our goal is to show that u and v lie in the same connected component of $G_{\mathcal{P}} \setminus X'_{\mathcal{P}}$ if and only if they lie in the same connected component of $G_{\mathcal{P}} \setminus X_{\mathcal{P}}$. We present the proof only in one direction, as the proofs in both directions are totally symmetric: we use only the facts that both $X_{\mathcal{P}} \cap W$ and $X'_{\mathcal{P}} \cap W$ are solutions to $(\mathcal{J}_b^*, \mathcal{P}^*)$ and that $X_{\mathcal{P}} \setminus V^* = X'_{\mathcal{P}} \setminus V^*$. The last equality holds because $Z_W \subseteq T_b^*$, so $Z_W \cap X_{\mathcal{P}} = Z_W \cap X_{\mathcal{P}^*}^*$.

Thus, assume that $u, v \in T \cup (T_b \setminus X_b)$ and u and v lie in the same connected component of $G_{\mathcal{P}} \setminus X_{\mathcal{P}}$. Let P be a path that connects u and v in $G_{\mathcal{P}} \setminus X_{\mathcal{P}}$ and let $u = v_0, v_1, v_2, \dots, v_r = v$ be a sequence of vertices that lie on the path P and belong to $D := T \cup (T_b \setminus X_b) \cup Z_W$, in the order they appear on P . First note that, since $X_{\mathcal{P}} \setminus V^* = X'_{\mathcal{P}} \setminus V^*$ and both $X_{\mathcal{P}} \cap W$ and $X'_{\mathcal{P}} \cap W$ are solutions to $(\mathcal{J}_b^*, \mathcal{P}^*)$, we have that $X_{\mathcal{P}} \cap D = X'_{\mathcal{P}} \cap D = X_b \cup X_b^*$. Thus, for each $0 \leq i \leq r$, we have $v_i \notin X'_{\mathcal{P}}$. To finish the proof of the lemma we need to show that for any $0 \leq i < r$, the vertices v_i and v_{i+1} lie in the same connected component of $G_{\mathcal{P}} \setminus X'_{\mathcal{P}}$.

Let P_i be the subpath of P between v_i and v_{i+1} . As $Z_W \subseteq D$, P_i is either a path in $G_{\mathcal{P}} \setminus V^*$ or a path in $G_{\mathcal{P}}[W]$. In the first case, since $X_{\mathcal{P}} \setminus V^* = X'_{\mathcal{P}} \setminus V^*$, we infer that the path P_i is present in $G_{\mathcal{P}} \setminus X'_{\mathcal{P}}$ and the claim is proven. In the second case, note that we have $(v_i, v_{i+1}) \in \mathcal{R}_b^*$. As $X'_{\mathcal{P}} \cap W$ is a solution to $(\mathcal{J}_b^*, \mathcal{P}^*)$, we infer that v_i and v_{i+1} are connected via a path P_i^* in $G_{\mathcal{P}^*}^* \setminus (X'_{\mathcal{P}} \cap W)$. However, by the definition of E_b^* , for any edge $w_1 w_2 \in E_b^*$ on P_i^* , the vertices w_1 and w_2 are in the same connected component of $G_{\mathcal{P}} \setminus ((V^* \setminus T_b) \cup X_{\mathcal{P}})$. Since $X_{\mathcal{P}} \setminus V^* = X'_{\mathcal{P}} \setminus V^*$ and $X_{\mathcal{P}} \cap T_b = X'_{\mathcal{P}} \cap T_b$, we have that $X_{\mathcal{P}} \setminus (V^* \setminus T_b) = X'_{\mathcal{P}} \setminus (V^* \setminus T_b)$ and the claim is proven. This finishes the proof of the lemma. \square

Note that in Lemma F.9 we have $|U(\mathcal{J}_b^*) \cap V^*| \leq q$.

A recursive call due to an application of Lemma F.9 allows us to reduce the number of nonterminal vertices in V^* to at most $q = 2^{O(k \log k)}$. To make the recursion work in FPT time, we need to reduce the number of terminals as well. Fortunately, this is quite easy, due to the identifying operation and Lemma F.7.

We are now ready to present the recursive step of the algorithm.

Step F.4. Assume we are given a BORDER N-MWCU instance $\mathcal{J}_b = (G, T, \mathcal{R}, k, T_b)$. Invoke first the algorithm of Lemma C.3 in a search for (q, k) -good node separation (with $V^\infty = T$). If it returns a good node separation (Z, V_1, V_2) , let $j \in \{1, 2\}$ be such that $|V_j \cap T_b| \leq k$ and denote $Z^* = Z$, $V^* = V_j$. Otherwise, if it returns that no such good node separation exists in G , invoke the algorithm of Lemma C.4 in a search for (q, k) -flower separation w.r.t. T_b (with $V^\infty = T$ again). If it returns that no such flower separation exists in G , pass the instance \mathcal{J}_b to the next step. Otherwise, if it returns a flower separation $(Z, (V_i)_{i=1}^\ell)$, denote $Z^* = Z$ and $V^* = \bigcup_{i=1}^\ell V_i$.

In the case we have obtained Z^* and V^* (either from Lemma C.3 or Lemma C.4), invoke the algorithm recursively for the BORDER N-MWCU instance \mathcal{J}_b^* defined as in the statement of Lemma F.9 for separator Z^* and set V^* , obtaining an output $(\mathbf{sol}_{\mathcal{P}}^*)_{\mathcal{P} \in \mathbb{P}(\mathcal{J}_b^*)}$. Compute the set $U(\mathcal{J}_b^*)$. Bypass (in an arbitrary order) all vertices of $V^* \setminus (T \cup U(\mathcal{J}_b^*))$. Recall that $T_b^* \subseteq U(\mathcal{J}_b^*)$, so no border terminal gets bypassed.

After all vertices of $V^* \setminus U(\mathcal{J}_b^*)$ are bypassed, perform the following operations on terminals of $V^* \cap T$:

1. As long as there exist two different $u, v \in V^* \cap T$ that satisfy $uv \in E(G)$ or $|N_G(u) \cap N_G(v)| > k$ do as follows: if $(u, v) \in \mathcal{R}$, identify u and v , and otherwise output \perp for all $\mathcal{P} \in \mathbb{P}(\mathcal{J}_b)$.
2. If the above is not applicable, then, as long as there exist three pairwise distinct terminals $u_1, u_2, u_3 \in T$ of the same equivalence class of \mathcal{R} that have the same neighborhood, delete u_3 from the graph (and delete all pairs containing u_3 from \mathcal{R}).

Let \mathcal{J}'_b be the outcome instance.

Finally, restart this step on the new instance \mathcal{J}'_b and obtain a family of solutions $(\mathbf{sol}'_{\mathcal{P}})_{\mathcal{P} \in \mathbb{P}(\mathcal{J}_b)}$ and return this family as an output to the instance \mathcal{J}_b .

Let us first verify that the application of Lemma F.9 is justified. Indeed, by the definitions of the good node separation and the flower separation, as well as the choice of V^* , we have in both cases $|V^* \cap T_b| \leq k$ and that $G[V^* \cup N_G(V^*)]$ is connected. Moreover, note that the recursive call is applied to a graph with strictly smaller number of vertices than G : in the case of a good node separation, V_2 is removed from the graph, and in the case of a flower separation, recall that the definition of the flower separation requires $Z \cup \bigcup_{i=1}^\ell V_i$ to be a proper subset of $V(G)$.

We have that, after the bypassing operations, V^* contains at most q vertices that are not terminals (at most k border terminals and at most $q - k$ vertices which are neither terminals nor border terminals). Let us now bound the number of terminal vertices once Step F.4 is applied. Note that, after Step F.4 is applied, for any $v \in T \cap V^*$, we have $N_G(v) \subseteq (V^* \setminus T) \cup Z$ and $|(V^* \setminus T) \cup Z| \leq (q + k)$. Due to the first rule in Step F.4, for any set $A \subseteq (V^* \setminus T) \cup Z$ of size $k + 1$, at most one terminal of $T \cap V^*$ is adjacent to all vertices of A . Due to the second rule in Step F.4, for any set $B \subseteq (V^* \setminus T) \cup Z$ of size at most k and for each equivalence class of \mathcal{R} , there are at most two terminals of this equivalence class with neighborhood exactly B . We infer that

$$|T \cap V^*| \leq (q + k)^{k+1} + 2\ell \sum_{i=1}^k (q + k)^i =: q'.$$

Note that $q' = 2^{O(k^2 \log k)}$.

The following lemma verifies the correctness of Step F.4.

Lemma F.10. *Assume we are given a BORDER N-MWCU instance $\mathcal{J}_b = (G, T, \mathcal{R}, k, T_b)$ on which Step F.4 is applied, and let \mathcal{J}'_b be an instance after Step F.4 is applied. Then any correct output to the instance \mathcal{J}'_b is a correct output to the instance \mathcal{J}_b as well. Moreover, if Step F.4 outputs \perp for all $\mathcal{P} \in \mathbb{P}(\mathcal{J}_b)$, then this is a correct output to \mathcal{J}_b .*

Proof. The lemma is a straightforward corollary of Lemma F.9, the properties of the bypassing operation described in Lemma F.5, and Lemmas F.7 and F.8. Lemma F.9 ensures us that each vertex not in $U(\mathcal{J}'_b)$ is omitted by some optimal solution for every $\mathcal{P} \in \mathbb{P}(\mathcal{J}_b)$, which enables us to use Lemma F.5. Finally, if for any terminals $u, v \in T$, we have $uv \in E(G)$ or $|N_G(u) \cap N_G(v)| > k$, then u and v are in the same connected component of $G \setminus X$ for any set X of at most k nonterminals and, if $(u, v) \notin \mathcal{R}$, for any $\mathcal{P} \in \mathbb{P}(\mathcal{J}_b)$, there is no solution to $(\mathcal{J}_b, \mathcal{P})$. \square

We are left with the analysis of the time complexity of Step F.4. The applications of Lemmas C.3 and C.4 use $O(2^{O(\min(q,k)\log(q+k))}n^3 \log n) = O(2^{O(k^2 \log k)}n^3 \log n)$ time. Let $n' = |V^*|$; the recursive step is applied to a graph with at most $n' + k$ vertices and, after bypassing, there are at most $\min(n - 1, n - n' + q + q')$ vertices left. Moreover, each bypassing operation takes $O(n^2)$ time, the computation of $U(\mathcal{J}'_b)$ takes $O(2^{O(k \log k)}n)$ time. Application of Lemma F.7 takes $O(kn^2)$ time per operation, which can be implemented by having a counter for each pair of terminals and increasing those counters accordingly by considering every pair of terminals of $N_G(x)$, for each $x \in V$. Since when a counter reaches value $k + 1$ for vertices u, v , we know that $|N_G(u) \cap N_G(v)| > k$, the total time consumed is bounded by $O(kn^2)$. Application of Lemma F.8 takes $O(n^2 \log n)$ time per one operation, since we can sort terminals from one equivalence class according to their sets of neighbours. Thus all applications of Lemmata F.7 and F.8 take $O(n^3(k + \log n))$ time in total. The value of k do not change in this step. Therefore, we have the following recursive formula for time complexity as a function of the number of vertices of G :

$$T(n) \leq \max_{q+1 \leq n' \leq n-q-1} \left(O(2^{O(k^2 \log k)}n^3 \log n) + T(n' + k) + T(\min(n - 1, n - n' + q + q')) \right). \quad (4)$$

Note that the function $p(t) = t^4 \log t$ is convex, so it is easy to see that the maximum is attained either when $n' = q + q' - 1$, or when $n' = n - q - 1$. A straightforward inductive check of both of the ends proves that we have indeed the claimed bound on the complexity, i.e., $T(n) = O(2^{O(k^2 \log k)}n^4 \log n)$.

We conclude this section with a note that Lemma C.5 asserts that, if Step F.4 is not applicable, then for any set $Z \subseteq V(G) \setminus T$ of size at most k , the graph $G \setminus Z$ contains at most $t := (2q + 2)(2^k - 1) + 2k + 1$ connected components containing a non-terminal, out of which at most one has more than q vertices not from T .

F.4 Brute force approach

If the graph output by Step F.4 has small number of vertices outside T , the algorithm may apply a straightforward brute-force approach to the BORDER N-MWCU problem. In this section we describe this method formally.

Lemma F.11. *A correct output to a BORDER N-MWCU instance $\mathcal{J}_b = (G, T, \mathcal{R}, k, T_b)$ can be computed in $O(2^{O(k \log k)}n^2 n_{-T}^k)$ time, where $n_{-T} = |V(G) \setminus T|$.*

Proof. Simply, for each $\mathcal{P} \in \mathbb{P}(\mathcal{J}_b)$ (at most $2^{O(k \log k)}$ choices) for each deletion set $X \subseteq V(G) \setminus T$ with $|X| \leq k$ (at most $(k+1)n_{-T}^k$ choices) we verify in $O(n^2)$ time if X is a solution to $(\mathcal{J}_b, \mathcal{P})$. \square

Step F.5. If $|V(G) \setminus T| \leq qt + k$, apply Lemma F.11 to find a correct output to a BORDER N-MWCU instance $\mathcal{J}_b = (G, T, \mathcal{R}, k, T_b)$.

Recall that $q, t \leq 2^{O(k \log k)}$. Thus, if Step F.5 is applicable, its running time is $O(2^{O(k^2 \log k)}n^2)$.

F.5 High connectivity phase

Assume we have a BORDER N-MWCU instance $\mathcal{J}_b = (G, T, \mathcal{R}, k, T_b)$ where Steps F.4 and F.5 are not applicable. In this section we show that high connectivity of G makes the problem much easier. To this end, fix $\mathcal{P} = (X_b, E_b, \mathcal{R}_b) \in \mathbb{P}(\mathcal{J}_b)$. We focus on finding the solution $\text{sol}_{\mathcal{P}}$; iterating through all the possible \mathcal{P} gives additional $2^{O(k \log k)}$ overhead to the running time. Recall that $G_{\mathcal{P}} = G \cup E_b$.

Note that, if $|V(G) \setminus T|$ is too large for Step F.5 to be applicable, for any set $Z \subseteq V(G) \setminus T$ of size at most k , the bound on the number of connected components from Lemma C.5 implies that there exists exactly one connected component of $G \setminus Z$ with more than q vertices outside T ; denote its vertex set by $\text{big}(Z)$.

We now use Lemma 1.1 to get some more structure of the graph G .

Definition F.12. Let $Z \subseteq V(G) \setminus T$ be a set of size at most k and let $S \subseteq V(G) \setminus T$. We say that S *interrogates* Z if the following holds:

1. $S \cap Z = \emptyset$;
2. for any connected component C of $G \setminus Z$ with at most q vertices outside T , all vertices of C belong to $S \cup T$.

Lemma F.13. Let \mathcal{F} be a family obtained by the algorithm of Lemma 1.1 for universe $U = V(G) \setminus T$ and constants $a = qt$ and $b = k$, Then, for any $Z \subseteq V(G) \setminus T$ with $|Z| \leq k$, there exists a set $S \in \mathcal{F}$ that interrogates Z .

Proof. Fix $Z \subseteq V(G) \setminus T$ with $|Z| \leq k$. Let A be the union of vertex sets of all connected components of $G \setminus Z$ that have at most q vertices outside T ; by Lemma C.5, $|A \setminus T| \leq qt$. By Lemma 1.1, as $|A \setminus T| \leq qt$ and $|Z| \leq k$, there exists a set $S \in \mathcal{F}$ that contains $A \setminus T$ and is disjoint with Z . By the construction of the set A , S interrogates Z and the lemma is proven. \square

Note that, as $q, t = 2^{O(k \log k)}$, the family \mathcal{F} of Lemma F.13 is of size $2^{O(k^2 \log k)} \log n$ and can be computed in $O(2^{O(k^2 \log k)} n \log n)$ time. Therefore we may branch, guessing a set S that interrogates a solution $\text{sol}_{\mathcal{P}} = X_{\mathcal{P}}$ we are looking for. Formally, we perform computations in each branch and return the minimum size solution from those obtained in the branches.

Step F.6. Compute the family \mathcal{F} from Lemma F.13 and branch into $|\mathcal{F}|$ subcases, indexed by sets $S \in \mathcal{F}$. In a branch S we seek for a set $X_{\mathcal{P}}$ with minimum possible $|X_{\mathcal{P}}|$ that not only is a solution to $(\mathcal{J}_b, \mathcal{P})$, but also is interrogated by S .

Lemma F.13 verifies the correctness of the branching of Step F.6; as discussed, the step is applied in $O(2^{O(k^2 \log k)} n \log n)$ time and leads to $O(2^{O(k^2 \log k)} \log n)$ subcases.

The following observation is crucial to for the final step.

Lemma F.14. Let $X_{\mathcal{P}}$ be a minimum size set that is a solution to $(\mathcal{J}_b, \mathcal{P})$ interrogated by S . Then there exists a set $T^{\text{big}} \subseteq T \cup (T_b \setminus X_b)$ that is empty or contains all vertices of exactly one equivalence class of \mathcal{R}_b , such that $X_{\mathcal{P}} = X_b \cup N_G(S(T^{\text{big}}))$, where $S(T^{\text{big}})$ is the union of vertex sets of all connected components of $G[S \cup T \cup (T_b \setminus X_b)]$ that contain a vertex of $(T \cup (T_b \setminus X_b)) \setminus T^{\text{big}}$.

Proof. Consider the graph $G_{\mathcal{P}} \setminus X_{\mathcal{P}}$ and let $\text{big}_{\mathcal{P}}(X_{\mathcal{P}})$ be the vertex set of the connected component of $G_{\mathcal{P}} \setminus X_{\mathcal{P}}$ that contain $\text{big}(X_{\mathcal{P}})$ (recall that $G_{\mathcal{P}}$ is the graph G with additional edges $E_{\mathcal{P}}$; thus $\text{big}_{\mathcal{P}}(X_{\mathcal{P}})$ may be significantly larger than $\text{big}(X_{\mathcal{P}})$). As $X_{\mathcal{P}}$ is a solution to $(\mathcal{J}_b, \mathcal{P})$, we have $X_{\mathcal{P}} \cap T_b = X_b$. Define $T^{\text{big}} = (T \cup (T_b \setminus X_b)) \cap \text{big}_{\mathcal{P}}(X_{\mathcal{P}})$; as $X_{\mathcal{P}}$ is a solution to $(\mathcal{J}_b, \mathcal{P})$, T^{big} is empty or contains vertices of exactly one equivalence class of \mathcal{R}_b .

Now let C be the vertex set of a connected component of $G \setminus X_{\mathcal{P}}$ that contains a vertex $v \in (T \cup (T_b \setminus X_b)) \setminus T^{\text{big}}$. Clearly, $v \notin \text{big}_{\mathcal{P}}(X_{\mathcal{P}})$. As S interrogates $X_{\mathcal{P}}$, $\text{big}_{\mathcal{P}}(X_{\mathcal{P}})$ contains $\text{big}(X_{\mathcal{P}})$ and $X_{\mathcal{P}} \cap (T \cup T_b) = X_b \subseteq T_b$, we infer that C is the vertex set of a connected component of $G[S \cup T \cup (T_b \setminus X_b)]$ as well. As $v \in C$, C is a connected component of $G[S(T^{\text{big}})]$. Since the choice of C was arbitrary, we infer that $N_G(S(T^{\text{big}})) \subseteq X_{\mathcal{P}}$. Denote $X'_{\mathcal{P}} = X_b \cup N_G(S(T^{\text{big}})) \subseteq X_{\mathcal{P}}$. To finish the proof of the lemma we need to show that $X'_{\mathcal{P}}$ is a solution to $(\mathcal{J}_b, \mathcal{P})$ as well.

Clearly, $X'_{\mathcal{P}} \cap (T \cup T_b) = X_b$, as $N_G(S(T^{\text{big}})) \cap (T \cup (T_b \setminus X_b)) = \emptyset$ by the definition of $S(T^{\text{big}})$. Moreover, as $X'_{\mathcal{P}} \subseteq X_{\mathcal{P}}$ and $X_{\mathcal{P}}$ is a solution to $(\mathcal{J}_b, \mathcal{P})$, if $(u, v) \in \mathcal{R}_b$ then u and v are in the same connected component of $G_{\mathcal{P}} \setminus X'_{\mathcal{P}}$. We now show that for any $(u, v) \notin \mathcal{R}_b$ the vertices u and v are in different connected components of $G_{\mathcal{P}} \setminus X'_{\mathcal{P}}$. Assume the contrary, and let $u, v \in T \cup (T_b \setminus X_b)$ be such that $(u, v) \notin \mathcal{R}_b$, u and v are in the same connected component of $G_{\mathcal{P}} \setminus X'_{\mathcal{P}}$ and that the distance between u and v in $G_{\mathcal{P}} \setminus X'_{\mathcal{P}}$ is minimum possible. Let P be a shortest path between u and v in $G_{\mathcal{P}} \setminus X'_{\mathcal{P}}$.

As $X_{\mathcal{P}}$ is a solution to (\mathcal{J}_b, X_b) , u and v are in different connected components of $G_{\mathcal{P}} \setminus X_{\mathcal{P}}$; without loss of generality assume $v \notin \text{big}_{\mathcal{P}}(X_{\mathcal{P}})$ and let C be the vertex set of the connected component of $G \setminus X_{\mathcal{P}}$ that contains v . Clearly, since $(u, v) \notin \mathcal{R}_b$, we have $u \notin C$. Moreover, $v \in (T \cup (T_b \setminus X_b)) \setminus T^{\text{big}}$ and C is a connected component of $G[S(T^{\text{big}})]$. Therefore $N_G(C) \subseteq X'_{\mathcal{P}}$. Since $u \notin C$, the path P needs to go via an edge $v_1 u_1 \in E_b$, where $v_1 \in C$ but $u_1 \notin C$. Note that then $u_1, v_1 \in T_b$. As $v_1 \in C$ and $X_{\mathcal{P}}$ is a solution to $(\mathcal{J}_b, \mathcal{P})$, we have $(v, v_1) \in \mathcal{R}_b$. As $E_b \subseteq \mathcal{R}_b$, we have that $(v, u_1) \in \mathcal{R}_b$. As $(u, v) \notin \mathcal{R}_b$, we infer that $(u_1, u) \notin \mathcal{R}_b$, but u_1 and u are connected via a proper subpath of P in $G_{\mathcal{P}} \setminus X'_{\mathcal{P}}$, a contradiction to the choice of u, v and P . This finishes the proof of the lemma. \square

Lemma F.14 justifies the final step.

Step F.7. In each branch, let S be the corresponding guess, for each set T^{big} that is empty or contains all vertices of one equivalence class of \mathcal{R}_b , check if $X_b \cup N_G(S(T^{\text{big}}))$ is a solution to $(\mathcal{J}_b, \mathcal{P})$ that is interrogated by S . For given \mathcal{P} , output the smallest solution to $(\mathcal{J}_b, \mathcal{P})$ found, or \perp if no solution is found for any choice of S and T^{big} .

Note that \mathcal{R} has at most $\ell = k^2 + k$ equivalence classes. As $|T_b| \leq 2k$, there are at most $1 + 3k + k^2$ choices of the set T^{big} . For each T^{big} , computing $X_b \cup N_G(S(T^{\text{big}}))$ and verifying if it is a solution to $(\mathcal{J}_b, \mathcal{P})$ interrogated by S takes $O(n^2)$ time. Therefore Step F.7 takes $O(2^{O(k^2 \log k)} n^2 \log n)$ time for all subcases.

This finishes the description of fixed-parameter algorithm for NODE MULTIWAY CUT-UNCUT.

G Lower bound for big alphabet size

In this section we prove that the dependence on s in the algorithm from Theorem 1.2 is probably essential, even for the edge deletion case and in the classical setting, when every vertex has a full list of possible labels and the partial permutations on edges are required to be permutations. We define formally the problem as follows:

<p>EDGE UNIQUE LABEL COVER (k)</p> <p>Input: An undirected graph G, a finite alphabet Σ of size s, an integer k, and: for each vertex $v \in V(G)$ a set $\phi_v \subseteq \Sigma$ and for each edge $e \in E(G)$ and each its endpoint v a partial permutation $\psi_{e,v}$ of Σ, such that if $e = uv$ then $\psi_{e,u} = \psi_{e,v}^{-1}$.</p> <p>Question: Does there exist a set $F \subseteq E(G)$ of size at most k and a function $\Psi : V(G) \rightarrow \Sigma$ such that for any $v \in V(G)$ we have $\Psi(v) \in \phi_v$ and for any $uv \in E(G) \setminus F$ we have $(\Psi(u), \Psi(v)) \in \psi_{uv,u}$?</p>	<p>Parameter: k</p>
--	---

Theorem G.1. EDGE UNIQUE LABEL COVER (k) is $W[1]$ -hard, even in the restricted case, when $\phi_v = \Sigma$ for all $v \in V(G)$ and $\psi_{uv,u}, \psi_{uv,v}$ are permutations for all $uv \in E(G)$.

Before we proceed to the proof, we state that this restricted case is not easier than the general one.

Lemma G.2. *There exists a polynomial time algorithm that, given an EDGE UNIQUE LABEL COVER instance $I = (G, \Sigma, k, (\phi_v)_{v \in V(G)}, (\psi_{e,v})_{e \in E(G), v \in e})$, outputs an equivalent instance $I' = (G', \Sigma', k', (\phi'_v)_{v \in V(G)}, (\psi'_{e,v})_{e \in E(G), v \in e})$ where $k' = k(k+2)$, $|\Sigma'| = |\Sigma| + k + 2$, $\phi'_v = \Sigma'$ for all $v \in V(G)$ and $\psi'_{e,v}$ is a permutation for all $e \in E(G), v \in e$.*

Proof. The graph G' we are going to construct will be a multigraph, possibly with loops. Note that we can easily get rid of multiple edges and loops by subdividing every edge and loop, and for each subdivision preserving the constraint on one of the obtained edges while setting the constraint on the other edge to be identity.

We start with setting $k' = k(k+2)$ and $\Sigma' = \Sigma \cup \Gamma$, where $\Gamma = \{\gamma_1, \gamma_2, \dots, \gamma_{k+2}\}$ is the set of $k+2$ new symbols that do not belong to Σ . Now we construct the multigraph G' as follows. Firstly, $V(G') = V(G)$. For every vertex $v \in V(G)$ we take an arbitrary permutation π_v of Σ' such that ϕ_v is exactly the set of labels that π_v stabilizes; note that this is possible due to $k+2 \geq 2$. We create $k'+1$ loops in v with π_v as the constraint. Then, for every edge $uv \in E(G)$ denote by $X_{uv,u}$ the set of labels from Σ that do not have an image in $\psi_{uv,u}$, and similarly denote by $X_{uv,v}$ the set of labels from Σ that do not have an image in $\psi_{uv,v}$. Let $\{\psi_{uv,u}^i\}_{i=1, \dots, k+2}$ be an arbitrary family of permutations of Σ' , such that:

- each $\psi_{uv,u}^i$ extends $\psi_{uv,u}$;
- each label $\alpha \in X_{uv,u} \cup \Gamma$ is mapped to pairwise different labels in $\psi_{uv,u}^i$ for $i = 1, 2, \dots, k+2$;
- each label $\beta \in X_{uv,v} \cup \Gamma$ is mapped to pairwise different labels in $\psi_{uv,v}^i = (\psi_{uv,u}^i)^{-1}$ for $i = 1, 2, \dots, k+2$.

Observe that as $|\Gamma| = k+2$, one can find such family $\{\psi_{uv,u}^i\}_{i=1, \dots, k+2}$ by enumerating $X_{uv,u} \cup \Gamma$ and $X_{uv,v} \cup \Gamma$ in arbitrary orders, fixing one bijection between them and shifting it cyclicly $k+1$ times. Between u and v we insert the set of $k+2$ edges $P_{uv} = \{uv^i\}_{i=1, 2, \dots, k+2}$, imposing the constraints $(\psi_{uv,u}^i, \psi_{uv,v}^i)$ on uv^i . Finally, we set $\phi'_v = \Sigma'$ for all $v \in V(G)$. This concludes the construction. We are left with a formal proof of the equivalence.

Assume first that there exists a set of edges $F \subseteq E(G)$, $|F| \leq k$, such that $G \setminus F$ admits a labeling Ψ respecting constraints in the input instance I . Let $F' = \{e^i : e \in F\}$; note that $|F'| = (k+2)|F| \leq k'$. A direct check shows that Ψ is also a correct labeling in $G' \setminus F'$, which proves that F' is a solution to the instance I' .

Now assume that there exists a set of edges $F' \subseteq E(G')$, $|F'| \leq k'$, such that $G' \setminus F'$ admits a labeling Ψ' respecting constraints in the output instance I' . Note that for each $v \in V(G)$ we have that $\Psi'(v) \in \phi_v$, as otherwise the set F' would need to contain $k'+1$ loops at v . Let $F \subseteq E(G)$ be the set of edges uv of G such that $(\Psi'(u), \Psi'(v)) \notin \psi_{uv,u}$. Clearly, F is a solution in the instance I as Ψ' is a correct labeling of $G \setminus F$. It remains to prove that $|F| \leq k$. Assume otherwise, i.e., $|F| \geq k+1$.

Consider an edge $uv \in E(G)$ such that $(\Psi'(u), \Psi'(v)) \notin \psi_{uv,u}$. We claim that $|F' \cap P_{uv}| \geq k+1$. If $\Psi'(u)$ belongs to the domain of $\psi_{uv,u}$, then all the constraints $\psi_{uv,u}^i$ map $\Psi'(u)$ to a label different than $\Psi'(v)$. Hence $P_{uv} \subseteq F'$ and the claim holds. Otherwise, $\Psi'(v)$ is mapped to $k+2$ different images in constraints $\psi_{uv,u}^i$, which means that at least $k+1$ of them must be different than $\Psi'(v)$. The corresponding edges have to be contained in F' and the claim holds in this case as well. As $|F| \geq k+1$, we have that $|F'| \geq (k+1)^2 = k'+1$, which is a contradiction. \square

We are now ready to prove Theorem G.1.

Proof of Theorem G.1. By Lemma G.2, we may consider the general problem definition, where we allow lists in vertices and partial permutations as constraints imposed on edges.

We provide a parameterized reduction from the MULTICOLORED CLIQUE problem, which is known to be W[1]-hard [17].

MULTICOLORED CLIQUE

Parameter: k

Input: An undirected graph H with vertices partitioned into k parts V_0, V_1, \dots, V_{k-1} , such that H does not contain edges connecting vertices from the same part V_i , for $i = 0, 1, \dots, k-1$.

Question: Is there a clique C in G of size k ?

Observe that by the assumption on the structure of H , the clique C has to contain exactly one vertex from each part V_i . Moreover, by adding independent vertices we can assume that each part V_i is of the same size n . In each part V_i fix an arbitrary ordering of vertices $v_0^i, v_1^i, \dots, v_{n-1}^i$.

Now, we are going to construct an instance $(G, \Sigma, k', (\phi_v)_{v \in V(G)}, (\psi_{e,v})_{e \in E(G), v \in e})$ that is a YES instance of EDGE UNIQUE LABEL COVER if and only if H contains a clique of size k . As the construction will be performed in polynomial time and $k' = k^2$, this gives the promised parameterized reduction.

We take $\Sigma = \{0, 1, 2, \dots, n\} \times \{0, 1, 2, \dots, n\}$, and let $\Lambda = \{0, 1, \dots, n-1\} \times \{0, 1, \dots, n-1\} \subseteq \Sigma$. For every part V_i we create a cycle C_i of length kn . Denote the vertices of C_i by $u_0^i, u_1^i, \dots, u_{kn-2}^i, u_{kn-1}^i$ in the order of their appearance on the cycle. For every vertex u_p^i let $\text{next}(u_p^i)$ be the next vertex on the cycle C_i , i.e., u_{p+1}^i if $p < kn-1$ and u_0^i if $p = kn-1$. Let $e(u_p^i)$ be the edge connecting u_p^i with $\text{next}(u_p^i)$.

On every edge of the cycle C_i we impose a constraint given by the permutation $\pi_0((a, b)) = (a-1, b)$, where the numbers behave cyclicly modulo $n+1$. More precisely, the constraint on the edge $e(u_p^i)$ states that the label of $\text{next}(u_p^i)$ has the first coordinate decremented by 1 modulo $n+1$ comparing to the label of u_p^i . Now, for every $i \neq j$, $0 \leq i, j < k$, we create an edge $u_{j \cdot n}^i u_{i \cdot n}^j$ with constraint given by the partial permutation $\sigma_{i,j} = \{(p, q), (q, p) \mid v_p^i v_q^j \in E(H)\}$. In other words, from the domain of the permutation $\sigma((a, b)) = (b, a)$ we remove out all the pairs that contain $n+1$ and all the pairs that correspond to nonedges between V_i and V_j . Finally, we set $\phi_v = \Lambda$ for every $v \in V(G)$ and $k' = k^2$. This concludes the construction.

Let us firstly assume that C is a clique of size k in H and let $\{v_{c_i}^i\} = V(C) \cap V_i$. We construct

- a set of edges $F = \{e(u_{j \cdot n + c_i}^i) \mid 1 \leq i, j < k\}$;
- a labeling $\Psi(u_p^i) = ((c_i - p) \bmod n, c_{q/n})$, where u_q^i is the closest next vertex on the cycle that has lower index being a multiplicity of n , i.e., $q/n = \lceil p/n \rceil \bmod n$.

Obviously, $|F| = k'$. Let us check that Ψ is a correct labeling of $G \setminus F$. Clearly, $\Psi(v) \in \Lambda = \phi_v$ for any $v \in V(G)$. Consider any edge $e(u_p^i) \notin F$. As $p \bmod n \neq c_i$, we have that $\Psi(u_p^i) = (x, y)$ for some $x > 0$ and $\Psi(\text{next}(u_p^i)) = (x-1, y)$; hence, these constraints are satisfied. Now consider any edge of the form $u_{j \cdot n}^i u_{i \cdot n}^j$ for $i \neq j$, $0 \leq i, j < k$. By the construction of Ψ we have that $\Psi(u_{j \cdot n}^i) = (c_i, c_j)$ and $\Psi(u_{i \cdot n}^j) = (c_j, c_i)$. Recall that C is a clique, so $v_{c_i}^i v_{c_j}^j \in E(H)$. Hence, (c_i, c_j) lies in the domain of $\sigma_{i,j}$ and the constraint imposed on this edge is satisfied as well.

Let us now assume that there is a set of edges $F \subseteq E(G)$, $|F| \leq k'$, such that there exists a correct labeling Ψ of $G \setminus F$. Firstly, we claim that for every n consecutive edges of every cycle C_i , F has to contain at least one of these edges. Otherwise there would be $n+1$ consecutive vertices $u_p^i, u_{p+1}^i, \dots, u_{p+n}^i$ such that edges $u_{p+i}^i u_{p+i+1}^i$ do not belong to F for $i = 0, 1, \dots, n-1$ (indices behave cyclicly). It follows that if $\Psi(u_p^i) = (\ell, d)$ for some $\ell < n$, then we would have $\Psi(u_{p+\ell+1}^i) = (n, d)$, but n is a forbidden value in a label for every vertex. As every cycle

C_i has length kn , it has to contain at least k edges from F . As $k' = k^2$, it has to contain exactly k edges from F . We can use again the claim to infer that between every two subsequent edges from F there must be exactly $n - 1$ edges not from F , as otherwise there would be n consecutive edges not belonging to F . Moreover, the same argumentation yields that the vertices of each interval on the cycle between the two subsequent edges from F have to be labeled with $(n - 1, d), (n - 2, d), \dots, (0, d)$, in this order, for some d depending on the interval, but constant within. Hence, for every cycle C_i we can find an integer $c_i \in \{0, 1, \dots, n - 1\}$, such that $F = \{e(u_{jn+c_i}^i) \mid 1 \leq i, j < k\}$ and $\Psi(u_{jn}^i) = (c_i, d_j^i)$ for all $j = 0, 1, \dots, k - 1$ and some numbers d_j^i .

We are going to prove that vertices $v_{c_i}^i$ for $i = 0, 1, \dots, k - 1$ induce a clique in H . Take parts V_i, V_j for $i \neq j$ and examine the edge $u_{j \cdot n}^i u_{i \cdot n}^j$ with constraint σ_{ij} . As $\Psi(u_{j \cdot n}^i) = (c_i, d_j^i)$, $\Psi(u_{i \cdot n}^j) = (c_j, d_i^j)$, and σ_{ij} swaps the elements of the pair, we find that $d_j^i = c_j$ and $d_i^j = c_i$. Moreover, (c_i, c_j) is in the domain of σ_{ij} if and only if $v_{c_i}^i v_{c_j}^j \in E(H)$. Therefore, $v_{c_i}^i$ and $v_{c_j}^j$ are adjacent for all $i \neq j$ and we are done. \square

H Weights

We would like to note that using our technique we can solve a more general problem, where the graph is edge-weighted (or vertex-weighted, in the vertex-deletion setting), and the goal is, instead of minimizing the cardinality of the cutset, to find a cutset of size at most k , having minimum sum of weights of the edges (or vertices) it contains. For example for the problem considered in Section 2, the formal definition is as follows.

WEIGHTED EDGE MULTIWAY CUT-UNCUT (W-E-MWCU) **Parameter:** k
Input: A graph $G = (V, E)$ together with a weight function $\omega : E \rightarrow \mathbb{R}_+$, set of terminals $T \subseteq V(G)$, an equivalence relation \mathcal{R} on the set T , and an integer k .
Question: What is the minimum weight set of edges $X \subseteq E(G)$ of cardinality at most k , such that for any $u, v \in T$, the vertices u and v belong to the same connected component of $G \setminus X$ if and only if $(u, v) \in \mathcal{R}$?

Note, that now we have to reformulate the bordered problem definition as well, because solutions to the bordered problem need to have a prescribed cardinality in order to make them comparable. Let us see it on the example of W-E-MWCU.

By $\mathbb{P}(\mathcal{J}_b)$ we define the set of all quadruples $\mathcal{P} = (X_b, E_b, \mathcal{R}_b, k_b)$, such that $X_b \subseteq T_b$, E_b is an equivalence relation on $T_b \setminus X_b$, \mathcal{R}_b is an equivalence relation on $T \cup (T_b \setminus X_b)$ such that $E_b \subseteq \mathcal{R}_b$ and $\mathcal{R}_b|_T = \mathcal{R}$, and $0 \leq k_b \leq k$. For a quadruple $\mathcal{P} = (X_b, E_b, \mathcal{R}_b, k_b)$, by $G_{\mathcal{P}}$ we denote the graph $G \cup E_b$, that is, the graph G with additional edges E_b .

We say that a set $X \subseteq V(G) \setminus T$ is a solution to $(\mathcal{J}_b, \mathcal{P})$ if $|X| \leq k_b$, $X \cap T_b = X_b$ and for any $u, v \in T \cup (T_b \setminus X_b)$, the vertices u and v are in the same connected component of the graph $G_{\mathcal{P}} \setminus X$ (i.e., we delete vertices X and add edges E_b) if and only if $(u, v) \in \mathcal{R}_b$, and the sum of weights of edges in X is minimum (comparing to all other sets X' satisfying the remaining constraints).

BORDER W-E-MWCU
Input: An W-E-MWCU instance $\mathcal{J} = (G, \omega, T, \mathcal{R}, k)$ with G being connected and a set $T_b \subseteq V(G) \setminus T$ of size at most $2k$; denote $\mathcal{J}_b = (G, T, \mathcal{R}, k, T_b)$
Output: For each $\mathcal{P} = (X_b, E_b, \mathcal{R}_b, k_b) \in \mathbb{P}(\mathcal{J}_b)$, output a $\text{sol}_{\mathcal{P}} = X_{\mathcal{P}}$ being a solution to $(\mathcal{J}_b, \mathcal{P})$ with minimum possible sum of weights of $X_{\mathcal{P}}$, or $\text{sol}_{\mathcal{P}} = \perp$ if such a solution does not exist.

Since, while finding a good separation, our algorithm does not perform any greedy choices,

we almost leave the algorithm unchanged. Similarly, the recursive understanding step in the node-deletion problems is not affected significantly by this change. However, when solving a weighted problem, we need to be more careful in the final, high connectivity phase, as the existence of weights limits our possibilities of being greedy. In the following paragraphs we shortly argue that the high connectivity phases of the algorithms presented in this paper can be adjusted to the weighted variants without greater effort.

NODE MULTIWAY CUT-UNCUT. The simplicity of the high connectivity phase of the N-MWCU algorithm allows us to solve the weighted variant with almost no changes. Recall that in this phase we first guess a set S that is (i) disjoint with the solution Z we are looking for, (ii) covers all nonterminal vertices of small connected components of $G \setminus Z$. Then we argue that any inclusion-wise minimal solution chooses at most one equivalence relation of \mathcal{R}_b to be the set of terminals contained in the big connected component of $G \setminus Z$, and takes as the solution the neighbourhood of all connected components of $G[S \cup T]$ that contain a terminal not contained in the selected equivalence class. The same argumentation holds in the case of nonnegative weights; note that in the border problem we require $|X| \leq k_b$ (instead of maybe more natural $|X| = k_b$), thus we may consider only inclusion-wise minimal solutions.

STEINER CUT. In the case of the STEINER CUT problem, we need to slightly change the final dynamic programming routine. Recall that in the high connectivity phase for this problem we first guess a set of edges S that is (i) disjoint with the solution Z we are looking for, (ii) contains a spanning tree of each small connected component of $G \setminus Z$, (iii) contains a large spanning tree with an endpoint of an edge of the solution Z , for each such endpoint contained in the large connected component of $G \setminus Z$. Then we obtain a graph H by contracting the edges of S and identifying the images of the large trees of S (assumed in point (iii)) into the core vertex b . For each connected component B'_i of $H \setminus b$ we have two choices: either we delete all edges, or no edges from $B'_i \cup \{b\}$. The choices between different components B'_i are independent, and we find the optimal solution via a simple dynamic programming routine. In the weighted case we need to add to the dynamic programming table one more dimension responsible for storing the cardinality of the constructed cutset, and the value in the table T is the minimum weight of a cutset of the prescribed cardinality.

NODE UNIQUE LABEL COVER. In the case of the NODE UNIQUE LABEL COVER problem, the high connectivity phase remains almost unchanged; however, we need to argue that all greedy steps used in this part of the algorithm are justified also in the weighted case. As in the case of the other problems, we start with guessing an interrogating set that is (i) disjoint with the solution Z we are looking for, (ii) contains all vertices of all small connected components of $G \setminus Z$, (iii) contains a large connected set adjacent to each vertex of Z that is adjacent to the large connected component of $G \setminus Z$. The algorithm performs now two simple greedy steps: it checks whether $Z = \emptyset$ is a solution, and looks for not forsaken vertices without neighbours in S . Both steps can be easily justified in the weighted case, as we assume nonnegative weights and we require only $|X| \leq k_b$ in the border problem definition. The crucial observation — that there are only at most s reasonable labelings of the big stains (big connected components) of S — does not interfere with weights. In the final bounded search tree algorithm we argue that there is a limited number of vertices, out of which we need to delete at least one (the Neighbourhood Branching Rule) or that there are only limited number of ways a small stain can be handled (the Small Stains Rule). Both argumentations are oblivious to weights; note that this is also true in the second part of Lemma D.18, where we argue about a greedy choice of a labeling in case when the chosen labeling of the big stains can be consistently extended to the closed neighbourhood of a connected component of $G \setminus (X_0 \cup N_G[S^{\text{big}}])$.

References

- [1] Amit Agarwal, Noga Alon, and Moses Charikar. Improved approximation for directed cut problems. In *Proc. of STOC'07*, pages 671–680, 2007.
- [2] Noga Alon, Raphy Yuster, and Uri Zwick. Color-coding: a new method for finding simple paths, cycles and other small subgraphs within large graphs. In *Proc. of STOC'94*, pages 326–335, 1994.
- [3] Sanjeev Arora, Boaz Barak, and David Steurer. Subexponential algorithms for unique games and related problems. In *Proc. of FOCS'10*, pages 563–572, 2010.
- [4] Vineet Bafna, Piotr Berman, and Toshihiro Fujito. A 2-approximation algorithm for the undirected feedback vertex set problem. *SIAM J. Discrete Math.*, 12(3):289–297, 1999.
- [5] Hans L. Bodlaender, Fedor V. Fomin, Daniel Lokshtanov, Eelko Penninkx, Saket Saurabh, and Dimitrios M. Thilikos. (meta) kernelization. In *Proc. of FOCS'09*, pages 629–638, 2009.
- [6] Nicolas Bousquet, Jean Daligault, and Stéphan Thomassé. Multicut is FPT. In *Proc. of STOC'11*, pages 459–468, 2011.
- [7] Moses Charikar, Konstantin Makarychev, and Yury Makarychev. Near-optimal algorithms for unique games. In *Proc. of STOC'06*, pages 205–214, 2006.
- [8] Chandra Chekuri, Sudipto Guha, and Joseph Naor. The Steiner k-Cut Problem. *SIAM J. Discrete Math.*, 20(1):261–271, 2006.
- [9] Jianer Chen, Yang Liu, and Songjian Lu. An improved parameterized algorithm for the minimum node multiway cut problem. *Algorithmica*, 55(1):1–13, 2009.
- [10] Jianer Chen, Yang Liu, Songjian Lu, Barry O’Sullivan, and Igor Razgon. A fixed-parameter algorithm for the directed feedback vertex set problem. In *Proc. of STOC'08*, pages 177–186, 2008.
- [11] Rajesh Hemant Chitnis, MohammadTaghi Hajiaghayi, and Dániel Marx. Fixed-parameter tractability of directed multiway cut parameterized by the size of the cutset. In *Proc. of SODA'12*, pages 1713–1725, 2012.
- [12] Marek Cygan, Marcin Pilipczuk, and Michał Pilipczuk. On group feedback vertex set parameterized by the size of the cutset. *CoRR*, abs/1112.6255, 2011.
- [13] Marek Cygan, Marcin Pilipczuk, Michał Pilipczuk, and Jakub Onufry Wojtaszczyk. Subset feedback vertex set is fixed-parameter tractable. In *Proc. of ICALP'11*, pages 449–461, 2011.
- [14] R. G. Downey and M. R. Fellows. *Parameterized Complexity*. Springer, 1999.
- [15] Guy Even, Joseph Naor, Baruch Schieber, and Madhu Sudan. Approximating minimum feedback sets and multi-cuts in directed graphs. In *Proc. of IPCO'95*, pages 14–28, 1995.
- [16] Guy Even, Joseph (Seffi) Naor, and Leonid Zosin. An 8-approximation algorithm for the subset feedback vertex set problem. In *Proc. of FOCS'99*, pages 310–319, 1996.
- [17] Micheal R. Fellows, Danny Hermelin, Frances A. Rosamond, and Stéphane Vialette. On the parameterized complexity of multiple-interval graph problems. *Theor. Comput. Sci.*, 410:53–61, 2009.

- [18] J. Flum and M. Grohe. *Parameterized Complexity Theory*. Texts in Theoretical Computer Science. An EATCS Series. Springer, 1 edition, March 2006.
- [19] Fedor V. Fomin, Daniel Lokshtanov, Saket Saurabh, and Dimitrios M. Thilikos. Linear kernels for (connected) dominating set on h -minor-free graphs. In *Proc. of SODA '12*, pages 82–93, 2012.
- [20] Naveen Garg, Vijay V. Vazirani, and Mihalis Yannakakis. Multiway cuts in directed and node weighted graphs. In *Proc. of ICALP'94*, pages 487–498, 1994.
- [21] Naveen Garg, Vijay V. Vazirani, and Mihalis Yannakakis. Approximate max-flow min-(multi)cut theorems and their applications. *SIAM J. Comput.*, 25(2):235–251, 1996.
- [22] F. Granot and R. Hassin. Multi-terminal maximum flows in node capacitated networks. *Disc. Appl. Math.*, 13:157–163, 1986.
- [23] Sylvain Guillemot. FPT algorithms for path-transversal and cycle-transversal problems. *Discrete Optimization*, 8(1):61–71, 2011.
- [24] Sylvain Guillemot. Parameterized complexity and approximability of the longest compatible sequence problem. *Discrete Optimization*, 8(1):50–60, 2011.
- [25] David R. Karger. Minimum cuts in near-linear time. *J. ACM*, 47(1):46–76, 2000.
- [26] David R. Karger, Philip N. Klein, Clifford Stein, Mikkel Thorup, and Neal E. Young. Rounding algorithms for a geometric embedding of minimum multiway cut. In *Proc. of STOC'99*, pages 668–678, 1999.
- [27] Ken-ichi Kawarabayashi and Mikkel Thorup. The minimum k -way cut of bounded size is fixed-parameter tractable. In *Proc. of FOCS'11*, pages 160–169, 2011.
- [28] Subhash Khot. On the power of unique 2-prover 1-round games. In *Proc. of STOC'02*, pages 767–775, 2002.
- [29] Subhash Khot. On the unique games conjecture (invited survey). In *IEEE Conference on Computational Complexity*, pages 99–121, 2010.
- [30] Stefan Kratsch, Marcin Pilipczuk, Michał Pilipczuk, and Magnus Wahlström. Fixed-parameter tractability of multicut in directed acyclic graphs. *CoRR*, abs/1202.5749, 2012.
- [31] Daniel Lokshtanov and Dániel Marx. Clustering with local restrictions. In *Proc. of ICALP'11*, pages 785–797, 2011.
- [32] Dániel Marx. Parameterized graph separation problems. *Theor. Comput. Sci.*, 351(3):394–406, 2006.
- [33] Dániel Marx, Barry O'Sullivan, and Igor Razgon. Finding small separators in linear time via treewidth reduction. *CoRR*, abs/1110.4765, 2012.
- [34] Dániel Marx and Igor Razgon. Fixed-parameter tractability of multicut parameterized by the size of the cutset. In *Proc. of STOC'11*, pages 469–478, 2011.
- [35] Hiroshi Nagamochi and Toshihide Ibaraki. A Linear-Time Algorithm for Finding a Sparse k -Connected Spanning Subgraph of a k -Connected Graph. *Algorithmica*, 7(5&6):583–596, 1992.

- [36] Joseph Naor and Yuval Rabani. Tree packing and approximating k-cuts. In *Proc. of SODA '01*, pages 26–27, 2001.
- [37] Moni Naor, Leonard J. Schulman, and Aravind Srinivasan. Splitters and near-optimal derandomization. In *Proc. of FOCS'95*, pages 182–191, 1995.
- [38] R. Ravi and Amitabh Sinha II. Approximating k-cuts via network strength. In *Proc. of SODA '02*, pages 621–622, 2002.
- [39] Igor Razgon. Large isolating cuts shrink the multiway cut. *CoRR*, abs/1104.5361, 2011.
- [40] Igor Razgon and Barry O'Sullivan. Almost 2-SAT is fixed-parameter tractable. *J. Comput. Syst. Sci.*, 75(8):435–450, 2009.
- [41] Paul D. Seymour. Packing directed circuits fractionally. *Combinatorica*, 15(2):281–288, 1995.