RATIONALITY OF ADMISSIBLE AFFINE VERTEX ALGEBRAS IN THE CATEGORY $\ensuremath{\mathcal{O}}$

TOMOYUKI ARAKAWA

ABSTRACT. We study the vertex algebras associated with modular invariant representations of affine Kac-Moody algebras at fractional levels, whose simple highest weight modules are classified by Joseph's characteristic varieties. We show that an irreducible highest weight representation of a non-twisted affine Kac-Moody algebra at an admissible level k is a module over the associated simple affine vertex algebra if and only if it is an admissible representation whose integral root system is isomorphic to that of the vertex algebra itself. This in particular proves the conjecture of Adamović and Milas [AM] on the rationality of admissible affine vertex algebras in the category \mathcal{O} .

1. INTRODUCTION

Let \mathfrak{g} be a complex simple Lie algebra, $\hat{\mathfrak{g}}$ the non-twisted affine Kac-Moody algebra associated with \mathfrak{g} , $V^k(\mathfrak{g})$ the universal affine vertex algebra associated with \mathfrak{g} at a non-critical level k, $L(k\Lambda_0)$ the unique simple quotient of $V^k(\mathfrak{g})$. The simple affine vertex algebra $L(k\Lambda_0)$ is called *admissible* if it is isomorphic to an admissible representation [KW2] as a $\hat{\mathfrak{g}}$ -module. The purpose of this article is to classify simple modules over admissible affine vertex algebras.

By a well-known result of Zhu [Zhu], there is a one-to-one correspondence between positively graded simple modules over a graded vertex algebra V and simple A(V)-modules, where A(V) is Zhu's algebra of V. In the case that V is an affine vertex algebra $L(k\Lambda_0)$, we have

(1)
$$A(L(k\Lambda_0)) \cong U(\mathfrak{g})/I_k$$

for some two sided-ideal I_k of the universal enveloping algebra $U(\mathfrak{g})$ of \mathfrak{g} . Since a simple $U(\mathfrak{g})$ -module M is an $A(L(k\Lambda_0))$ -module if and only if the annihilating ideal of M in $U(\mathfrak{g})$ contains I_k , our problem amounts to classify the primitive ideals of $U(\mathfrak{g})$ containing I_k . Because any primitive ideal of $U(\mathfrak{g})$ is the annihilating ideal of a highest weight representation of \mathfrak{g} [Duf], it suffices to classify simple highest weight representations of $A(L(k\Lambda_0))$, or equivalently, to classify simple $L(k\Lambda_0)$ -modules in the category \mathcal{O} of $\hat{\mathfrak{g}}$.

Let $L(\lambda)$ be the irreducible highest weight representation of $\hat{\mathfrak{g}}$ with highest weight λ .

Main Theorem. Let k be an admissible number¹, λ a weight of $\hat{\mathfrak{g}}$ of level k. Then $L(\lambda)$ is a module over $L(k\Lambda_0)$ if and only if it is an admissible representation whose

²⁰⁰⁰ Mathematics Subject Classification. 17B69, 17B67, 17B08, 17B55.

This work is partially supported by the JSPS Grant-in-Aid for Scientific Research (B) No. 20340007 and JSPS Grant-in-Aid for Challenging Exploratory Research No. 23654006.

¹A complex number k is called admissible if $L(k\Lambda_0)$ is an admissible representation.

integral root system is isomorphic to that of $L(k\Lambda_0)$. In particular any $\hat{\mathfrak{g}}$ -module from the category \mathcal{O} is an $L(k\Lambda_0)$ -module if and only if it is a direct sum of such admissible representations of $\hat{\mathfrak{g}}$ of level k.

The second statement of Main Theorem was conjectured by Adamović and Milas [AM, Conjecture 3.5.7]. It has been proved in some special cases: for type $C_{\ell}^{(1)}$ admissible half integer levels by Adamović [Ada]; for $\widehat{\mathfrak{sl}}_2$ by Adamović and Milas [AM], Dong, Li and Mason [DLM] and Feigin and Malikov [FeM]; for some cases in type $A_{\ell}^{(1)}$, $B_{\ell}^{(1)}$ admissible half integer levels by Perše [Per1, Per2]; for a few cases in type $G_2^{(1)}$ admissible one-third integer levels by Axtell and Lee [AL]. These works are based on the explicit computation of the singular vector of $V^k(\mathfrak{g})$ which generates the maximal ideal. Our method in this article is completely different.

Let us explain the outline of the proof of Main Theorem briefly. By (1) simple highest weight $L(k\Lambda_0)$ -modules are classified by Joseph's *characteristic variety* [Jos] $\mathcal{V}(I_k)$ of I_k , which is a Zariski closed subset of the dual \mathfrak{h}^* of the Cartan subalgebra \mathfrak{h} of \mathfrak{g} (Proposition 2.5). We deduce the "if" part of Main Theorem from a result of Frenkel and Malikov [FrM, MF] which states that every *G*-integrable admissible representation at level *k* is a module over $L(k\Lambda_0)$, together with an affine analogue of the Duflo-Joseph Lemma (Lemma 2.8) for characteristic varieties. We prove the "only if" part of Main Theorem by reducing to the $\widehat{\mathfrak{sl}}_2$ -cases [AM] using the *semiinfinite restriction functor* $H^{\frac{\infty}{2}+i}(\mathfrak{m}^{(i)}[t,t^{-1}],?)$ studied in [A3], where $\mathfrak{m}^{(i)}$ is the nilradical of a minimal parabolic subalgebra of \mathfrak{g} , see §4.3 for the details.

It should be mentioned that there is another variety naturally associated with $A(L(k\Lambda_0))$, that is, the zero set $V(\operatorname{gr} I_k)$ of the associated graded ideal $\operatorname{gr} I_k$ of $\mathbb{C}[\mathfrak{g}^*]$. We have a surjection

(2)
$$R_{L(k\Lambda_0)} \twoheadrightarrow \operatorname{gr} A(L(k\Lambda_0)) = \mathbb{C}[\mathfrak{g}^*]/\operatorname{gr} I_k$$

of Poisson algebras, where $R_{L(k\Lambda_0)}$ is Zhu's C_2 -algebra of $L(k\Lambda_0)$, see [ALY, Proposition 3.3]. Hence $V(\text{gr } I_k)$ is contained in the associated variety [A1] $X_{L(k\Lambda_0)} = \operatorname{Specm} R_{L(k\Lambda_0)}$. Although (2) is not an isomorphism in general, in a subsequent paper [A4] we prove the following:

Theorem 1.1. Let k be an admissible number, with $k + h^{\vee} = p/q$, $p, q \in \mathbb{N}$, (p,q) = 1. Then (2) induces the isomorphism of varieties

$$V(\operatorname{gr} I_k) \cong X_{L(k\Lambda_0)}.$$

Namely [A2], we have

$$V(\operatorname{gr} I_k) \cong \begin{cases} \{x \in \mathfrak{g}; (\operatorname{ad} x)^{2q} = 0\} & \text{if } (r^{\vee}, q) = 1, \\ \{x \in \mathfrak{g}; \pi_{\theta_s}(x)^{2q/r^{\vee}} = 0\} & \text{if } (r^{\vee}, q) = r^{\vee}, \end{cases}$$

which is an irreducible G-invariant subvariety of $\mathfrak{g}^* \cong \mathfrak{g}$. Here r^{\vee} is the lacing number of \mathfrak{g} , θ_s is the highest short root of \mathfrak{g} and π_{θ_s} is the irreducible finite-dimensional representation of \mathfrak{g} with highest weight θ_s .

Although they themselves are not rational in the usual sense² it has been conjectured [FKW, KW3] that admissible affine vertex algebras produce rational W-algebras in many cases by the method of the (generalized) quantum Drinfeld-Sokolov reduction. By applying the result in this article in a subsequent paper [A4] we prove the rationality of all the minimal series principal W-algebras [FKW].

This paper is organized as follows. In §2 we establish the relationship between Joseph's characteristic varieties and the representation theory of affine vertex algebras and prove an affine analogue of the Duflo-Joseph Lemma (Lemma 2.8). In §3 we prove the "if part" of Main Theorem. In §4 we prove the "only if part" of Main Theorem. In appendix we apply Main Theorem to prove a semi-infinite analogue of Kostant's generalized Borel-Weil-Bott Theorem [Kos] for admissible representations announced in [A3].

Acknowledgments. Some part of this work was done while the author was visiting Weizmann Institute, Israel, in May 2011, Emmy Noether Center in Erlangen, Germany in June 2011, Isaac Newton Institute for Mathematical Sciences, UK, in 2011, The University of Manchester, University of Birmingham, The University of Edinburgh, Lancaster University, York University, UK, in November 2011, Academia Sinica, Taiwan, in December 2011. He is grateful to those institutes for their hospitality.

2. Affine vertex algebras and Joseph's characteristic varieties

Let \mathfrak{g} be a complex simple Lie algebra of rank l. Fix a triangular decomposition

$$\mathfrak{g} = \mathfrak{n}_- \oplus \mathfrak{h} \oplus \mathfrak{n}_+,$$

with a Cartan subalgebra \mathfrak{h} of \mathfrak{g} . We will often identify \mathfrak{h} with \mathfrak{h}^* using the normalized invariant bilinear form (|) of \mathfrak{g} . Let Δ , Δ_+ , Δ_- , $\Pi = \{\alpha_1, \ldots, \alpha_l\}$ be the sets of the roots, positive roots, negative roots, and simple roots of \mathfrak{g} , respectively. Also, let θ be the highest root of \mathfrak{g} , θ_s the highest short root, ρ the half sum of the positive roots, $W = \langle s_\alpha; \alpha \in \Delta \rangle \subset \operatorname{Aut} \mathfrak{h}^*$ the Weyl group of \mathfrak{g} . Here $s_\alpha(\lambda) = \lambda - \langle \lambda, \alpha^{\vee} \rangle \alpha$, $\alpha^{\vee} = 2\alpha/(\alpha|\alpha)$. We set $s_i = s_{\alpha_i}$ for $i = 1, \ldots, l$. Let $w \circ \lambda = w(\lambda + \rho) - \rho$ for $w \in W, \lambda \in \mathfrak{h}^*$.

Denote by $\mathcal{O}^{\mathfrak{g}}$ the Bernstein-Gelfand-Gelfand category \mathcal{O} of \mathfrak{g} , by $L^{\mathfrak{g}}_{\lambda}$ the irreducible highest weight representation of \mathfrak{g} with highest weight $\lambda \in \mathfrak{h}^*$.

Set $U(\mathfrak{g})^{\mathfrak{h}} := \{ u \in U(\mathfrak{g}); [h, u] = 0 \text{ for all } h \in \mathfrak{h} \}$, and let

$$\Upsilon: U(\mathfrak{g})^{\mathfrak{g}} \to U(\mathfrak{h})$$

be the restriction of the projection $U(\mathfrak{g}) = U(\mathfrak{h}) \oplus (\mathfrak{n}_{-}U(\mathfrak{g}) + U(\mathfrak{g})\mathfrak{n}_{+}) \to U(\mathfrak{h})$ to $U(\mathfrak{g})^{\mathfrak{h}}$. One knows that Υ is an algebra homomorphism.

For a two-sided ideal I of $U(\mathfrak{g})$, the *characteristic variety* [Jos] (without the ρ -shift) of I is defined as

$$\mathcal{V}(I) = \{\lambda \in \mathfrak{h}^*; p(\lambda) = 0 \text{ for all } p \in \Upsilon(I^{\mathfrak{h}})\},\$$

where $I^{\mathfrak{h}} = I \cap U(\mathfrak{g})^{\mathfrak{h}}$.

Lemma 2.1. For $\lambda \in \mathfrak{h}^*$, $\lambda \in \mathcal{V}(I)$ if and only if $IL_{\lambda}^{\mathfrak{g}} = 0$.

²However Main Theorem implies that admissible affine vertex algebras are rational in the sense of [DLM]. Also, the fact [A2] that $X_{L(k\Lambda_0)}$ is contained in the nilpotent cone of \mathfrak{g} implies that they are C_2 -cofinite in the sense of [DLM].

Proof. The "if part" is obvious. To see the converse, suppose that $IL_{\lambda}^{\mathfrak{g}} \neq 0$. Then $IL_{\lambda}^{\mathfrak{g}} = L_{\lambda}^{\mathfrak{g}}$ since $L_{\lambda}^{\mathfrak{g}}$ is simple. It follows that there exists $a \in I^{\mathfrak{h}}$ such that $v_{\lambda} = av_{\lambda} = \Upsilon(a)(\lambda)v_{\lambda}$, where v_{λ} is the highest weight vector of $L_{\lambda}^{\mathfrak{g}}$. Hence $\lambda \notin \mathcal{V}(I)$. \Box

By Lemma 2.1, the characteristic variety $\mathcal{V}(I)$ of I classifies simple $U(\mathfrak{g})/I$ -module in the category $\mathcal{O}^{\mathfrak{g}}$.

The following fact is useful for us.

Lemma 2.2 ([Duf], [Jos, Lemma 2]). Let I be a two-sided ideal of $U(\mathfrak{g}), \lambda \in \mathcal{V}(I)$. Suppose that $\langle \lambda + \rho, \alpha_i^{\vee} \rangle \notin \mathbb{N}$ for $\alpha_i \in \Pi$. Then $s_i \circ \lambda \in \mathcal{V}(I)$.

Let

$$\widehat{\mathfrak{g}} = \mathfrak{g}[t, t^{-1}] \oplus \mathbb{C}K$$

be the affine Kac-Moody algebra associated with \mathfrak{g} as in Introduction. The commutation relations of $\widehat{\mathfrak{g}}$ are given by

$$[xt^{m}, yt^{n}] = [x, y]t^{m+n} + m(x|y)\delta_{m+n,0}K, \quad [K, \hat{g}] = 0$$

for $x, y \in \mathfrak{g}, m, n \in \mathbb{Z}$. Let

$$\widehat{\mathfrak{g}} = \widehat{\mathfrak{n}}_{-} \oplus \widehat{\mathfrak{h}} \oplus \widehat{\mathfrak{n}}_{+}$$

be the triangular decomposition of $\hat{\mathfrak{g}}$, where $\hat{\mathfrak{n}}_{-} = \mathfrak{n}_{-} \oplus \mathfrak{g}[t^{-1}]t^{-1}$, $\hat{\mathfrak{h}} = \mathfrak{h} \oplus \mathbb{C}K$, $\hat{\mathfrak{n}}_{+} = \mathfrak{n}_{+} \oplus \mathfrak{g}[t]t$. Let $\hat{\mathfrak{h}}^{*} = \mathfrak{h}^{*} \oplus \mathbb{C}\Lambda_{0}$ be the dual of $\hat{\mathfrak{h}}$, where $\Lambda_{0}(K) = 1$, $\Lambda_{0}(\mathfrak{h}) = 0$. For $\lambda \in \hat{\mathfrak{h}}^{*}$, denote by $\bar{\lambda} \in \mathfrak{h}^{*}$ the restriction of λ to \mathfrak{h} .

Let $\widehat{\Delta}^{re}$ be the set of real roots of $\widehat{\mathfrak{g}}$ in the dual $\widetilde{\mathfrak{h}}^* = \widehat{\mathfrak{h}}^* \oplus \mathbb{C}\delta$ of the extended Cartan subalgebra $\widetilde{\mathfrak{h}} = \widehat{\mathfrak{h}} \oplus \mathbb{C}D$, $\widehat{\Delta}^{re}_+$ the set of positive real roots, $\widehat{\Pi} = \{\alpha_0, \alpha_1, \ldots, \alpha_l\}$ the the set of simple roots of $\widehat{\mathfrak{g}}$, where $\alpha_0 = -\theta + \delta$.

Let $\widehat{W} = W \ltimes Q^{\vee}$, the Weyl group of $\widehat{\mathfrak{g}}$, which is generated by s_0, s_1, \ldots, s_l . Here $Q^{\vee} = \sum_{\alpha \in \Delta} \mathbb{Z} \alpha^{\vee}$, the coroot lattice of \mathfrak{g} , and s_0 is the reflection corresponding to α_0 . Let $\widetilde{W} = W \ltimes P^{\vee}$, the extended Weyl group of $\widehat{\mathfrak{g}}$, where P^{\vee} is the coweight lattice of \mathfrak{g} . We write t_{λ} for the element of \widehat{W} corresponding to $\lambda \in P^{\vee}$. It holds that

$$\widetilde{W} = \widetilde{W}_+ \ltimes \widehat{W},$$

where \widetilde{W}_{+} is the subgroup of \widetilde{W} consisting of elements which fix the set $\widehat{\Pi}$. Write $\theta = \sum_{i=1}^{l} a_i \alpha_i$ and set $J = \{i \in \{0, 1, \ldots, l\}; a_i = 1\}$. The group \widetilde{W}_{+} is described as

(3)
$$\widetilde{W}_{+} = \{t_{\overline{\Lambda}_{i}} w_{j}; j \in J\},\$$

where $\bar{\Lambda}_j$ is the *j*-th fundamental weight of \mathfrak{g} and w_j is the unique element of Wwhich fixes the set $\{\alpha_1, \ldots, \alpha_l, -\theta\}$ and $w_j(-\theta) = \alpha_j$. We have the isomorphism $\widetilde{W}_+ \cong P^{\vee}/Q^{\vee}, t_{\bar{\Lambda}_j} w_j \mapsto \bar{\Lambda}_j + Q^{\vee}.$

Let $k \in \mathbb{C}$. Denote by $U_k(\hat{\mathfrak{g}})$ the quotient of the universal enveloping algebra $U(\hat{\mathfrak{g}})$ by the ideal generated by K - k id, and let

$$\widetilde{U}_k(\widehat{\mathfrak{g}}) := \lim_{\stackrel{\leftarrow}{\underset{N}{\longleftarrow}}} U_k(\mathfrak{g})/(U_k(\mathfrak{g})\mathfrak{g}[t]t^N),$$

the completed universal enveloping algebra of $\hat{\mathfrak{g}}$ ([FBZ, 4.2.1]) at level k.

A $\hat{\mathfrak{g}}$ -module M is said to be of level k if K acts as the scalar k. It is called smooth if $(xt^n)m = 0$ for any $x \in \hat{\mathfrak{g}}, m \in M$ and a sufficiently large n. A smooth $\hat{\mathfrak{g}}$ -module M of level k is naturally considered as a continuous $\widetilde{U}_k(\hat{\mathfrak{g}})$ -module.

Let $V^k(\mathfrak{g})$ be the universal affine vertex algebra associated with \mathfrak{g} at level k as in Introduction. By definition,

$$V^k(\mathfrak{g}) = U(\widehat{\mathfrak{g}}) \otimes_{U(\mathfrak{g}[t] \oplus \mathbb{C}K)} \mathbb{C}_k$$

as a $\widehat{\mathfrak{g}}$ -module, where \mathbb{C}_k is the one-dimensional representation of $\mathfrak{g}[t] \oplus \mathbb{C}K$ on which $\mathfrak{g}[t]$ acts trivially and K acts as a multiplication by k. Note that by the PBW theorem we have $V^k(\mathfrak{g}) \cong U(\mathfrak{g}[t^{-1}]t^{-1})$ as vectors spaces. Define the linear map

$$\widetilde{Y}: V^k(\mathfrak{g}) \to \widetilde{U}_k(\widehat{\mathfrak{g}})[[z, z^{-1}], \quad a \mapsto \widetilde{Y}(a, z) = \sum_{n \in \mathbb{Z}} a_{(n)} z^{-n-1}$$

by

$$\tilde{Y}((x_1t^{-n_1-1})\dots(x_rt^{-n_r-1})\mathbf{1},z) = \frac{1}{n_1!\dots n_r!} : \partial_z^{n_1}x_1(z)\dots\partial_z^{n_r}x_r(z):,$$

where $\mathbf{1} = 1 \otimes 1$, $x(z) = \sum_{n \in \mathbb{Z}} (xt^n) z^{-n-1}$ for $x \in \mathfrak{g}$ and : : denotes the normally ordered product (see [Kac, 3.1]). The space $V^k(\mathfrak{g})$ is equipped with the vertex algebra structure whose state-field correspondence $Y: V^k(\mathfrak{g}) \to (\operatorname{End} V^k(\mathfrak{g}))[[z, z^{-1}]]$ is given by the composition of \tilde{Y} with the action map $\widetilde{U}_k(\hat{\mathfrak{g}}) \to \operatorname{End} V^k(\mathfrak{g})$, see [Kac, FBZ] for the details. We have

(4)
$$[a_{(m)}, b_{(n)}] = \sum_{i \ge 0} \binom{m}{i} (a_{(i)}b)_{(m+n-i)},$$

(5)
$$(a_{(m)}b)_{(n)} = \sum_{i\geq 0} (-1)^i \binom{m}{i} (a_{(m-i)}b_{(n+i)} - (-1)^r b_{(m+n-i)}a_{(i)})$$

for $a, b \in V^k(\mathfrak{g}), m, n \in \mathbb{Z}$, in $\widetilde{U}_k(\widehat{\mathfrak{g}})$ by [FBZ, Lemma 4.3.2]. For a smooth $\widehat{\mathfrak{g}}$ -module of level k, let $Y^M : V^k(\mathfrak{g}) \to (\operatorname{End} M)[[z, z^{-1}]]$ be the composition of \tilde{Y} with the action map $\widetilde{U}_k(\hat{\mathfrak{g}}) \to \operatorname{End} M$. This gives $M \neq V^k(\mathfrak{g})$ module structure. Conversely any $V^k(\mathfrak{g})$ -module can be considered as a smooth $\hat{\mathfrak{g}}$ -module of level k. Hence a $V^k(\mathfrak{g})$ -module is the same as a smooth $\hat{\mathfrak{g}}$ -module of level k.

We assume that the level k is non-critical, that is, $k \neq -h^{\vee}$, where h^{\vee} is the dual Coxeter number of \mathfrak{g} . The vertex algebra $V^k(\mathfrak{g})$ is conformal by the Sugawara construction; let

$$L(z) = \sum_{n \in \mathbb{Z}} L_n z^{-n-2} = \tilde{Y}(\omega, z),$$

where $\omega = \frac{1}{2(k+h^{\vee})} \sum_{i=1}^{\dim \mathfrak{g}} x_i t^{-1} x^i t^{-1} \mathbf{1}, \{x_i\}$ is a basis of $\mathfrak{g}, \{x^i\}$ is the dual basis, and $\mathbf{1} = 1 \otimes 1 \in V^k(\mathfrak{g})$. We have

$$[L_m, L_n] = (m-n)L_{m+n} + \frac{m^3 - m}{12} \delta_{n+m,0} \frac{k \dim \mathfrak{g}}{k+h^{\vee}},$$
$$[L_m, xt^n] = -nxt^{n-m} \quad \text{for } x \in \mathfrak{g}, \ n \in \mathbb{Z}.$$

For a smooth $\widehat{\mathfrak{g}}$ -module M of level k and $d \in \mathbb{C}$, set

$$M_d = \{ m \in M; (L_0 - d)^r m = 0 \text{ for } r \gg 0 \}.$$

Note that each M_d is a \mathfrak{g} -submodule of M. We say M is graded if $M = \bigoplus_{d \in \mathbb{C}} M_d$; positively graded if there exists $d_{top} \in \mathbb{C}$ such that $M_{d_{top}} \neq 0$ and $M = \bigoplus_{n \in \mathbb{Z}_{\geq 0}} M_{d_{top}+n}$. If this is the case we often write M_{top} for $M_{d_{top}}$.

The module $V^k(\mathfrak{g})$ is positively graded:

$$V^k(\mathfrak{g}) = \bigoplus_{\Delta=0}^{\infty} V^k(\mathfrak{g})_{\Delta}.$$

For a homogeneous element $a \in V^k(\mathfrak{g})$, the L_0 -eigenvalue of a is called the *conformal* weight of a and denoted by Δ_a .

We denote by $\mathcal{L}(V^k(\mathfrak{g}))$ Borcherds' Lie algebra associated with $V^k(\mathfrak{g})$:

$$\mathcal{L}(V^{k}(\mathfrak{g})) = V^{k}(\mathfrak{g}) \otimes \mathbb{C}[t, t^{-1}]/(L_{-1} \otimes 1 + 1 \oplus \frac{d}{dt})(V^{k}(\mathfrak{g}) \otimes \mathbb{C}[t, t^{-1}]).$$

The Lie bracket of $\mathcal{L}(V^k(\mathfrak{g}))$ is given by

$$[a_{\{m\}}, b_{\{n\}}] = \sum_{i \ge 0} \binom{m}{i} (a_{(i)}b)_{\{m+n-i\}}$$

for $a, b \in V$, $m, n \in \mathbb{Z}$, where $a_{\{n\}}$ denotes the image of $a \otimes t^n$ in $\mathcal{L}(V^k(\mathfrak{g}))$. The Lie algebra $\mathcal{L}(V^k(\mathfrak{g}))$ is graded:

$$\mathcal{L}(V^k(\mathfrak{g})) = \bigoplus_{\Delta} \mathcal{L}(V^k(\mathfrak{g}))_{\Delta},$$

where deg $a_{\{m\}} = m - \Delta_a + 1$ for a homogeneous vector $a \in V^k(\mathfrak{g})$.

Note that $(xt^{-1}\mathbf{1})_{\{m\}}$, $x \in \mathfrak{g}$, $m \in \mathbb{Z}$, generates the Kac-Moody algebra $\hat{\mathfrak{g}}$ at level k, and hence, there is an adjoint action of $\hat{\mathfrak{g}}$ on $\mathcal{L}(V^k(\mathfrak{g}))$.

By (4), there is a natural Lie algebra homomorphism

$$\mathcal{L}(V^k(\mathfrak{g})) \to U_k(\widehat{\mathfrak{g}}), \quad a_{\{m\}} \mapsto a_{(m)},$$

which is known to be injective ([FBZ, 4.2.7]). Below we consider $\mathcal{L}(V^k(\mathfrak{g}))$ as a graded Lie subalgebra of $\widetilde{U}_k(\widehat{\mathfrak{g}})$ and write $a_{(m)}$ for $a_{\{m\}}$. Clearly, $\mathcal{L}(V^k(\mathfrak{g}))$ is a ad $\widehat{\mathfrak{g}}$ -submodule of $\widetilde{U}_k(\widehat{\mathfrak{g}})$.

For a Z-graded vertex algebra V, let A(V) be Zhu's algebra of V. By [Zhu, Theorem 3.1.1], one knows that $A(V^k(\mathfrak{g})) \xrightarrow{\sim} U(\mathfrak{g})$ as algebras. This isomorphism may be constructed as follows: Consider the decomposition $U_k(\widehat{\mathfrak{g}}) = U(\mathfrak{g}) \oplus (\mathfrak{g}[t^{-1}]t^{-1}U_k(\widehat{\mathfrak{g}}) + U_k(\widehat{\mathfrak{g}})\mathfrak{g}[t]t)$. This induces the decomposition

(6)
$$\widetilde{U}_k(\widehat{\mathfrak{g}}) = U(\mathfrak{g}) \oplus \overline{(\mathfrak{g}[t^{-1}]t^{-1}U_k(\widehat{\mathfrak{g}}) + U_k(\widehat{\mathfrak{g}})\mathfrak{g}[t]t)},$$

where, for a subspace M of $\widetilde{U}_k(\widehat{\mathfrak{g}})$, \overline{M} denotes the closure of M. Let $\Phi : \widetilde{U}_k(\widehat{\mathfrak{g}}) \to U(\mathfrak{g})$ be the projection with respect to the above decomposition.

Lemma 2.3. The correspondence $[a] \mapsto \Phi(o(a))$ gives the algebra isomorphism $A(V^k(\mathfrak{g})) \xrightarrow{\sim} U(\mathfrak{g})$, where $o: V^k(\mathfrak{g}) \to \widetilde{U}_k(\widehat{\mathfrak{g}})$ is the linear map defined by $o(a) = a_{(\Delta_a-1)}$ for a homogeneous element a of V.

Proof. The fact that the above map is an algebra homomorphism follows from the formula

(7)
$$o(a)o(b) \equiv o(a * b) \pmod{(\mathfrak{g}[t^{-1}]t^{-1}U_k(\widehat{\mathfrak{g}}) + U_k(\widehat{\mathfrak{g}})\mathfrak{g}[t]t)},$$

where $a * b = \sum_{i \ge 0} {\Delta_a \choose i} a_{(i-1)} b$ for a homogeneous element $a \in V^k(\mathfrak{g})$ (see [Zhu, Theorem 2.1.2]). The assertion follows by recalling the proof of [FZ, Theorem 3.1.1].

Let \mathcal{O}_k be the full subcategory of $\hat{\mathfrak{g}}$ -modules of level k consisting of objects Mon which (1) L_0 acts locally finitely, (2) $\hat{\mathfrak{n}}_+$ acts locally nilpotently, (3) $\hat{\mathfrak{h}}$ acts semisimply. An object of \mathcal{O}_k is obviously smooth. Note that $V^k(\mathfrak{g})$ is an object of \mathcal{O}_k . Let $L(\lambda)$ be the irreducible highest weight representation of $\hat{\mathfrak{g}}$ with highest weight $\lambda \in \hat{\mathfrak{h}}^*$. Clearly, $L(\lambda)$ is positively graded and

$$L(\lambda)_{top} \cong L^{\mathfrak{g}}_{\overline{\lambda}}$$

as \mathfrak{g} -modules. The category \mathcal{O}_k is naturally regarded as a full subcategory of the category of $V^k(\mathfrak{g})$ -modules.

Let N_k be the unique maximal ideal of $V^k(\mathfrak{g})$. Then

$$V^k(\mathfrak{g})/N_k \cong L(k\Lambda_0)$$

as $\widehat{\mathfrak{g}}$ -modules, where $\Lambda_0 \in \widehat{\mathfrak{h}}^*$ is the 0-th fundamental weight of $\widehat{\mathfrak{g}}$: $\Lambda_0(K) = 1$, $\Lambda_0(\mathfrak{h}) = 0$. The vertex algebra $L(k\Lambda_0)$ is called the *(simple) affine vertex algebra* associated with \mathfrak{g} at level k. By definition, a $V^k(\mathfrak{g})$ -module M is a $L(k\Lambda_0)$ -module if and only if $a_{(n)}$ annihilates M for all $a \in N_k$, $n \in \mathbb{Z}$.

Let $\mathcal{L}(N_k)$ be the image of N_k in $\mathcal{L}(V^k(\mathfrak{g}))$. Then $\mathcal{L}(N_k)$ is a graded ideal of $\mathcal{L}(V^k(\mathfrak{g}))$: $\mathcal{L}(N_k) = \bigoplus_{\Delta \in \mathbb{Z}} \mathcal{L}(N_k)_{\Delta}$, where $\mathcal{L}(N_k)_{\Delta} = \mathcal{L}(N_k) \cap \mathcal{L}(V^k(\mathfrak{g}))_{\Delta}$. We have $\mathcal{L}(L(k\Lambda_0)) \cong \mathcal{L}(V^k(\mathfrak{g}))/\mathcal{L}(N_k)$. Set

$$I_k := \Phi(\mathcal{L}(N_k)_0) \subset U(\mathfrak{g}).$$

By (7), it follows that I_k is an two-sided ideal of $U(\mathfrak{g})$.

The following follows immediately from the definition of Zhu's algebra [FZ].

Lemma 2.4. The isomorphism $A(V^k(\mathfrak{g})) \xrightarrow{\sim} U(\mathfrak{g})$ in Lemma 2.3 induces the isomorphism $A(L(k\Lambda_0)) \xrightarrow{\sim} U(\mathfrak{g})/I_k$.

By Zhu's theorem [Zhu], the correspondence

$$M \mapsto M_{top}$$

gives a bijection between the set of isomorphism classes of simple positively graded $L(k\Lambda_0)$ -modules and that of simple $A(L(k\Lambda_0))$ -modules. Hence Lemma 2.1 and Lemma 2.4 give the following assertion.

Proposition 2.5. For a weight $\lambda \in \hat{\mathfrak{h}}^*$ of level k, $L(\lambda)$ is a module over $L(k\Lambda_0)$ if and only if $\overline{\lambda} \in \mathcal{V}(I_k)$. (Recall that $\overline{\lambda}$ denotes the restriction of λ to \mathfrak{h} .)

Lemma 2.6. Let $\lambda \in \widehat{\mathfrak{h}}^*$ be a weight of level k such that $\overline{\lambda} \in \mathcal{V}(I_k)$. Suppose that $\langle \lambda + \widehat{\rho}, \alpha_i^{\vee} \rangle \notin \mathbb{N}$ for $\alpha_i \in \widehat{\Pi}$. Then $\overline{s_i \circ \lambda} \in \mathcal{V}(I_k)$.

Proof. The case i = 1, ..., l is the statement of Lemma 2.2. We shall prove the assertion for i = 0 based on the argument in the proof of [Jos, Lemma 2]. Let $\mathfrak{sl}_2^{(0)} = \operatorname{span}_{\mathbb{C}} \{e_0, f_0, \alpha_0^{\vee}\} \subset \widehat{\mathfrak{g}}$ be the copy of $\mathfrak{sl}_2(\mathbb{C})$, where e_0 and f_0 are root vectors of root α_0 and $-\alpha_0$, respectively. Set

$$\widehat{\mathfrak{b}}^{(0)}=\mathfrak{sl}_2^{(0)}+(\widehat{\mathfrak{h}}\oplus\widehat{\mathfrak{n}}_+)=\widehat{\mathfrak{l}}\oplus\widehat{\mathfrak{m}},$$

the minimal parabolic subalgebra of $\hat{\mathfrak{g}}$, where $\hat{\mathfrak{l}}$ is its Levi subalgebra and $\hat{\mathfrak{m}}$ is its nilradical. We have $\hat{\mathfrak{l}} = \mathfrak{sl}_2^{(0)} + \hat{\mathfrak{h}}_0^{\perp}$ and $\hat{\mathfrak{m}} = \bigoplus_{\alpha \in \widehat{\Delta}_1^{re} \atop \alpha \neq \alpha_0} \widehat{\mathfrak{g}}_{\alpha}$, where $\hat{\mathfrak{h}}_0^{\perp}$ is the orthogonal

complement of $\mathbb{C}\alpha_0^{\vee}$ in $\hat{\mathfrak{h}}$ and $\hat{\mathfrak{g}}_{\alpha}$ denotes the root space of $\hat{\mathfrak{g}}$ of root α . Denote by $\hat{\mathfrak{m}}_-$ the opposite subalgebra of $\hat{\mathfrak{m}}$, so that $\hat{\mathfrak{g}} = \hat{\mathfrak{m}}_- \oplus \hat{\mathfrak{l}} \oplus \hat{\mathfrak{m}}_+$. Then we have the decomposition $U_k(\hat{\mathfrak{g}}) = U_k(\hat{\mathfrak{l}}) \oplus (\hat{\mathfrak{m}}_- U_k(\hat{\mathfrak{g}}) + U_k(\hat{\mathfrak{g}})\hat{\mathfrak{m}})$, where $U_k(\hat{\mathfrak{l}}) = U(\hat{\mathfrak{l}})/(K - k)U(\hat{\mathfrak{l}})$. This induces the decomposition

$$\widetilde{U}_k(\widehat{\mathfrak{g}}) = U_k(\widehat{\mathfrak{l}}) \oplus (\widehat{\mathfrak{m}}_- \widetilde{U}_k(\widehat{\mathfrak{g}}) \oplus \widetilde{U}_k(\widehat{\mathfrak{g}}) \widehat{\mathfrak{m}}),$$

Let $\widehat{\Upsilon}^{(0)}: \widetilde{U}_k(\widehat{\mathfrak{g}}) \to U_k(\widehat{\mathfrak{l}})$ be the projection with respect to this decomposition. Set

$$I := \widehat{\Upsilon}^{(0)}(\mathcal{L}(N_k)_0),$$

which is an ad $\hat{\mathfrak{l}}$ -invariant subspace of $U_k(\hat{\mathfrak{l}})$. Note that we have

(8)
$$\Upsilon(I_k) = \gamma^{(0)}(I)$$

where $\gamma^{(0)} : U(\hat{\mathfrak{l}}) \to U(\mathfrak{h})$ is the projection defined by the decomposition $U_k(\hat{\mathfrak{l}}) = U(\mathfrak{h}) \oplus (f_0 U_k(\hat{\mathfrak{l}}) + U_k(\hat{\mathfrak{l}})e_0).$

Since it is a direct sum of semisimple finite-dimensional representations I is generated by $I^{e_0} := \{u \in I; [e_0, u] = 0\}$ as an ad \hat{I} -module, and by Kostant's separation theorem I^{e_0} is spanned by the vectors of the form

$$e_0^n b_n$$
 with $n \in \mathbb{Z}_{\geq 0}$, $b_n \in Z_0$,

where Z_0 is the subalgebra of $U_k(\hat{\mathfrak{l}})$ generated by the quadratic Casimir element $\Omega_0 \in U(\mathfrak{sl}_2^{(0)})$ and $\hat{\mathfrak{h}}_0^{\perp}$.

Now suppose that $e_0^n b_n \in I^e$, $n \in \mathbb{Z}_{\geq 0}$, $b_k \in Z_0$, annihilates $L(\lambda)$. Then $e_0^n f_0^n b_n$ also annihilates $L(\lambda)$. Since

$$e_0^n f_0^n b_n v_{\lambda} = \lambda(\alpha_0^{\vee})(\lambda(\alpha_0^{\vee}) - 1) \dots (\lambda(\alpha_0^{\vee}) - n + 1) b_n v_{\lambda},$$

where v_{λ} is the highest weight vector of $L(\lambda)$, the assumption implies that λ is a zero of $\gamma^{(0)}(b_n)$, and so is $s_0 \circ \lambda$. It follows that I^{e_0} is spanned by the vectors $e_0^n b_n$, $n \in \mathbb{Z}_{\geq 0}, b_n \in Z_0$, such that $\gamma^{(0)}(b_n)(s_0 \circ \lambda) = 0$. By (8), we conclude that $s_0 \circ \lambda$ is a zero of $\Upsilon(I_k)$.

Lemma 2.7. Let $\lambda \in \hat{\mathfrak{h}}^*$ be a weight of level k such that $\overline{\lambda} \in \mathcal{V}(I_k)$. Then $\overline{w \circ \lambda} \in \mathcal{V}(I_k)$ for any $w \in \widetilde{W}_+$.

Proof. Let $w \in \widetilde{W}_+$. Then $w^{-1} = t_{\overline{\Lambda}_j} w_j$ for some $j \in J$, see (3). Let \widetilde{w}_j be a Tits lifting of w_j to the adjoint group of \mathfrak{g} . Also, let $\Delta(\overline{\Lambda}_j, z)$ be Li's delta operator [Li] corresponding to $\overline{\Lambda}_j$. One can obtain a new simple $L(k\Lambda_0)$ -module from $L(\lambda)$ by twisting the action $a \mapsto Y^{L(k\Lambda_0)}(a, z)$ on $L(\lambda)$ by $a \mapsto Y^{L(k\Lambda_0)}(\Delta(\overline{\Lambda}_j, z)\widetilde{w}_j a, z)$. Since the twisting the action by $\Delta(\bar{\Lambda}_j, z)$ induces the automorphism of $\hat{\mathfrak{g}}$ corresponding to $t_{\bar{\Lambda}_j}$ (see [Li, (3.15)–(3.17)]), one finds that the resulting $L(k\Lambda_0)$ -module is in the category \mathcal{O}_k and isomorphic to $L(w \circ \lambda)$ as $\hat{\mathfrak{g}}$ -modules. This completes the proof.

Lemma 2.8. Let $\lambda \in \widehat{\mathfrak{h}}^*$ be a weight of level k such that $\overline{\lambda} \in \mathcal{V}(I_k)$, and let $w \in \widetilde{W}$. Suppose that $\langle \lambda + \widehat{\rho}, \alpha^{\vee} \rangle \notin \mathbb{N}$ for all $\alpha \in \widehat{\Delta}^{re}_+ \cap w^{-1}(\widehat{\Delta}^{re}_-)$. Then $\overline{w \circ \lambda} \in \mathcal{V}(I_k)$.

Proof. By Lemma 2.7 we may assume that $w \in \widehat{W}$. We proceed by induction on the length $\ell(w)$ of w. The case that $\ell(w) = 1$ has been proved in Lemma 2.6. So let $\ell(w) > 1$. Write $w = s_i y$ with $i \in \{0, 1, \ldots, l\}, y \in \widehat{W}, \ell(w) = \ell(y) + 1$. Then

$$\widehat{\Delta}^{re}_+ \cap w^{-1}(\widehat{\Delta}^{re}_-) = \{y^{-1}(\alpha_i)\} \sqcup (\widehat{\Delta}^{re}_+ \cap y^{-1}(\widehat{\Delta}^{re}_-)).$$

Hence by the induction hypothesis we have $\overline{y \circ \lambda} \in \mathcal{V}(I_k)$. Since $\underline{\langle y \circ \lambda + \hat{\rho}, \alpha_i^{\vee} \rangle} = \langle y(\lambda + \hat{p}), \alpha_i^{\vee} \rangle = \langle \lambda + \hat{\rho}, y^{-1}(\alpha_i^{\vee}) \rangle \notin \mathbb{N}$, Lemma 2.6 gives that $\overline{w \circ \lambda} \in \mathcal{V}(I_k)$ as required.

3. KAC-WAKIMOTO ADMISSIBLE REPRESENTATIONS

For $\lambda \in \widehat{\mathfrak{h}}^*$, let $\widehat{\Delta}(\lambda)$ and $\widehat{W}(\lambda)$ be its integral root system and its integral Weyl group, respectively:

$$\widehat{\Delta}(\lambda) = \{ \alpha \in \widehat{\Delta}^{re}; \langle \lambda + \widehat{\rho}, \alpha^{\vee} \rangle \in \mathbb{Z} \}, \qquad \widehat{W}(\lambda) = \langle s_{\alpha}; \alpha \in \widehat{\Delta}(\lambda) \rangle.$$

Let $\widehat{\Delta}(\lambda)_+ = \widehat{\Delta}(\lambda) \cap \widehat{\Delta}_+^{re}$, the set of positive roots of $\widehat{\Delta}(\lambda)$ and $\Pi(\lambda) \subset \widehat{\Delta}(\lambda)_+$, the set of simple roots.

A weight $\lambda \in \mathfrak{h}^*$ is called *admissible* if

- (i) λ is regular dominant, that is, $\langle \lambda + \hat{\rho}, \alpha^{\vee} \rangle \notin \{0, -1, -2, \dots\},\$
- (ii) $\mathbb{Q}\widehat{\Delta}(\lambda) = \mathbb{Q}\widehat{\Delta}^{re}$.

An *admissible number* (for $\hat{\mathfrak{g}}$) is a complex number k such that $k\Lambda_0$ is admissible.

Proposition 3.1 ([KW2, KW3]). A complex number k is admissible if and only if

$$k+h^{\vee}=\frac{p}{q} \quad \text{with } p,q\in\mathbb{N}, \ (p,q)=1, \ p\geq \begin{cases} h^{\vee} & \text{if } (r^{\vee},q)=1\\ h & \text{if } (r^{\vee},q)=r^{\vee}, \end{cases}$$

where h is the the Coxeter number of \mathfrak{g} and r^{\vee} is the lacing number of \mathfrak{g} , that is, the maximal number of the edges in the Dynkin diagram of \mathfrak{g} . If this is the case we have $\widehat{\Pi}(k\Lambda_0) = {\dot{\alpha}_0, \alpha_1, \alpha_2, \ldots, \alpha_l}$, where

$$\dot{\alpha_0} = \begin{cases} -\theta + q\delta & \text{if } (r^{\vee}, q) = 1\\ -\theta_s + \frac{q}{r^{\vee}}\delta & \text{if } (r^{\vee}, q) = r^{\vee}. \end{cases}$$

For an admissible number k let Pr^k be the set of admissible weights λ such that $\widehat{\Delta}(\lambda) \cong \widehat{\Delta}(k\Lambda_0)$ as root systems. Set

$$Pr_{\mathbb{Z}}^{k} = Pr^{k} \cap \mathfrak{h}_{k,\mathbb{Z}}^{*},$$

where

(9)
$$\widehat{\mathfrak{h}}_{k,\mathbb{Z}}^* = \{\lambda \in \widehat{\mathfrak{h}}^*; \lambda(K) = k, \ \langle \lambda, \alpha_i^{\vee} \rangle \in \mathbb{Z} \text{ for all } i = 1, \dots, l\}.$$

Then (10)

$$Pr_{\mathbb{Z}}^{k} = \begin{cases} \{\lambda \in \widehat{\mathfrak{h}}_{k,\mathbb{Z}}^{*}; \langle \lambda, \alpha_{i}^{\vee} \rangle \geq 0 \text{ for } i = 1, \dots, l, \, \langle \lambda, \theta \rangle \leq p - h^{\vee} \} & \text{if } (r^{\vee}, q) = 1, \\ \{\lambda \in \widehat{\mathfrak{h}}_{k,\mathbb{Z}}^{*}; \langle \lambda, \alpha_{i}^{\vee} \rangle \geq 0 \text{ for } i = 1, \dots, l, \, \langle \lambda, \theta_{s}^{\vee} \rangle \leq p - h \} & \text{if } (r^{\vee}, q) = r^{\vee}. \end{cases}$$

We have [KW2]

(11)
$$Pr^{k} = \bigcup_{\substack{y \in \widetilde{W} \\ y(\widehat{\Delta}(k\Lambda_{0})_{+}) \subset \widehat{\Delta}_{+}^{re}}} Pr_{y}^{k}, \quad Pr_{y}^{k} := y \circ Pr_{\mathbb{Z}}^{k}.$$

Note that

$$Pr_{\mathbb{Z}}^{k} \cong \begin{cases} \widehat{P}_{+}^{p-h^{\vee}} & \text{if } (r^{\vee},q) = 1, \\ {}^{L}\widehat{P}_{+}^{\vee,p-h} & \text{if } (r^{\vee},q) = r^{\vee}. \end{cases}$$

where \widehat{P}^m_+ is the set of level *m* integral dominant weights of $\widehat{\mathfrak{g}}$, and ${}^L \widehat{P}^{\vee,m}_+$ is the set of level *m* integral dominant coweights of the affine Kac-Moody algebra $\widehat{{}^L \mathfrak{g}}$ associated with the Langlands dual Lie algebra ${}^L \mathfrak{g}$. Note also that for $\lambda \in Pr_{\mathbb{Z}}^k$ we have

$$\widehat{\Delta}(\lambda) = \begin{cases} \{\alpha + nq\delta; \alpha \in \Delta, \ n \in \mathbb{Z}\} & \text{if } (q, r^{\vee}) = 1, \\ \{\alpha + nq\delta; \alpha \in \Delta_{long}\} \sqcup \{\alpha + \frac{nq}{r^{\vee}}\delta; \alpha \in \Delta_{short}, n \in \mathbb{Z}\} & \text{if } (q, r^{\vee}) = r^{\vee}, \end{cases}$$

where Δ_{long} (resp. Δ_{short}) is the sets of long roots (resp. short roots) of \mathfrak{g} . It follows that

$$\widehat{W}(\lambda) = \begin{cases} W \ltimes q Q^{\vee} & \text{if } (q, r^{\vee}) = 1, \\ W \ltimes q Q & \text{if } (q, r^{\vee}) = r^{\vee} \end{cases}$$

for $\lambda \in Pr_{\mathbb{Z}}^k$. In particular

$$\widehat{W}(\lambda) \cong \begin{cases} \widehat{W} & \text{if } (q, r^{\vee}) = 1, \\ {}^{L}\widehat{W} & \text{if } (q, r^{\vee}) = r^{\vee}, \end{cases}$$

for $\lambda \in Pr^k$, where ${}^{L}\widehat{W}$ is the Weyl group of $\widehat{{}^{L}\mathfrak{g}}$.

Proposition 3.2 ([FKW, 1.5]). Let k be an admissible number for $\widehat{\mathfrak{g}}$ with denominator q as in Proposition 3.1. Suppose that $(q, r^{\vee}) = 1$, and let \widetilde{W}_+ acts on $Pr_{\mathbb{Z}}^k$ by $t_{\overline{\Lambda}_j}w_j \mapsto t_{q\overline{\Lambda}_j}w_j$, on P^{\vee}/qQ^{\vee} be the identification $\widetilde{W}_+ = P^{\vee}/Q^{\vee} = qP^{\vee}/qQ^{\vee}$, and on $Pr_{\mathbb{Z}}^k \times (P^{\vee}/qQ^{\vee})$ diagonally. We have a bijection

$$\begin{array}{ccc} (Pr_{\mathbb{Z}}^{k} \times (P^{\vee}/qQ^{\vee}))/\widetilde{W}_{+} & \stackrel{\sim}{\to} & Pr^{k}, \\ & & & & \\ [(\lambda,\mu)] & \mapsto & t_{-\mu}w \circ \lambda \end{array}$$

where w is the element of $W \ltimes qQ^{\vee}$ such that $t_{-\mu}w(\widehat{\Delta}(k\Lambda_0)_+) \subset \widehat{\Delta}_+^{re}$, which exists uniquely. In particular

$$|Pr^{k}| = |P^{\vee}/qP^{\vee}||\widehat{P}_{+}^{p-h^{\vee}}| = q^{l}|\widehat{P}_{+}^{p-h^{\vee}}|.$$

Let ${}^{L}\widetilde{W}_{+}$ denote the group \widetilde{W}_{+} for $\widehat{L}\mathfrak{g}$, that is, the subgroup of the extended Weyl group ${}^{L}\widetilde{W}$ of $\widehat{L}\mathfrak{g}$ consisting of elements of length zero. It acts on $Pr_{\mathbb{Z}}^{k}$ by the identification $Pr_{\mathbb{Z}}^{k} \cong {}^{L}\widehat{P}_{+}^{\vee,p-h}$ if r^{\vee} divides the denominator of k. It acts also

on the set P^{\vee}/qQ by the identification ${}^{L}\widetilde{W}_{+} = P/Q = qP/qQ$, and hence on $Pr_{\mathbb{Z}}^{k} \times (P^{\vee}/qQ)$ diagonally. The following assertion can be proved in the same manner as Proposition 3.2.

Proposition 3.3. Let k be an admissible number for $\hat{\mathfrak{g}}$ with denominator q as in Proposition 3.1. Suppose that $(q, r^{\vee}) = r^{\vee}$. We have a bijection

$$\begin{array}{rcl} (Pr_{\mathbb{Z}}^k \times (P^{\vee}/qQ))/{}^L \widetilde{W}_+ & \xrightarrow{\sim} & Pr^k, \\ & & & & \\ [(\lambda,\mu)] & \mapsto & t_{-\mu} w \circ \lambda, \end{array}$$

where w is the element of $W \ltimes qQ$ such that $t_{-\mu}w(\widehat{\Delta}(k\Lambda_0)_+) \subset \widehat{\Delta}_+^{re}$, which exists uniquely. In particular

$$|Pr^k| = |P^{\vee}/qP||^L \widehat{P}_+^{\vee,p-h}| = \frac{q^l|^L \widehat{P}_+^{\vee,p-h}|}{(r^{\vee})^{|short\ simple\ roots\ of\ \mathfrak{g}|}}.$$

Now recall the following important result [MF] (see also [FrM]).

Theorem 3.4 ([MF]). Let k be an admissible number and $\lambda \in Pr_{\mathbb{Z}}^k$. Then $L(\lambda)$ is a module over $L(k\Lambda_0)$.

The following assertion follows immediately from Proposition 2.5 and Theorem 3.4.

Corollary 3.5. We have $\overline{\lambda} \in \mathcal{V}(I_k)$ for $\lambda \in Pr_{\mathbb{Z}}^k$.

Proof of the "if part" of Main Theorem. Let $\mu \in Pr_y^k$. Then there exist $\lambda \in Pr_{\mathbb{Z}}^k$ and $w \in \widetilde{W}$ such that $\mu = y \circ \lambda$ and $y(\widehat{\Delta}(\lambda)_+) \subset \widehat{\Delta}_+^{re}$. (Note that $\widehat{\Delta}(\lambda)_+ = \Delta(k\Lambda_0)_+$.) The last condition is equivalent to that if $\alpha \in \widehat{\Delta}_+^{re}$ such that $\langle \lambda + \rho, \alpha^{\vee} \rangle \in \mathbb{Z}$ then $y(\alpha) \in \widehat{\Delta}_+^{re}$. Or equivalently, $\langle \lambda + \rho, \alpha^{\vee} \rangle \notin \mathbb{Z}$ for all $\alpha \in \widehat{\Delta}_+^{re} \cap y^{-1}(\widehat{\Delta}_-^{re})$. Since $\overline{\lambda}$ belongs to $\mathcal{V}(I_k)$ by Theorem 3.4, so does $\overline{\mu}$ by Lemma 2.8. By Proposition 2.5, this completes the proof of the "if part" of Main Theorem. \Box

4. Semi-infinite restriction functors

In this section we prove the "only if" part of Main Theorem by reducing to the \mathfrak{sl}_2 -cases. (Recall that statement of Main Theorem has been proved for $\mathfrak{g} = \mathfrak{sl}_2$ by Adamović and Milas [AM].)

Let k be an admissible number for $\hat{\mathfrak{g}}$, so that

$$k + h^{\vee} = \frac{p}{q}, \ p, q \in \mathbb{N}, \ (p,q) = 1, \ p \ge \begin{cases} h^{\vee} & \text{if } (r^{\vee},q) = 1, \\ h & \text{if } (r^{\vee},q) = r^{\vee}. \end{cases}$$

Let $\{e_i, h_i, f_i; i = 1, ..., l\}$ be a set of Chevalley generators of \mathfrak{g} . For i = 1, ..., l, let \mathfrak{p}_i be the minimal parabolic subalgebra $\mathbb{C}f_i \oplus \mathfrak{h} \oplus \mathfrak{n}$ of \mathfrak{g} , \mathfrak{l}_i its Levi subalgebra, \mathfrak{m}_i its nil-radical. We have

$$\mathfrak{p}_i = \mathfrak{l}_i \oplus \mathfrak{m}_i, \quad \mathfrak{l}_i = \mathfrak{sl}_2^{(i)} \oplus \mathfrak{h}_i^{\perp},$$

where $\mathfrak{sl}_2^{(i)}$ is the copy of \mathfrak{sl}_2 spanned by e_i, h_i and f_i , and \mathfrak{h}_i^{\perp} is the orthogonal component of $\mathbb{C}h_i$ in \mathfrak{h} . We have $\mathfrak{m}_i = \bigoplus_{\alpha \in \Delta(\mathfrak{m}_i)} \mathbb{C}x_{\alpha}$, where $\Delta(\mathfrak{m}_i) = \Delta_+ \setminus \{\alpha_i\}$ and x_{α} is a root vector of \mathfrak{g} of root α .

 Set

$$L\mathfrak{m}_i = \mathfrak{m}_i[t, t^{-1}] \subset \widehat{\mathfrak{g}}.$$

We shall consider the semi-infinite $L\mathfrak{m}_i$ -cohomology $H^{\frac{\infty}{2}+\bullet}(L\mathfrak{m}_i, M)$ with coefficient in $M \in \mathcal{O}_k$. This is defined by Feigin's standard complex $(C^{\bullet}(L\mathfrak{m}_i, M), d)$ ([Feĭ]); $H^{\frac{\infty}{2}+\bullet}(L\mathfrak{m}_i, M) = H^i(C^{\bullet}(L\mathfrak{m}_i, M), d), C^{\bullet}(L\mathfrak{m}_i, M) = M \otimes \bigwedge^{\frac{\infty}{2}+\bullet} \mathfrak{m}_i$, where $\bigwedge^{\frac{\infty}{2}+\bullet} \mathfrak{m}_i$ is a vertex (super)algebra generated by the odd field $\psi_{\alpha}(z) = \sum_{n \in \mathbb{Z}} \psi_{\alpha,n} z^{-n-1}$, $\psi^*_{\alpha}(z) = \sum_{n \in \mathbb{Z}} \psi^*_{\alpha,n} z^{-n}, \alpha \in \Delta(\mathfrak{m}_i)$, satisfying the OPEs

$$\psi_{\alpha}(z)\psi_{\beta}^{*}(w) \sim \frac{\delta_{\alpha,\beta}}{z-w}, \quad \psi_{\alpha}(z)\psi_{\beta}(w) \sim \psi_{\alpha}^{*}(z)\psi_{\beta}^{*}(w) \sim 0.$$

Differential d is the coefficient of z^{-1} of the field

$$d(z) = \sum_{\alpha \in \Delta(\mathfrak{m}_i)} x_{\alpha}(z) \psi_{\alpha}^*(z) - \frac{1}{2} \sum_{\alpha, \beta, \gamma \in \Delta(\mathfrak{m}_i)} c_{\alpha, \beta}^{\gamma} : \psi_{\alpha}^*(z) \psi_{\beta}^*(z) \psi_{\gamma}(z) :,$$

where $c_{\alpha,\beta}^{\gamma}$ is the structure constant: $[x_{\alpha}, x_{\beta}] = \sum_{\gamma} c_{\alpha,\beta}^{\gamma} x_{\gamma}$ and we have omitted the tensor product symbol.

The space $H^{\bullet}(L\mathfrak{m}_i, V^k(\mathfrak{g}))$ inherits the vertex (super)algebra structure from $C^{\bullet}(L\mathfrak{m}_i, V^k(\mathfrak{g}))$. It follows that the space $H^{\frac{\infty}{2}+\bullet}(L\mathfrak{m}_i, M), M \in \mathcal{O}_k$, is a module over the vertex algebra $H^{\frac{\infty}{2}+\bullet}(L\mathfrak{m}_i, M)$. Similarly, $H^{\bullet}(L\mathfrak{m}_i, L(k\Lambda_0))$ is naturally a vertex algebra, and if $M \in \mathcal{O}_k$ is a $L(k\Lambda_0)$ -module then $H^{\frac{\infty}{2}+\bullet}(L\mathfrak{m}_i, M)$ is a module over $H^{\frac{\infty}{2}+\bullet}(L\mathfrak{m}_i, L(k\Lambda_0))$.

Define the rational number k_i by the formula

(13)
$$k_i + 2 = \frac{2}{(\alpha_i | \alpha_i)} (k + h^{\vee}) = \begin{cases} \frac{p}{q} & \text{if } \alpha_i \text{ is a long root,} \\ \frac{r^{\vee} p}{q} & \text{if } \alpha_i \text{ is a short root.} \end{cases}$$

Note that k_i is an admissible number for \mathfrak{sl}_2 .

Let

$$V^{k_i}(\mathfrak{l}_i) = V^{k_i}(\mathfrak{sl}_2^{(i)}) \otimes \pi,$$

where π is the Heisenberg vertex algebra associated with $\mathfrak{h}_i^{\perp}[t, t^{-1}] \oplus \mathbb{C}K \subset \hat{\mathfrak{g}}$ at level $k + h^{\vee}$:

$$\pi = U(\mathfrak{h}_i^{\perp}[t, t^{-1}] \oplus \mathbb{C}K) \otimes_{U(\mathfrak{h}_i^{\perp}[t] \oplus \mathbb{C}K)} \mathbb{C}_{k+h^{\vee}},$$

where $\mathbb{C}_{k+h^{\vee}}$ is the one-dimensional representation of $\mathfrak{h}_i^{\perp}[t] \oplus \mathbb{C}K$ on which $\mathfrak{h}_i^{\perp}[t]$ acts trivially and K acts as the scalar $k + h^{\vee}$.

For $x \in \mathfrak{l}_i$, set

$$\widehat{x}(z) = x(z) - \sum_{\beta,\gamma \in \Delta(\mathfrak{m}_i)} (x_{-\gamma} | [x, x_\beta]) : \psi_\beta^*(z) \psi_\gamma(z) :,$$

where $x_{-\gamma}$ is a root vector of root $-\gamma$ such that $(x_{-\gamma}|x_{\gamma}) = 1$. Then $x(z) \mapsto \hat{x}(z)$, $x \in \mathfrak{l}_i$, gives a vertex algebra homomorphism $V^{k_i}(\mathfrak{l}_i) \to C^{\bullet}(L\mathfrak{m}_i, V^k(\mathfrak{g}))$. Since $\hat{x}(z)$, $x \in \mathfrak{l}_i$, commutes with the action of d, it induces a vertex algebra homomorphism

(14)
$$V^{k_i}(\mathfrak{l}_i) \to H^{\underline{\mathfrak{s}}_2+0}(L\mathfrak{m}_i, V^k(\mathfrak{g})),$$

see e.g. [HT] for the details of the above facts. Thus, $H^{\frac{\infty}{2}+0}(L\mathfrak{m}_i, M)$, $M \in \mathcal{O}_k$, is a module over $V^{k_i}(\mathfrak{l}_i)$. In particular, it is a module over $\widehat{\mathfrak{sl}}_2^{(i)}$ of level k_i , which belongs to the category \mathcal{O}_{k_i} of $\widehat{\mathfrak{sl}}_2^{(i)}$.

Denote by $V_{k_i}(\mathfrak{l}_i)$ the simple quotient of $V^{k_i}(\mathfrak{l}_i)$. We have

$$V_{k_i}(\mathfrak{l}_i) = L_{\widehat{\mathfrak{sl}}_0^{(i)}}(k_i \Lambda_0) \otimes \pi,$$

where $L_{\widehat{\mathfrak{sl}}_2^{(i)}}(\mu)$ denotes the irreducible representation of $\widehat{\mathfrak{sl}}_2^{(i)}$ with highest weight μ . The quotient map $V^k(\mathfrak{g}) \to L(k\Lambda_0)$ induces the vertex algebra homomorphism $H^{\frac{\infty}{2}+0}(L\mathfrak{m}_i, V^k(\mathfrak{g})) \to H^{\frac{\infty}{2}+0}(L\mathfrak{m}_i, L(k\Lambda_0))$. Hence we have the vertex algebra homomorphism

(15)
$$V^{k_i}(\mathfrak{l}_i) \to H^{\frac{\infty}{2}+0}(L\mathfrak{m}_i, L(k\Lambda_0)).$$

Theorem 4.1 ([A3, Theorem 7.5]). The vertex algebra homomorphism (15) factors through the vertex algebra homomorphism

$$V_{k_i}(\mathfrak{l}_i) \to H^{\frac{\infty}{2}+0}(L\mathfrak{m}_i, L(k\Lambda_0))$$

In particular, the fields $\hat{e}_i(z)$, $\hat{h}_i(z)$, $\hat{f}_i(z)$ generate the admissible affine vertex algebra $L_{\widehat{\mathfrak{sl}}^{(i)}}(k_i\Lambda_0)$ in $H^{\frac{\infty}{2}+0}(L\mathfrak{m}_i, L(k\Lambda_0))$.

Proof. For reader's convenience we shall sketch the proof here (see [A3] for the details). For a weight $\mu \in \hat{\mathfrak{h}}^*$ of level k, let $W(\mu)$ be the Wakimoto module of $\hat{\mathfrak{g}}$ with highest weight μ ([FF, Fre]). We have

(16)
$$H^{\frac{\infty}{2}+i}(L\mathfrak{m}_i, W(\mu)) \cong \begin{cases} W_{\mathfrak{l}_i}(\mu) & \text{for } i=0, \\ 0 & \text{for } i\neq 0, \end{cases}$$

where $W_{\mathfrak{l}_i}(\mu) = W_{\widehat{\mathfrak{sl}}_2^{(i)}}(\mu^{(i)}) \otimes \pi_{\mu|_{\mathfrak{h}_i^{\perp}}}, \, \mu^{(i)}$ is a weight of $\widehat{\mathfrak{sl}}_2^{(i)}$ defined by

$$\mu^{(i)} = \mu|_{\mathbb{C}h_i} + k_i \Lambda_0,$$

 $W_{\widehat{\mathfrak{sl}}_{2}^{(i)}}(\mu^{(i)})$ is the Wakimoto module of $\widehat{\mathfrak{sl}}_{2}^{(i)}$ with highest weight $\mu^{(i)}$ and π_{γ} is the simple π -module with highest weight γ . In particular $W(\mu)$ is acyclic with respect to the functor $H^{\frac{\infty}{2}+i}(L\mathfrak{m}_{i},?)$. Hence one may compute the cohomology $H^{\frac{\infty}{2}+i}(L\mathfrak{m}_{i},M)$ by using a resolution of M in terms of Wakimoto modules.

Set $\lambda = k\Lambda_0$. For the admissible representation $L(\lambda)$ there is a natural choice for such a resolution, which is a two-sided analogue of the usual BGG resolution: there exists a complex

(17)
$$C^{\bullet}: \dots \to C^{-1} \xrightarrow{d_{-1}} C^0 \xrightarrow{d_0} C^1 \xrightarrow{d_1} \cdots$$

of $\widehat{\mathfrak{g}}$ -modules such that

$$C^{i} = \bigoplus_{\substack{w \in \widehat{W}(\lambda) \\ \ell_{\lambda}^{\frac{\infty}{2}}(w) = i}} W(w \circ \lambda), \quad d_{i} = \sum_{\substack{w, w' \in \widehat{W}(\lambda) \\ \ell_{\lambda}^{\frac{\infty}{2}}(w) = i, w \succ w'}} d_{w, w'},$$

and $H^{i}(C^{\bullet}) = \begin{cases} L(\lambda) & \text{for } i = 0, \\ 0 & \text{for } i \neq 0, \end{cases}$

where $\ell_{\lambda}^{\frac{\infty}{2}}(w)$ denotes the semi-infinite length of $w \in \widehat{W}(\lambda)$, the symbol \triangleright denotes the covering in the semi-infinite Bruhat order, and $d_{w,w'}$ for $w \triangleright w'$ is a non-trivial $\widehat{\mathfrak{g}}$ module homomorphism $W(w \circ \lambda) \to W(w' \circ \lambda)$, which is unique up to multiplication by a nonzero constant. The existence of such a resolution was conjectured in [FKW], and was proved in [A3] by applying a result [Fie] of Fiebig and a method [Ark] of Arkhipov: Fiebig's equivalence between different blocks of categories \mathcal{O} shows the existence of the usual BGG resolution of $L(\lambda)$, and Arkhipov's method enables us to transform the usual BGG resolution to the two-sided BGG resolution, see [A3] for the details.

Now the space $H^{\frac{\infty}{2}+\bullet}(L\mathfrak{m}_i, L(\lambda))$ is isomorphic to the cohomology of the complex $H^{\frac{\infty}{2}+0}(L\mathfrak{m}_i, C^{\bullet})$ obtained from (17) by applying the functor $H^{\frac{\infty}{2}+0}(L\mathfrak{m}_i, ?)$, which has the following form:

$$\cdots \to \bigoplus_{\substack{w \in \widehat{W}(k\Lambda_0) \\ \ell_{\lambda}^{\frac{\infty}{2}}(w) = -1}} W_{\mathfrak{l}_i}(w \circ \lambda) \xrightarrow{d'_{-1}} \bigoplus_{\substack{w \in \widehat{W}(k\Lambda_0) \\ \ell_{\lambda}^{\frac{\infty}{2}}(w) = 0}} W_{\mathfrak{l}_i}(w \circ \lambda) \xrightarrow{d'_{-1}} \bigoplus_{\substack{w \in \widehat{W}(k\Lambda_0) \\ \ell_{\lambda}^{\frac{\infty}{2}}(w) = 1}} W_{\mathfrak{l}_i}(w \circ \lambda) \xrightarrow{d'_{-1}} \cdots,$$

where $d'_i = \sum_{\substack{w,w' \in \widehat{W}(k\Lambda_0)\\\ell^{\frac{\infty}{2}}(w)=i, w \vDash w'}} d'_{w,w'}$ and $d'_{w,w'}$ is the homomorphism $W_{\mathfrak{l}_i}(w \circ \lambda) \to W_{\mathfrak{l}_i}(w)$

 $W_{\mathfrak{l}_i}(w' \circ \lambda)$ induced by $d_{w,w'}$. Observe that in this realization of $H^{\frac{\infty}{2}+\bullet}(L\mathfrak{m}_i, L(\lambda))$ the vertex algebra homomorphism (15) is obtained from the vertex algebra embedding

$$V^{k_i}(\mathfrak{l}_i) \hookrightarrow W_{\mathfrak{l}_i}(k\Lambda_0) \subset H^{\frac{\infty}{2}+0}(L\mathfrak{m}_i, C^0)$$

described in [Fre, Theorem 5.1]. Therefore in order to prove the assertion it remains to show that the maximal proper submodule N of $V^{k_i}(\mathfrak{l}_i)$ is contained in the image of d'_{-1} . Note that N is tensor product of the maximal proper submodule of $V^{k_i}(\mathfrak{sl}_2^{(i)})$ and π .

 Set

$$\dot{\alpha}_{0}^{(i)} = \begin{cases} -\alpha_{i} + q\delta & \text{if } \alpha_{i} \text{ is a long root or } r^{\vee} \not| q, \\ -\alpha_{i} + \frac{q}{r^{\vee}}\delta & \text{if } \alpha_{i} \text{ is a short root and } r^{\vee} | q, \end{cases}$$

and put $\dot{s}_{0}^{(i)} = s_{\dot{\alpha}_{0}^{(i)}}$. Then $\dot{\alpha}_{0}^{(i)} \in \Delta(k\Lambda_{0}), \ \ell_{\lambda}^{\frac{\infty}{2}}(\dot{s}_{0}^{(i)}) = -1, \ \dot{s}_{0}^{(i)} \triangleright 1$, and the maximal submodule of $V^{k_{i}}(\mathfrak{sl}_{2})$ is generated by a singular vector of weight $(\dot{s}_{0}^{(i)} \circ k\Lambda_{0})^{(i)}$ ([KW1]). One sees that the image of the highest weight vector of $W_{\mathfrak{l}_{i}}(\dot{s}_{0}^{(i)} \circ k\Lambda_{0})$ in $W_{\mathfrak{l}_{i}}(k\Lambda_{0})$ by the homomorphism $d'_{\dot{s}_{0}^{(i)},1}$ is nonzero and generates the maximal submodule of $V^{k_{i}}(\mathfrak{l}_{i}) \subset W_{\mathfrak{l}_{i}}(k\Lambda_{0})$. This completes the proof.

Theorem 4.2 ([A3, Theorem 7.6]). If M is a module over $L(k\Lambda_0)$, the space $H^{\frac{\infty}{2}+r}(L\mathfrak{m}_i, M), r \in \mathbb{Z}$, is a direct sum of irreducible admissible representations of $\widehat{\mathfrak{sl}}_2^{(i)}$ of level k_i .

Proof. By Theorem 4.1, $H^{\frac{\infty}{2}+\bullet}(L\mathfrak{m}_i, M)$ is a module over $L_{\mathfrak{sl}_2^{(i)}}(k_i\Lambda_0) \subset V_{k_i}(\mathfrak{l}_i)$, which belongs to the category \mathcal{O} of $\mathfrak{sl}_2^{(i)}$ of level k_i . Hence the assertion follows immediately from [AM].

Let

$$L^{tot}(z) = \sum_{n \in \mathbb{Z}} L_n^{tot} z^{-n-2} := L(z) + : \sum_{\alpha \in \Delta(\mathfrak{m}_i)} : \psi_\alpha(z) \partial_z \psi_\alpha^*(z) : .$$

Then $[L_0^{tot}, xt^n] = -nxt^n$ for $x \in \mathfrak{g}$, $[L_0^{tot}, \psi_{\alpha,n}] = -n\psi_{\alpha,n}$, $[L_0^{tot}, \psi_{\alpha,n}^*] = -n\psi_{\alpha,n}^*$. It follows that L^{tot} commutes with d, and thus, acts on the space $H^{\frac{\infty}{2}+0}(L\mathfrak{m}_i, M)$.

For $M \in \mathcal{O}_k$, let $C^{\bullet}(L\mathfrak{m}_i, M)_d \subset C^{\bullet}(L\mathfrak{m}_i, M), H^{\frac{\infty}{2} + \bullet}(L\mathfrak{m}_i, M)_d \subset H^{\frac{\infty}{2} + \bullet}(L\mathfrak{m}_i, M)$, be the generalized L_0^{tot} -eigenspaces of eigenvalue d. Then

$$C^{\bullet}(L\mathfrak{m}_{i}, M) = \bigoplus_{d \in \mathbb{C}} C^{\bullet}(L\mathfrak{m}_{i}, M)_{d}$$
$$H^{\frac{\infty}{2} + \bullet}(L\mathfrak{m}_{i}, M) = \bigoplus_{d \in \mathbb{C}} H^{\frac{\infty}{2} + \bullet}(L\mathfrak{m}_{i}, M)_{d}$$

Note that $C^{\bullet}(L\mathfrak{m}_i, M)_d$ is a subcomplex of $C^{\bullet}(L\mathfrak{m}_i, M)$ and $H^{\frac{\infty}{2}+\bullet}(L\mathfrak{m}_i, M)_d$ is the cohomology of the subcomplex $C^{\bullet}(L\mathfrak{m}_i, M)_d$.

Lemma 4.3. Let M be a positively graded $\widehat{\mathfrak{g}}$ -module in the category \mathcal{O}_k , and let d_{top} be the L_0 -eigenvalue on M_{top} . Then $H^{\frac{\infty}{2}+\bullet}(L\mathfrak{m}_i, M) = \bigoplus_{n \in \mathbb{Z}_{\geq 0}} H^{\frac{\infty}{2}+\bullet}(L\mathfrak{m}_i, M)_{d_{top}+n}$

and

$$H^{\frac{\infty}{2}+r}(L\mathfrak{m}_i, M)_{d_{top}} \cong \begin{cases} H^r(\mathfrak{m}_i, M_{top}) & \text{if } r \ge 0, \\ 0 & \text{otherwise} \end{cases}$$

where $H^{\bullet}(\mathfrak{m}_i, N)$ denotes the (usual) Lie algebra \mathfrak{m}_i -cohomology with coefficient in an \mathfrak{m}_i -module N.

Proof. The first assertion is easy to see. The second assertion follows by observing that $C^{\bullet}(L\mathfrak{m}_i, M)_{d_{top}} = M_{top} \otimes \bigwedge^{\bullet}(\mathfrak{m}_i)$ and the restriction of the differential to $C^{\bullet}(L\mathfrak{m}_i, M)_{d_{top}}$ coincides with the differential of the Chevalley complex for calculating $H^{\bullet}(\mathfrak{m}_i, M_{top})$.

Lemma 4.4. Let k be an admissible number and suppose that $L(\lambda)$ is a module over $L(k\Lambda_0)$. Then $\lambda^{(i)} = \lambda|_{\mathbb{C}h_i} \oplus k_i\Lambda_0$ is an admissible weight for $\widehat{\mathfrak{sl}}_2$ for all $i = 1, \ldots, l$. *Proof.* Let d_{top} be the top weight of $L(\lambda)$. By Lemma 4.3, $H^{\frac{\infty}{2}+0}(L\mathfrak{m}_i, L(\lambda)) = \bigoplus_{d \geq d_{top}} H^{\frac{\infty}{2}+0}(L\mathfrak{m}_i, L(\lambda))_d$ and

$$H^{\frac{\infty}{2}+0}(L\mathfrak{m}_i, L(\lambda))_{d_{top}} = H^0(\mathfrak{m}_i, L^{\mathfrak{g}}_{\overline{\lambda}}).$$

It follows that the image $[v_{\lambda}]$ of the highest weight vector v_{λ} of $L(\lambda)$ is a non-zero singular vector of $\widehat{\mathfrak{sl}}_{2}^{(i)}$ of weight $\lambda^{(i)}$ in $H^{\frac{\infty}{2}+0}(L\mathfrak{m}_{i}, L(\lambda))$. Hence $[v_{\lambda}]$ generates a (nonzero) highest weight $\widehat{\mathfrak{sl}}_{2}^{(i)}$ -submodule with highest weight $\lambda^{(i)}$. But according to Theorem 4.2 such a module must be admissible. This completes the proof. \Box

Proposition 4.5. Suppose that $L(\lambda)$ is a $L(k\Lambda_0)$ -module. Then $\widehat{\Delta}(\lambda) \cong \widehat{\Delta}(k\Lambda_0)$.

Proof. By Lemma 4.4, $\langle \lambda + \rho, \alpha_i^{\vee} \rangle \in \frac{2}{(\alpha_i | \alpha_i)q} \mathbb{Z}$ for all $i = 1, \dots l$. It follows that there exists $n_i \in \mathbb{Z}$ for each $i = 1, \dots, l$ such that $\alpha_i + n_i \delta \in \widehat{\Delta}(\lambda)$. Hence

(18) there exists $n_{\alpha} \in \mathbb{Z}$ for each $\alpha \in \Delta$ such that $\alpha + n_{\alpha}\delta \in \widehat{\Delta}(\lambda)$.

This implies that $\mathbb{Q}\widehat{\Delta}(\lambda) = \mathbb{Q}\widehat{\Delta}^{re}$. Therefore [KW2] $\widehat{\Delta}(\lambda)$ is isomorphic to the integral root system of some admissible weight of level k. But according to the classification of admissible weights by Kac and Wakimoto [KW2], the integral roots of admissible weights satisfying the property (18) must be isomorphic to $\widehat{\Delta}(k\Lambda_0)$.

Proposition 4.6. Let k be an admissible number and $\lambda \in \hat{\mathfrak{h}}_{k,\mathbb{Z}}^*$ (see (9)). Suppose that $L(\lambda)$ is a module over $L(k\Lambda_0)$. Then $L(\lambda)$ is admissible.

Proof. First, it follows that

$$\langle \lambda, \alpha_i^{\vee} \rangle \ge 0 \text{ for } i = 1, \dots, l$$

from Lemma 4.4. Therefore the \mathfrak{g} -submodule $L^{\mathfrak{g}}_{\overline{\lambda}}$ generated by the highest weight vector of $L(\lambda)$ is finite-dimensional. By [Kos] we have

$$H^{r}(\mathfrak{m}_{i}, L^{\mathfrak{g}}_{\bar{\lambda}})) \cong \bigoplus_{\substack{w \in W^{(i)}\\\ell(w) = r}} L^{\mathfrak{l}_{i}}_{w \circ \bar{\lambda}}$$

for $r \geq 0$ as l_i -modules, where $L^{l_i}_{\bar{\mu}}$ is the irreducible highest weight representation of l_i with highest weight $\bar{\mu}$ and

$$W^{(i)} = \{ w \in W; w^{-1}(\alpha_i) \in \Delta_+ \}.$$

Therefore

$$H^{\frac{\infty}{2}+r}(L\mathfrak{m}_i, L(\lambda))_{d_{top}} \cong \bigoplus_{\substack{w \in W^{(i)}\\\ell(w)=r}} L^{\mathfrak{l}_i}_{w \circ \bar{\lambda}}$$

by Lemma 4.3. Since each l_i -highest weight vector of $H^{\frac{\infty}{2}+r}(L\mathfrak{m}_i, L(\lambda))_{d_{top}}$ generates a highest weight representation of $\widehat{\mathfrak{sl}}_{2}^{(i)}$, the weights $(w \circ \lambda)^{(i)}$ must be admissible by Theorem 4.2, that is,

(19)
$$(w \circ \lambda)^{(i)} \in Pr_{\mathbb{Z}}^{(i),k_i} \text{ for all } w \in W^{(i)}, i = 1, \dots, l.$$

where $Pr_{\mathbb{Z}}^{(i),k_i}$ denotes the set $Pr_{\mathbb{Z}}^{k_i}$ for $\widehat{\mathfrak{sl}}_2^{(i)}$. Now first consider the case that $(r^{\vee},q) = 1$. It remains to show that $\langle \lambda,\theta \rangle \leq \frac{1}{2}$ $p - h^{\vee}$, see (10). Let α_i be any simple long root of \mathfrak{g} , so that there exists $w \in W$ such that $\theta = w^{-1}(\alpha_i)$. Since $(w \circ \lambda)^{(i)} \in Pr_{\mathbb{Z}}^{(i),k_i}$ and $k_i + 2 = p/q$, we have

$$p-2 \ge \langle w \circ \lambda, \alpha_i^{\vee} \rangle = \langle w(\lambda+\rho), \alpha_i^{\vee} \rangle - 1 = \langle \lambda+\rho, \theta \rangle - 1 = \langle \lambda, \theta \rangle + h^{\vee} - 2.$$

We have shown that $\lambda \in Pr_{\mathbb{Z}}^k$.

Next consider the case that $(r^{\vee}, q) = r^{\vee}$. We need to show that $\langle \lambda, \theta_s^{\vee} \rangle \leq p - h$. Let α_i be any simple short root of \mathfrak{g} . Then there exists $w \in W$ such that $\theta_s = w^{-1}(\alpha_i)$. Since $(w \circ \lambda)^{(i)} \in Pr_{\mathbb{Z}}^{(i),k_i}$ and $k_i + 2 = r^{\vee}p/q = p/(q/r^{\vee})$, we have

$$p-2 \ge \langle w \circ \lambda, \alpha_i^{\vee} \rangle = \langle w(\lambda+\rho), \alpha_i^{\vee} \rangle - 1 = \langle \lambda+\rho, \theta_s^{\vee} \rangle - 1 = \langle \lambda, \theta_s^{\vee} \rangle + h - 2.$$

Hence $\lambda \in Pr_{\mathbb{Z}}^k$ as required.

Proof of the "only if" part of Main Theorem. Suppose that $L(\lambda)$ is a module over $L(k\Lambda_0)$. By Proposition 4.5, $\widehat{\Delta}(\lambda) \cong \widehat{\Delta}(k\Lambda_0)$. Therefore by [KW2, Lemma 2.1] there exist $y \in \widetilde{W}$ such that $\widehat{\Delta}(\lambda)_+ = y(\widehat{\Delta}(k\Lambda_0)_+)$. Let $\mu = y^{-1} \circ \lambda$. Then

$$\widehat{\Delta}(\mu)_{+} = \widehat{\Delta}(k\Lambda_{0})_{+} = y^{-1}(\widehat{\Delta}(\lambda)_{+}).$$

This gives on the one hand that $\mu \in \widehat{\mathfrak{h}}_{k,\mathbb{Z}}^*$ and on the other hand $\langle \mu + \rho, \alpha^{\vee} \rangle \notin \mathbb{Z}$ for $\alpha \in \widehat{\Delta}^{re}_+ \cap y(\widehat{\Delta}^{re}_-)$. The latter condition implies that $L(\mu)$ is also a module over $L(k\Lambda_0)$ by Lemma 2.8. But then Proposition 4.6 implies that μ must be an admissible weight since $\mu \in \widehat{\mathfrak{h}}_{k,\mathbb{Z}}^*$. Therefore λ is also an admissible weight. This completes the proof of the "only if" part of Main Theorem.

The last assertion of Main Theorem now follows immediately from the fact that $\operatorname{Ext}_{\mathcal{O}_k}^1(L(\lambda), L(\mu)) = 0$ for any admissible weights λ, μ of level k since they are regular dominant weights. In fact it is known by [GK, Theorem 0.2] that $\operatorname{Ext}_{\widehat{\mathfrak{G}}}^1(L(\lambda), L(\mu)) = 0$ for any admissible weights λ, μ . (Note that it is obvious that $\operatorname{Ext}_{\mathcal{O}_k}^1(L(\lambda), L(\lambda)) = 0$, but it is highly non-trivial that $\operatorname{Ext}_{\widehat{\mathfrak{g}}}^1(L(\lambda), L(\lambda)) = 0$.) Hence we have the following slightly more general assertion than the last assertion of Main Theorem.

Proposition 4.7. Let M be a finitely generated $L(k\Lambda_0)$ -module on which $\hat{\mathfrak{n}}_+$ locally nilpotently and $\hat{\mathfrak{h}}$ acts locally finitely. Then M is a direct sum of admissible representations of $\hat{\mathfrak{g}}$ whose integral Weyl groups are isomorphic to that of $L(k\Lambda_0)$.

Appendix A. Generalized semi-infinite Borel-Weil-Bott Theorem for admissible representations

Let \mathfrak{p} be a parabolic subalgebra of \mathfrak{g} containing \mathfrak{b}_- , and let $\mathfrak{p} = \mathfrak{l} \oplus \mathfrak{r}_-$ be the direct sum decomposition of \mathfrak{p} with the Levi subalgebra \mathfrak{l} containing \mathfrak{h} and the nilpotent radical \mathfrak{m}_- . Denote by $\mathfrak{m} \subset \mathfrak{n}$ the opposite algebra of \mathfrak{m}_- , so that $\mathfrak{g} = \mathfrak{p} \oplus \mathfrak{m}$. Let

$$\mathfrak{l} = \mathfrak{l}_0 \oplus \bigoplus_{i=1}^s \mathfrak{l}_i$$

be the decomposition of \mathfrak{l} into direct sum of simple Lie subalgebras \mathfrak{l}_i , $i = 1, \ldots, s$, and its center \mathfrak{l}_0 of \mathfrak{l} . Let $\mathfrak{h}_i = \mathfrak{l} \cap \mathfrak{h}$, the Cartan subalgebra of \mathfrak{l}_i , and denote by $\Delta_i \subset \Delta$ the subroot system of \mathfrak{g} corresponding to \mathfrak{l}_i , $\Delta_{i,+} = \Delta_i \cap \Delta_+$, $\Pi_i = \Pi \cap \Delta_i$. Let h_i^{\vee} be the dual Coxeter number of \mathfrak{l}_i (with a convention that $h_0^{\vee} = 0$), θ_i the highest root of Δ_i , $\theta_{i,s}$ the highest short roof of Δ_i .

Let $\hat{\mathfrak{l}}_i = \mathfrak{l}_i[t, t^{-1}] \oplus \mathbb{C}K \subset \hat{\mathfrak{g}}$ for $i = 0, 1, \dots, s$. Set

$$K_i = \frac{2}{(\theta_i | \theta_i)} K,$$

and we consider K_i as an element of \hat{l}_i . Thus,

$$\widehat{\mathfrak{l}}_i = \mathfrak{l}_i[t, t^{-1}] \oplus \mathbb{C}K_i.$$

Then $\widehat{\mathfrak{h}}_i = \mathfrak{h}_i \oplus \mathbb{C}K_i$ is a Cartan subalgebra of $\widehat{\mathfrak{l}}_i$. Set $\widehat{\mathfrak{h}}_i^* = \mathfrak{h}_i^* \oplus \mathbb{C}\Lambda_{0,i} \subset \widehat{\mathfrak{h}}^*$, where $\Lambda_{0,i} = \frac{(\theta_i|\theta_i)}{2}\Lambda_0$, and we regard $\widehat{\mathfrak{h}}_i^*$ as the dual of $\widehat{\mathfrak{h}}_i$.

$$\widehat{\Delta}_{\mathfrak{l}_i}^{re} = \{ \alpha + n\delta \in \widehat{\Delta}^{re}; \alpha_i \in \Delta_i \}$$

Then $\widehat{\Delta}_{\mathfrak{l}_i}^{re}$ is a subroot system of $\widehat{\Delta}^{re}$. We regard $\widehat{\Delta}_{\mathfrak{l}_i}^{re}$ as a real root system of $\widehat{\mathfrak{l}}_i$. In particular subgroup of \widehat{W} generated $s_{\alpha}, \alpha \in \widehat{\Delta}_+^{re}$, is identified with the Weyl group \widehat{W}_i of $\widehat{\mathfrak{l}}_i$.

For $\lambda \in \widehat{\mathfrak{h}}^*$ define $\lambda_i \in \widehat{\mathfrak{h}}_i^*, i = 0, 1, \dots, s$, by

$$\langle \lambda_i + \widehat{\rho}_i, \alpha^{\vee} + nK_i \rangle = \langle \lambda + \widehat{\rho}, \alpha^{\vee} + \frac{2n}{(\theta_i | \theta_i)} K \rangle \quad \text{for } \alpha^{\vee} \in \mathfrak{h}_i,$$

where $\hat{\rho}_i = \rho_i + h_i^{\vee} \Lambda_{0,i}$, $\rho_i = \sum_{\alpha \in \Delta_{i,+}} \alpha/2$. Observe that if λ is an admissible weight of $\hat{\mathfrak{g}}$ level k, then λ_i is an admissible weight of $\hat{\mathfrak{l}}_i$ of level k_i for $i = 1, 2, \ldots, s$, where

$$k_i + h_i^{\vee} = \frac{2}{(\theta_i | \theta_i)} (k + h^{\vee}).$$

Define Lie algebras

$$\widehat{\mathfrak{l}} = \bigoplus_{i=0}^{s} \widehat{\mathfrak{l}}_{i}, \quad \widehat{\mathfrak{t}} = \bigoplus_{i=0}^{s} \widehat{\mathfrak{h}}_{i} \subset \widehat{\mathfrak{l}}.$$

For $\lambda \in \widehat{\mathfrak{h}}^*$, set

$$\lambda_{\mathfrak{l}} = \sum_{i=0}^{s} \lambda_i \in \widehat{\mathfrak{t}}^*,$$

and let $L_{\mathfrak{l}}(\lambda_{\mathfrak{l}})$ be the irreducible highest weight representation of $\hat{\mathfrak{l}}$ with highest weight $\lambda_{\mathfrak{l}}$. Then $L_{\mathfrak{l}}(\lambda_{\mathfrak{l}}) = \bigotimes_{i=0}^{s} L_{\mathfrak{l}_{i}}(\lambda_{i})$, where $L_{\mathfrak{l}_{i}}(\lambda_{i})$ is the irreducible highest weight representation of $\hat{\mathfrak{l}}_{i}$ with highest weight λ_{i} . The weight $\lambda_{\mathfrak{l}} \in \hat{\mathfrak{t}}^{*}$ is called admissible if λ_{i} is admissible for all $i \neq 0$, and if this is the case the $\hat{\mathfrak{l}}$ -module $L_{\mathfrak{l}}(\lambda)$ is called admissible.

Now let k be an admissible number for $\hat{\mathfrak{g}}$, and let $\lambda \in Pr_{\mathbb{Z}}^k$. Set

$$\widehat{\Delta}_{\mathfrak{l}}(\lambda) = \bigcup_{i=1}^{s} \widehat{\Delta}_{\mathfrak{l}_{i}}(\lambda), \quad \widehat{\Delta}_{\mathfrak{l}_{i}}(\lambda) = \widehat{\Delta}_{\mathfrak{l}_{i}}^{re} \cap \widehat{\Delta}(\lambda),$$

and let $\widehat{W}_{\mathfrak{l}}(\lambda)$ be the subgroup of $\widehat{W}(\lambda)$ generated by $s_{\alpha}, \alpha \in \widehat{\Delta}_{\mathfrak{l}}(\lambda)$. Define

$$\widehat{W}^{\mathfrak{l}}(\lambda) = \{ w \in \widehat{W}(\lambda); w^{-1}(\widehat{\Delta}(\lambda)_{+}) \subset \widehat{\Delta}_{+}^{re} \}.$$

Then the multiplication map $\widehat{W}_{\mathfrak{l}}(\lambda) \times \widehat{W}^{\mathfrak{l}}(\lambda) \to \widehat{W}(\lambda), (u, v) \mapsto uv$, is a bijection. Moreover we have

$$\ell_{\lambda}^{\frac{\infty}{2}}(uv) = \ell_{\lambda}^{\frac{\infty}{2}}(u) + \ell_{\lambda}^{\frac{\infty}{2}}(v)$$

for $u \in \widehat{W}_{\mathfrak{l}}(\lambda), v \in \widehat{W}^{\mathfrak{l}}(\lambda)$ ([Pet], see [A3, Theorem 3.3]).

Lemma A.1. (i) For $w \in \widehat{W}(\lambda)$, $(w \circ \lambda)_{\mathfrak{l}}$ is admissible if and only if $w \in \widehat{W}^{\mathfrak{l}}(\lambda)$.

(ii) For $w, w' \in \widehat{W}^{\mathfrak{l}}(\lambda)$, $(w \circ \lambda)_{\mathfrak{l}} = (w' \circ \lambda)_{\mathfrak{l}}$ if and only if w = w'.

Let $L\mathfrak{m} = \mathfrak{m}[t, t^{-1}] \subset \hat{\mathfrak{g}}$. The semi-infnite $L\mathfrak{m}$ -cohomology $H^{\frac{\infty}{2}+0}(L\mathfrak{m}, L(k\Lambda_0))$ is naturally an $\hat{\mathfrak{l}}$ -module. In fact, as in the case that \mathfrak{p} is a minimal parabolic subalgebra (see Theorem 4.1), there is an injective vertex algebra homomorphism

(20)
$$L_{\mathfrak{l}}((k\Lambda_0)_{\mathfrak{l}}) = \bigotimes_{i=0}^{s} L_{\mathfrak{l}_i}(k_i\Lambda_{i,0}) \hookrightarrow H^{\frac{\infty}{2}+0}(L\mathfrak{m}, L(k\Lambda_0))$$

provided that k is admissible ([A3, Theorem 7.5]). Since $L(\lambda)$ is a module over $L(k\Lambda_0)$, $H^{\frac{\infty}{2}+p}(L\mathfrak{m}, L(\lambda))$ is a module over the vertex algebra $H^{\frac{\infty}{2}+0}(L\mathfrak{m}, L(k\Lambda_0))$, and hence, is a module over $L_{\mathfrak{l}}((k\Lambda_0)_{\mathfrak{l}})$. Since each k_i is an admissible number for $\hat{\mathfrak{l}}_i$ for all $i \neq 0$, it follows from Main Theorem that $H^{\frac{\infty}{2}+p}(L\mathfrak{m}, L(\lambda))$ is a direct sum of admissible representations of $\hat{\mathfrak{l}}$ for each $p \in \mathbb{Z}$.

The following assertion can be proved in the same manner as in [A3, Theorem 7.7].

Theorem A.2. Let k be an admissible number for $\hat{\mathfrak{g}}$, $\lambda \in Pr_{\mathbb{Z}}^k$. For each $p \in \mathbb{Z}$ we have the $\hat{\mathfrak{l}}$ -module isomorphism

$$H^{\frac{\infty}{2}+p}(L\mathfrak{m},L(\lambda)) \cong \bigoplus_{\substack{w \in \widehat{W}^{\mathfrak{l}}(\lambda) \\ \ell_{\lambda}^{\frac{\infty}{2}}(w)=p}} L_{\mathfrak{l}}((w \circ \lambda)_{\mathfrak{l}}).$$

Remark A.3. Although an admissible affine vertex algebra has only finitely many isomorphism classes of simple modules in the category \mathcal{O} the sum in Theorem A.2 is infinite since $\hat{\mathfrak{l}}_0$ is a Heisenberg algebra which has infinitely many isomorphism classes of simple modules in the category \mathcal{O} .

Theorem A.2 should be regarded as an analogue of Kostant's generalized Borel-Weil-Bott Theorem [Kos] appeared in the proof of Proposition 4.6. It has been proved in [HT] in the case that $L(\lambda)$ is an integrable representation of $\hat{\mathfrak{g}}$ using the fact that integrable representations are unitarizable, following the original idea of Kostant's proof. Their proof does not apply to our case since non-integrable admissible representations are not unitarizable.

We also note that in the above formula it was essential that $L(\lambda)$ is an admissible representations in order to to apply Main Theorem for $\hat{\mathfrak{l}}_i, i \neq 0$. Otherwise, the complete reducibility does not necessarily hold and the cohomology $H^{\frac{\infty}{2}+p}(L\mathfrak{m}, L(\lambda))$ may be very complicated.

References

- [Ada] Dražen Adamović. Some rational vertex algebras. Glas. Mat. Ser. III, 29(49)(1):25–40, 1994.
- [ALY] Tomoyuki Arakawa, Ching Hung Lam, and Hiromichi Yamada. Zhu's algebra, C₂-algebra and C₂-cofiniteness of parafermion vertex operator algebras. Adv. Math., 264:261–295, 2014.
- [AM] Dražen Adamović and Antun Milas. Vertex operator algebras associated to modular invariant representations for $A_1^{(1)}$. Math. Res. Lett., 2(5):563–575, 1995.
- [A1] Tomoyuki Arakawa. A remark on the C_2 cofiniteness condition on vertex algebras. Math. Z., 270(1-2):559–575, 2012.
- [A2] Tomoyuki Arakawa. Associated varieties of modules over Kac-Moody algebras and C₂cofiniteness of W-algebras. arXiv:1004.1554[math.QA].
- [A3] T. Arakawa. Two-sided BGG resolution of admissible representations. Represent. Theory, 19(3):183–222, 2014.
- [A4] Tomoyuki Arakawa. Rationality of W-algebras; principal nilpotent cases. arXiv:1211.7124[math.QA].
- [Ark] Sergey Arkhipov. A new construction of the semi-infinite BGG resolution. math.QA/9605043.
- [DLM] Chongying Dong, Haisheng Li, and Geoffrey Mason. Vertex operator algebras associated to admissible representations of \hat{sl}_2 . Comm. Math. Phys., 184(1):65–93, 1997.
- [Duf] Michel Duflo. Sur la classification des idéaux primitifs dans l'algèbre enveloppante d'une algèbre de Lie semi-simple. Ann. of Math. (2), 105(1):107–120, 1977.
- [FBZ] Edward Frenkel and David Ben-Zvi. Vertex algebras and algebraic curves, volume 88 of Mathematical Surveys and Monographs. American Mathematical Society, Providence, RI, second edition, 2004.
- [Feĭ] B. L. Feĭgin. Semi-infinite homology of Lie, Kac-Moody and Virasoro algebras. Uspekhi Mat. Nauk, 39(2(236)):195–196, 1984.

- [FF] Boris L. Feigin and Edward V. Frenkel. Affine Kac-Moody algebras and semi-infinite flag manifolds. Comm. Math. Phys., 128(1):161–189, 1990.
- $\begin{array}{ll} \mbox{[Fie]} & \mbox{Peter Fiebig. The combinatorics of category \mathcal{O} over symmetrizable Kac-Moody algebras.} \\ & Transform. \ Groups, 11(1):29-49, 2006. \end{array}$
- [FKW] Edward Frenkel, Victor Kac, and Minoru Wakimoto. Characters and fusion rules for Walgebras via quantized Drinfel'd-Sokolov reduction. Comm. Math. Phys., 147(2):295–328, 1992.
- [FeM] Boris Feigin and Fyodor Malikov. Modular functor and representation theory of sl₂ at a rational level. In Operads: Proceedings of Renaissance Conferences (Hartford, CT/Luminy, 1995), volume 202 of Contemp. Math., pages 357–405, Providence, RI, 1997. Amer. Math. Soc.
- [FrM] Igor Frenkel and Fyodor Malikov. Kazhdan-Lusztig tensoring and Harish-Chandra categories. arXiv:q-alg/9703010.
- [Fre] Edward Frenkel. Wakimoto modules, opers and the center at the critical level. Adv. Math., 195(2):297–404, 2005.
- [FZ] Igor B. Frenkel and Yongchang Zhu. Vertex operator algebras associated to representations of affine and Virasoro algebras. Duke Math. J., 66(1):123–168, 1992.
- [GK] Maria Gorelik and Victor Kac. On complete reducibility for infinite-dimensional Lie algebras. Adv. Math., 226(2):1911–1972, 2011.
- [HT] Shinobu Hosono and Akihiro Tsuchiya. Lie algebra cohomology and N = 2 SCFT based on the GKO construction. *Comm. Math. Phys.*, 136(3):451–486, 1991.
- [Jos] A. Joseph. A characteristic variety for the primitive spectrum of a semisimple Lie algebra. In Non-commutative harmonic analysis (Actes Colloq., Marseille-Luminy, 1976), pages 102–118. Lecture Notes in Math., Vol. 587. Springer, Berlin, 1977.
- [Kac] Victor Kac. Vertex algebras for beginners, volume 10 of University Lecture Series. American Mathematical Society, Providence, RI, second edition, 1998.
- [Kos] Bertram Kostant. Lie algebra cohomology and the generalized Borel-Weil theorem. Ann. of Math. (2), 74:329–387, 1961.
- [KW1] Victor G. Kac and Minoru Wakimoto. Modular invariant representations of infinitedimensional Lie algebras and superalgebras. Proc. Nat. Acad. Sci. U.S.A., 85(14):4956– 4960, 1988.
- [KW2] V. G. Kac and M. Wakimoto. Classification of modular invariant representations of affine algebras. In Infinite-dimensional Lie algebras and groups (Luminy-Marseille, 1988), volume 7 of Adv. Ser. Math. Phys., pages 138–177. World Sci. Publ., Teaneck, NJ, 1989.
- [KW3] Victor G. Kac and Minoru Wakimoto. On rationality of W-algebras. Transform. Groups, 13(3-4):671–713, 2008.
- [Li] Haisheng Li. The physics superselection principle in vertex operator algebra theory. J. Algebra, 196(2):436–457, 1997.
- [MF] F. G. Malikov and I. B. Frenkel'. Annihilating ideals and tilting functors. Funktsional. Anal. i Prilozhen., 33(2):31–42, 95, 1999.
- [Per1] Ozren Perše. Vertex operator algebras associated to type B affine Lie algebras on admissible half-integer levels. J. Algebra, 307(1):215–248, 2007.
- [Per2] Ozren Perše. Vertex operator algebras associated to certain admissible modules for affine Lie algebras of type A. Glas. Mat. Ser. III, 43(63)(1):41–57, 2008.
- [Pet] D. Peterson. Quantum cohomology of G/P,. Lecture Notes, Cambridge, MA, Spring: Massachusetts Institute of Technology, 1997.
- [Zhu] Yongchang Zhu. Modular invariance of characters of vertex operator algebras. J. Amer. Math. Soc., 9(1):237–302, 1996.

Research Institute for Mathematical Sciences, Kyoto University, Kyoto 606-8502 JAPAN

E-mail address: arakawa@kurims.kyoto-u.ac.jp