COMPLEX CROSS-RATIOS AND THE PTOLEMAEAN INEQUALITY

IOANNIS D. PLATIS

ABSTRACT. We use Korányi–Reimann complex cross–ratios to prove the Ptolemaean inequality and the Theorem of Ptolemaeus in the setting of the boundary of complex hyperbolic space and the first Heisenberg group.

1. INTRODUCTION

The Theorem of Ptolemaeus in planar Euclidean geometry states that the product of the euclidean lengths of the diagonals of an inscribed quadrilateral equals to the sum of the products of the euclidean lengths of its opposite sides. When one vertex of the quadrilateral does not lie on the circle passing from the other three vertices, then we have inequality, known as the Ptolemaean inequality.

The Ptolemaean inequality and its generalisation to various spaces has recently been the study of many authors, for example see the innovative paper of S. Buyalo and V. Schroeder, [1] and the work of S.M. Buckley, K. Falk and D.J. Wraith in CAT(0) spaces, [2]. In the present paper we use the Korányi–Reimann complex cross–ratios to give a proof of the Ptolemaean Inequality and the Theorem of Ptolemaeus in the boundary of complex hyperbolic space $\partial \mathbf{H}_{\mathbb{C}}^2$ (Theorem 3.1) and accordingly in the Heisenberg group (Theorem 3.2).

2. Preliminaries

2.1. Complex Hyperbolic Space. Let $\mathbb{C}^{2,1}$ be the vector space \mathbb{C}^3 with the Hermitian form of signature (2,1) given by

$$\langle \mathbf{z}, \mathbf{w} \rangle = \mathbf{w}^* J \mathbf{z} = z_1 \overline{w}_3 + z_2 \overline{w}_2 + z_2 \overline{w}_1$$

with matrix

$$J = \left[\begin{array}{rrr} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{array} \right].$$

We consider the following subspaces of $\mathbb{C}^{2,1}$:

$$V_{-} = \left\{ \mathbf{z} \in \mathbb{C}^{2,1} : \langle \mathbf{z}, \mathbf{z} \rangle < 0 \right\},$$
$$V_{0} = \left\{ \mathbf{z} \in \mathbb{C}^{2,1} - \{\mathbf{0}\} : \langle \mathbf{z}, \mathbf{z} \rangle = 0 \right\}.$$

Let $\mathbb{P} : \mathbb{C}^{2,1} - \{0\} \longrightarrow \mathbb{C}P^2$ be the canonical projection onto complex projective space. Then complex hyperbolic space $\mathbf{H}^2_{\mathbb{C}}$ is defined to be $\mathbb{P}V_-$ and its boundary $\partial \mathbf{H}^2_{\mathbb{C}}$ is $\mathbb{P}V_0$. Specifically, $\mathbb{C}^{2,1} - \{\mathbf{0}\}$ may be covered with three charts H_1 , H_2 , H_3 where H_j comprises those points in $\mathbb{C}^{2,1} - \{\mathbf{0}\}$ for which $z_j \neq 0$. It is clear that V_- is contained in H_3 . The canonical projection from H_3 to \mathbb{C}^2 is given by $\mathbb{P}(\mathbf{z}) = (z_1/z_3, z_2/z_3) = z$. Therefore we can write $\mathbf{H}^2_{\mathbb{C}} = \mathbb{P}(V_-)$ as

$$\mathbf{H}_{\mathbb{C}}^{2} = \left\{ (z_{1}, z_{2}) \in \mathbb{C}^{2} : 2\Re(z_{1}) + |z_{2}|^{2} < 0 \right\}.$$

Date: 12 May 2012.

I.D. PLATIS

There are distinguished points in V_0 which we denote by **o** and ∞ :

$$\mathbf{o} = \begin{bmatrix} 0\\0\\1 \end{bmatrix}, \quad \infty = \begin{bmatrix} 1\\0\\0 \end{bmatrix}.$$

Then $V_0 - \{\infty\}$ is contained in H_3 and $V_0 - \{\mathbf{o}\}$ (in particular ∞) is contained in H_1 . Let $\mathbb{P}\mathbf{o} = o$ and $\mathbb{P}\infty = \infty$. Then we can write $\partial \mathbf{H}_{\mathbb{C}}^2 = \mathbb{P}(V_0)$ as

$$\partial \mathbf{H}_{\mathbb{C}}^2 - \{\infty\} = \{(z_1, z_2) \in \mathbb{C}^2 : 2\Re(z_1) + |z_2|^2 = 1\}.$$

In particular $o = (0,0) \in \mathbb{C}^2$. In this manner, $\mathbf{H}^2_{\mathbb{C}}$ is the Siegel domain in \mathbb{C}^2 ; see [6].

Conversely, given a point z of $\mathbb{C}^2 = \mathbb{P}(H_3) \subset \mathbb{C}P^2$ we may lift $z = (z_1, z_2)$ to a point z in $H_3 \subset \mathbb{C}^{2,1}$, called the standard lift of z, by writing z in non-homogeneous coordinates as

$$\mathbf{z} = \begin{bmatrix} z_1 \\ z_2 \\ 1 \end{bmatrix}.$$

The Bergman metric on $\mathbf{H}^2_{\mathbb{C}}$ is defined by the distance function ρ given by the formula

$$\cosh^2\left(\frac{\rho(z,w)}{2}\right) = \frac{\langle \mathbf{z}, \mathbf{w} \rangle \langle \mathbf{w}, \mathbf{z} \rangle}{\langle \mathbf{z}, \mathbf{z} \rangle \langle \mathbf{w}, \mathbf{w} \rangle} = \frac{\left| \langle \mathbf{z}, \mathbf{w} \rangle \right|^2}{|\mathbf{z}|^2 |\mathbf{w}|^2}$$

where \mathbf{z} and \mathbf{w} in V_{-} are the standard lifts of z and w in $\mathbf{H}^{2}_{\mathbb{C}}$ and $|\mathbf{z}| = \sqrt{-\langle \mathbf{z}, \mathbf{z} \rangle}$. Alternatively,

$$ds^{2} = -\frac{4}{\langle \mathbf{z}, \mathbf{z} \rangle^{2}} \det \begin{bmatrix} \langle \mathbf{z}, \mathbf{z} \rangle & \langle d\mathbf{z}, \mathbf{z} \rangle \\ \langle \mathbf{z}, d\mathbf{z} \rangle & \langle d\mathbf{z}, d\mathbf{z} \rangle \end{bmatrix}.$$

The holomorphic sectional curvature of $\mathbf{H}_{\mathbb{C}}^2$ equals to -1 and its real sectional curvature is pinched between -1 and -1/4.

2.1.1. Isometries, complex lines, Lagrangian planes. Let U(2, 1) be the group of unitary matrices for the Hermitian form $\langle \cdot, \cdot \rangle$. Each such matrix A satisfies the relation $A^{-1} = JA^*J$ where A^* is the Hermitian transpose of A.

The full group of holomorphic isometries of complex hyperbolic space is the projective unitary group PU(2,1) = U(2,1)/U(1), where $U(1) = \{e^{i\theta}I, \theta \in [0,2\pi)\}$ and I is the 3×3 identity matrix. We find sometimes convenient to consider instead the group SU(2,1) of matrices which are unitary with respect to $\langle \cdot, \cdot \rangle$, and have determinant 1. Therefore $PU(2,1) = SU(2,1)/\{I, \omega I, \omega^2 I\}$, where ω is a non real cube root of unity, and so SU(2,1) is a 3-fold covering of PU(2,1).

A complex line is an isometric image of the embedding of $\mathbf{H}^1_{\mathbb{C}}$ into $\mathbf{H}^2_{\mathbb{C}}$. A Lagrangian plane is an isometric image of $\mathbf{H}^2_{\mathbb{R}}$ into $\mathbf{H}^2_{\mathbb{C}}$.

2.2. The boundary–Heisenberg group. A finite point z is in the boundary of the Siegel domain if its standard lift to $\mathbb{C}^{2,1}$ is z where

$$\mathbf{z} = \begin{bmatrix} z_1 \\ z_2 \\ 1 \end{bmatrix}$$
 where $z_1 + \bar{z}_1 + |z_2|^2 = 0.$

We write $z = z_2/\sqrt{2} \in \mathbb{C}$ and this condition becomes $2\Re(z_1) = -2|z|^2$. Hence we may write $z_1 = -|z|^2 + it$ for $t \in \mathbb{R}$. That is for $z \in \mathbb{C}$ and $t \in \mathbb{R}$:

$$\mathbf{z} = \begin{bmatrix} -|z|^2 + it \\ \sqrt{2}z \\ 1 \end{bmatrix}$$

Therefore we may identify the boundary of the Siegel domain with the one point compactification of $\mathbb{C} \times \mathbb{R}$.

The action of the stabiliser of infinity $\operatorname{Stab}(\infty)$ gives to the set of these points the structure of a non Abelian group. This is the Heisenberg group \mathfrak{H} which is $\mathbb{C} \times \mathbb{R}$ with group law

$$(z,t) * (w,s) = (z+w,t+s+2\Im(\overline{w}z)).$$

The Heisenberg norm (Korányi gauge) is given by

$$|(z,t)| = ||z|^2 - it|^{1/2}.$$

From this norm we obtain a metric, the Korányi–Cygan (K–C) metric, on \mathfrak{H} by the relation

$$d_K((z_1,t_1),(z_2,t_2)) = |(z_1,t_1)^{-1} * (z_2,t_2)|$$

Or, in other words

$$d_K\left((z_1,t_1),\,(z_2,t_2)\right) = \left||z_1-z_2|^2 - it_1 + it_2 - 2i\Im(z_1\bar{z}_2)\right|^{1/2}.$$

By taking the standard lift of points on $\partial \mathbf{H}_{\mathbb{C}}^2 - \{\infty\}$ to $\mathbb{C}^{2,1}$ we can write the K–C metric as:

$$d_K\left((z_1, t_1), (z_2, t_2)\right) = \left| \left\langle \begin{bmatrix} -|z_1|^2 + it_1\\ \sqrt{2}z_1\\ 1 \end{bmatrix}, \begin{bmatrix} -|z_2|^2 + it_2\\ \sqrt{2}z_2\\ 1 \end{bmatrix} \right\rangle \right|^{1/2}$$

The K–C metric is invariant under

- (1) the left action of \mathfrak{H} , $(z,t) \to (\zeta, s) * (z,t)$;
- (2) the Heisenberg translations $(z,t) \mapsto (z,t+s), s \in \mathbb{R};$
- (3) the rotations about the vertical axis $(z,t) \mapsto (ze^{i\phi},t), \phi \in \mathbb{R}$.

The group $\text{Isom}(\mathfrak{H}, d_K)$ of *Heisenberg isometries*, is thus represented by the group consisting of matrices of the form

(2.1)
$$\begin{bmatrix} 1 & -\sqrt{2}\zeta e^{i\phi} & -|\zeta|^2 + is \\ 0 & e^{i\phi} & \sqrt{2}\zeta \\ 0 & 0 & 1 \end{bmatrix}.$$

The K–C metric is also scaled up to multiplicative constants by the action of Heisenberg dilations $(z,t) \mapsto (rz, r^2t), r \in \mathbb{R}_*$. The group

$$\operatorname{Sim}(\mathfrak{H}, d_K) = \mathbb{R} \times \mathrm{U}(1) \times \mathfrak{H}$$

acting on \mathfrak{H} is called the group of Heisenberg similarities.

2.2.1. \mathbb{R} -circles and \mathbb{C} -circles. \mathbb{R} -circles are boundaries of Lagrangian planes and \mathbb{C} -circles are boundaries of complex lines. They come in two flavours, infinite ones (i.e. containing the point at infinity) and finite ones. We refer to [6] for more more details about these curves.

I.D. PLATIS

2.2.2. Cross-ratios. Given a quadruple of distinct points $\mathbf{p} = (p_1, p_2, p_3, p_4)$ in $\partial \mathbf{H}^2_{\mathbb{C}}$, their complex cross-ratio as defined by Korányi and Reimann in [7] is

$$\mathbb{X}(p_1, p_2, p_3, p_4) = \frac{\langle \mathbf{p}_3, \mathbf{p}_1 \rangle \langle \mathbf{p}_4, \mathbf{p}_2 \rangle}{\langle \mathbf{p}_4, \mathbf{p}_1 \rangle \langle \mathbf{p}_3, \mathbf{p}_2 \rangle},$$

where \mathbf{p}_i are lifts of p_i , i = 1, 2, 3, 4. The cross-ratio is independent of the choice of lifts and remains invariant under the diagonal action of PU(2, 1). We stress here that for points in the Heisenberg group, the square root of its absolute value is

$$|\mathbb{X}(p_1, p_2, p_3, p_4)|^{1/2} = \frac{d_K(p_4, p_2) \cdot d_K(p_3, p_1)}{d_K(p_4, p_1) \cdot d_K(p_3, p_2)}$$

2.3. Cross-ratio variety. Given a quadruple $\mathfrak{p} = (p_1, p_2, p_3, p_4)$ of distinct points in the boundary, all possible permutations of points gives us 24 complex cross-ratios corresponding to \mathfrak{p} . Due to symmetries, see [4], Falbel showed that all cross-ratios corresponding to a quadruple of points depend on three cross-ratios which satisfy two real equations. The following proposition holds.

Proposition 2.1. Let $\mathfrak{p} = (p_1, p_2, p_3, p_4)$ be any quadruple of distinct points in $\partial \mathbf{H}^2_{\mathbb{C}}$. Let

$$X_1(\mathbf{p}) = X(p_1, p_2, p_3, p_4), \quad X_2(\mathbf{p}) = X(p_1, p_3, p_2, p_4), \quad X_3(\mathbf{p}) = X(p_2, p_3, p_1, p_4)$$

Then

(2.2)
$$|\mathbb{X}_2| = |\mathbb{X}_1| |\mathbb{X}_3|,$$

(2.3)
$$2|\mathbb{X}_1|^2 \Re(\mathbb{X}_3) = |\mathbb{X}_1|^2 + |\mathbb{X}_2|^2 - 2\Re(\mathbb{X}_1 + \mathbb{X}_2) + 1$$

For the proof, see for instance in [8]. The above equations define a 4–dimensional real subvariety of $\mathbb{C}^{3,1}$

Definition 2.2. The *cross-ratio variety* \mathfrak{X} is the subset of \mathbb{C}^3 at which Eqs. 2.2 and 2.3 are satisfied.

3. PTOLEMAEAN INEQUALITY AND PTOLEMAEUS' THEOREM

In this section we prove an S^3 version of the Ptolemaean inequality and Ptolemaeus Theorem respectively, which are derived almost immediately from the properties of complex cross-ratios.

Theorem 3.1. (Ptolemaean inequality and Ptolemaeus' Theorem in $\partial \mathbf{H}^2_{\mathbb{C}}$) Let $\mathfrak{p} = (p_1, p_2, p_3, p_4)$ a quadruple of distinct points in $\partial \mathbf{H}^2_{\mathbb{C}}$ and $\mathbb{X}_i = \mathbb{X}_i(\mathfrak{p})$, i = 1, 2, 3, its corresponding complex cross-ratios. Then the following inequalities hold:

(3.1)
$$|\mathbb{X}_1|^{1/2} + |\mathbb{X}_2|^{1/2} \ge 1$$
, and $-1 \le |\mathbb{X}_1|^{1/2} - |\mathbb{X}_2|^{1/2} \le 1$.

Each inequality 3.1 holds if and only if all four points of \mathfrak{p} lie in an \mathbb{R} -circle. Then, $\mathbb{X}_i > 0$, i = 1, 2 and

(1) $\mathbb{X}_{1}^{1/2} - \mathbb{X}_{2}^{1/2} = 1$ if p_{1} and p_{3} separate p_{2} and p_{4} ; (2) $\mathbb{X}_{2}^{1/2} - \mathbb{X}_{1}^{1/2} = 1$ if p_{1} and p_{2} separate p_{3} and p_{4} ; (3) $\mathbb{X}_{1}^{1/2} + \mathbb{X}_{2}^{1/2} = 1$ if p_{1} and p_{4} separate p_{2} and p_{3} .

¹We note here that Equalities 2.2 and 2.3 also hold in the case of quadruples of distinct points in the boundary of the quaternionic hyperbolic space $\partial \mathbf{H}_{\mathbb{H}}^2$. This is identified with the one point compactification of the quaternionic Heisenberg group, see [9].

Proof. From the defining equations of \mathfrak{X} we have

Therefore,

 $(|\mathbb{X}_1|^{1/2} - |\mathbb{X}_2|^{1/2} - 1) \cdot (|\mathbb{X}_1|^{1/2} - |\mathbb{X}_2|^{1/2} + 1) \cdot (|\mathbb{X}_1|^{1/2} + |\mathbb{X}_2|^{1/2} - 1) \cdot (|\mathbb{X}_1|^{1/2} + |\mathbb{X}_2|^{1/2} + 1) \le 0$ and since $|\mathbb{X}_1|^{1/2} + |\mathbb{X}_2|^{1/2} + 1 > 0$, this reduces to

$$(|\mathbb{X}_1|^{1/2} - |\mathbb{X}_2|^{1/2} - 1) \cdot (|\mathbb{X}_1|^{1/2} - |\mathbb{X}_2|^{1/2} + 1) \cdot (|\mathbb{X}_1|^{1/2} + |\mathbb{X}_2|^{1/2} - 1) \le 0.$$

Suppose that $(|\mathbb{X}_1|^{1/2} - |\mathbb{X}_2|^{1/2})^2 > 1$; then applying triangle inequality we are reduced to a contradiction since we must also have $(|\mathbb{X}_1|^{1/2} - |\mathbb{X}_2|^{1/2})^2 > 1$. Therefore, $(|\mathbb{X}_1|^{1/2} - |\mathbb{X}_2|^{1/2})^2 \leq 1$ and $(|\mathbb{X}_1|^{1/2} - |\mathbb{X}_2|^{1/2})^2 \geq 1$ which proves the Ptolemaean inequality.

To prove Ptolemaeus' Theorem, suppose first that one of the inequalities holds as an equality. Then,

 $(|\mathbb{X}_1|^{1/2} - |\mathbb{X}_2|^{1/2} - 1) \cdot (|\mathbb{X}_1|^{1/2} - |\mathbb{X}_2|^{1/2} + 1) \cdot (|\mathbb{X}_1|^{1/2} + |\mathbb{X}_2|^{1/2} - 1) \cdot (|\mathbb{X}_1|^{1/2} + |\mathbb{X}_2|^{1/2} + 1) = 0,$ which is equivalent to

$$(|\mathbb{X}_1| - |\mathbb{X}_2|)^2 = 2(|\mathbb{X}_1| + |\mathbb{X}_2|) - 1.$$

Since $(|\mathbb{X}_1| - |\mathbb{X}_2|)^2 \le 2\Re(\mathbb{X}_1 + \mathbb{X}_2) - 1$, we have $|\mathbb{X}_1| + |\mathbb{X}_2| \le \Re(\mathbb{X}_1 + \mathbb{X}_2)$ and therefore $0 \ge \Re(\mathbb{X}_2) - |\mathbb{X}_2| \ge |\mathbb{X}_1| - \Re(\mathbb{X}_1) \ge 0.$

Thus X_1, X_2 are positive. Now from Equation 2.3 it follows

usi

$$2\mathbb{X}_{1}^{2}\Re(\mathbb{X}_{3}) = \mathbb{X}_{1}^{2} + \mathbb{X}_{2}^{2} - 2\mathbb{X}_{1} - 2\mathbb{X}_{2} + 1$$
$$= \mathbb{X}_{1}^{2} + \mathbb{X}_{2}^{2} - (\mathbb{X}_{1} - \mathbb{X}_{2})^{2} = 2\mathbb{X}_{1}\mathbb{X}_{2}$$
ng Eq. 2.2 = $2\mathbb{X}_{1}^{2}|\mathbb{X}_{3}|,$

and thus $X_3 > 0$. By Proposition 5.12 (ii) in [8] all points have to lie in a Lagrangian plane.

Conversely, if all points lie on an \mathbb{R} -circle, then we have equality in one of the inequalities, see Proposition 5.14 of [8]. The last statement is just Corollary 5.15 of [8].

Let now $\mathfrak{p} = (p_1, p_2, p_3, p_4)$ a quadruple of distinct points in the Heisenberg group \mathfrak{H} . Inequalities 3.1 of Theorem 3.1 can be written as

$$(3.2) d_K(p_2, p_3) \cdot d_K(p_1, p_4) \le d_K(p_2, p_4) \cdot d_K(p_1, p_3) + d_K(p_1, p_2) \cdot d_K(p_3, p_4),$$

$$(3.3) d_K(p_1, p_3) \cdot d_K(p_2, p_4) \le d_K(p_1, p_2) \cdot d_K(p_3, p_4) + d_K(p_2, p_3) \cdot d_K(p_1, p_4)$$

$$(3.4) d_K(p_1, p_2) \cdot d_K(p_3, p_4) \le d_K(p_1, p_3) \cdot d_K(p_2, p_4) + d_K(p_2, p_3) \cdot d_K(p_1, p_4)$$

As a corollary of the above discussion we derive the following.

I.D. PLATIS

Theorem 3.2. (Ptolemaean inequality and Ptolemaeus' Theorem in the Heisenberg group) The Korányi–Cygan metric d_K in the Heisenberg group \mathfrak{H} satisfies the Ptolemaean inequality: for each quadruple of points $\mathfrak{p} = (p_1, p_2, p_3, p_4)$, inequalities 3.2, 3.3 and 3.4 hold. Moreover, each of these inequalities hold as an equality if and only if all points lie in an \mathbb{R} -circle. Explicitly,

- (1) $d_K(p_2, p_3) \cdot d_K(p_1, p_4) = d_K(p_2, p_4) \cdot d_K(p_1, p_3) + d_K(p_1, p_2) \cdot d_K(p_3, p_4)$ if and only if all points lie in an \mathbb{R} -circle and p_1 and p_4 separate p_2 and p_3 ;
- (2) $d_K(p_1, p_3) \cdot d_K(p_2, p_4) = d_K(p_1, p_2) \cdot d_K(p_3, p_4) + d_K(p_2, p_3) \cdot d_K(p_1, p_4)$ if and only if all points lie in an \mathbb{R} -circle and p_1 and p_3 separate p_2 and p_4 ;
- (3) $d_K(p_1, p_2) \cdot d_K(p_3, p_4) = d_K(p_1, p_3) \cdot d_K(p_2, p_4) + d_K(p_2, p_3) \cdot d_K(p_1, p_4)$ if and only if all points lie in an \mathbb{R} -circle and p_1 and p_2 separate p_3 and p_4 .

References

- S. Buyalo & V. Schroeder; Möbius structures and Ptolemy spaces: boundary at innity of complex hyperbolic spaces. arXiv:1012.1699v1 [math.MG]
- [2] S.M. Buckley, K. Falk & D.J. Wraith; Ptolemaic inequality and CAT(0) Glasgow Math. J. 51 (2009) 301314.
- [3] H. Cunha & N. Gusevskii; On the moduli space of quadruples of points in the boundary of complex hyperbolic space. Transform. Groups 15 (2010), no. 2, 261283.
- [4] E. Falbel; Geometric structures associated to triangulations as fixed point sets of involutions. Topol. and its Appl. 154 (2007), no. 6, 1041-1052. Corrected version in www.math.jussieu.fr/~falbel
- [5] E. Falbel & I.D. Platis; The PU(2,1) configuration space of four points in S³ and the cross-ratio variety. Math. Ann. 340 (2008), no. 4, 935–962.
- [6] W. Goldman; Complex hyperbolic geometry. Oxford Mathematical Monographs. Oxford Science Publications. The Clarendon Press, Oxford University Press, New York, 1999.
- [7] A. Korányi & H.M. Reimann; The complex cross ratio on the Heisenberg group. Enseign. Math. (2) 33 (1987), no. 3-4, 291–300.
- [8] J.R. Parker & I.D. Platis; Complex hyperbolic Fenchel-Nielsen coordinates. Topology (47) (2008), 101–135.
- [9] I.D. Platis; The Psp(2,1)-configuration space of four points in S^7 . In preparation.

DEPARTMENT OF MATHEMATICS, UNIVERITY OF CRETE, GREECE *E-mail address*: jplatis@math.uoc.gr