Non-finite axiomatizability of Dynamic Topological Logic

David Fernández-Duque

Sevilla

November 23, 2018

Abstract

Dynamic topological logic (\mathcal{DTL}) is a polymodal logic designed for reasoning about *dynamic topological systems*. These are pairs $\langle X, f \rangle$, where X is a topological space and $f : X \to X$ is continuous. \mathcal{DTL} uses a language L which combines the topological $S4$ modality \square with temporal operators from linear temporal logic.

Recently, I gave a sound and complete axiomatization DTL^* for an extension of the logic to the language L^* , where \Diamond is allowed to act on finite sets of formulas and is interpreted as a tangled closure operator. No complete axiomatization is known over L, although one proof system, which we shall call KM, was conjectured to be complete by Kremer and Mints.

In this paper we show that, given any language L' such that $\mathsf{L} \subseteq \mathsf{L}' \subseteq$ L^* , the set of valid formulas of L' is not finitely axiomatizable. It follows, in particular, that KM is incomplete.

1 Introduction

Finding a transparent axiomatization for Dynamic Topological Logic (\mathcal{DTL}) has been an elusive open problem since 2005, when one (which we shall call KM) was proposed by Kremer and Mints in [\[13\]](#page-21-0) without establishing its completeness. In [\[9\]](#page-20-0) I offered a complete axiomatization, not over the language L used in [\[13\]](#page-21-0), but rather in an extended language L^* which allowed the modal \Diamond to be applied to finite sets of formulas. It was then interpreted as a 'tangled closure' operator (see Section [3\)](#page-3-0). The resulting logic is called DTL^* .

However, the fact that DTL^* used the unfamiliar 'tangled closure' operation and was substantially less intuitive than KM left the completeness of the latter as a relevant open problem. Actually, the only motivation given in [\[9\]](#page-20-0) for passing to an extended language was, There is a completeness proof which works in the extended language but not in the original one; a valid, but not terribly compelling, reason.

The results in this paper will show that indeed the use of the tangled closure is an essential part of this axiomatization, and cannot be removed without extending KM, although it is not clear what such an extension should look like. In fact, we prove more. We show that, given $k < \omega$, there is a formula Trouble^k \in L such that Trouble^k is derivable in DTL^{*} only by using formulas of the form $\Diamond \Gamma$, where Γ has at least k elements. This shows that \mathcal{DTL} can be written as a strictly increasing sequence of theories and hence is not finitely axiomatizable; it follows, in particular, that KM is incomplete.

1.1 Previous work on $DT\mathcal{L}$

Dynamic topological logic (\mathcal{DTL}) combines the topological S4 with Linear Temporal Logic. The 'topological interior' interpretation of modal logic was already studied by Tarski, McKinsey and others around the 1940s [\[17\]](#page-21-1) and more recently in works like [\[2,](#page-20-1) [12,](#page-21-2) [15\]](#page-21-3). Temporal logic also has a long history, having been studied by Prior before 1960 [\[16\]](#page-21-4) and received substantial attention since; see [\[14\]](#page-21-5) for a nice overview.

The purpose of $DT\mathcal{L}$ is to reason about *dynamic topological systems* (dts's); these are pairs $\langle X, f \rangle$, where X is a topological space and $f : X \to X$ is a continuous function, and shall be discussed in greater detail in Section [4.](#page-3-1) Dynamic Topological Logic was originally introduced as a bimodal logic in [\[1\]](#page-20-2), where it was called S4C. In our notation, it uses the 'interior' modality \Box , interpreted topologically, and 'next-time' modality f , interpreted as a preimage operator; see Section [4](#page-3-1) for details. The logic S4C is a rather well-behaved modal logic; it is decidable, axiomatizable and has the finite model property, all of which was established in [\[1\]](#page-20-2). Later, [\[13\]](#page-21-0) showed that a variant, called S4H, is complete for the class of dynamical systems where f is a homeomorphism.

Also in [\[13\]](#page-21-0), it was noted that by adding the infinitary temporal modality 'henceforth' (here denoted [f]), one could reason about long-term behavior of dts's, capturing phenomena such as topological recurrence. Thus they introduced an extension of S4C, which we denote $\mathcal{D}\mathcal{TL}$. $\mathcal{D}\mathcal{TL}$ turned out to behave much worse than S4C; it was proven to be undecidable in [\[10\]](#page-20-3), and in [\[11\]](#page-20-4) it was also shown that, if we restrict to the case where f is a homeomorphism, then the logic becomes non-axiomatizable. Fortunately, with arbitrary continuous functions the logic turned out to be recursively enumerable [\[6\]](#page-20-5), but the only currently known axiomatization is over L^* [\[9\]](#page-20-0).

This axiomatization uses the fact, first observed in [\[4\]](#page-20-6), that L^* is more expressive than L. There, it is shown that, over the class of finite S4 models, L^{*}_○ (i.e., the fragment of L^{*} without temporal modalities) is equally expressive to the bisimulation-invariant fragment of both first-order logic and monadic second-order logic, while L_{\lozenge} is weaker. This added expressive power is used in an important way in [\[9\]](#page-20-0), although it is not proven that such an extension is necessary.

1.2 Layout of the paper

Sections [2-](#page-2-0)[5](#page-4-0) give a general review of dynamic topological logic and the known results relevant for this paper. In Section [2,](#page-2-0) we introduce topological spaces and show how one can see preorders as a special case. Section [3](#page-3-0) gives the main properties of the tangled closure operator, an important part of \mathcal{DTL}^* , introduced in Section [4.](#page-3-1) Section [5](#page-4-0) then reviews the axiomatization from [\[9\]](#page-20-0) and defines some important sublogics.

Section [6](#page-7-0) introduces tangled bisimulations, which are based on those pre-sented in [\[4\]](#page-20-6). These are used to show that $\left\langle \right\rangle_{i=1}^{k+1}$ $\sum_{i=1}^{n+1} \gamma_i$ cannot in general be defined using exclusively k-adic occurrences of \Diamond .

Section [7](#page-10-0) defines the formulas $\texttt{Trouble}^k$ which are derivable in \texttt{DTL}^k , as well as other formulas which are useful for our purposes. Finally, Section [8](#page-12-0) shows that, indeed, DTL^k is consistent with $\neg \text{Trouble}^{k+1}$, thus stratifying Dynamic Topological Logic into a strictly increasing sequence of theories, from which it follows that it is not finitely axiomatizable.

2 Topologies and preorders

In this section we shall very briefly review some basic notions from topology. As is well-known, topological spaces provide an interpretation of the modal logic S4, generalizing its familiar Kripke semantics.

Let us recall the definition of a topological space:

Definition 2.1. A topological space is a pair $\mathfrak{X} = \langle |\mathfrak{X}|, \mathcal{T}_{\mathfrak{X}} \rangle$, where $|\mathfrak{X}|$ is a set and $\mathcal{T}_{\mathfrak{X}}$ a family of subsets of $|\mathfrak{X}|$ satisfying

- 1. $\varnothing, |\mathfrak{X}| \in \mathcal{T}_{\mathfrak{X}};$
- 2. if $U, V \in \mathcal{T}_{\mathfrak{X}}$ then $U \cap V \in \mathcal{T}_{\mathfrak{X}}$ and
- 3. if $\mathcal{O} \subseteq \mathcal{T}_{\mathfrak{X}}$ then $\bigcup \mathcal{O} \in \mathcal{T}_{\mathfrak{X}}$.

The elements of $\mathcal{T}_{\mathfrak{X}}$ are called open sets. Complements of open sets are closed sets.

Given a set $A \subseteq |\mathfrak{X}|$, its *interior*, denoted A° , is defined by

$$
A^{\circ} = \bigcup \{ U \in \mathcal{T}_{\mathfrak{X}} : U \subseteq A \}.
$$

Dually, we define the closure \overline{A} as $|\mathfrak{X}| \setminus (|\mathfrak{X}| \setminus A)^\circ$; this is the smallest closed set containing A.

Topological spaces generalize transitive, reflexive Kripke frames. Recall that these are pairs $\mathfrak{W} = \langle |\mathfrak{W}|, \preccurlyeq_{\mathfrak{W}} \rangle$, where $\preccurlyeq_{\mathfrak{W}}$ is a preorder on the set $|\mathfrak{W}|$. We will write \preccurlyeq instead of $\preccurlyeq_{\mathfrak{M}}$ whenever this does not lead to confusion.

To see a preorder as a special case of a topological space, define

$$
\downarrow w = \{v : v \preccurlyeq w\}.
$$

Then consider the topology $\mathcal{T}_{\preccurlyeq}$ on $|\mathfrak{W}|$ given by setting $U \subseteq |\mathfrak{W}|$ to be open if and only if, whenever $w \in U$, we have $\downarrow w \subseteq U$ (so that the sets of the form $\downarrow w$ provide a basis for $\mathcal{T}_{\preccurlyeq}$). A topology of this form is a preorder topology^{[1](#page-3-2)}.

Throughout this text we will often identify preorders with their corresponding topologies, and many times do so tacitly.

We will also use the notation

- $w \prec v$ for $w \preccurlyeq v$ but $v \not\preccurlyeq w$ and
- $w \approx v$ for $w \preccurlyeq v$ and $v \preccurlyeq w$.

The relation \approx is an equivalence relation; the equivalence class of a point $x \in |\mathfrak{W}|$ is usually called a *cluster*, and we will denote it by [x].

3 The tangled closure

The tangled closure is an important component of DTL^{*}. It was introduced in [\[4\]](#page-20-6) for Kripke frames and has also appeared in [\[7,](#page-20-7) [8,](#page-20-8) [9,](#page-20-0) [5\]](#page-20-9).

Definition 3.1. Let \mathfrak{X} be a topological space and $\mathcal{S} \subseteq 2^{|\mathfrak{X}|}$.

Given $E \subseteq |\mathfrak{X}|$, we say S is tangled in E if, for all $A \in \mathcal{S}$, $A \cap E$ is dense in E.

We define the tangled closure of S, denoted S^* , to be the union of all sets E such that S is tangled in E .

It is important for us to note that the tangled closure is defined over any topological space; however, we will often be concerned with locally finite preorders in this paper. Here, the tangled closure is relatively simple.

Lemma 3.1. Let $\langle S, \preccurlyeq \rangle$ be a finite preorder, $x \in S$ and $\mathcal{O} \subseteq 2^S$. Then, $x \in \mathcal{O}^*$ if and only if there exist $\langle y_A \rangle_{A \in \mathcal{O}}$ such that $y_A \in A$, $y_A \preccurlyeq x$ for all $A \in \mathcal{O}$ and $y_A \approx y_B$ for all $A, B \in \mathcal{O}$.

Proof. A proof can be found in any of [\[7,](#page-20-7) [8,](#page-20-8) [9,](#page-20-0) [5\]](#page-20-9).

 \Box

4 Dynamic Topological Logic

The language L^{*} is built from propositional variables in a countably infinite set PV using the Boolean connectives \wedge and \neg (all other connectives are to be defined in terms of these), the unary modal operators f ('next') and $[f]$ ('henceforth'), along with a polyadic modality \Diamond which acts on finite sets, so that if Γ is a finite set of formulas then $\Diamond \Gamma$ is also a formula. Note that this is a modification of the usual language of \mathcal{DTL} , where \diamondsuit acts on single formulas only. We write \square as a shorthand for $\neg \Diamond \neg$; similarly, $\langle f \rangle$ denotes the dual of $[f]$.

 1 Or, more specifically, a *downset topology*. Note that I stray from convention, since most authors use the upset topology here, but I find this presentation more natural. This will later be reflected in the semantics for $\Box.$

We also write $\Diamond \gamma$ instead of $\Diamond \{\gamma\}$; its meaning is identical to that of the usual S4 modality [\[5\]](#page-20-9). We will often write $\left\langle \right\rangle_{n}^{N}$ $\sum_{n=1}^{N} \gamma_i$ instead of $\sqrt{\{\gamma_i\}}_{1 \leq i \leq N}$.

Given a formula ϕ , the *depth* of ϕ , denoted dpt(ϕ), is the modal nesting depth of ϕ , while its width, wdt (ϕ) , denotes the maximal k such that ϕ has a subformula of the form $\left\langle \right\rangle _{k}^{k}$ $\sum_{i=1}^k \gamma_i$. For $k < \omega$, L^k denotes the sublanguage of L^* where all formulas have width at most k. Thus, in particular, $L = L¹$.

Formulas of L [∗] are interpreted on dynamical systems over topological spaces, or dynamic topological systems.

Definition 4.1. A weak dynamic topological system (dts) is a triple

$$
\mathfrak{X}=\left\langle |\mathfrak{X}|,\mathcal{T}_{\mathfrak{X}},f_{\mathfrak{X}}\right\rangle ,
$$

where $\langle |\mathfrak{X}|, \mathcal{T}_{\mathfrak{X}} \rangle$ is a topological space and

$$
f_{\mathfrak{X}}:|\mathfrak{X}|\to|\mathfrak{X}|.
$$

If further $f_{\mathfrak{X}}$ is continuous^{[2](#page-4-1)}, we say \mathfrak{X} is a dynamical system.

Definition 4.2. Given a (weak) dynamic topological system \mathfrak{X} , a valuation on $\mathfrak X$ is a function

 $\llbracket \cdot \rrbracket : \mathsf{L}^* \to 2^{|\mathfrak{X}|}$

satisfying

$$
\begin{aligned}\n\llbracket \alpha \wedge \beta \rrbracket_{\mathfrak{X}} &= \llbracket \alpha \rrbracket_{\mathfrak{X}} \cap \llbracket \beta \rrbracket_{\mathfrak{X}} \\
\llbracket \neg \alpha \rrbracket_{\mathfrak{X}} &= |\mathfrak{X}| \setminus \llbracket \alpha \rrbracket_{\mathfrak{X}} \\
\llbracket f \alpha \rrbracket_{\mathfrak{X}} &= f^{-1} \llbracket \alpha \rrbracket_{\mathfrak{X}} \\
\llbracket [f] \alpha \rrbracket_{\mathfrak{X}} &= \bigcap_{n \geq 0} f^{-n} \llbracket \alpha \rrbracket_{\mathfrak{X}} \\
\llbracket \Diamond \{ \alpha_1, ..., \alpha_n \} \rrbracket_{\mathfrak{X}} &= \{ \llbracket \alpha_1 \rrbracket_{\mathfrak{X}}, ..., \llbracket \alpha_n \rrbracket_{\mathfrak{X}} \}^*\n\end{aligned}.
$$

A (weak) dynamic topological model (wdtm/dtm) is a (weak) dynamic topological system \mathfrak{X} equipped with a valuation $[\![\cdot]\!]_{\mathfrak{X}}$. We say a formula ϕ is valid
on \mathfrak{X} if $[\![\phi]\!] = |\mathfrak{X}|$ and write $\mathfrak{X} \models \phi$. If a formula ϕ is valid on every dynamic on \mathfrak{X} if $\llbracket \phi \rrbracket_{\mathfrak{X}} = |\mathfrak{X}|$, and write $\mathfrak{X} \models \phi$. If a formula ϕ is valid on every dynamic topological model, then we write $\models \phi$. \mathcal{DTL} is the set of valid formulas of L under this interpretation, while \mathcal{DTL}^* denotes the set of valid formulas of L^* .

We will often write $\langle \mathfrak{X}, x \rangle \models \phi$ instead of $x \in [\![\phi]\!]_{\mathfrak{X}}$.

5 The axiomatization

We shall distinguish \mathcal{DTL}^* , which is defined semantically, from DTL^* , which is a proof system. The two have the same set of theorems, but we will be interested in natural subsystems of DTL[∗] which are defined syntactically.

²That is, whenever $U \subseteq |\mathfrak{X}|$ is open, then so is $f^{-1}(U)$

Below, note that the modality f is unary, and $f\Gamma$ is merely a shorthand for ${f \gamma : \gamma \in \Gamma};$ p denotes a propositional variable and P a finite set of propositional variables. Then, the axiomatization DTL[∗] consists of the following:

Taut All propositional tautologies.

Topological axioms

$$
\begin{aligned}\n\mathsf{K} \quad & \Box(p \to q) \to (\Box p \to \Box q) \\
\mathsf{T} \quad & \Box p \to p \\
\mathsf{4} \quad & \Box p \to \Box \Box p \\
\mathsf{Fix}_{\Diamond} \quad & \Diamond P \to \bigwedge_{q \in P} \Diamond(q \land \Diamond P) \\
\mathsf{Ind}_{\Diamond} \quad & \Box \quad \bigwedge_{q \in P} (p \to \Diamond(q \land p)) \to (p \to \Diamond P) \\
& \mathsf{q} \in P\n\end{aligned}
$$

Temporal axioms

$$
\begin{aligned}\n\text{Neg}_f &\neg fp \leftrightarrow f \neg p \\
\text{And}_f &\ f(p \land q) \leftrightarrow fp \land fq \\
\text{Fix}_{[f]} &\ [f]p \to p \land f[f]p \\
\text{Ind}_{[f]} &\ [f](p \to fp) \to (p \to [f]p)\n\end{aligned}
$$

Cont^{*} $\Diamond fP \rightarrow f\Diamond P$.

Rules

MP Modus ponens

Subs
$$
\frac{\phi}{\phi[\vec{p}/\vec{\psi}]}
$$

\n $N_{\Box} \frac{\phi}{\Box \phi}$ $N_{f} \frac{\phi}{f\phi}$ $N_{[f]} \frac{\phi}{[f]\phi}$

This axiomatization is sound and complete, as proven in [\[9\]](#page-20-0):

Theorem 5.1. DTL^{*} is sound and complete for the class of dynamic topological models.

There are many subtleties in our proof system, so before continuing we should make a few remarks.

First, let us say a few words about the substitution rule. It is to be understood as 'simulataneous substitution', where \vec{p} represents a finite sequence of variables, $\bar{\psi}$ a finite sequence of formulas and each variable is replaced by the respective formula. By standard arguments, this rule preserves validity, as there is nothing in our semantics distinguishing atomic facts from complex propositions.

Further, since we are concerned with finite axiomatizability of a logic it is important to include it; otherwise, each substitution instance of any of the axioms would have to be regarded as a new axiom and the finite axiomatizability would fail for obvious reasons. Of course this is not the only possible presentation, as one can also consider axiomatizations by finitely many schemas, but here we shall consider different formulas to be different also as axioms.

With this in mind, we should note that the above axiomatization is not finite, nor can it be modified into a finite version in an obvious way. Evidently the set of all propositional tautologies can be replaced by finitely many axioms, but this is not what concerns us. Much more importantly, we need infinitely many axioms for \Diamond , and it is only in the metalanguage that we can give them a uniform presentation. In fact, the symbol P representing a finite set of propositional variables is not a symbol of L^* , where we would have to write out explicitly $\{p_1, \ldots, p_k\}$ for each given value of k.

Of particular interest is the schema Cont[∗] . This was originally named TCont; we adopt the new notation to stress that the standard 'continuity' axiom,

$$
\mathsf{Cont}^1:\Diamond fp\to f\Diamond p,
$$

is indeed a special case.

Cont^{*} is really an infinite collection of axioms. To be precise, for $k < \omega$ let

$$
\mathsf{Cont}^k = \bigotimes_{i \in [1,k]} fp_i \to f \bigotimes_{i \in [1,k]} p_i.
$$

Note that Cont^{k+1} extends Cont^k since we can always substitute p_{k+1} by p_k .

We then let DTL^k be the variant of DTL^* where $Cont^*$ is replaced by $Cont^k$. We denote derivability in DTL^k by \vdash^k . DTL^0 denotes the system with no continuity axiom.

Our goal will be to show that $\langle DTL^k \rangle_{k \leq \omega}$ gives a sequence of theories of strictly increasing strength. Since DTL^{*} is the union of these theories, it will follow as a straightforward consequence that DTL[∗] is not finitely axiomatizable. However, to do this we will need a second refinement, this time of each DTL^k .

For $n, k < \omega$, we let DTL_n^k be the subtheory of DTL^k which restricts the substitution rule in the following ways:

- 1. Subs may only be applied immediately to axioms and
- 2. if Subs is applied to Cont^k, then each p_i must be replaced by a formula with modal depth at most n .

A very easy induction on derivations shows that any proof in DTL^k may be transformed into one satisfying the above two conditions for some value of n and hence $DTL^k = \bigcup_{n < \omega} DTL_n^k$. We denote derivability in DTL_n^k by \vdash_n^k .

The reason for passing to DTL_n^k is that the substitution rule, while preserving validity, does not preserve model validity; if $\mathfrak{M} \models \phi$, it does not always follow that $\mathfrak{M} \models \phi[p/\psi]$. Later we wish to build specific models of fragments of DTL^{*}, and to check soundness for these models, DTL_n^k has the advantage that we only need to focus on substitution instances of axioms. This will become relevant in Section [8.](#page-12-0)

DTL^{*} is an extension of KM, which can be defined as follows:

Definition 5.1. The calculus KM is the restriction^{[3](#page-7-1)} of DTL^{*} to L^1 .

In KM, all appearances of \Diamond must be applied to a single formula; in particular, the axioms Fix_{\Diamond} and Ind_{\Diamond} are not present, and Cont^{*} becomes Cont¹. We should note that DTL^1 is very similar, but not identical, to KM. DTL^1 allows formulas of the form $\Diamond \Gamma$ within derivations for Γ arbitrarily large, but Cont^{*} is also replaced by Cont¹. We do have, however, that $KM \subseteq DTL^{1}$.

Later we shall show that the sequence $\langle DTL^k \rangle_{k \lt \omega}$ is strictly increasing in strength, even over L; i.e., there are formulas $\text{Trouble}^k \in L$ such that \vdash^{k+1} Trouble^{k+1} but $\not\vdash^k$ Trouble^{k+1}. These are defined in Section [7;](#page-10-0) but first, we need to define partial tangled bisimulations, the fundamental tool we shall use to prove our main results.

6 Tangled bisimulations

Our main results are based on partial bisimulation techniques. As we will be working in a polyadic system, we shall need a notion of partial bisimulation which preserves the polyadic \Diamond . Such a notion was already introduced in [\[4\]](#page-20-6); here we present a slight generalization which is more sensitive to the width of formulas. For more information on partial bisimulations, we refer the reader to a text such as [\[3\]](#page-20-10).

Definition 6.1 (Tangled partial bisimulation). Given models $\mathfrak{X}, \mathfrak{Y}, n < \omega$ and $k \leq \omega$, we define a binary relation $\omega_k^{\Omega} \subseteq |\mathfrak{X}| \times |\mathfrak{Y}|$ by inducion on n as follows.
For $n = 0$, $x \leftrightarrow 0$ *u* if and only if x and *u* satisfy the same set of atoms.

For $n = 0$, $x \triangleq_k^0 y$ if and only if x and y satisfy the same set of atoms.
Otherwise $x \triangle^{n+1} y$ if x y satisfy the same set of atoms and Otherwise, $x \triangleq_k^{n+1} y$ if x, y satisfy the same set of atoms and

- Forth whenever $m < k$ and $x_1 \approx x_2 \approx \ldots \approx x_m \preccurlyeq x$, there are $y_1 \approx y_2 \approx$ $\ldots y_m \preccurlyeq y$ such that $x_i \triangleq_k^n y_i$ for all $i \leq m$,
- Back \leq whenever $m < k$ and $y_1 \approx y_2 \approx \ldots \approx y_m \leq y$, there are $x_1 \approx x_2 \approx \ldots \approx x_m$ $x_m \preccurlyeq x$ such that $x_i \triangleq_k^n y_i$ for all $i \leq m$,

Forth $f f_{\mathfrak{X}}(x) \rightleftharpoons_k^n f_{\mathfrak{Y}}(y)$

Forth_[f] for every $m < \omega$ there is $m' < \omega$ such that $f_{\mathfrak{X}}^m(x) \rightleftharpoons_k^n f_{\mathfrak{Y}}^{m'}(y)$ and

 $\mathsf{Back}_{[f]}$ for every $m < \omega$ there is $m' < \omega$ such that $f_{\mathfrak{Y}}^{m}(y) \rightleftharpoons_{k}^{n} f_{\mathfrak{X}}^{m'}(x)$.

³Of course this description is anachronical, and it would be more accurate to think of DTL[∗] as an extension of KM.

We will write \Rightarrow^n_k when $k = \omega$; in this case there are no bounds on the clauses \rightarrow Note that there is no 'book' clause for f as it would be identical to **Forth**. for \leq . Note that there is no 'back' clause for f as it would be identical to Forth f. For purely topological structures (i.e., without the function $f_{\mathfrak{X}}$), we shall also use the analogous notion of partial bisimulation, simply removing the clauses for f , $[f]$.

When the respective structures are clear from context, we may write $x \triangleq_k^n y$ instead of $\langle \mathfrak{X}, x \rangle \Leftrightarrow_k^n \langle \mathfrak{Y}, y \rangle$.

Lemma 6.1. If φ is a formula with $\text{dpt}(\varphi) \leq n$ and $\text{wdt}(\varphi) < k$ and $\mathfrak{X}, \mathfrak{Y}$ are finite dtm's, then whenever $\langle \mathfrak{X}, x \rangle \triangleq_k^n \langle \mathfrak{Y}, y \rangle$, we have that $x \in [\![\varphi]\!]_{\mathfrak{X}}$ if and only if $y \in [\![\varphi]\!]_{\mathfrak{N}}$.

Proof. The proof proceeds by a standard induction on $\mathrm{dpt}(\varphi)$ and we omit it. \Box

Below and throughout the text, \prod denotes a disjoint union.

Definition 6.2 (Simple models). Let $k < \omega$.

A preordered model \mathfrak{S} is k-simple (or merely simple) if

$$
|\mathfrak{S}| = \coprod_{i=1}^k [p_i]_{\mathfrak{S}}.
$$

If \mathfrak{S} is k-simple and $x \in |\mathfrak{S}|$, we write $p_{\mathfrak{S}}(x)$ for the unique $p \in \{p_1, \ldots, p_k\}$ such that $x \in [p]_{\mathfrak{S}}$.

As always, we will drop subindices when it does not lead to confusion, writing $p(x)$ instead of $p_{\mathfrak{S}}(x)$.

Before continuing, let us establish a notational convention. Given natural numbers n, k, we will denote by $|n|_k$ the unique element m of $\{1,\ldots,k\}$ such that $n \equiv m \pmod{k}$. Note that this strays from the standard remainder in that $|k|_k = k$, but it shall simplify several expressions later on. Intervals shall be assumed to be intervals of natural numbers, i.e.

$$
[a, b] = \{ n \in \mathbb{N} : a \le n \le b \}.
$$

Further, it will be convenient to assume that the set of propositional variables is enumerated by $\langle p_k \rangle_{k \langle \omega}$.

In the remainder of this section, we shall use partial tangled bisimulations to show that L^{k+1} is more expressive than L^k . This might not be too surprising given results in [\[4,](#page-20-6) [7\]](#page-20-7), but to the best of my knowledge this has not been stated explicitly before and will provide a good warm-up for the techniques we shall use later on.

To be precise, by more expressive we mean the following: given languages $\lambda, \lambda' \subseteq L^*$ and a class of models X, we say λ' is at least as expressive as λ over X if, given $\varphi \in \lambda$, there is $\varphi' \in \lambda'$ such that, for every $\mathfrak{X} \in \mathcal{X}$, $[\![\varphi]\!]_{\mathfrak{X}} = [\![\varphi']\!]_{\mathfrak{X}}$.
The language λ' is more expressive than λ if λ' is at least as expressive as λ . The language λ' is *more expressive* than λ if λ' is at least as expressive as λ , The language λ' is *more expressive* than λ if λ' is at least as expressive as λ , but not vice-versa.

The following structures will be useful in proving our expressiveness result:

Figure 1: The models $\mathfrak{A}(2,2)$ (above) and $\mathfrak{A}(1,3)$. Points marked by a triangle satisfy p_1 , circles p_2 and squares p_3 . Ellipses indicate clusters, and the numbers shown represent the h-coordinate.

Definition 6.3 ($\mathfrak{A}(N, K)$). Given natural numbers N, K we define a K-simple structure $\mathfrak{A} = \mathfrak{A}(N, K)$ as follows:

- 1. $|\mathfrak{A}|$ is the set of all pairs (h, k) such that $h \in [0, NK], k \in [1, K]$ and either
- (a) $h = 0$ or (b) $k \neq |h|_K;$ 2. $(h, k) \preccurlyeq_{\mathfrak{A}} (h', k')$ if and only if $h \geq h'$; 3. $p(h, k) = p_k$.

Lemma 6.2. Given natural numbers $N, K, m, k \in [1, K]$ and

$$
h \in [1, (N-m)K]
$$

such that $h \not\equiv k \pmod{K}$,

$$
\langle \mathfrak{A}(N,K),(0,k)\rangle \cong_K^m \langle \mathfrak{A}(N,K),(h,k)\rangle.
$$

Proof. Let $\mathfrak{A} = \mathfrak{A}(N, K), k \in [1, K]$ and $1 \leq h \leq (N - m)K$.

We proceed by induction on m. The base case, when $m = 0$, is simple, since $(0, k)$ and (h, k) satisfy the same set of propositional variables (namely, $\{p_k\}$).

For the inductive step, we assume that $1 \leq h \leq (N - (m + 1))K$. Let us first check that $\mathsf{Forth}_{\preccurlyeq}$ holds.

Suppose that $x_1 \approx \ldots \approx x_{K-1} \preccurlyeq (0, k)$. Write $x_i = (\ell, k_i)$ and consider two cases. If $\ell \geq h$, we may set $y_i = x_i$ which clearly satisfy the conditions of Forth \preccurlyeq .

Otherwise, there exists a value $k_* \in [1, K]$ such that $k_i \neq k_*$ for all $i \in$ [1, K − 1]. Pick $h' \in [(N - (m + 1))K, (N – m)K]$ such that $h' \equiv k_* \pmod{K}$. For all $i, (h', k_i) \in |\mathfrak{A}|$, and by induction on m we have that

$$
y_i = (h', k_i) \rightleftharpoons_K^m (0, k_i) \rightleftharpoons_K^m (\ell, k_i),
$$

while clearly $y_1 \approx \ldots y_{K-1} \preccurlyeq (h, k)$, hence satisfying Forth \Diamond .

For Back_{\diamond}, suppose $y_1 \approx \ldots \approx y_{K-1} \preccurlyeq (h, k)$; then clearly $y_i \preccurlyeq (0, k)$ for all i, and we can set $x_i = y_i$. \Box

With this, we can show that $\left\langle \right\rangle_{k=1}^{k+1}$ $\prod_{i=1}^{n+1} \gamma_i$ cannot generally be defined by formulas of smaller tangled width:

Theorem 6.1. L^{k+1} is strictly more expressive than L^k for all k over the class of all finite topological models.

Proof. Let $\eta^k = \left\langle \right\rangle_{i=1}^{k+1}$ $\sum_{i=1}^{k+1} p_i$. Suppose $\varphi \in L^k$ has depth n and consider $\mathfrak{A} =$ $\mathfrak{A}(n+1,k+1).$

Then, by Lemma [6.2,](#page-9-0) $(1, 2) \rightleftarrows_{k+1}^{n} (0, 2)$, so that by Lemma [6.1,](#page-8-0) $\langle \mathfrak{A}, (1, 2) \rangle \rightleftarrows_{k}$ φ if and only if $\langle \mathfrak{A},(0, 2)\rangle \models \varphi$. However, it is easy to check that $\langle \mathfrak{A},(1, 2)\rangle \not\models \eta^k$ yet $\langle \mathfrak{A}, (0, 2) \rangle \models \eta^k$; hence φ cannot be equivalent to η^k .

Since $\varphi \in L^k$ was arbitrary, we conclude that η^k is not expressible in L^k over the class of finite topological models. \Box

7 Trouble formulas

In this section we shall introduce a sequence of formulas \langle Trouble $k \rangle_{k \leq \omega}$ with the property that \vdash^k Trouble^k. As we shall see later, $\not\vdash^k$ Trouble^{k+1}, thus establishing that DTL^{k+1} is stronger than DTL^k . The formulas Trouble^k will all be in L^1 .

Definition 7.1. The following abbreviations shall be used throughout the text.

$$
\operatorname{Cycle}^k = \Diamond p_k \to \bigwedge_{i=1}^k (p_i \to fp_{|i+1|_k})
$$
\n
$$
\operatorname{Start}_i^k = p_i \land [f] \operatorname{Cycle}^k
$$
\n
$$
\operatorname{Bundle}^k = \Box \bigwedge_{i=1}^k \Diamond \operatorname{Start}_i^k
$$
\n
$$
\operatorname{Tangle}^k = \bigotimes_{i \in [1,k]} \operatorname{Start}_i^k
$$
\n
$$
\operatorname{rankle}^k = \operatorname{Rundle}^k \to [f] \Diamond p_k
$$

Trouble^{κ} = Bundle^{κ} \rightarrow $|J|\sqrt{p_k}$

Before continuing, let us give some intuition for these formulas. The formula Cycle^k states that f 'cycles' the values of $p(x)$; if $p(x) = p_i$, $p(f(x)) = p_{i+1}$, unless $i = k$ in which case $p(f(x)) = p_1$. The formula $\Diamond p_k$ is used as a sort of trigger for this cycling behavior; when $\Diamond p_k$ fails, $p(f(x))$ is unspecified.

Start $_i^k$ is used to begin the cycling behavior described by Cycle^k at p_i ; it says that, initially, p_i holds, and from then on, f cycles the values of $p(f^n(x))$, provided that $\Diamond p_k$ holds at each step.

<code>Bundle</code> k and <code>Tangle</code> k are similar, but <code>Bundle</code> k is stronger. As we will mainly be interpreting these formulas over finite Kripke models, let us restrict the discussion to this setting. Here, the meaning of \texttt{Tangle}^k should be familiar; it says there is a cluster where there is a point x_i satisfying each Start^k.

The formula Bundle^k, meanwhile, says that each \texttt{Start}_i^k is dense near x; in particular, each minimal cluster must have one point satisfying each B undle^k. But such a cluster would be a witness to \texttt{Tangle}^k .

Thus we have that \models <code>Bundle</code>^{k} \rightarrow <code>Tangle</code>^{k}; but note that the former formula is in L^1 , while the latter is not.

Meanwhile, we should also expect \models Bundle^k \rightarrow fBundle^k; this is because, if $x_1 \approx x_2 \approx \ldots \approx x_k$ is a cluster with x_i satisfying Start^k_i, then clearly each x_i satsifies $\Diamond p_k$ (since $x_k \preccurlyeq x_i$), so that $f(x_i)$ satisfies $p_{|i+1|_k}$.

Thus also $f(x_1) \approx f(x_2) \approx \ldots \approx f(x_k)$ is a cluster of points satisfying each \texttt{Start}_i^k (although 'shifted' one step). It then follows that these points also satisfy \texttt{Tangle}^k . By induction, we see that $\models \texttt{Tangle}^k \rightarrow [f] \texttt{Tangle}^k$; but this clearly makes $\texttt{Trouble}^k$ true, since $\models \texttt{Tangle}^k \rightarrow \Diamond p_k.$

The reasoning we have just made is easy to formalize in DTL^k , as we show below. Later we shall also see that it is impossible to formalize in DTL^{k-1} .

 $\bf Proposition~7.1.}$ $Given$ $k<\omega,$ \vdash^k Trouble $^k.$

Proof. Reasoning within 54 one readily sees that, for any $i < k$, \vdash^0 Bundle^k \rightarrow \Diamond (Start $_k^k \land$ Bundle^k); thus we may apply necessitation and Ind_{\Diamond} to derive $\texttt{Bundle}^k \rightarrow \bigwedge^k_k$ $\sum_{i=1}^{\kappa}$ Start $\frac{k}{i}$ and obtain

$$
\vdash^{0} \text{Bundle}^{k} \to \text{Tangle}^{k}.\tag{1}
$$

Further, we note that

$$
\vdash^0 \mathtt{Tangle}^k \to \Diamond p_k,\tag{2}
$$

since this is a consequence of the axiom Fix_{\Diamond} .

For any $i \in [1, k]$ we may use Fix_{\diamond} to see that

$$
\vdash^0\mathtt{Tangle}^k\to \Diamond(\mathtt{Start}_i^k\wedge\mathtt{Tangle}^k).
$$

Using [\(2\)](#page-11-0), this imples

 \overline{a}

$$
\vdash^0\mathtt{Tangle}^k\to \Diamond(\mathtt{Start}_i^k\wedge \Diamond p_k\wedge\mathtt{Tangle}^k),
$$

i.e.

$$
{}^{-0}\text{ Tangle}^k \rightarrow \Diamond (p_i \wedge [f]\text{Cycle}^k \wedge \Diamond p_k \wedge \text{Tangle}^k).
$$

Now, by ${\sf Fix}_{[f]},$ $\vdash^0 [f]{\sf Cycle}^k \to ({\sf Cycle}^k_i \wedge f[f]{\sf Cycle}^k),$ whereas $\vdash^0 \Diamond p_k \wedge \texttt{Cycle}^k \to (p_i \to fp_{|i+1|_k}),$

i.e. $\vdash^0 p_i \wedge \Diamond p_k \wedge \text{Cycle}^k \rightarrow fp_{|i+1|k}$. From this we conclude that

$$
\vdash^0 \mathtt{Tangle}^k \to \Diamond (f p_{|i+1|_k} \land f[f] \mathtt{Cycle}^k \land \mathtt{Tangle}^k),
$$

and since it holds for all $i \in [1, k]$ we can use Ind_\Diamond to obtain

$$
\vdash^0 \mathtt{Tangle}^k \to \bigotimes_{i \in [1,k]} (fp_{|i+1|_k} \wedge f[f] \mathtt{Cycle}^k),
$$

which, rearranging indices and pulling out f , shows that

$$
\vdash^0 \mathtt{Tangle}^k \to \bigotimes_{i \in [1,k]} f(p_i \wedge [f] \mathtt{Cycle}^k).
$$

Now, we may use Cont^k to obtain

$$
\vdash^k \mathtt{Tangle}^k \to f \bigotimes_{i \in [1,k]} (p_i \wedge [f] \mathtt{Cycle}^k);
$$

by necessiation and $\text{Ind}_{[f]}$ this yields

$$
\vdash^k \mathtt{Tangle}^k \to [f] \bigotimes_{i \in [1,k]} (p_i \wedge [f] \mathtt{Cycle}^k),
$$

i.e. $\vdash^k \mathtt{Tangle}^k \to [f]\mathtt{Tangle}^k.$

Putting this together with [\(2\)](#page-11-0) we see that

$$
\vdash^k \mathtt{Tangle}^k \to [f] \Diamond p_k,\tag{3}
$$

which together with [\(1\)](#page-11-1) gives us

$$
\vdash^k\mathtt{Bundle}^k\to [f]\Diamond p_k,
$$

i.e. \vdash^k Trouble^k, as claimed.

8 Incompleteness of finite fragments

The formula $\texttt{Trouble}^k$ is derivable in \texttt{DTL}^k ; let us now see that $\texttt{Trouble}^{k+1}$ is not. To prove this, we shall introduce models $\mathfrak{D}(N, K)$. They will be composed of two submodels; $\mathfrak{C}(K)$, defined later, and $\mathfrak{B}(N, K)$, defined below.

The general idea is that the models $\mathfrak{D}(n+1, k+1)$ will satisfy

$$
\mathsf{DTL}_n^k \cup \{\neg \mathtt{Trouble}^{k+1}\},
$$

thus showing that ψ_n^k Trouble^{k+1} for all *n*. From this we may conclude that $\not\vdash^k$ Trouble $^{k+1}.$

 \Box

Figure 2: The model $\mathfrak{D}(2,2)$, described in Definition [8.3.](#page-16-0) $\mathfrak{B}(2,2)$, as in Defini-tion [8.1,](#page-14-0) is the submodel on the right-hand side of the dotted line and $\mathfrak{C}(2)$ is the submodel on its left. Arrows indicate $f_{\mathcal{D}}$, while \preccurlyeq is the transitive, reflexive closure of the relation represented by \prec together with the ellipse on the left, which represents \approx . Points represented by a circle satisfy p_1 , by a triangle, p_2 .

Before defining our structures formally, let us give a general idea. Consider the model $\mathfrak{D} = \mathfrak{D}(2, 2)$ depicted in Figure [2.](#page-13-0) We will name a point x using triples $(h(x), t(x), k(x))$, where $h(x)$ is the 'spatial' (vertical) coordinate, $t(x)$ the 'temporal' (horizontal) coordinate and $k(x)$ is the index of $p(x)$, which in this case is 1 for points represented by a circle and 2 for triangles. The points on the left of the dotted line will be written $(0, -1, k(x))$.

First, let us observe that $\mathfrak{D} \models \text{Cycle}^2$, since $f_{\mathfrak{D}}$ alternates between circles (which satisfy p_1) and triangles (which satisfy p_2). The exception for this are the points on the main diagonal $h + t = 6$ and on the 'tail' $t \geq 4$, but these points do not satisfy $\Diamond p_2$ and thus they also satisfy Cycle². From this, one can easily check that $(0, -1, 1)$ satisfies \neg Trouble².

Meanwhile, the key aspect of the model is that $f_{\mathfrak{D}}$ is discontinuous, since $(0, -1, 2) \preccurlyeq (0, -1, 1)$ yet

$$
f_{\mathfrak{D}}(0, -1, 2) = (0, 0, 1) \nless (1, 0, 2) = f_{\mathfrak{D}}(0, -1, 1).
$$

This discontinuity is easily seen to make the following instance of Cont^2 fail on $(0, -1, 1)$:

$$
\Diamond f\{p_1, p_2\} \to f\Diamond \{p_1, p_2\}.
$$

However, instances of Cont^1 of small modal depth do hold. Consider, for example,

$$
\Diamond fp_1 \to f\Diamond p_1.
$$

Here we see that $f_{\mathfrak{D}}(0, -1, 2)$ satisfies fp_1 , so that $(0, -1, 1)$ satisfies $\Diamond fp_1$. If $f_{\mathfrak{D}}$ were continuous, we would be able to use $f_{\mathfrak{D}}(0, -1, 2)$ as a witness that $(0, -1, 1)$ satisfies $f \Diamond p_1$, but in this case we cannot. However, we do have a different witness, namely $(2, 0, 1)$. More generally, as we shall see in Lemma [8.1,](#page-16-1) $(2, 0, 1) \rightleftharpoons^1 (0, 0, 1)$ so the two satisfy the same formulas of modal depth one.
Thus Ω extisfies $DT^{1,1}$ as well as π **reauble**² from which we conclude the

Thus \mathfrak{D} satisfies DTL¹ as well as \neg **Trouble**², from which we conclude that \forall_1^1 Trouble². To see that \forall_n^1 Trouble², we need to consider a larger model, $\mathfrak{D}(n, 2)$, which is built much like $\mathfrak{D}(2, 2)$ but is deeper. By varying n, we conclude that $\not\vdash^1$ Trouble².

Now, let us give the formal definition of $\mathfrak{B}(N, K)$, which is the submodel of $\mathfrak{D}(N, K)$ on the right of the dotted lines in Figure [2.](#page-13-0)

Definition 8.1. Given $N, K < \omega$, we define a K-simple dynamic model $\mathfrak{B} =$ $\mathfrak{B}(N,K)$ by letting

1. $|\mathfrak{B}|$ be the set of all triples of natural numbers (h, t, k) such that either

(a)
$$
h + t \le NK
$$
, $k \in [1, K]$ and $k \not\equiv h + t \pmod{K}$ or
(b) $h = 0$, $t \in [NK + 1, N(K + 1)]$ and $k \ne K$.

2. $(h_1, t_1, k_1) \preceq_{\mathfrak{B}} (h_2, t_2, k_2)$ if and only if $t_1 = t_2$ and $h_1 \geq h_2$;

$$
3. \ f_{\mathfrak{B}}(h,t,k) = \begin{cases} (h, t+1, |k+1|_K) & \text{if } h+t < NK \\ (h-1, t+1, k) & \text{if } h+t = NK \text{ and } h > 0 \\ (h, t+1, k) & \text{if } t \in [NK+1, N(K+1)) \\ (0, 0, |k+1|_K) & \text{if } t = N(K+1) \end{cases}
$$

4. $p(h, t, k) = p_k$.

We will write points as $x = (h(x), k(x), t(x))$. We will also write $s(x) =$ $h(x) + t(x)$.

It will be convenient to describe the ω_n^m -equivalence classes over $\mathfrak{B}(N, K)$.
shall do this using the relations ω_n^m defined below. We shall do this using the relations \sim^m , defined below.

Definition 8.2. For $m < N$, say $x \sim^m y$ if $p(x) = p(y)$ and one (or more) of the following occurs:

- 1. $s(x) = s(y)$,
- 2. $s(x), s(y) \leq K(N-m)$ or
- 3. $s(x), s(y) \in [NK, N(K+1) m].$

The models $\mathfrak{B}(k,n)$ are designed to be very homogeneous, so that different points are hard to distinguished using L^* . The relations ∼^m are representative of this.

First, note that every point is \sim^m -similar to another on the t-axis; this can be seen in Figure [3,](#page-15-0) where every point can be 'slid' down the diagonals to one

Figure 3: The relation \sim ¹ on $\mathfrak{B}(2, 2)$. Points inside the large lower-left triangle are ∼¹ -related if and only if they satisfy the same propositional variable, which in this figure occurs when they have the same shape. This triangle would grow if we were to consider \sim^0 , or shrink if we were to consider \sim^2 .

on the line $h = 0$. Points in the large left-hand triangle may have more than one such representative.

Another useful property is that all points have a very similar orbit; namely, if x is any point and y satisfies $h(y) = 0$, then y lies in the orbit of x. This is easily seen in Figure [2,](#page-13-0) where if we follow the $f_{\mathcal{D}}$ -arrows starting anywhere, we eventually reach $(0, 0, 1)$, and from here $f_{\mathfrak{D}}$ simply cycles around the t-axis indefinitely.

The situation is slightly more involved for larger values of k , where we may turn to Figure [4.](#page-17-0) Consider, for example, the point $(0, 0, 2)$; in this example, circles have third coordinate 1, triangles have 2, squares 3. Here, notice that after four iterations of $f_{\mathfrak{B}}$ we reach $(0, 4, 1)$, which then maps to $(0, 0, 1)$. Afterwards, $f_{\mathfrak{B}}$ will cycle through the second row of the t-axis, and then return again to $(0, 0, 2)$.

Let us collect these observations into a lemma:

Lemma 8.1. For every $x \in [\mathfrak{B}(N,K)]$ and $m < N$,

- 1. there is $y \sim^m x$ with $h(y) = 0$ and
- 2. if $h(y) = 0$ there is $n < \omega$ such that $f_{\mathfrak{B}}^n(x) = y$.

Proof. The first claim is obvious if we notice that

 $(h, t, k) \sim^m (0, h + t, k).$

For the second, first we observe that $h(f^{N(K+1)+1}(x))=0$ independently of x; then note that $f_{\mathfrak{B}}$ is clearly transitive on those elements z with $h(z) = 0$, given that

$$
f^{N(K+1)+1}(0,0,k) = (0,0,|k+1|_K),
$$

thus 'rotating' $k(z)$.

Now, let us see that \sim^m does, indeed, guarantee partial bisimulation.

 \Box

Proposition 8.1. If $x \sim^m y$ then $x \approx^m y$.

Proof. We work by induction on m , considering each clause of a tangled bisimulation. Note that \sim^m preserves atoms, in particular covering the case $m = 0$.

Otherwise, suppose $x \sim^{m+1} y$. Clearly we only need to prove the 'forth' clauses, since the 'back' clauses are symmetric.

Forth_{\leq} We shall only consider the case where $s(x), s(y) < NK$; the other case is similar and easier.

Suppose $x_0 \approx x_1 \approx \ldots \approx x_{I-1} \preccurlyeq x$; note that we can assume $I \leq K$, since **B** has cluster width K. Note also that each x_i has $h(x_i) \geq h(x)$ and $t(x_i) = t(x).$

Consider $h' = h(x_i) + t(y) - t(x)$. If $h' \ge h(y)$, set $h = h'$; otherwise, let h be the least value such that $h \geq h(y)$ and $h + t(y) \equiv h(x_i) + t(x)$ (mod K). Then, set $y_i = (h, t(y), k(x_i))$.

First, note that $s(y_i) \equiv s(x_i) \pmod{K}$, so that all y_i are elements of $|\mathfrak{B}|$. Now, we further have that $s(y_i) = s(x_i)$ except in the case that $h' < h(y)$, in which it easily follows that $s(x) \neq s(y)$, so $s(x)$, $s(y) < K(N - (m + 1))$ and thus $s(x_i), s(y_i) < K(N-m)$.

In either case we use our induction hypothesis to see that $y_i \trianglelefteq^m x_i$, as also claimed.

- Forth f This follows from observing that the required (in)equalities are preserved by $f_{\mathfrak{B}}$ and we skip it.
- Forth $[f]$ Let $n < \omega$ and consider $z = f_{\mathfrak{B}}^n(x)$. Then, by Lemma [8.1.](#page-15-1)[1,](#page-15-2) there is $z' \sim m$ z with $h(z') = 0$, while by Lemma [8.1,](#page-15-1)[2,](#page-15-3) there is n' such that $f_{\mathfrak{B}}^{n'}(y) = z'$, as required.

 \Box

Now that we have studied the models $\mathfrak{B}(N, K)$, let us add the 'head' $\mathfrak{C}(K)$, which is where the trouble really lurks. The resulting model will be called $\mathfrak{D}(N, K)$, where points in $\mathfrak{C}(K)$ will map discontinuously onto $\mathfrak{B}(N, K)$. However, these discontinuities will require large formulas to capture in L^{K-1} , given that $\mathfrak{C}(K)$ will consist of a cluster with K points.

Definition 8.3. We define a model $\mathfrak{C} = \mathfrak{C}(K)$ where

- $|\mathfrak{C}| = \{0\} \times \{-1\} \times [1, K]$
- $\preccurlyeq_{\mathfrak{C}}$ is total (i.e., \mathfrak{C} consists of a single cluster)
- $p(0, -1, k) = p_k$.

We define a model $\mathfrak{D} = \mathfrak{D}(N, K)$ based on $|\mathfrak{C}(K)| \cup |\mathfrak{B}(N, K)|$ with

$$
\preccurlyeq_{\mathfrak{D}} = \preccurlyeq_{\mathfrak{C}(K)} \cup \preccurlyeq_{\mathfrak{B}(N,K)}
$$

Figure 4: The model $\mathfrak{D}(1,3)$. Notation is similar to that for Figure [2,](#page-13-0) but this time we have only partially indicated the relations; points marked with a square now satisfy p_3 .

and

$$
f_{\mathfrak{D}}(0,-1,k) = \begin{cases} (0,0,|k+1|_K) & \text{if } k \neq K-1 \\ (1,0,K) & \text{if } k = K-1. \end{cases}
$$

Our strategy now is to show that $\mathfrak{D}(N+1, K+1)$ is a model of DTL_N^K $\{\neg \text{Trouble}^{K+1}\}$; from this we may conclude that \forall^{K} Trouble^{K+1}, given that $\dot{\mathsf{D}}\mathsf{T}\mathsf{L}^K = \bigcup_{n<\omega} \mathsf{D}\mathsf{T}\mathsf{L}_n^K.$

Lemma 8.2. $\mathfrak{D}(N+1,K+1) \models \mathsf{DTL}_N^K$.

Proof. All the rules of DTL_N^K preserve model validity, so it suffices to check that $\mathfrak{D}(N+1,K+1)$ satisfies all axioms of DTL_N^K ; that is, all permitted substitution instances of axioms of DTL^K .

Since $\mathfrak{D}(N+1, K+1)$ is a weak dynamical system, it satisfies every axiom of DTL_N^K except possibly for instances of Cont^K.

So, let

$$
\sigma = \bigotimes_{i \leq K} f \delta_i \to f \bigotimes_{i \leq K} \delta_i
$$

be a substitution instance of Cont^K where each δ_i has modal depth at most N. Let $x \in |\mathfrak{D}|$ and assume that

$$
x \in \left[\left[\bigotimes_{i \leq K} f \delta_i \right]_{\mathfrak{D}};
$$

since $f_{\mathfrak{D}} \restriction |\mathfrak{B}|$ is continuous, we can suppose that $x \in |\mathfrak{C}|$, for otherwise $x \in$ $[\![\sigma]\!]_{\mathfrak{D}}.$

Then, given $i < K$ there is $x_i = (0, -1, k_i) \approx x$ such that $x_i \in [f\delta_i]_{\mathcal{D}}$. For at least one value of $k_* \in [1, K + 1]$ we have that $k_* \neq |k_i + 1|_K$ for all i; we then have that $y_i = (k_*, 0, |k_i + 1|_K)$ is an element of $|\mathfrak{B}|$ and by Proposition [8.1](#page-16-1)

$$
y_i \triangleq^N_* f_{\mathfrak{D}}(x_i).
$$

Meanwhile, $y_i \approx y_j \preccurlyeq f_{\mathfrak{D}}(x)$ for all i, j , so that

$$
x \in \left[f \bigotimes_{i \leq K} \delta_i \right]_{\mathfrak{D}},
$$

as required.

Lemma 8.3. Given $K, N \leq \omega$ and $k \in [1, k]$,

$$
\langle \mathfrak{D}(N,K), (0,-1,k) \rangle \models \neg \texttt{Trouble}^K.
$$

Proof. Let $\mathfrak{D} = \mathfrak{D}(N, K)$.

First, let us show that every $x \in |\mathfrak{D}|$ satisfies

$$
\operatorname{Cycle}^K = \Diamond p_K \to \bigwedge_{k \le K} (p_k \to fp_{|k+1|_K}).
$$

If $s(x) \ge NK$, then $x \notin [\![\Diamond p_K]\!]_{\mathfrak{D}}$ and thus $x \notin [\![\Diamond p_K]\!]_{\mathfrak{D}}$. This shows that $x \in \llbracket \texttt{Cycle}^K \rrbracket_{\mathfrak{D}}, \text{ as required.}$

Otherwise, letting $y = f_{\mathfrak{D}}(x)$, we note by case-by-case inspection that $k(y) =$ $|k(x) + 1|_K$, so that x satisfies $p_{k(x)} \to fp_{|k(x)+1|_K}$, whereas for $k \neq k(x)$, x satisfies $p_k \to fp_{|k+1|_K}$ trivially. Thus Cycle^K holds everywhere, as claimed.

It follows from this, in particular, that $(0, -1, k)$ satisfies $p_k \wedge [f]$ Cycle^K, i.e. Start^K; this shows that $(0, -1, k)$ satisfies $\bigwedge_{1 \leq i \leq K} \Diamond$ Start^K and, given that $k \in [1, K]$ was arbitrary,

$$
\langle \mathfrak{D}, (0,-1,k) \rangle \models \Box \bigwedge_{i=1}^k \Diamond \mathrm{Start}^K_i = \mathrm{Bundle}^K.
$$

It remains to show that $(0, -1, k)$ satisfies $\langle f \rangle \Box \neg p_K$; but this follows from the observation that

$$
f_{\mathfrak{D}}^{NK+1}(0,-1,k)=(0,NK,k') \notin [\![\Diamond p_K]\!]_{\mathfrak{D}}.
$$

We conclude that

$$
\langle \mathfrak{D}, (0,-1,k) \rangle \models \texttt{Bundle}^K \land \neg[f] \Diamond p_K \equiv \neg \texttt{Trouble}^K,
$$

as claimed.

The following lemma summarizes our results so far:

 \Box

 \Box

Lemma 8.4. For all $k < \omega$, the formula Trouble^{k+1} $\in L^1$ is derivable in DTL^{k+1} , but not in DTL^k .

Proof. By Proposition [7.1,](#page-11-2) \vdash^{k+1} Trouble^{k+1}; meanwhile, if \vdash^{k} Trouble^{k+1}, we would have that \vdash_n^k Trouble^{k+1} for some *n*.

But this cannot be, since we have seen that

$$
\mathfrak{D}(n+1,k+1) \models \mathsf{DTL}_n^k \cup \{\neg \mathsf{Trouble}^{k+1}\},
$$

and thus $\not\vdash_n^k$ Trouble^{k+1}.

With this, we may easily prove our main result.

Theorem 8.1. Let λ be any language such that $L \subseteq \lambda \subseteq L^*$, and let $\mathcal{DTL}[\lambda] =$ $\mathcal{DTL}^* \cap \lambda$.

Similarly, for $k < \omega$, define $DTL^k[\lambda] = DTL^k \cap \lambda$.

Then, given any natural number k, $DT{\mathcal{L}}[\lambda]$ is not finitely axiomatizable^{[4](#page-19-0)} *over* $DTL^{k}[\lambda]$.

Proof. Let T be any sound, finite extension of $DTL^k[\lambda]$, so that without loss of generality we may assume $\mathsf{T} = \mathsf{DTL}^k[\lambda] + \phi$ for some valid formula ϕ .

Since DTL^{*} is complete, we would have that that DTL^{*} $\vdash \phi$, and hence, for some value of K, $DTL^{\tilde{K}} \vdash \phi$; obviously, we may take $K \geq k$.

But then, we have by Lemma [8.4](#page-19-1) that DTL^K \neq Trouble^{K+1}, and hence

$$
\mathsf{DTL}^k[\lambda] + \phi \not\vdash \mathsf{Trouble}^{K+1} \in \mathcal{DTL}[\lambda].
$$

Meanwhile, T was arbitrary, so we conclude that $\mathcal{DTL}[\lambda]$ is not finitely axiomatizable over $DTL^k[\lambda]$. \Box

This result is quite general, so it may be convenient to explicitly mention some special cases. The following corollary states some immediate consequences of Theorem [8.1;](#page-19-2) below, recall that $KM \subseteq DTL^1[L]$.

Corollary 8.1. $DT\mathcal{L}$ and $DT\mathcal{L}^*$ are not finitely axiomatizable. In particular, KM is incomplete for the class of dynamic topological models.

9 Concluding remarks

The axiomatization DTL^{*} introduced the tangled modality to Dynamic Topological Logic as a sort of scaffolding, to be removed once the appropriate techniques were available. However, I believe the present work to be a convincing argument that indeed it is a central element of the logic; tangled sets affect the behavior of dynamical systems and to be unable to reason about them directly gives a logical formalism an unnecessary handicap.

 \Box

⁴Observe that DTL^{k+1} is finitely axiomatizable over DTL^k, but it does not necessarily follow from this that $DTL^{k+1}[\lambda]$ is finitely axiomatizable over $DTL^{k}[\lambda]$ for all λ .

Of course, none of the results presented here show that a reasonable axiomatization within L^1 is impossible to find. I am not sure how relevant such an axiomatization would be at this point, but it remains an interesting problem.

Meanwhile, I believe a more fruitful direction is to analyze other logics which are hard to axiomatize because of a similar lack in expressive power. In particular, there are many products of modal logics which have very similar models to those of Dynamic Topological Logic; perhaps they too would benefit from a polyadic variant?

References

- [1] S.N. Artemov, J.M. Davoren, and A. Nerode. Modal logics and topological semantics for hybrid systems. Technical Report MSI 97-05, 1997.
- [2] Guram Bezhanishvili and Mai Gehrke. Completeness of S4 with respect to the real line: revisited. Annals of Pure and Applied Logic, 131(1–3):287 – 301, 2005.
- [3] P. Blackburn, M. de Rijke, and Y. Venema. Modal Logic. Cambridge University Press, 2001.
- [4] A. Dawar and M. Otto. Modal characterisation theorems over special classes of frames. Annals of Pure and Applied Logic, 161:1–42, 2009. Extended journal version LICS 2005 paper.
- [5] David Fern´andez Duque. Tangled modal logic for topological dynamics. Annals of Pure and Applied Logic, 163(4):467–481, 2012.
- [6] D. Fern´andez-Duque. Non-deterministic semantics for dynamic topological logic. Annals of Pure and Applied Logic, 157(2-3):110–121, 2009. Kurt Gödel Centenary Research Prize Fellowships.
- [7] D. Fernández-Duque. On the modal definability of simulability by finite transitive models. Studia Logica, 98:347–373, August 2011.
- [8] D. Fern´andez-Duque. Tangled modal logic for spatial reasoning. In T. Walsh, editor, Proceedings of IJCAI, pages 857–862, 2011.
- [9] D. Fernández-Duque. A sound and complete axiomatization for dynamic topological logic. Journal of Symbolic Logic, 2012. forthcoming.
- [10] B. Konev, R. Kontchakov, F. Wolter, and M. Zakharyaschev. Dynamic topological logics over spaces with continuous functions. In G. Governatori, I. Hodkinson, and Y. Venema, editors, Advances in Modal Logic, volume 6, pages 299–318, London, 2006. College Publications.
- [11] B. Konev, R. Kontchakov, F. Wolter, and M. Zakharyaschev. On dynamic topological and metric logics. Studia Logica, 84:129–160, 2006.
- [12] P. Kremer. Strong completeness of S4 wrt the real line. 2012.
- [13] P. Kremer and G. Mints. Dynamic topological logic. Annals of Pure and Applied Logic, 131:133–158, 2005.
- [14] O. Lichtenstein and A. Pnueli. Propositional temporal logics: Decidability and completeness. Logic Jounal of the IGPL, 8(1):55–85, 2000.
- [15] G. Mints and T. Zhang. Propositional logic of continuous transformations in Cantor space. Archive for Mathematical Logic, 44:783–799, 2005.
- [16] A. Prior. Time and modality. 1957.
- [17] A. Tarski. Der Aussagenkalkül und die Topologie. Fundamenta Mathematica, 31:103–134, 1938.