

# Rational Calogero-Moser model: Explicit forms and $r$ -matrix structure of the second Poisson structure

J. Avan<sup>a</sup> and E. Ragoucy<sup>b</sup>

<sup>a</sup> *LPTM, CNRS and Université de Cergy-Pontoise,  
2 avenue A. Chauvin, 95302 Cergy-Pontoise Cedex, France  
E-mail: avan@u-cergy.fr*

<sup>b</sup> *LAPTh, CNRS and Université de Savoie,  
9 chemin de Bellevue, BP 110, 74941, Annecy-Le-Vieux Cedex, France  
E-mail: ragoucy@lapp.in2p3.fr*

## Abstract

We compute the full expression of the second Poisson bracket structure for  $N = 2$  and  $N = 3$  site rational classical Calogero-Moser model. We propose an  $r$ -matrix formulation for  $N = 2$ . It is identified with the classical limit of the second dynamical boundary algebra previously built by the authors.

MSC: 81R12, 16T15, 16T25

Keywords: Integrable systems; Dynamical algebras; Reflection algebras

July 2012  
LAPTH-036/12

# 1 Introduction

## 1.1 Dynamical $r$ -matrices

The Calogero-Moser model [1, 2] provided [3] a textbook example of a classical dynamical  $r$ -matrix structure for a Lax representation [4] of a classical Liouville integrable system. Remember that Liouville integrability for a  $N$ -dimensional Hamiltonian system is characterized (see [5] and references therein) by the existence of  $N$  independent Poisson-commuting quantities  $H^{(i)}, i = 1 \cdots N$  including the original Hamiltonian. The system admits a Lax representation when the equations of motion can be represented as a spectrum-preserving evolution of a  $n \times n$  matrix  $L$  encapsulating the  $2N$  dynamical variables  $\{p_i, q_i\}, i = 1 \cdots N$ :

$$\frac{dL}{dt} = [L, M]. \quad (1.1)$$

The quantities  $TrL^k, k = 1 \cdots n$  may then provide the Liouville Hamiltonians if they Poisson-commute and they are in sufficient number (e.g. if  $n \geq N$  or there exists a spectral-parameter dependence). They build in this case a so-called Hamiltonian hierarchy of mutually compatible equations of motion. This Poisson-commuting property is equivalent [6] to rewriting the Poisson brackets of the Lax matrix coefficients in a specific algebraic form involving a matrix  $r$ , living in the tensor product  $M_n(\mathbb{C}) \otimes M_n(\mathbb{C})$ ,  $M_n(\mathbb{C})$  being self-explanatorily the algebra of complex  $n \times n$  matrices:

$$\{L_1, L_2\} = [r_{12}, L_1] - [r_{21}, L_2]. \quad (1.2)$$

Expression (1.2) is the so-called linear form of  $r$ -matrix structure. Associativity of the Poisson bracket form (1.2) is guaranteed by the consistency equation:

$$[r_{12}, r_{13}] + [r_{12}, r_{23}] + [r_{32}, r_{13}] + \{r_{12}, L_3\} - \{r_{13}, L_2\} = 0 \quad (1.3)$$

generically known as “dynamical Yang-Baxter equation” (dYB), see e.g. [28]. If  $r$  does not depend on the dynamical variables (1.3) becomes a purely algebraic “classical Yang-Baxter equation”. If not the issue arises of an available algebraic reexpression of the dynamical contribution  $\{r_{12}, L_3\} - \{r_{13}, L_2\}$ . At least one such form is available in the literature and occurs in the Calogero-Moser model, where one identifies:

$$\{r_{12}, L_3\} = \sum_{k=1}^n h_s^{(3)} \frac{d}{dq_s} r_{12} \quad (1.4)$$

where  $h_s, s = 1 \cdots n$  is a representation of a Cartan subalgebra<sup>1</sup>.

Equation (1.4) now allows to identify the generic dYB equation (1.3) with a specific equation known as the classical Gervais-Neveu-Felder equation [8, 18, 19]. The problem of

---

<sup>1</sup>In a more general, abstract context the abelian property is dropped and one is lead to consider “non-abelian dynamical algebras” [7]

realizing an explicit algebraic form for the dynamical terms remains however open in many interesting cases.

It is crucial to underline here that in the Lax representation (1.1) there exists a one-to-one algebraic correspondence, parametrized by the linear  $r$  matrix, between the Hamiltonian  $H(L)$  triggering a specific time evolution, and the associated  $M$  matrix. Namely [12] the  $M$  matrix is given by:

$$M = Tr_1(r_{12} dH(L)_2). \quad (1.5)$$

The Lax matrix  $L$  however is characteristic of the whole Hamiltonian hierarchy and remains therefore unique.

## 1.2 Hierarchy of Poisson brackets: Magri construction

A dual formulation of Liouville integrability was proposed by Magri [27] where the hierarchy of Hamiltonians  $H^{(k)} = TrL^k, k = 1 \cdots n$  acting simultaneously on the dynamical variables  $p_i, q_j$  through a single Poisson bracket structure (e.g. the canonical one  $\{p_i, q_j\} = \delta_{ij}$ ), is substituted by a hierarchy of mutually compatible Poisson brackets  $\{\}_{(i)}$ , where  $i = 1$  corresponds to the above canonical “first” bracket. The duality between the two formulations is summarized by the identity:

$$\{H^{(k)}, X(p, q)\}_{(l)} = \{H^{(k')}, X(p, q)\}_{(l')} \quad \text{for } k + l = k' + l'. \quad (1.6)$$

The explicit construction of the higher ( $l \geq 2$ ) Poisson brackets uses the so-called “recursion operator” (see e.g. [9]).

It then follows that the Lax matrix for a classically Liouville integrable system exhibits a corresponding hierarchy of  $r$ -matrix structures associated to each Poisson bracket in the Magri hierarchy. Such structures are explicitly known when the  $r$ -matrix for the first (canonical) bracket is non-dynamical. For instance the second Poisson bracket is the famous quadratic Sklyanin bracket [10, 11, 28]:

$$\{L_1, L_2\} = [a_{12}, L_1 L_2] + L_1 s_{12} L_2 - L_2 s_{12} L_1 \quad (1.7)$$

where  $a$  and  $s$  are respectively the skew-symmetric,  $a_{12} = \frac{1}{2}(r_{12} - r_{21})$ , and symmetric,  $s_{12} = \frac{1}{2}(r_{12} + r_{21})$ , part of the  $r$ -matrix. The Sklyanin bracket, properly said, corresponds to the case  $s = 0$ .

Note that a quadratic Poisson bracket (1.7) takes in any case the general linear form (1.2) with a linear  $r$  matrix defined as

$$r_{12} \equiv \frac{1}{2}(a_{12} L_2 + L_2 a_{12}) - L_2 s_{12}. \quad (1.8)$$

The third bracket has a more complicated form, cubic in terms of  $L$ , derived explicitly in [11].

When  $r$  is dynamical no such general formulation is available. Our purpose here is therefore to give a first example of an explicit  $r$ -matrix structure for the simplest example of the rational  $N$ -site Calogero-Moser Lax matrix. Its second Poisson structure was recently derived explicitly [26, 27] which makes it possible in principle to define an associated  $r$ -matrix structure.

The Poisson bracket hierarchy however is only formulated in terms of the adjoint invariants of the Lax matrix  $I_k = Tr L^k$  and mixed Lax-position matrix  $J_k = Tr L^k Q$  where  $Q$  is the diagonal position matrix  $Diag (q_1 \cdots q_N)$ . Inverting this form to reexpress the Poisson brackets of the original first-bracket canonical variables (in which the Lax matrix elements have simple expressions) is our first aim, partially achieved in Section 3.

We first of all (Section 2) derive some general Poisson bracket identities valid for all values of  $N$ , which allows us to reobtain the full  $N = 2$  second Poisson bracket structure derived in [26, 27] and construct explicitly the  $N = 3$  Poisson brackets (Section 3). Formulae for  $N = 3$  are considerably more complicated and do not yield at this time an obvious generalization to any value of  $N \geq 4$ .

We then (Section 4) propose a completely explicit form for second Poisson bracket of the  $N = 2$  site Lax matrix. This structure is defined as an explicit quadratic Sklyanin form (1.7) in terms of the Lax matrix  $L$ . It is identified with the classical limit of the Second Dynamical Boundary Algebra, recently built in [25]. The case  $N = 3$  seems at this time too complex and not clearly enough understood to allow for a reasonable attempt at building an  $r$  matrix structure. We conclude with some remarks and proposals.

## 2 General properties of $N$ -Calogero

The Lax matrix and position matrix for the rational Calogero-Mpser model are defined as:

$$L = \begin{pmatrix} p_1 & \frac{1}{q_{12}} & \cdots & \frac{1}{q_{1N}} \\ \frac{-1}{q_{12}} & p_2 & \ddots & \vdots \\ \vdots & \ddots & \ddots & \frac{1}{q_{N-1,N}} \\ \frac{-1}{q_{1N}} & \cdots & \frac{-1}{q_{N-1,N}} & p_N \end{pmatrix} \quad \text{and} \quad Q = \begin{pmatrix} q_1 & 0 & \cdots & 0 \\ 0 & q_2 & \ddots & \vdots \\ \vdots & \ddots & \ddots & 0 \\ 0 & \cdots & 0 & q_N \end{pmatrix} \quad (2.1)$$

where  $q_{ij} = q_i - q_j$ .

The second Poisson structure is expressed [26] in the basis

$$I_n = \frac{1}{n} Tr L^n \quad \text{and} \quad J_{n+1} = Tr(Q L^n). \quad (2.2)$$

It reads

$$\{I_n, I_m\} = 0 \quad ; \quad \{J_n, I_m\} = (m+n-1)I_{m+n-1} \quad ; \quad \{J_n, J_m\} = (n-m)J_{m+n-1}. \quad (2.3)$$

We call this (PB-)algebra  $\mathcal{A}_N$ . Remark that the first generators  $I_1$  and  $J_1$  correspond to the center of mass position and impulsion,

$$p_0 = I_1 = \sum_{j=1}^N p_j \quad \text{and} \quad q_0 = J_1 = \sum_{j=1}^N q_j, \quad (2.4)$$

while  $I_2$  is the Calogero Hamiltonian

$$H = \sum_{j=1}^N p_j^2 - 2 \sum_{j \neq i}^N \frac{1}{(q_i - q_j)^2}.$$

## 2.1 Decoupling of the center of mass position

We extract consistently the centre-of-mass variable  $q_0 \equiv J_1$  by setting:

$$\tilde{Q} = Q - \frac{\text{Tr}(Q)}{N} \mathbb{I} = Q - \frac{q_0}{N} \mathbb{I} \quad \text{and} \quad \tilde{J}_{n+1} = \text{Tr}(\tilde{Q} I_n) = J_{n+1} - \frac{n}{N} q_0 I_n \quad (2.5)$$

with the convention  $\lim_{n \rightarrow 0} (n I_n) = N$ .

**Proposition 2.1** *The algebra  $\mathcal{A}_N$  is the semi-direct sum of the subalgebra  $\tilde{\mathcal{A}}_N$ , generated by  $I_n$  and  $\tilde{J}_n$  and the (PB-)commutative algebra  $\{q_0\}$ .*

$$\{I_n, I_m\} = 0 \quad ; \quad \{\tilde{J}_{n+1}, I_m\} = (m+n) I_{m+n} - \frac{mn}{N} I_m I_n \quad (2.6)$$

$$\{\tilde{J}_{n+1}, \tilde{J}_{m+1}\} = (n-m) \tilde{J}_{m+n+1} + \frac{mn}{N} (\tilde{J}_{m+1} I_n - \tilde{J}_{n+1} I_m) \quad (2.7)$$

$$\{q_0, I_m\} = m I_m \quad ; \quad \{q_0, \tilde{J}_{n+1}\} = n \tilde{J}_{n+1} \quad (2.8)$$

*The structure of the algebra  $\tilde{\mathcal{A}}_N$  is entirely determined by the PBs of the  $p_i$ 's and the  $q_i$ 's and do not depend on the PBs of  $q_0$ .*

Proof: The PBs are obtained by direct calculation. It follows that, since  $\tilde{Q}$  and  $L$  do not depend on  $q_0$ , the PBs of  $q_0$  with  $I$  and  $J$  variables are not relevant in the calculation of the  $L$  Poisson brackets and the associated  $r$ -matrix structure. They shall be considered separately. ■

## 2.2 PBs of the center of mass momentum

**Proposition 2.2** *The Poisson brackets of  $p_0$  are given by*

$$\{p_0, q_j\} = -p_j \quad \text{and} \quad \{p_0, p_j\} = -2 \sum_{n \neq j} q_{jn}^{-3} \quad (2.9)$$

Proof: Let

$$K = \begin{pmatrix} a_1 & q_{12}^{-2} & \cdots & q_{1N}^{-2} \\ q_{12}^{-2} & a_2 & \ddots & \vdots \\ \vdots & \ddots & \ddots & q_{N-1,N}^{-2} \\ q_{1N}^{-2} & \cdots & q_{N-1,N}^{-2} & a_N \end{pmatrix} \quad \text{with} \quad a_j = - \sum_{n \neq j} q_{nj}^{-2} \quad (2.10)$$

then (2.9) is equivalent to

$$\{p_0, L\} = [L, K] \quad \text{and} \quad \{p_0, Q\} = [Q, K] - L$$

This matricial form of the PBs implies

$$\{p_0, I_n\} = \sum_{m=0}^n \text{Tr} L^m [L, K] L^{n-m-1} = \text{Tr} (L^{n-1} K - K L^{n-1}) = 0$$

and

$$\{p_0, J_{n+1}\} = \text{Tr} ([Q, K] - L) L^n + \sum_{m=0}^{n+1} \text{Tr} Q L^m [L, K] L^{n-m} \quad (2.11)$$

$$= \text{Tr} (K Q L^n - Q L^n K) - \text{Tr} L^{n+1} = -(n+1) I_{n+1} \quad (2.12)$$

Hence, (2.9) reproduce the PBs of  $p_0$  with all the generators of  $\mathcal{A}_N$ . Since the correspondence between  $I, J$  and  $p, q$  variables is one-to-one [26], the PB's of  $p_0$  with  $q_i, p_j$  are univocally determined by the PB's of  $p_0$  with  $I_i, J_j$  establishing that (2.9) is correct. ■

It is interesting to remark that this matrix  $K$  (used above) is such that the sum of elements on every line or every column yields 0. It implies that  $K$  commutes with the matrix  $\mu \equiv \sum_{i \neq j} e_{ij}$  which is the moment map used to define the Calogero-Moser model by Hamiltonian reduction of a free motion on the cotangent bundle of the Lie algebra  $M_n(\mathbb{C})$  [13]. However the corresponding Poisson structure is the *first*, not *second* one and this property of  $K$  is therefore not easy to interpret.

**Corollary 2.3** *For any function of  $\vec{q}$  and  $\vec{p}$ , we have*

$$\{p_0, f(\vec{q}, \vec{p})\} = \mathcal{D}f(\vec{q}, \vec{p}) \quad \text{with} \quad \mathcal{D} = \sum_{n=1}^N \left\{ p_n \frac{\partial}{\partial q_n} - 2 \sum_{j \neq n} q_{jn}^{-3} \frac{\partial}{\partial q_j} \right\}$$

Proof: Direct calculation using (2.9). ■

### 3 Second Calogero Poisson brackets

#### 3.1 $N = 2$ Poisson brackets

These second PBs have been already calculated in [26]. We recall them for the sake of completion and show on a simple case the method we use for  $N = 3$ .

From the calculation of section 2.2, one deduces that

$$\{p_1, p_2\} = -\frac{1}{q_{12}^3} \quad \text{and} \quad \{p_j, q_k\} = -\delta_{jk} p_j + z_{jk} \quad \text{with} \quad z_{1j} + z_{2j} = 0 \quad \forall j$$

which leaves us with three unknowns,  $z_{11}$ ,  $z_{22}$  and the PB  $\{q_1, q_2\}$ .

Plugging this partial result into the PBs given in (2.3), one deduce the final form, parametrized as:

$$\begin{aligned} z_{11} = -z_{22} = -z_{21} = z_{12} &= \frac{(p_1 - p_2)}{(q_1 - q_2)^2} \frac{1}{\frac{4}{(q_1 - q_2)^2} - (p_1 - p_2)^2} \\ \{q_1, q_2\} &= \frac{1}{(q_1 - q_2)} \frac{1}{\frac{4}{(q_1 - q_2)^2} - (p_1 - p_2)^2}. \end{aligned} \quad (3.1)$$

As already mentioned, these expressions were calculated in [26], directly from the PBs of the  $I_n$ 's and  $J_m$ 's. However, a direct calculation becomes highly complicated for larger values of  $N$ , and one needs to use the results obtained in sections 2.1 and 2.2. That defines the strategy we will adopt in the next subsection.

#### 3.2 $N = 3$ Calogero

We set the following forms for the  $\{p, p\}$  and  $\{p, q\}$  brackets:

$$\{p_i, p_j\} = -\frac{2}{q_{ij}^3} + x_{ij} \quad \text{and} \quad \{p_i, q_{jk}\} = (\delta_{ik} - \delta_{ji})p_i + z_{i;jk} \quad (3.2)$$

$$\text{with} \quad x_{ij} = -x_{ji} \quad \text{and} \quad z_{i;jk} = -z_{i;kj} \quad (3.3)$$

A careful study of Jacobi identity for the triplet  $p_0, p_1, p_2$  finally yields:

$$x_{12} = x_{23} = x_{31} \equiv x_0 \quad (3.4)$$

Consistency conditions on the  $z$  coefficients read:

$$z_{1;jk} + z_{2;jk} + z_{3;jk} = 0, \quad \forall j \neq k \quad (3.5)$$

$$z_{i;12} + z_{i;23} + z_{i;31} = 0, \quad \forall i \quad (3.6)$$

Explicit resolution of the Poisson bracket structure can then be achieved and yields:

$$z_{i;jk} = - \left( \frac{q_{jk}}{q_{i_2} q_{i_3}} \right)^3 \frac{n_{i;jk}}{\mathbf{d}} \quad \text{with } (i, i_2, i_3) = \text{circ.perm.}(1, 2, 3) \quad (3.7)$$

Two forms of  $n$  coefficients are defined depending on which independent indices are present. One gets:

$$\begin{aligned} n_{i;jk} = & -q_{ij}q_{ik} \left\{ q_{ij}^2 q_{ik}^2 (-q_{ij}^3 p_j^2 + q_{ik}^3 p_k^2) + p_i(p_j - p_k) q_{ij}^2 q_{ik}^2 (q_{ij} + \frac{1}{2} q_{ik}) \mathbf{q}^2 \right. \\ & \left. - p_j p_k q_{ij}^2 q_{ik}^2 q_{jk} (q_{jk}^2 + 3q_{ik} q_{jk}) + \frac{1}{2} \mathbf{q}^2 q_{jk} (q_{ij} + q_{ik})^2 \right\} x_0(i, j, k) \\ & - \mathbf{q}^2 (q_{ij} + q_{ik})^2 \end{aligned} \quad (3.8)$$

$$\begin{aligned} n_{i;ij} = & \left\{ - (q_{ij} q_{jk} q_{ki})^3 (p_j - p_i)(p_j - p_k) + \frac{1}{2} \mathbf{q}^2 q_{ij} q_{ik} q_{jk} (q_{ij} + q_{ik})(q_{ik} + q_{jk}) \right\} x_0(i, j, k) \\ & + 4q_{ik}^3 - 2q_{ij} q_{jk} (q_{ik}^2 + q_{ij} q_{jk}) \end{aligned} \quad (3.9)$$

Here it is understood that  $(i, j, k)$  is any permutation of the three indices 1, 2, 3. In addition one defines:

$$\mathbf{d} = -\mathbf{q}^2 \left( p_1(q_{12} + q_{13}) + p_2(q_{23} + q_{21}) + p_3(q_{31} + q_{32}) \right) \quad (3.10)$$

$$\mathbf{q}^2 = q_{12}^2 + q_{23}^2 + q_{13}^2. \quad (3.11)$$

Finally the single, pure  $q_{ij}$  bracket reads:

$$\begin{aligned} \{q_{12}, q_{23}\} &= -\frac{n_{12} n_{23}}{2\mathbf{d}} \\ n_{12} n_{23} &= q_{12} q_{13} q_{23} x_0 \left\{ (q_{12} q_{13} q_{23})^2 (p_1 p_2 (p_1 - p_2) + p_2 p_3 (p_2 - p_3) + p_1 p_3 (p_1 - p_3)) \right. \\ & \quad \left. + q_{23}^2 (q_{12} - q_{13}) p_1 - q_{13}^2 (q_{23} - q_{12}) p_2 - q_{12}^2 (q_{13} + q_{23}) p_3 \right\} \\ & \quad + \mathbf{q}^2 \left\{ p_1 q_{23} (q_{12} + q_{13}) + p_2 q_{13} (q_{12} + q_{32}) - p_3 q_{12} (q_{13} + q_{23}) \right\} \end{aligned} \quad (3.12)$$

The quantity  $x_0$  is explicitly obtained as a very complicated rational function of all dynamical variables  $p$  and  $q_i - q_j$ . It can be characterized however as the single solution of the differential equation ( $\mathcal{D}$  being defined in Corollary 2.3):

$$\begin{aligned} \mathcal{D}x_0 = & \frac{x_0}{q_{12} q_{23} q_{31} \mathbf{d}} \left\{ q_{23} (q_{12}^4 + q_{13}^4) p_1^2 + q_{31} (q_{21}^4 + q_{23}^4) p_2^2 + q_{12} (q_{31}^4 + q_{31}^4) p_3^2 \right. \\ & + p_2 p_3 q_{23} (2q_{23}^4 + 3q_{12} q_{23}^2 q_{13} - q_{12}^2 q_{13}^2) + p_1 p_2 q_{12} (2q_{12}^4 + 3q_{23} q_{12}^2 q_{13} - q_{23}^2 q_{13}^2) \\ & \left. + p_1 p_3 q_{31} (2q_{13}^4 - 3q_{23} q_{13}^2 q_{12} - q_{23}^2 q_{12}^2) + \frac{(\mathbf{q}^2)^3}{4q_{12} q_{23} q_{31}} \right\} \\ & + \frac{2(q_{13}^6 + q_{12}^6 + q_{23}^6) - 6q_{23}^2 q_{12}^2 q_{13}^2}{(q_{12} q_{23} q_{31})^3 \mathbf{d}} \end{aligned} \quad (3.13)$$

with the particular value at  $q_{12} = q_{23}$

$$x_0 \Big|_{q_{12}=q_{23}} = -18 \frac{81 - 3q_{12}^2(p_3 + p_1 - 2p_2)^2 - 4q_{12}^4(p_2 - p_1)(p_1 - p_3)^2(p_3 - p_2)}{q_{13} q_{12}^2 g(q_{12}) g(-q_{12})} \quad (3.14)$$

with

$$g(q_{12}) = 27 + 27q_{12}(p_3 - p_1) + 4q_{12}^2(2p_3 - p_2 - p_1)(p_2 + p_3 - 2p_1) + 4q_{12}^3(p_2 - p_1)(p_3 - p_1)(p_3 - p_2).$$

Note also that

$$x_0 \sim \frac{-2}{q_{23} q_{13} q_{12}} \quad \text{for } q_{12} \rightarrow 0$$

for  $q_{23}$  finite.

### 3.3 Particular cases

We present here some particular cases where the PBs simplify drastically. They correspond to particular positions of the particles. Remark that in some cases, the particular positions make the Lax formalism ill-defined, but the PBs are themselves well-defined.

#### 3.3.1 Three free particles

If we consider  $q_{12} = q_{23} \rightarrow \infty$ , the three particles are far away one from each other, so that they can be considered as decoupled. Indeed, in that case, the PBs simplify to

$$\{q_i, q_j\} = 0, \quad \{p_i, q_j\} = -\delta_{ij}p_i, \quad \{p_i, p_j\} = -\frac{2}{q_{ij}^3}. \quad (3.15)$$

One recognizes the second PB structure of free particles.

Let us note that to take properly the limit, one has to explicitly use the behavior

$$x_0 \sim \frac{9}{4(p_2 - p_1)(p_2 - p_3)q_{12}^5} \quad \text{when } q_{23} = q_{12} \rightarrow \infty.$$

#### 3.3.2 One free particle

If now we consider that only one particle, say particle 1, is far from the two others, one has to take the limit  $q_{12} \rightarrow \infty$  keeping  $q_{23}$  finite. The first particle (associated to the index 1) decouples while the particles 2 and 3 still interact. Indeed, in that case, the PBs simplify to

$$\{q_1, q_j\} = 0, \quad \{p_j, q_1\} = -\delta_{1j}p_1, \quad \{p_1, q_j\} = -\delta_{1j}p_1, \quad \{p_1, p_j\} = -\frac{2}{q_{1j}^3} \quad \forall j \quad (3.16)$$

$$\{q_2, q_3\} =, \quad \{p_2, p_3\} = -\frac{2}{q_{23}^3} + o(1/q_1) \quad (3.17)$$

$$\{p_j, q_k\} = -\delta_{kj}p_j + z_{kj}, \quad j, k = 2, 3 \quad (3.18)$$

$$z_{2;12} = -2q_{23}/((-q_{23}p_3 + 2 + q_{23}p_2)(-q_{23}p_3 - 2 + q_{23}p_2)) \quad (3.19)$$

$$z_{1;12} = 0, \quad z_{1;23} = 0, \quad \{q_{12}, q_{23}\} = 0. \quad (3.20)$$

One recovers indeed  $z_{jk}$  such as computed in (3.1). Again, one needs to know that for  $q_{12} \rightarrow \infty$ ,

$$x_0 \sim \frac{2}{q_{23}^2(p_1^2 - p_1(p_3 - p_2) + p_2p_3) + 1} \left( \frac{-1/q_{23}}{(q_{23} + q_{12})q_{12}} + \frac{(p_3 - p_2)(2p_1 - p_3 - p_2)q_{23}^2}{(q_{23}^2(p_3 - p_2)^2 - 4)(q_{23} + q_{12})q_{12}^2} \right)$$

to get a correct answer.

## 4 Dynamical r-matrix algebra for $N = 2$ Calogero

### 4.1 The quadratic algebra

Let us now formulate the Poisson bracket structure (3.1) in terms of an  $r$ -matrix structure. We postulate that a quadratic formulation mimicking (1.7) will be adequate for this second Poisson bracket, although both  $a$  and  $s$  matrices will be expected to be dynamical. We recall that given a classical Lax matrix  $\ell$ , the most general quadratic form for the associated Poisson structure is

$$\{\ell_1, \ell_2\} = a_{12} \ell_1 \ell_2 + \ell_1 b_{12} \ell_2 - \ell_2 c_{12} \ell_1 - \ell_1 \ell_2 d_{12} \quad (4.1)$$

where consistency conditions imply that  $a_{12} = -a_{21}$ ,  $d_{12} = -d_{21}$ ,  $b_{12} = c_{21}$ . Note that (4.1) implies that the functions  $\{\text{tr } \ell^m, m \in \mathbb{Z}_+\}$  Poisson-commute if  $a + b = c + d$ . A more general trace formula,  $\text{tr}(\gamma^{-1} \ell)^m$ , occurs whenever a scalar matrix  $\gamma$  exists such that

$$a_{12} \gamma_1 \gamma_2 + \gamma_1 b_{12} \gamma_2 - \gamma_2 c_{12} \gamma_1 - \gamma_1 \gamma_2 d_{12} = 0, \quad (4.2)$$

see [17].

Dynamical dependence of  $abcd$  now is assumed to be solely on coordinates  $q_i$ ,  $i = 1, \dots, n$ , on a dual  $\mathfrak{h}^*$  of the Cartan subalgebra  $\mathfrak{h}$  in  $sl(n, \mathbb{C})$ .

In the 2 sites case we get

$$\ell = \begin{pmatrix} p_1 & \frac{1}{q_{12}} \\ -1 & p_2 \\ \frac{1}{q_{12}} & \end{pmatrix} \quad (4.3)$$

The PB deduced from (3.1) reads

$$\{\ell_1, \ell_2\} = \frac{1}{q_{12}} \begin{pmatrix} 0 & \frac{p_1}{q_{12}} & -\frac{p_1}{q_{12}} & 0 \\ -\frac{p_1}{q_{12}} & -\frac{2}{(q_{12})^2} & 0 & \frac{p_2}{q_{12}} \\ \frac{p_1}{q_{12}} & 0 & \frac{2}{(q_{12})^2} & -\frac{p_2}{q_{12}} \\ 0 & -\frac{p_2}{q_{12}} & \frac{p_2}{q_{12}} & 0 \end{pmatrix} \quad (4.4)$$

As in the linear case (1.3)-(1.4), associativity for the PB structure (4.1) is implied by algebraic consistency conditions (Yang-Baxter classical equations) for  $a, b, c, d$ , provided the *a priori* undetermined bracket  $\{r_{12}, \ell_3\}$ ,  $r = a, b, c, d$ , be of an algebraic form. We postulate here the following form for this PB

$$\{r_{12}, \ell_3\} = \epsilon_R (h_3 \partial r_{12}) \ell_3 + \epsilon_L \ell_3 h_3 \partial r_{12} \quad (4.5)$$

$$h \partial = \sum_{i=1}^n \mu e_{ii} \otimes \frac{\partial}{\partial q_i}, \quad (4.6)$$

where  $e_{ii} \in \mathfrak{h}$ ,  $\epsilon_R, \epsilon_L$  are c-numbers to be determined. Notice the difference in homogeneity factors in  $l$  with respect to the linear case (1.4).

We will see below that this postulate is consistent and the correct choice of parameters  $\epsilon$  is

$$\epsilon_L = \epsilon_R = \frac{1}{2}. \quad (4.7)$$

A solution to express the PB (4.4) as a quadratic form is given by

$$a_{12} = d_{12} = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & \frac{-w_1}{2q_{12}} & \frac{1}{2q_{12}} & 0 \\ 0 & \frac{-1}{2q_{12}} & \frac{w_1}{2q_{12}} & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \quad (4.8)$$

$$b_{12} = c_{21} = c_{12} = \begin{pmatrix} \frac{-w_1}{2q_{12}} & 0 & 0 & \frac{1}{2q_{12}} \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ \frac{-1}{2q_{12}} & 0 & 0 & \frac{w_1}{2q_{12}} \end{pmatrix} \quad (4.9)$$

$$(4.10)$$

where  $w_1$  is a free parameter.

Jacobi identity for the quadratic PB is ensured by the following (sufficient) classical dynamical Yang-Baxter equations:

$$[a_{12}, a_{13}] + [a_{12}, a_{23}] + [a_{32}, a_{13}] + \frac{1}{2} (h_3 \partial a_{12} + h_1 \partial a_{23} + h_2 \partial a_{31}) = 0, \quad (4.11)$$

$$[d_{12}, d_{13}] + [d_{12}, d_{23}] + [d_{32}, d_{13}] + \frac{1}{2} (h_3 \partial d_{12} + h_1 \partial d_{23} + h_2 \partial d_{31}) = 0, \quad (4.12)$$

$$[a_{12}, c_{13} + c_{23}] + [c_{13}, c_{23}] + \frac{1}{2} (-h_3 \partial a_{12} + h_1 \partial c_{23} - h_2 \partial c_{13}) = 0, \quad (4.13)$$

$$[d_{12}, b_{13} + b_{23}] + [b_{13}, b_{23}] + \frac{1}{2} (-h_3 \partial d_{12} + h_1 \partial b_{23} - h_2 \partial b_{13}) = 0. \quad (4.14)$$

In the absence of dynamical term, one would recover the usual classical quadratic algebra [17].

A connection between the classical DYB equations associated with quadratic and linear Poisson brackets is established as follows: In the simplest case of Poisson commutation of traces when  $a + b = c + d$  in (4.1) the matrix  $r \equiv a + b$  obeys the linear dynamical Yang Baxter equation obtained from (1.3), (1.4), that is, with two derivative terms provided that  $a, b, c, d$  obey (4.11)-(4.14), coupled DYB with three derivative terms.

This result however does *not* imply that (1.3) and (4.11)-(4.14) are identified as first and second Poisson structure since the algebraic forms thus obtained require respectively the additional identification of the Poisson brackets of  $l$  with  $r$  or  $a, b, c, d$  as (1.4) or (4.5) which is not implied in any way by the form of PB's (1.3) and (4.11)-(4.14). In other words, contrary to the non-dynamical case (Sklyanin bracket) one cannot establish that the quadratic form (4.11)-(4.14) be a consistent Poisson structure *solely* from the fact that (1.3) be one such structure.

One then immediately observes that (4.11)-(4.14) is a classical limit ( $\hbar \rightarrow 0$ ) of a set of 4 dynamical Yang-Baxter equations first formulated in [25]

$$A_{12}(\mathbf{q}) A_{13}(\mathbf{q} - \epsilon_R h^{(2)}) A_{23}(\mathbf{q}) = A_{23}(\mathbf{q} - \epsilon_R h^{(1)}) A_{13}(\mathbf{q}) A_{12}(\mathbf{q} - \epsilon_R h^{(3)}), \quad (4.15)$$

$$D_{12}(\mathbf{q} + \epsilon_L h^{(3)}) D_{13}(\mathbf{q}) D_{23}(\mathbf{q} + \epsilon_L h^{(1)}) = D_{23}(\mathbf{q}) D_{13}(\mathbf{q} + \epsilon_L h^{(2)}) D_{12}(\mathbf{q}), \quad (4.16)$$

$$A_{12}(\mathbf{q}) C_{13}(\mathbf{q} - \epsilon_R h^{(2)}) C_{23}(\mathbf{q}) = C_{23}(\mathbf{q} - \epsilon_R h^{(1)}) C_{13}(\mathbf{q}) A_{12}(\mathbf{q} + \epsilon_L h^{(3)}), \quad (4.17)$$

$$D_{12}(\mathbf{q} - \epsilon_R h^{(3)}) B_{13}(\mathbf{q}) B_{23}(\mathbf{q} + \epsilon_L h^{(1)}) = B_{23}(\mathbf{q}) B_{13}(\mathbf{q} + \epsilon_L h^{(2)}) D_{12}(\mathbf{q}), \quad (4.18)$$

with the particular choice of "weight parameters"  $\epsilon_L = \epsilon_R = \frac{1}{2}$  and the classical limit is defined by setting

$$R(\mathbf{q}) = \mathbb{I} + \hbar r(\mathbf{q}) + o(\hbar^2), \quad R = A, B, C, D \quad \text{and} \quad r = a, b, c, d \quad (4.19)$$

$$h^{(i)} = \hbar h_i + o(\hbar^3), \quad (4.20)$$

and keeping the order  $\hbar^2$  in (4.15)-(4.18), orders 1 and  $\hbar$  being trivial.

These 4 equations are in turn characterized as sufficient conditions for associativity of a quantum quadratic dynamical exchange algebra defined generically in [25]:

$$A_{12}(\mathbf{q}) K_1(\mathbf{q} - \epsilon_R h^{(2)}) B_{12}(\mathbf{q}) K_2(\mathbf{q} + \epsilon_L h^{(1)}) = K_2(\mathbf{q} - \epsilon_R h^{(1)}) C_{12}(\mathbf{q}) K_1(\mathbf{q} + \epsilon_L h^{(2)}) D_{12}(\mathbf{q}) \quad (4.21)$$

assuming a set of zero-weight conditions

$$\epsilon_R [h^{(1)} + h^{(2)}, A_{12}] = \epsilon_L [h^{(1)} + h^{(2)}, D_{12}] = 0 \quad (4.22)$$

$$[\epsilon_R h^{(1)} - \epsilon_L h^{(2)}, C_{12}] = [\epsilon_L h^{(1)} - \epsilon_R h^{(2)}, B_{12}] = 0, \quad (4.23)$$

and unitary hypothesis

$$A_{12} A_{21} = D_{12} D_{21} = \mathbb{I} \otimes \mathbb{I} \quad ; \quad C_{12} = B_{21}. \quad (4.24)$$

Altogether, these relations ensure associativity of the product in the dynamical algebra.

Note that the free parameter  $w_1$  is the signature in this classical limit of one gauge covariance of the dynamical Yang Baxter equation for  $A$ , pointed out in e.g. [14] under which diagonal coordinates  $d_{ij}$  of  $R$ -matrix on basis elements  $e_{ij} \otimes e_{ji}$  in  $M_n(\mathbb{C}) \otimes M_n(\mathbb{C})$  contain constant parameters  $g_{ij} \equiv f_i - f_j$  and  $f_i$  are arbitrary non-dynamical c-numbers. Here  $\hbar w_1 \equiv f_1 - f_2$ .

## 4.2 Comparison with the first Poisson bracket $r$ -matrix

We recall that the first Poisson bracket of rational Calogero-Moser model is expressed linearly in terms of the Lax matrix  $L$  following the formulation (1.2). The  $r$ -matrix takes the form:

$$r = \sum_{i \neq j} \frac{1}{q_i - q_j} e_{ij} \otimes e_{ji} + \sum_{k \neq j} \frac{1}{q_k - q_j} e_{kk} \otimes e_{kj} \quad (4.25)$$

It is interpreted [20–22, 24] as a combination  $r_{12} = d_{12} + c_{12}$  of two matrices realizing with  $b_{12} \equiv c_{21}$  and  $a = d + c - b$  a classical semi-dynamical reflection algebra corresponding to the choice  $\epsilon_L = 0, \epsilon_R = 1$  in (4.15)-(4.18). In particular the symmetric part of  $r$  is now a sum  $\frac{1}{2}b + c$  of two matrices with respective weights  $(0, 1)$  and  $(1, 0)$  under adjoint action of  $\mathfrak{h} \oplus \mathfrak{h}$ . It is therefore *not* related with the  $abcd$  quadruplet realizing the second Poisson structure of the rational CM Lax matrix, although the  $d$  matrices themselves are identical. The second CM bracket is therefore not realized as a strict Sklyanin-type quadratization of the first CM bracket. This counterexample arising in the simplest available situation for dynamical  $r$ -matrices thus eliminates any possibility of extending directly the Sklyanin-Li-Parmentier procedure for higher Poisson brackets to the case of dynamical  $r$ -matrices.

Curiously enough a quadratic Poisson bracket involving the components  $abcd$  deduced from the  $r$ -matrix of the *first* CM bracket does exist: it arises in the formulation of the *first* PB of the Ruijsenaar Schneider rational Lax matrix [20, 23].

## 5 Open questions

The formulation of a simple algebraic relation à la Sklyanin between first and second Poisson bracket structure in the context of Lax matrices where dynamical  $r$ -matrices arise in the linear expression of their first PB seems thus, if not altogether excluded, at least unreachable at the moment. Other issues remain open at this time:

- Define the crossed  $r$ -matrix formulation for  $L$  and  $Q$ , and the  $r$ -matrix structure for  $Q$  when  $N = 2$ . In this way a complete understanding of the second PB structure including the  $\{I, J\}$  and  $\{J, J\}$  brackets, will be achieved.

- Define the  $r$ -matrix formulation describing the second PB structure for  $N = 2$  Ruijsenaar-Schneider model (starting with the rational case). In fact a form for this second PB structure has been conjectured [15] but not explicitly built in terms of Lax matrix (only in terms of

trace invariants extending naturally the variables  $I$  and  $J$ ). One conjectures here that some cubic- $l$  dependent form will be relevant.

- Define the  $r$ -matrix formulation for  $N = 3$ . The complexity of the expressions for the coordinate Poisson brackets seems to present a difficult technical challenge here. This technical complexity indicates in any case that the  $r$ -matrix quadruplet in a postulated quadratic form will exhibit a dependence on both  $p$  and  $q$  variables, suggesting that the dynamical dependence here goes beyond the Gervais-Neveu-Felder formulation (1.4). A similar issue arose some time ago [16] for  $N \geq 4$  elliptic Calogero-Moser Lax formulation without spectral parameter, and has not been satisfactorily solved since. The issue of  $p, q$  dependant  $r$ -matrices is in any case a yet mostly unexplored one which we hope to come back to in a near future.

## Acknowledgements

This work was sponsored by CNRS, Université de Cergy-Pontoise, Université de Savoie and ANR Project DIADEMS (Programme Blanc ANR SIMI1 2010-BLAN-0120-02). J.A. wishes to thank LAPTh for their kind hospitality.

## References

- [1] F. Calogero, *Exactly solvable one-dimensional Many-body problems*, Lett. Nuov. Cim. **13** (1975) 411; F. Calogero, *On a functional equation connected with integrable Many-body problems*, Lett. Nuov. Cim. **16** (1976) 77.
- [2] J.Moser, *Three integrable Hamiltonian systems connected with isospectral deformations*, Adv. Math. **16** (1975), 1.
- [3] J. Avan, M. Talon, *Classical  $r$ -matrix structure for the Calogero model*, Phys. Lett. **B303** (1993), 33.
- [4] P.D.Lax, *Integrals of non linear equations and solitary waves*, Comm. Pure Appl. Math. **21** (1968), 467
- [5] O. Babelon, D. Bernard, M. Talon, *Introduction to classical integrable systems*, Cambridge Monographs on Mathematical Physics, Cambridge University Press (2003).
- [6] O. Babelon, C.M. Viallet, *Hamiltonian structures and Lax equations*, Phys. Lett. **B237** (1989), 411.
- [7] Ping Xu, *Quantum dynamical yang Baxter equation over a non abelian base*, Comm. Math. Phys. **226** (2002), 475.

- [8] J. Balog, L. Dabrowski, L. Feher, *Classical r-matrix and exchange algebra in WZNW and Toda theories*, Phys. Lett. **B244** (1990), 227.
- [9] W. Oevel, O. Ragnisco, *R-matrices and higher Poisson brackets for integrable systems*, Physica **A161** (1989), 181.
- [10] E.K. Sklyanin, *On some algebraic structures linked with the YB equation*, Funct. Anal. Appl. **16**, 4 (1982), 27.
- [11] L.C. Li, S.Parmentier, *Non linear Poisson structures and r-matrices*, Comm. Math. Phys. **125** (1989), 545.
- [12] M.A. Semenov-Tjan-Shanskii, *What is a classical r-matrix*, Funct. Anal. Appl. **17** (1983), 259;
- [13] M.A. Olshanetski, A.M. Peremolov, *Classical integrable Finite-dimensional Systems related to Lie algebras*, Physics Reports **71** (1981), 313.
- [14] J. Avan, B. Billaud, G. Rollet, *Classification of non-affine non-Hecke dynamical R-matrices*, arXiv:1204.2746
- [15] . I. Aniceto, J. Avan, A. Jevicki, *Poisson Structures of Calogero Moser and Ruijsenaar Schneider models*, J. Phys. **A 43** (2010), 185201
- [16] H.W. Braden, T. Suzuki, *R-matrices for elliptic Calogero-Moser models*, Lett. Math. Phys. **30** (1994), 147
- [17] L. Freidel and J.M. Maillet, *Quadratic algebras and integrable systems*, Phys. Lett. **B262** (1991) 278.
- [18] J.L. Gervais, A. Neveu, *Novel triangle relation and absence of tachyons in Liouville string field theory*, Nucl. Phys. **B238** (1984) 125.
- [19] G. Felder, *Elliptic quantum groups*, Proc. ICMP Paris 1994, pp 211 and arXiv:hep-th/9412207.
- [20] Yu. B. Suris, *Why are the Elliptic Ruijsenaar Schneider and Calogero-Moser hierarchies governed by the same r-matrix ?*, Phys. Lett. **A225** (1997), 253
- [21] G.E. Arutyunov and S.A. Frolov, *Comm. Math. Phys.* **191** (1998) 15-29 and arXiv:q-alg/9610009.
- [22] G.E. Arutyunov, L.O. Chekhov and S.A. Frolov, *Comm. Math. Phys* **192** (1998) 405-432, arXiv:q-alg/9612032.

- [23] S.N.M. Ruijsenaars and H. Schneider, *A new class of integrable systems and its relation to solitons*, Ann. Phys. **170** (1986) 370.
- [24] Z. Nagy, J. Avan and G. Rollet, *Construction of dynamical quadratic algebras*, Lett. Math. Phys. **67** (2004) 1-11 and [arXiv:math/0307026](#).
- [25] J. Avan and E. Ragoucy, *New Dynamical Reflection Algebra and Related Quantum Integrable Systems*, Lett. Math. Phys. **101** (2012) 85 and [arXiv:1106.3264](#).
- [26] C. Bartocci, G. Falqui, I. Mencattini, G. Ortenzi, M. Pedroni *On the geometric origin of the bi-Hamiltonian structure of the Calogero-Moser system* Int. Math. Res. Not. **2010** (2010) 279-296 and [arXiv:0902.0953v2](#).
- [27] F. Magri, P. Casati, G. Falqui, M. Pedroni, *Eight lectures on Integrable Systems*, In: Integrability of Nonlinear Systems (Y. Kosmann-Schwarzbach et al. eds.), Lecture Notes in Physics **495** (2nd edition), 2004, pp. 209-250.
- [28] J.-M. Maillet, *New integrable canonical structures in two-dimensional models*, Nucl.Phys **B269** (1986) 54.
- [29] J.-M. Maillet, *Lax equations and quantum groups*, Phys. Lett. **B245** (1990) 480.