

# Generalisation Bounds for Metric and Similarity Learning \*

Qiong Cao, Zheng-Chu Guo and Yiming Ying

College of Engineering, Mathematics and Physical Sciences  
University of Exeter, Harrison Building, EX4 4QF, UK

## Abstract

Recently, metric learning and similarity learning have attracted a large amount of interest. Many models and optimisation algorithms have been proposed. However, there is relatively little work on the generalisation analysis of such methods. In this paper, we derive novel generalisation bounds of metric and similarity learning. In particular, we first show that the generalisation analysis reduces to the estimation of the Rademacher average over “sums-of-i.i.d.” sample-blocks related to the specific matrix norm. Then, we derive generalisation bounds for metric/similarity learning with different matrix-norm regularisers by estimating their specific Rademacher complexities. Our analysis indicates that sparse metric/similarity learning with  $L^1$ -norm regularisation could lead to significantly better bounds than those with Frobenius-norm regularisation. Our novel generalisation analysis develops and refines the techniques of U-statistics and Rademacher complexity analysis.

## 1 Introduction

The success of many machine learning algorithms (e.g. the nearest neighborhood classification and k-means clustering) depends on the concepts of distance metric and similarity. For instance, k-nearest-neighbor (kNN) classifier depends on a distance function to identify the nearest neighbors for classification; k-means algorithms depend on the pairwise distance measurements between examples for clustering. Kernel methods and information retrieval methods rely on a similarity measure between samples. Many existing studies have been devoted to learning a metric or similarity automatically from data, which is usually referred to as *metric learning* and *similarity learning*, respectively.

---

\*Corresponding author: Yiming Ying. Email: y.ying@exeter.ac.uk

Most work in metric learning focuses on learning a (squared) Mahalanobis distance defined, for any  $x, t \in \mathbb{R}^d$ , by  $d_M(x, t) = (x - t)M(x - t)^\top$  where  $M$  is a positive semi-definite matrix, see e.g. [1, 8, 9, 10, 23, 25, 26, 27, 28]. Concurrently, the pairwise similarity defined by  $s_M(x, t) = xMt^\top$  was studied in [6, 14, 18, 22]. These methods have been successfully applied to various real-world problems including information retrieval and face verification [6, 11, 12, 29]. Although there are a large number of studies devoted to supervised metric/similarity learning based on different objective functions, few studies address the generalisation analysis of such methods. The recent work [13] pioneered the generalisation analysis for metric learning using the concept of uniform stability [4]. However, this approach only works for the strongly convex norm, e.g. the Frobenius norm, and the offset term is fixed which makes the generalisation analysis essentially different.

In this paper, we develop a novel approach for generalisation analysis of metric learning and similarity learning which can deal with general matrix regularisation terms including Frobenius norm [13], sparse  $L^1$ -norm [21], mixed  $(2, 1)$ -norm [28] and trace-norm [28, 23]. In particular, we first show that the generalisation analysis for metric/similarity learning reduces to the estimation of the Rademacher average over “sums-of-i.i.d.” sample-blocks related to the specific matrix norm, which we refer to as the *Rademacher complexity for metric (similarity) learning*. Then, we show how to estimate the Rademacher complexities with different matrix regularisers. Our analysis indicates that sparse metric/similarity learning with  $L^1$ -norm regularisation could lead to significantly better generalisation bounds than that with Frobenius norm regularisation, especially when the dimension of the input data is high. This is nicely consistent with the rationale that sparse methods are more effective for high-dimensional data analysis. Our novel generalisation analysis develops and extends Rademacher complexity analysis [2, 15] to the setting of metric/similarity learning by using techniques of U-statistics [7, 20].

The paper is organized as follows. The next section reviews the models of metric/similarity learning. Section 3 establishes the main theorems. In Section 4, we derive and discuss generalisation bounds for metric/similarity learning with various matrix-norm regularisation terms. Section 5 concludes the paper.

**Notation:** Let  $\mathbb{N}_n = \{1, 2, \dots, n\}$  for any  $n \in \mathbb{N}$ . For any  $X, Y \in \mathbb{R}^{d \times n}$ ,  $\langle X, Y \rangle = \mathbf{Tr}(X^\top Y)$  where  $\mathbf{Tr}(\cdot)$  denotes the trace of a matrix. The space of symmetric  $d$  times  $d$  matrices will be denoted by  $\mathbb{S}^d$ . We equip  $\mathbb{S}^d$  with a general matrix norm  $\|\cdot\|$ ; it can be a Frobenius norm, trace-norm and mixed norm. Its associated dual norm is denoted, for any  $M \in \mathbb{S}^d$ , by  $\|M\|_* = \sup\{\langle X, M \rangle : X \in \mathbb{S}^d, \|X\| \leq 1\}$ . The Frobenius norm on matrices or vector is always denoted by  $\|\cdot\|_F$ . Later on we use the conventional notation that  $X_{ij} = (x_i - x_j)(x_i - x_j)^\top$  and  $\tilde{X}_{ij} = x_i x_j^\top$ .

## 2 Metric/Similarity Learning Formulation

In our learning setting, we have an input space  $\mathcal{X} \subseteq \mathbb{R}^d$  and an output (label) space  $\mathcal{Y}$ . Denote  $\mathcal{Z} = \mathcal{X} \times \mathcal{Y}$  and suppose  $\mathbf{z} := \{z_i = (x_i, y_i) \in \mathcal{Z} : i \in \mathbb{N}_n\}$  an i.i.d. training set according to an unknown distribution  $\rho$  on  $\mathcal{Z}$ . Denote the  $d \times n$  input data matrix by  $\mathbf{X} = (x_i : i \in \mathbb{N}_n)$  and the  $d \times d$  distance matrix by  $M = (M_{\ell k})_{\ell, k \in \mathbb{N}_d}$ . Then, the (pseudo-) distance between  $x_i$  and  $x_j$  is measured by

$$d_M(x_i, x_j) = (x_i - x_j)^\top M (x_i - x_j).$$

The goal of metric learning is to identify a distance function  $d_M(x_i, x_j)$  such that it yields a small value for a similar pair and a large value for a dissimilar pair. The bilinear similarity function is defined by

$$s_M(x_i, x_j) = x_i^\top M x_j.$$

Similarly, the target of similarity learning is to learn  $M \in \mathbb{S}^d$  such that it reports a large similarity value for a similar pair and a small similarity value for a dissimilar pair. It is worth pointing out that we do not require the positive semi-definiteness of the matrix  $M$  throughout this paper. However, we do assume  $M$  to be symmetric, since this will guarantee the distance (similarity) between  $x_i$  and  $x_j$  ( $d_M(x_i, x_j)$ ) is equivalent to that between  $x_j$  and  $x_i$  ( $d_M(x_j, x_i)$ ).

There are two main terms in the metric/similarity learning model: *empirical error* and *matrix regularisation term*. The empirical error function is to employ the similarity and dissimilarity information provided by the label information and the appropriate matrix regularisation term is to avoid overfitting and improve generalisation performance.

For any pair of samples  $(x_i, x_j)$ , let  $r(y_i, y_j) = 1$  if  $y_i = y_j$  otherwise  $r(y_i, y_j) = -1$ . It is expected that there exists an offset term  $b \in \mathbb{R}$  such that  $d_M(x_i, x_j) \leq b$  for  $r(y_i, y_j) = 1$  and  $d_M(x_i, x_j) > b$  otherwise. This naturally leads to the empirical error [13] defined by

$$\frac{1}{n(n-1)} \sum_{i, j \in \mathbb{N}_n, i \neq j} I[r(y_i, y_j)(d_M(x_i, x_j) - b) > 0]$$

where the indicator function  $I[x]$  equal 1 if  $x$  is true and zero otherwise.

Due to the indicator function, the above empirical error is not convex which is difficult to do optimisation. A usual way to overcome this shortcoming is to upper-bound it with a convex loss function. For instance, we can use the hinge loss to upper-bound the indicator function which leads to the following empirical error:

$$\mathcal{E}_{\mathbf{z}}(M, b) := \frac{1}{n(n-1)} \sum_{i, j \in \mathbb{N}_n, i \neq j} [1 + r(y_i, y_j)(d_M(x_i, x_j) - b)]_+ \quad (1)$$

In order to avoid overfitting, we need to enforce a regularisation term denoted by  $\|M\|$ , which will restrict the complexity of the distance matrix. We emphasize here  $\|\cdot\|$  denotes a general matrix norm in the linear space  $\mathbb{S}^d$ . Putting the regularisation term and the empirical error term together yields the following metric learning model:

$$(M_{\mathbf{z}}, b_{\mathbf{z}}) = \arg \min_{M \in \mathbb{S}^d, b \in \mathbb{R}} \{\mathcal{E}_{\mathbf{z}}(M, b) + \lambda \|M\|^2\}, \quad (2)$$

where  $\lambda > 0$  is a trade-off parameter.

Different regularisation terms lead to different metric learning formulations. For instance, the Frobenius norm  $\|M\|_F$  is used in [13]. To favor the element-sparsity, [21] introduced the  $L^1$ -norm regularisation  $\|M\| = \sum_{\ell, k \in \mathbb{N}_d} |M_{\ell k}|$ . [28] proposed the mixed  $(2, 1)$ -norm  $\|M\| = \sum_{\ell \in \mathbb{N}_d} \left( \sum_{k \in \mathbb{N}_d} |M_{\ell k}|^2 \right)^{\frac{1}{2}}$  to encourage the column-wise sparsity of the distance matrix. The trace-norm regularisation  $\|M\| = \sum_{\ell} \sigma_{\ell}(M)$  was also considered by [28, 23]. Here,  $\{\sigma_{\ell} : \ell \in \mathbb{N}_d\}$  denote the singular values of a matrix  $M \in \mathbb{S}^d$ . Since  $M$  is symmetric, the singular values of  $M$  are identical to the absolute values of its eigenvalues.

In analogy to the formulation of metric learning, we consider the following empirical error for similarity learning [18, 6]:

$$\tilde{\mathcal{E}}_{\mathbf{z}}(M, b) := \frac{1}{n(n-1)} \sum_{i, j \in \mathbb{N}_n, i \neq j} [1 - r(y_i, y_j)(s_M(x_i, x_j) - b)]_+. \quad (3)$$

This leads to the regularised formulation for similarity learning defined as follows:

$$(\tilde{M}_{\mathbf{z}}, \tilde{b}_{\mathbf{z}}) = \arg \min_{M \in \mathbb{S}^d, b \in \mathbb{R}} \{\tilde{\mathcal{E}}_{\mathbf{z}}(M, b) + \lambda \|M\|^2\}. \quad (4)$$

[18] used the Frobenius-norm regularisation for similarity learning. The trace-norm regularisation has been used by [22] to encourage a low-rank similarity matrix  $M$ .

### 3 Statistical Generalisation Analysis

In this section, we mainly give a detailed proof of generalisation bounds for metric and similarity learning. In particular, we develop a novel line of generalisation analysis for metric and similarity learning with general matrix regularisation terms. The key observation is that the empirical data term  $\mathcal{E}_{\mathbf{z}}(M, b)$  for metric learning is a modification of U-statistics and it is expected to converge to its expected form defined by

$$\mathcal{E}(M, b) = \iint (1 + r(y, y')(d_M(x, x') - b))_+ d\rho(x, y) d\rho(x', y'). \quad (5)$$

The empirical term  $\tilde{\mathcal{E}}_{\mathbf{z}}(M, b)$  for similarity learning is expected to converge to

$$\tilde{\mathcal{E}}(M, b) = \iint (1 - r(y, y')(s_M(x, x') - b))_+ d\rho(x, y) d\rho(x', y'). \quad (6)$$

The target of generalisation analysis is to bound the true error  $\mathcal{E}(M_{\mathbf{z}}, b_{\mathbf{z}})$  by the empirical error  $\mathcal{E}_{\mathbf{z}}(M_{\mathbf{z}}, b_{\mathbf{z}})$  for metric learning and  $\tilde{\mathcal{E}}(\tilde{M}_{\mathbf{z}}, \tilde{b}_{\mathbf{z}})$  by the empirical error  $\tilde{\mathcal{E}}_{\mathbf{z}}(\tilde{M}_{\mathbf{z}}, \tilde{b}_{\mathbf{z}})$  for similarity learning.

In the sequel, we provide a detailed proof for generalisation bounds of metric learning. Since the proof for similarity learning is exactly the same as that for metric learning, we only mention the results followed with some brief comments.

### 3.1 Bounding the Solutions

By the definition of  $(M_{\mathbf{z}}, b_{\mathbf{z}})$ , we know that

$$\mathcal{E}_{\mathbf{z}}(M_{\mathbf{z}}, b_{\mathbf{z}}) + \lambda \|M_{\mathbf{z}}\|^2 \leq \mathcal{E}_{\mathbf{z}}(0, 0) + \lambda \|0\|^2 = 1$$

which implies that

$$\|M_{\mathbf{z}}\| \leq \frac{1}{\sqrt{\lambda}}. \quad (7)$$

Now we turn our attention to deriving the bound of the offset term  $b_{\mathbf{z}}$  by modifying the techniques in [5] which was originally developed to estimate the offset term of the soft-margin SVM.

**Lemma 1.** *For any samples  $\mathbf{z}$  and  $\lambda > 0$ , there exists a minimizer  $(M_{\mathbf{z}}, b_{\mathbf{z}})$  of problem (2) such that*

$$\min_{i \neq j} [d_{M_{\mathbf{z}}}(x_i, x_j) - b_{\mathbf{z}}] \leq 1, \quad \max_{i \neq j} [d_{M_{\mathbf{z}}}(x_i, x_j) - b_{\mathbf{z}}] \geq -1. \quad (8)$$

*Proof.* Firstly we prove the inequality  $\min_{i \neq j} [d_{M_{\mathbf{z}}}(x_i, x_j) - b_{\mathbf{z}}] \leq 1$ . To this end, we first consider the special case where the training set  $\mathbf{z}$  only contains two examples  $z_1 = (x_1, y_1)$  and  $z_2 = (x_2, y_2)$  with distinct labels, i.e.  $y_1 \neq y_2$ . For any  $\lambda > 0$ , let  $(M_{\mathbf{z}}, b_{\mathbf{z}}) = (\mathbf{0}, -1)$ , and observe that  $\mathcal{E}_{\mathbf{z}}(\mathbf{0}, -1) + \lambda \|\mathbf{0}\|^2 = 0$ . This observation implies that  $(M_{\mathbf{z}}, b_{\mathbf{z}})$  is a minimizer of problem (2). Consequently, we have the desired result since  $\min_{i \neq j} [d_{M_{\mathbf{z}}}(x_i, x_j) - b_{\mathbf{z}}] = d_{M_{\mathbf{z}}}(x_1, x_2) - b_{\mathbf{z}} = 1$ .

Now let us consider the general case where the training set  $\mathbf{z}$  has at least two examples with the same label. In this case, we prove the inequality by contradiction. Suppose that  $r = \min_{i \neq j} [d_{M_{\mathbf{z}}}(x_i, x_j) - b_{\mathbf{z}}] > 1$  which equivalently implies that  $d_{M_{\mathbf{z}}}(x_i, x_j) - (b_{\mathbf{z}} + r - 1) \geq 1$  for any  $i \neq j$ . Hence, for any  $i \neq j$  and any pair of examples  $(x_i, x_j)$  with distinct labels, i.e.  $y_i \neq y_j$  (equivalently  $r(y_i, y_j) = -1$ ), there holds

$$(1 + r(y_i, y_j)(d_{M_{\mathbf{z}}}(x_i, x_j) - b_{\mathbf{z}} - r + 1))_+ = (1 - (d_{M_{\mathbf{z}}}(x_i, x_j) - b_{\mathbf{z}} - r + 1))_+ = 0.$$

Consequently,

$$\begin{aligned}
\mathcal{E}_{\mathbf{z}}(M_{\mathbf{z}}, b_{\mathbf{z}} + r - 1) &= \frac{1}{n(n-1)} \sum_{i \neq j} \left( 1 + r(i, j)(d_{M_{\mathbf{z}}}(x_i, x_j) - b_{\mathbf{z}} - r + 1) \right)_+ \\
&= \frac{1}{n(n-1)} \sum_{i \neq j, y_i = y_j} (1 + d_{M_{\mathbf{z}}}(x_i, x_j) - b_{\mathbf{z}} - (r - 1))_+ \\
&< \frac{1}{n(n-1)} \sum_{i \neq j, y_i = y_j} (1 + d_{M_{\mathbf{z}}}(x_i, x_j) - b_{\mathbf{z}})_+ \leq \mathcal{E}_{\mathbf{z}}(M_{\mathbf{z}}, b_{\mathbf{z}}).
\end{aligned}$$

The above estimation implies that  $\mathcal{E}_{\mathbf{z}}(M_{\mathbf{z}}, b_{\mathbf{z}} + r - 1) + \lambda \|M_{\mathbf{z}}\| < \mathcal{E}_{\mathbf{z}}(M_{\mathbf{z}}, b_{\mathbf{z}}) + \lambda \|M_{\mathbf{z}}\|$  which contradicts the definition of the minimizer  $(M_{\mathbf{z}}, b_{\mathbf{z}})$ . Hence,  $r = \min_{i \neq j} [d_{M_{\mathbf{z}}}(x_i, x_j) - b_{\mathbf{z}}] \leq 1$ .

Secondly, we prove the inequality  $\max_{i \neq j} [d_{M_{\mathbf{z}}}(x_i, x_j) - b_{\mathbf{z}}] \geq -1$  in analogy to the above argument. Consider a special case where the training set  $\mathbf{z}$  contains only two examples  $z_1 = (x_1, y_1)$  and  $z_2 = (x_2, y_2)$  with the same label, i.e.  $y_1 = y_2$ . For any given  $\lambda > 0$ , let  $(M_{\mathbf{z}}, b_{\mathbf{z}}) = (\mathbf{0}, 1)$ . Since  $\mathcal{E}_{\mathbf{z}}(\mathbf{0}, 1) + \lambda \|\mathbf{0}\|^2 = 0$ ,  $(\mathbf{0}, 1)$  is a minimizer of problem (2). The desired estimation follows from the fact that  $\max_{i \neq j} d_{M_{\mathbf{z}}}(x_i, x_j) - b_{\mathbf{z}} = 0 - 1 = -1$ .

Now let us consider the general case where the training set  $\mathbf{z}$  has at least two examples with distinct labels. We prove the estimation by contradiction. Assume  $r = \max_{i \neq j} d_{M_{\mathbf{z}}}(x_i, x_j) - b_{\mathbf{z}} < -1$ , then  $d_{M_{\mathbf{z}}}(x_i, x_j) - (b_{\mathbf{z}} + r + 1) \leq -1$  holds for any  $i \neq j$ . This implies, for any pair of examples  $(x_i, x_j)$  with the same label, i.e.  $r(i, j) = 1$ , that  $\left( 1 + r(i, j)(d_{M_{\mathbf{z}}}(x_i, x_j) - b_{\mathbf{z}} - r - 1) \right)_+ = 0$ . Hence,

$$\begin{aligned}
\mathcal{E}_{\mathbf{z}}(M_{\mathbf{z}}, b_{\mathbf{z}} + r + 1) &= \frac{1}{n(n-1)} \sum_{i \neq j} \left( 1 + r(i, j)(d_{M_{\mathbf{z}}}(x_i, x_j) - b_{\mathbf{z}} - r - 1) \right)_+ \\
&= \frac{1}{n(n-1)} \sum_{i \neq j, y_i \neq y_j} \left( 1 - d_{M_{\mathbf{z}}}(x_i, x_j) + b_{\mathbf{z}} + (r + 1) \right)_+ \\
&< \frac{1}{n(n-1)} \sum_{i \neq j, y_i \neq y_j} (1 - d_{M_{\mathbf{z}}}(x_i, x_j) + b_{\mathbf{z}})_+ \leq \mathcal{E}_{\mathbf{z}}(M_{\mathbf{z}}, b_{\mathbf{z}}).
\end{aligned}$$

The above estimation yields that  $\mathcal{E}_{\mathbf{z}}(M_{\mathbf{z}}, b_{\mathbf{z}} + r + 1) + \lambda \|M_{\mathbf{z}}\|^2 < \mathcal{E}_{\mathbf{z}}(M_{\mathbf{z}}, b_{\mathbf{z}}) + \lambda \|M_{\mathbf{z}}\|^2$  which contradicts the definition of the minimizer  $(M_{\mathbf{z}}, b_{\mathbf{z}})$ . Hence, we have the desired inequality  $\max_{i \neq j} d_{M_{\mathbf{z}}}(x_i, x_j) - b_{\mathbf{z}} \geq -1$  which completes the proof of the lemma.  $\square$

**Corollary 2.** *For any samples  $\mathbf{z}$  and  $\lambda > 0$ , there exists a minimizer  $(M_{\mathbf{z}}, b_{\mathbf{z}})$  of problem (2) such that*

$$|b_{\mathbf{z}}| \leq 1 + \left( \max_{i \neq j} \|X_{ij}\|_* \right) \|M_{\mathbf{z}}\|. \quad (9)$$

*Proof.* Recall that  $X_{ij} = (x_i - x_j)(x_i - x_j)^\top$  and observe, by the definition of the dual norm  $\|\cdot\|_*$ , that

$$d_M(x_i, x_j) = \langle X_{ij}, M \rangle \leq \|X_{ij}\|_* \|M\|.$$

Using the above observation, estimation (9) follows directly from inequality (8). This completes the proof.  $\square$

Denote

$$\mathcal{F} = \left\{ (M, b) : \|M\| \leq 1/\sqrt{\lambda}, \quad |b| \leq 1 + X_* \|M\| \right\}, \quad (10)$$

where

$$X_* = \sup_{x, x' \in \mathcal{X}} \|(x - x')(x - x')^\top\|_*.$$

From the above corollary, for any samples  $\mathbf{z}$  we can easily see that the optimal solution  $(M_{\mathbf{z}}, b_{\mathbf{z}})$  of formulation (2) belongs to the bounded set  $\mathcal{F} \subseteq \mathbb{S}^d \times \mathbb{R}$ .

We end this subsection with two remarks. Firstly, in what follows, we restrict our attention to the minimizer  $(M_{\mathbf{z}}, b_{\mathbf{z}})$  of formulation (2) which satisfies inequality (9). Secondly, our formulation (2) for metric learning focused on the hinge loss which is widely used in the community of metric learning, see e.g [13, 25, 29]. Similar results to those in the above corollary can easily be obtained for  $q$ -norm loss given, for any  $x \in \mathbb{R}$ , by  $(1 - x)_+^q$  with  $q > 1$ . However, it still remains a question to us on how to estimate the term  $b$  for general loss functions.

### 3.2 Generalisation Bounds

Before stating the generalisation bounds, we introduce some notations. For any  $z = (x, y), z' = (x', y') \in \mathcal{Z}$ , let  $\Phi_{M,b}(z, z') = (1 + r(y, y')(d_M(x, x') - b))_+$ . Hence, for any  $(M, b) \in \mathcal{F}$ ,

$$\sup_{z, z'} \sup_{(M,b) \in \mathcal{F}} \Phi_{M,b}(z, z') \leq B_\lambda := 2(1 + X_*/\sqrt{\lambda}). \quad (11)$$

Let  $\lfloor \frac{n}{2} \rfloor$  denote the largest integer less than  $\frac{n}{2}$  and recall the definition that  $X_{ij} = (x_i - x_j)(x_i - x_j)^\top$ . We now define Rademacher average over sums-of-i.i.d. sample-blocks related to the dual matrix norm  $\|\cdot\|_*$  by

$$\widehat{R}_n = \frac{1}{\lfloor \frac{n}{2} \rfloor} \mathbb{E}_\sigma \left\| \sum_{i=1}^{\lfloor \frac{n}{2} \rfloor} \sigma_i X_{i(\lfloor \frac{n}{2} \rfloor + i)} \right\|_*, \quad (12)$$

and its expectation is denoted by  $R_n = \mathbb{E}_{\mathbf{z}} [\widehat{R}_n]$ . Our main theorem below shows that the generalisation bounds for metric learning critically depend on the quantity of  $R_n$ . For this reason, we refer to  $R_n$  as the *Rademacher complexity for metric learning*. It is worth mentioning that metric learning formulation (2) depends on the norm  $\|\cdot\|$  of the linear space  $\mathbb{S}^d$  and the Rademacher complexity  $R_n$  is related to its dual norm  $\|\cdot\|_*$ .

**Theorem 3.** Let  $(M_{\mathbf{z}}, b_{\mathbf{z}})$  be the solution of formulation (2). Then, for any  $0 < \delta < 1$ , with probability  $1 - \delta$  we have that

$$\begin{aligned} \mathcal{E}(M_{\mathbf{z}}, b_{\mathbf{z}}) - \mathcal{E}_{\mathbf{z}}(M_{\mathbf{z}}, b_{\mathbf{z}}) &\leq \sup_{(M,b) \in \mathcal{F}} \left[ \mathcal{E}(M, b) - \mathcal{E}_{\mathbf{z}}(M, b) \right] \\ &\leq \frac{4R_n}{\sqrt{\lambda}} + \frac{4(3+2X_*/\sqrt{\lambda})}{\sqrt{n}} + 2(1 + X_*/\sqrt{\lambda}) \left( \frac{2\ln(\frac{1}{\delta})}{n} \right)^{\frac{1}{2}}. \end{aligned} \quad (13)$$

*Proof.* The proof of the theorem can be divided into three steps as follows.

**Step 1:** Let  $\mathbb{E}_{\mathbf{z}}$  denote the expectation with respect to samples  $\mathbf{z}$ . Observe that  $\mathcal{E}(M_{\mathbf{z}}, b_{\mathbf{z}}) - \mathcal{E}_{\mathbf{z}}(M_{\mathbf{z}}, b_{\mathbf{z}}) \leq \sup_{(M,b) \in \mathcal{F}} \left[ \mathcal{E}(M, b) - \mathcal{E}_{\mathbf{z}}(M, b) \right]$ . For any  $\mathbf{z} = (z_1, \dots, z_{k-1}, z_k, z_{k+1}, \dots, z_n)$  and  $\mathbf{z}' = (z_1, \dots, z_{k-1}, z'_k, z_{k+1}, \dots, z_n)$  we know from inequality (11) that

$$\begin{aligned} &\left| \sup_{(M,b) \in \mathcal{F}} \left[ \mathcal{E}(M, b) - \mathcal{E}_{\mathbf{z}}(M, b) \right] - \sup_{(M,b) \in \mathcal{F}} \left[ \mathcal{E}(M, b) - \mathcal{E}_{\mathbf{z}'}(M, b) \right] \right| \\ &\leq \sup_{(M,b) \in \mathcal{F}} |\mathcal{E}_{\mathbf{z}}(M, b) - \mathcal{E}_{\mathbf{z}'}(M, b)| \\ &= \frac{1}{n(n-1)} \sup_{(M,b) \in \mathcal{F}} \sum_{j \in \mathbb{N}_n, j \neq k} |\Phi_{M,b}(z_k, z_j) - \Phi_{M,b}(z'_k, z_j)| \\ &\leq \frac{1}{n(n-1)} \sup_{(M,b) \in \mathcal{F}} \sum_{j \in \mathbb{N}_n, j \neq k} |\Phi_{M,b}(z_k, z_j)| + |\Phi_{M,b}(z'_k, z_j)| \\ &\leq 4(1 + X_*/\sqrt{\lambda})/n. \end{aligned}$$

Applying McDiarmid's inequality [19] (see Lemma 6 in the Appendix) to the term  $\sup_{(M,b) \in \mathcal{F}} \left[ \mathcal{E}(M, b) - \mathcal{E}_{\mathbf{z}}(M, b) \right]$ , with probability  $1 - \delta$  there holds

$$\begin{aligned} \sup_{(M,b) \in \mathcal{F}} \left[ \mathcal{E}(M, b) - \mathcal{E}_{\mathbf{z}}(M, b) \right] &\leq \mathbb{E}_{\mathbf{z}} \sup_{(M,b) \in \mathcal{F}} \left[ \mathcal{E}(M, b) - \mathcal{E}_{\mathbf{z}}(M, b) \right] \\ &\quad + 2(1 + X_*/\sqrt{\lambda}) \left( \frac{2\ln(\frac{1}{\delta})}{n} \right)^{\frac{1}{2}}. \end{aligned} \quad (14)$$

Now we only need to estimate the first term in the expectation form on the right-hand side of the above equation by symmetrization techniques.

**Step 2:** To estimate  $\mathbb{E}_{\mathbf{z}} \sup_{(M,b) \in \mathcal{F}} \left[ \mathcal{E}(M, b) - \mathcal{E}_{\mathbf{z}}(M, b) \right]$ , applying Lemma 7 with  $q_{(M,b)}(z_i, z_j) = \mathcal{E}(M, b) - (1 + r(y_i, y_j)(d_M(x_i, x_j) - b))_+$  implies that

$$\mathbb{E}_{\mathbf{z}} \sup_{(M,b) \in \mathcal{F}} \left[ \mathcal{E}(M, b) - \mathcal{E}_{\mathbf{z}}(M, b) \right] \leq \mathbb{E}_{\mathbf{z}} \sup_{(M,b) \in \mathcal{F}} \left[ \mathcal{E}(M, b) - \bar{\mathcal{E}}_{\mathbf{z}}(M, b) \right], \quad (15)$$

where  $\bar{\mathcal{E}}_{\mathbf{z}}(M, b) = \frac{1}{\lfloor \frac{n}{2} \rfloor} \sum_{i=1}^{\lfloor \frac{n}{2} \rfloor} \Phi_{M,b}(z_i, z_{\lfloor \frac{n}{2} \rfloor + i})$ . Now let  $\bar{\mathbf{z}} = \{\bar{z}_1, \bar{z}_2, \dots, \bar{z}_n\}$  be i.i.d. samples which are independent of  $\mathbf{z}$ , then

$$\begin{aligned} \mathbb{E}_{\mathbf{z}} \sup_{(M,b) \in \mathcal{F}} \left[ \mathcal{E}(M, b) - \bar{\mathcal{E}}_{\mathbf{z}}(M, b) \right] &= \mathbb{E}_{\mathbf{z}} \sup_{(M,b) \in \mathcal{F}} \left[ \mathbb{E}_{\bar{\mathbf{z}}} \left[ \bar{\mathcal{E}}_{\bar{\mathbf{z}}}(M, b) \right] - \bar{\mathcal{E}}_{\mathbf{z}}(M, b) \right] \\ &\leq \mathbb{E}_{\mathbf{z}, \bar{\mathbf{z}}} \sup_{(M,b) \in \mathcal{F}} \left[ \bar{\mathcal{E}}_{\bar{\mathbf{z}}}(M, b) - \bar{\mathcal{E}}_{\mathbf{z}}(M, b) \right] \end{aligned} \quad (16)$$

By standard symmetrization techniques (see e.g. [2]), for i.i.d. Rademacher variables  $\{\sigma_i \in \{\pm 1\} : i \in \mathbb{N}_{\lfloor \frac{n}{2} \rfloor}\}$ , we have that

$$\begin{aligned} &\mathbb{E}_{\mathbf{z}, \bar{\mathbf{z}}} \sup_{(M,b) \in \mathcal{F}} \left[ \bar{\mathcal{E}}_{\bar{\mathbf{z}}}(M, b) - \bar{\mathcal{E}}_{\mathbf{z}}(M, b) \right] \\ &= \mathbb{E}_{\mathbf{z}, \bar{\mathbf{z}}} \frac{1}{\lfloor \frac{n}{2} \rfloor} \sup_{(M,b) \in \mathcal{F}} \sum_{i=1}^{\lfloor \frac{n}{2} \rfloor} \sigma_i \left[ \Phi_{M,b}(\bar{z}_i, \bar{z}_{\lfloor \frac{n}{2} \rfloor + i}) - \Phi_{M,b}(z_i, z_{\lfloor \frac{n}{2} \rfloor + i}) \right] \\ &= 2 \mathbb{E}_{\mathbf{z}, \sigma} \frac{1}{\lfloor \frac{n}{2} \rfloor} \sup_{(M,b) \in \mathcal{F}} \sum_{i=1}^{\lfloor \frac{n}{2} \rfloor} \sigma_i \Phi_{M,b}(\bar{z}_i, \bar{z}_{\lfloor \frac{n}{2} \rfloor + i}) \\ &\leq 2 \mathbb{E}_{\mathbf{z}, \sigma} \frac{1}{\lfloor \frac{n}{2} \rfloor} \sup_{(M,b) \in \mathcal{F}} \left| \sum_{i=1}^{\lfloor \frac{n}{2} \rfloor} \sigma_i \Phi_{M,b}(\bar{z}_i, \bar{z}_{\lfloor \frac{n}{2} \rfloor + i}) \right|. \end{aligned} \quad (17)$$

Applying the contraction property of Rademacher averages (see Lemma 8 in the Appendix) with  $\Psi_i(t) = (1 + r(y_i, y_{\lfloor \frac{n}{2} \rfloor + i})t)_+ - 1$ , we have the following estimation for the last term on the righthand side of the above inequality:

$$\begin{aligned} &\mathbb{E}_{\sigma} \frac{1}{\lfloor \frac{n}{2} \rfloor} \sup_{(M,b) \in \mathcal{F}} \left| \sum_{i=1}^{\lfloor \frac{n}{2} \rfloor} \sigma_i \Phi_{M,b}(\bar{z}_i, \bar{z}_{\lfloor \frac{n}{2} \rfloor + i}) \right| \\ &\leq \mathbb{E}_{\sigma} \frac{1}{\lfloor \frac{n}{2} \rfloor} \sup_{(M,b) \in \mathcal{F}} \left| \sum_{i=1}^{\lfloor \frac{n}{2} \rfloor} \sigma_i (\Phi_{M,b}(\bar{z}_i, \bar{z}_{\lfloor \frac{n}{2} \rfloor + i}) - 1) \right| + \frac{1}{\lfloor \frac{n}{2} \rfloor} \mathbb{E}_{\sigma} \left| \sum_{i=1}^{\lfloor \frac{n}{2} \rfloor} \sigma_i \right| \\ &\leq \frac{2}{\lfloor \frac{n}{2} \rfloor} \mathbb{E}_{\sigma} \sup_{(M,b) \in \mathcal{F}} \left| \sum_{i=1}^{\lfloor \frac{n}{2} \rfloor} \sigma_i (d_M(x_i, x_{\lfloor \frac{n}{2} \rfloor + i}) - b) \right| + \frac{1}{\lfloor \frac{n}{2} \rfloor} \mathbb{E}_{\sigma} \left| \sum_{i=1}^{\lfloor \frac{n}{2} \rfloor} \sigma_i \right| \\ &\leq \frac{2}{\lfloor \frac{n}{2} \rfloor} \mathbb{E}_{\sigma} \sup_{\|M\| \leq \frac{1}{\sqrt{\lambda}}} \left| \sum_{i=1}^{\lfloor \frac{n}{2} \rfloor} \sigma_i d_M(x_i, x_{\lfloor \frac{n}{2} \rfloor + i}) \right| + \frac{(3 + 2X_*/\sqrt{\lambda})}{\lfloor \frac{n}{2} \rfloor} \mathbb{E}_{\sigma} \left| \sum_{i=1}^{\lfloor \frac{n}{2} \rfloor} \sigma_i \right| \end{aligned} \quad (18)$$

**Step 3** : It remains to estimate the terms on the righthand side of inequality (18). To this end, observe that

$$\mathbb{E}_{\sigma} \left| \sum_{i=1}^{\lfloor \frac{n}{2} \rfloor} \sigma_i \right| \leq \left( \mathbb{E}_{\sigma} \left| \sum_{i=1}^{\lfloor \frac{n}{2} \rfloor} \sigma_i \right|^2 \right)^{\frac{1}{2}} \leq \sqrt{\lfloor \frac{n}{2} \rfloor}.$$

Moreover,

$$\begin{aligned} \mathbb{E}_\sigma \sup_{\|M\| \leq \frac{1}{\sqrt{\lambda}}} \left| \sum_{i=1}^{\lfloor \frac{n}{2} \rfloor} \sigma_i d_M(x_i, x_{\lfloor \frac{n}{2} \rfloor + i}) \right| &= \mathbb{E}_\sigma \sup_{\|M\| \leq \frac{1}{\sqrt{\lambda}}} \left| \left\langle \sum_{i=1}^{\lfloor \frac{n}{2} \rfloor} \sigma_i (x_i - x_{\lfloor \frac{n}{2} \rfloor + i})(x_i - x_{\lfloor \frac{n}{2} \rfloor + i})^\top, M \right\rangle \right| \\ &\leq \frac{1}{\sqrt{\lambda}} \mathbb{E}_\sigma \left\| \sum_{i=1}^{\lfloor \frac{n}{2} \rfloor} \sigma_i X_{i(\lfloor \frac{n}{2} \rfloor + i)} \right\|_*. \end{aligned}$$

Putting the above estimations and inequalities (17), (18) together yields that

$$\mathbb{E}_{\mathbf{z}, \bar{\mathbf{z}}} \sup_{(M, b) \in \mathcal{F}} \left[ \bar{\mathcal{E}}_{\bar{\mathbf{z}}}(M, b) - \bar{\mathcal{E}}_{\mathbf{z}}(M, b) \right] \leq \frac{2(3 + 2X_*/\sqrt{\lambda})}{\sqrt{\lfloor \frac{n}{2} \rfloor}} + \frac{4R_n}{\sqrt{\lambda}} \leq \frac{4(3 + X_*/\sqrt{\lambda})}{\sqrt{n}} + \frac{2R_n}{\sqrt{\lambda}}.$$

Consequently, combining this with inequalities (15), (16) implies that

$$\mathbb{E}_{\mathbf{z}} \sup_{(M, b) \in \mathcal{F}} \left[ \mathcal{E}(M, b) - \mathcal{E}_{\mathbf{z}}(M, b) \right] \leq \frac{4(3 + 2X_*/\sqrt{\lambda})}{\sqrt{n}} + \frac{4R_n}{\sqrt{\lambda}}.$$

Putting this estimation with (14) completes the proof the theorem.  $\square$

In the setting of similarity learning,  $X_*$  and  $R_n$  are replaced by

$$\tilde{X}_* = \sup_{x, t \in \mathcal{X}} \|xt^\top\|_* \quad \text{and} \quad \tilde{R}_n = \frac{1}{\lfloor \frac{n}{2} \rfloor} \mathbb{E}_{\mathbf{z}} \mathbb{E}_\sigma \left\| \sum_{i=1}^{\lfloor \frac{n}{2} \rfloor} \sigma_i \tilde{X}_{i(\lfloor \frac{n}{2} \rfloor + i)} \right\|_*, \quad (19)$$

where  $\tilde{X}_{i(\lfloor \frac{n}{2} \rfloor + i)} = x_i x_{\lfloor \frac{n}{2} \rfloor + i}^\top$ . Let  $\tilde{\mathcal{F}} = \left\{ (M, b) : \|M\| \leq 1/\sqrt{\lambda}, |b| \leq 1 + \tilde{X}_* \|M\| \right\}$ . Using the exactly same argument as above, we can prove the following bound for similarity learning formulation (4).

**Theorem 4.** *Let  $(\tilde{M}_{\mathbf{z}}, \tilde{b}_{\mathbf{z}})$  be the solution of formulation (4). Then, for any  $0 < \delta < 1$ , with probability  $1 - \delta$  we have that*

$$\begin{aligned} \tilde{\mathcal{E}}(\tilde{M}_{\mathbf{z}}, \tilde{b}_{\mathbf{z}}) - \tilde{\mathcal{E}}_{\mathbf{z}}(\tilde{M}_{\mathbf{z}}, \tilde{b}_{\mathbf{z}}) &\leq \sup_{(M, b) \in \tilde{\mathcal{F}}} \left[ \tilde{\mathcal{E}}(M, b) - \tilde{\mathcal{E}}_{\mathbf{z}}(M, b) \right] \\ &\leq \frac{4\tilde{R}_n}{\sqrt{\lambda}} + \frac{4(3 + 2\tilde{X}_*/\sqrt{\lambda})}{\sqrt{n}} + 2(1 + \tilde{X}_*/\sqrt{\lambda}) \left( \frac{2 \ln(\frac{1}{\delta})}{n} \right)^{\frac{1}{2}}. \end{aligned} \quad (20)$$

## 4 Estimation of $R_n$ and Discussion

From Theorem 3, we need to estimate the Rademacher average for metric learning, i.e.  $R_n$ , and the quantity  $X_*$  for different matrix regularisation terms. Without loss of generality, we only focus on popular matrix norms such as the Frobenius norm [13],  $L^1$ -norm [21], trace-norm [28, 23] and mixed (2, 1)-norm [28].

**Example 1** (Frobenius norm). Let the matrix norm be the Frobenius norm i.e.  $\|M\| = \|M\|_F$ , then the quantity  $X_* = \sup_{x, x' \in \mathcal{X}} \|x - x'\|_F^2$  and the Rademacher complexity is estimated as follows:

$$R_n \leq \frac{2X_*}{\sqrt{n}} = \frac{2 \sup_{x, x' \in \mathcal{X}} \|x - x'\|_F^2}{\sqrt{n}}.$$

Let  $(M_{\mathbf{z}}, b_{\mathbf{z}})$  be a solution of formulation (2) with Frobenius norm regularisation. For any  $0 < \delta < 1$ , with probability  $1 - \delta$  there holds

$$\begin{aligned} \mathcal{E}(M_{\mathbf{z}}, b_{\mathbf{z}}) - \mathcal{E}_{\mathbf{z}}(M_{\mathbf{z}}, b_{\mathbf{z}}) &\leq 2 \left( 1 + \frac{\sup_{x, x' \in \mathcal{X}} \|x - x'\|_F^2}{\sqrt{\lambda}} \right) \sqrt{\frac{2 \ln(\frac{1}{\delta})}{n}} \\ &\quad + \frac{16 \sup_{x, x' \in \mathcal{X}} \|x - x'\|_F^2}{\sqrt{n\lambda}} + \frac{12}{\sqrt{n}}. \end{aligned} \quad (21)$$

*Proof.* Note that the dual norm of the Frobenius norm is itself. The estimation of  $X_*$  is straightforward. The Rademacher complexity  $R_n$  is estimated as follows:

$$\begin{aligned} R_n &= \frac{1}{\lfloor \frac{n}{2} \rfloor} \mathbb{E} \left( \sum_{i, j=1}^{\lfloor \frac{n}{2} \rfloor} \sigma_i \sigma_j \langle x_i - x_{\lfloor \frac{n}{2} \rfloor + i}, x_j - x_{\lfloor \frac{n}{2} \rfloor + j} \rangle^2 \right)^{\frac{1}{2}} \\ &\leq \frac{1}{\lfloor \frac{n}{2} \rfloor} \mathbb{E}_{\sigma} \left( \sum_{i, j=1}^{\lfloor \frac{n}{2} \rfloor} \sigma_i \sigma_j \langle x_i - x_{\lfloor \frac{n}{2} \rfloor + i}, x_j - x_{\lfloor \frac{n}{2} \rfloor + j} \rangle^2 \right)^{\frac{1}{2}} \\ &= \frac{1}{\lfloor \frac{n}{2} \rfloor} \mathbb{E}_{\mathbf{z}} \left( \sum_{i=1}^{\lfloor \frac{n}{2} \rfloor} \|x_i - x_{\lfloor \frac{n}{2} \rfloor + i}\|_F^4 \right)^{\frac{1}{2}} \\ &\leq X_* / \sqrt{\lfloor \frac{n}{2} \rfloor} \leq \frac{2X_*}{\sqrt{n}}. \end{aligned}$$

Putting this estimation back into equation (13) completes the proof of Example 1.  $\square$

Other popular matrix norms for metric learning are the  $L^1$ -norm, trace-norm and mixed  $(2, 1)$ -norm. The dual norms are respectively  $L^\infty$ -norm, spectral norm (i.e. the maximum of singular values) and mixed  $(2, \infty)$ -norm. All these dual norms mentioned above are less than the Frobenius norm. Hence, the following estimation always holds true for all the norms mentioned above:

$$X_* \leq \sup_{x, x' \in \mathcal{X}} \|x - x'\|_F^2, \quad \text{and} \quad R_n \leq \frac{2 \sup_{x, x' \in \mathcal{X}} \|x - x'\|_F^2}{\sqrt{n}}.$$

Consequently, the generalisation bound (21) holds true for metric learning formulation (2) with  $L^1$ -norm, or trace-norm or mixed  $(2, 1)$ -norm regularisation. However, in some cases, the above upper-bounds are too conservative. For instance, in the following examples we can show that more refined estimation of  $R_n$  can be obtained by applying the Khinchin inequalities for Rademacher averages [20].

**Example 2** (Sparse  $L^1$ -norm). Let the matrix norm be the  $L^1$ -norm i.e.  $\|M\| = \sum_{\ell, k \in \mathbb{N}_d} |M_{\ell k}|$ . Then,  $X_* = \sup_{x, x' \in \mathcal{X}} \|x - x'\|_\infty^2$  and

$$R_n \leq 4 \sup_{x, x' \in \mathcal{X}} \|x - x'\|_\infty^2 \sqrt{\frac{e \log d}{n}}.$$

Let  $(M_{\mathbf{z}}, b_{\mathbf{z}})$  be a solution of formulation (2) with  $L^1$ -norm regularisation. For any  $0 < \delta < 1$ , with probability  $1 - \delta$  there holds

$$\begin{aligned} \mathcal{E}(M_{\mathbf{z}}, b_{\mathbf{z}}) - \mathcal{E}_{\mathbf{z}}(M_{\mathbf{z}}, b_{\mathbf{z}}) &\leq 2 \left( 1 + \frac{\sup_{x, x' \in \mathcal{X}} \|x - x'\|_\infty^2}{\sqrt{\lambda}} \right) \sqrt{\frac{2 \ln(\frac{1}{\delta})}{n}} \\ &\quad + \frac{8 \sup_{x, x' \in \mathcal{X}} \|x - x'\|_\infty^2 (1 + 2\sqrt{e \log d})}{\sqrt{n\lambda}} + \frac{12}{\sqrt{n}}. \end{aligned} \quad (22)$$

*Proof.* The dual norm of the  $L^1$ -norm is the  $L^\infty$ -norm. Hence,  $X_* = \sup_{x, x' \in \mathcal{X}} \|x - x'\|_\infty^2$ . To estimate  $R_n$ , we observe, for any  $1 < q < \infty$ , that

$$\begin{aligned} R_n &= \frac{1}{\lfloor \frac{n}{2} \rfloor} \mathbb{E}_{\mathbf{z}} \mathbb{E}_{\sigma} \left\| \sum_{i=1}^{\lfloor \frac{n}{2} \rfloor} \sigma_i X_{i(\lfloor \frac{n}{2} \rfloor + i)} \right\|_\infty \leq \frac{1}{\lfloor \frac{n}{2} \rfloor} \mathbb{E}_{\mathbf{z}} \mathbb{E}_{\sigma} \left\| \sum_{i=1}^{\lfloor \frac{n}{2} \rfloor} \sigma_i X_{i(\lfloor \frac{n}{2} \rfloor + i)} \right\|_q \\ &:= \frac{1}{\lfloor \frac{n}{2} \rfloor} \mathbb{E}_{\mathbf{z}} \mathbb{E}_{\sigma} \left( \sum_{\ell, k \in \mathbb{N}_d} \left| \sum_{i=1}^{\lfloor \frac{n}{2} \rfloor} \sigma_i (x_i^k - x_{\lfloor \frac{n}{2} \rfloor + i}^k) (x_i^\ell - x_{\lfloor \frac{n}{2} \rfloor + i}^\ell) \right|^q \right)^{\frac{1}{q}} \\ &\leq \frac{1}{\lfloor \frac{n}{2} \rfloor} \mathbb{E}_{\mathbf{z}} \left( \sum_{\ell, k \in \mathbb{N}_d} \mathbb{E}_{\sigma} \left| \sum_{i=1}^{\lfloor \frac{n}{2} \rfloor} \sigma_i (x_i^k - x_{\lfloor \frac{n}{2} \rfloor + i}^k) (x_i^\ell - x_{\lfloor \frac{n}{2} \rfloor + i}^\ell) \right|^q \right)^{\frac{1}{q}} \end{aligned} \quad (23)$$

where  $x_i^k$  represents the  $k$ -th coordinate element of vector  $x_i \in \mathbb{R}^d$ . To estimate the term on the right-hand side of inequality (23), we apply the Khinchin-Kahane inequality (See Lemma 9 in the Appendix) with  $p = 2 < q < \infty$  yields that

$$\begin{aligned} &\mathbb{E}_{\sigma} \left| \sum_{i=1}^{\lfloor \frac{n}{2} \rfloor} \sigma_i (x_i^k - x_{\lfloor \frac{n}{2} \rfloor + i}^k) (x_i^\ell - x_{\lfloor \frac{n}{2} \rfloor + i}^\ell) \right|^q \\ &\leq q^{\frac{q}{2}} \left( \mathbb{E}_{\sigma} \left| \sum_{i=1}^{\lfloor \frac{n}{2} \rfloor} \sigma_i (x_i^k - x_{\lfloor \frac{n}{2} \rfloor + i}^k) (x_i^\ell - x_{\lfloor \frac{n}{2} \rfloor + i}^\ell) \right|^2 \right)^{\frac{q}{2}} \\ &= q^{\frac{q}{2}} \left( \sum_{i=1}^{\lfloor \frac{n}{2} \rfloor} (x_i^k - x_{\lfloor \frac{n}{2} \rfloor + i}^k)^2 (x_i^\ell - x_{\lfloor \frac{n}{2} \rfloor + i}^\ell)^2 \right)^{\frac{q}{2}} \leq \sup_{x, x' \in \mathcal{X}} \|x - x'\|_\infty^{2q} \left( \lfloor \frac{n}{2} \rfloor \right)^{\frac{q}{2}} q^{\frac{q}{2}}. \end{aligned} \quad (24)$$

Putting the above estimation back into (23) and letting  $q = 4 \log d$  implies that

$$\begin{aligned} R_n &\leq \sup_{x, x' \in \mathcal{X}} \|x - x'\|_\infty^2 d^{\frac{2}{q}} \sqrt{q} / \sqrt{\lfloor \frac{n}{2} \rfloor} = 2 \sup_{x, x' \in \mathcal{X}} \|x - x'\|_\infty^2 \sqrt{e \log d / \lfloor \frac{n}{2} \rfloor} \\ &\leq 4 \sup_{x, x' \in \mathcal{X}} \|x - x'\|_\infty^2 \sqrt{e \log d / n}. \end{aligned}$$

Putting the estimation for  $X_*$  and  $R_n$  into Theorem 13 yields inequality (22). This completes the proof of Example 2.  $\square$

**Example 3** (Mixed  $(2, 1)$ -norm). Consider  $\|M\| = \sum_{\ell \in \mathbb{N}_d} \sqrt{\sum_{k \in \mathbb{N}_d} |M_{\ell k}|^2}$ . Then, we have  $X_* = [\sup_{x, x' \in \mathcal{X}} \|x - x'\|_F] [\sup_{x, x' \in \mathcal{X}} \|x - x'\|_\infty]$ , and

$$R_n \leq 4 \left[ \sup_{x, x' \in \mathcal{X}} \|x - x'\|_\infty \right] \left[ \sup_{x, x' \in \mathcal{X}} \|x - x'\|_F \right] \sqrt{\frac{e \log d}{n}}.$$

Let  $(M_{\mathbf{z}}, b_{\mathbf{z}})$  be a solution of formulation (2) with mixed  $(2, 1)$ -norm. For any  $0 < \delta < 1$ , with probability  $1 - \delta$  there holds

$$\begin{aligned} \mathcal{E}(M_{\mathbf{z}}, b_{\mathbf{z}}) - \mathcal{E}_{\mathbf{z}}(M_{\mathbf{z}}, b_{\mathbf{z}}) &\leq 2 \left( 1 + \frac{\left[ \sup_{x, x' \in \mathcal{X}} \|x - x'\|_{\infty} \right] \left[ \sup_{x, x' \in \mathcal{X}} \|x - x'\|_F \right]}{\sqrt{\lambda}} \right) \sqrt{\frac{2 \ln \left( \frac{1}{\delta} \right)}{n}} \\ &\quad + \frac{8 \left[ \sup_{x, x' \in \mathcal{X}} \|x - x'\|_{\infty} \right] \left[ \sup_{x, x' \in \mathcal{X}} \|x - x'\|_F \right] (1 + 2\sqrt{e \log d})}{\sqrt{n\lambda}} + \frac{12}{\sqrt{n}}. \end{aligned} \quad (25)$$

*Proof.* The estimation of  $X_*$  is straightforward and we estimate  $R_n$  as follows. For any  $q > 1$ , there holds

$$\begin{aligned} R_n &= \frac{1}{\lfloor \frac{n}{2} \rfloor} \mathbb{E}_{\mathbf{z}} \mathbb{E}_{\sigma} \left\| \sum_{i=1}^{\lfloor \frac{n}{2} \rfloor} \sigma_i X_{i(\lfloor \frac{n}{2} \rfloor + i)} \right\|_{(2, \infty)} \\ &= \frac{1}{\lfloor \frac{n}{2} \rfloor} \mathbb{E}_{\mathbf{z}} \mathbb{E}_{\sigma} \sup_{\ell \in \mathbb{N}_d} \left( \sum_{k \in \mathbb{N}_d} \left| \sum_{i=1}^{\lfloor \frac{n}{2} \rfloor} \sigma_i (x_i^k - x_{\lfloor \frac{n}{2} \rfloor + i}^k) (x_i^{\ell} - x_{\lfloor \frac{n}{2} \rfloor + i}^{\ell}) \right|^2 \right)^{\frac{1}{2}} \\ &\leq \frac{1}{\lfloor \frac{n}{2} \rfloor} \mathbb{E}_{\mathbf{z}} \left( \sum_{k \in \mathbb{N}_d} \mathbb{E}_{\sigma} \sup_{\ell \in \mathbb{N}_d} \left| \sum_{i=1}^{\lfloor \frac{n}{2} \rfloor} \sigma_i (x_i^k - x_{\lfloor \frac{n}{2} \rfloor + i}^k) (x_i^{\ell} - x_{\lfloor \frac{n}{2} \rfloor + i}^{\ell}) \right|^2 \right)^{\frac{1}{2}}. \end{aligned} \quad (26)$$

It remains to estimate the terms inside the parenthesis on the right-hand side of the above inequality. To this end, we observe, for any  $q' > 1$ , that

$$\begin{aligned} &\mathbb{E}_{\sigma} \sup_{\ell \in \mathbb{N}_d} \left| \sum_{i=1}^{\lfloor \frac{n}{2} \rfloor} \sigma_i (x_i^k - x_{\lfloor \frac{n}{2} \rfloor + i}^k) (x_i^{\ell} - x_{\lfloor \frac{n}{2} \rfloor + i}^{\ell}) \right|^2 \\ &\leq \mathbb{E}_{\sigma} \left( \sum_{\ell \in \mathbb{N}_d} \left| \sum_{i=1}^{\lfloor \frac{n}{2} \rfloor} \sigma_i (x_i^k - x_{\lfloor \frac{n}{2} \rfloor + i}^k) (x_i^{\ell} - x_{\lfloor \frac{n}{2} \rfloor + i}^{\ell}) \right|^{2q'} \right)^{\frac{1}{q'}} \\ &\leq \left( \sum_{\ell \in \mathbb{N}_d} \mathbb{E}_{\sigma} \left| \sum_{i=1}^{\lfloor \frac{n}{2} \rfloor} \sigma_i (x_i^k - x_{\lfloor \frac{n}{2} \rfloor + i}^k) (x_i^{\ell} - x_{\lfloor \frac{n}{2} \rfloor + i}^{\ell}) \right|^{2q'} \right)^{\frac{1}{q'}}. \end{aligned}$$

Applying the Khinchin-Kahane inequality (Lemma 9 in the Appendix) with  $q = 2q' = 4 \log d$  and  $p = 2$  to the above inequality yields that

$$\begin{aligned} &\mathbb{E}_{\sigma} \sup_{\ell \in \mathbb{N}_d} \left| \sum_{i=1}^{\lfloor \frac{n}{2} \rfloor} \sigma_i (x_i^k - x_{\lfloor \frac{n}{2} \rfloor + i}^k) (x_i^{\ell} - x_{\lfloor \frac{n}{2} \rfloor + i}^{\ell}) \right|^2 \\ &\leq \left( \sum_{\ell \in \mathbb{N}_d} (2q')^{q'} \left[ \mathbb{E}_{\sigma} \left| \sum_{i=1}^{\lfloor \frac{n}{2} \rfloor} \sigma_i (x_i^k - x_{\lfloor \frac{n}{2} \rfloor + i}^k) (x_i^{\ell} - x_{\lfloor \frac{n}{2} \rfloor + i}^{\ell}) \right|^2 \right]^{q'} \right)^{\frac{1}{q'}} \\ &= \left( \sum_{\ell \in \mathbb{N}_d} (2q')^{q'} \left[ \sum_{i=1}^{\lfloor \frac{n}{2} \rfloor} (x_i^k - x_{\lfloor \frac{n}{2} \rfloor + i}^k)^2 (x_i^{\ell} - x_{\lfloor \frac{n}{2} \rfloor + i}^{\ell})^2 \right]^{q'} \right)^{\frac{1}{q'}} \\ &\leq 2q' \sup_{x, x' \in \mathcal{X}} \|x - x'\|_{\infty}^2 d^{\frac{1}{q'}} \left[ \sum_{i=1}^{\lfloor \frac{n}{2} \rfloor} (x_i^k - x_{\lfloor \frac{n}{2} \rfloor + i}^k)^2 \right] \\ &\leq 4e(\log d) \sup_{x, x' \in \mathcal{X}} \|x - x'\|_{\infty}^2 \left[ \sum_{i=1}^{\lfloor \frac{n}{2} \rfloor} (x_i^k - x_{\lfloor \frac{n}{2} \rfloor + i}^k)^2 \right] \end{aligned}$$

Putting the above estimation back into (26) implies that

$$\begin{aligned} R_n &\leq \sqrt{4e \log d} \left[ \sup_{x, x' \in \mathcal{X}} \|x - x'\|_{\infty} \right] \mathbb{E}_{\mathbf{z}} \left( \sum_{i=1}^{\lfloor \frac{n}{2} \rfloor} \|x_i - x_{\lfloor \frac{n}{2} \rfloor + i}\|_F^2 \right)^{\frac{1}{2}} / \lfloor \frac{n}{2} \rfloor \\ &\leq \sqrt{4e \log d} \left[ \sup_{x, x' \in \mathcal{X}} \|x - x'\|_{\infty} \right] \left[ \sup_{x, x' \in \mathcal{X}} \|x - x'\|_F \right] / \sqrt{\lfloor \frac{n}{2} \rfloor} \\ &\leq 4\sqrt{e \log d} \left[ \sup_{x, x' \in \mathcal{X}} \|x - x'\|_{\infty} \right] \left[ \sup_{x, x' \in \mathcal{X}} \|x - x'\|_F \right] / \sqrt{n}. \end{aligned}$$

Combining this with Theorem 3 implies the inequality (25). This completes the proof of the example.  $\square$

In the Frobenius-norm case, the main term of the bound (21) is  $\mathcal{O}\left(\frac{\sup_{x,x' \in \mathcal{X}} \|x-x'\|_F^2}{\sqrt{n\lambda}}\right)$ . This bound is consistent with that given by [13] where  $\sup_{x \in \mathcal{X}} \|x\|_F$  is assumed to be bounded by some constant  $B$ . Comparing the generalisation bounds in the above examples. The key terms  $X_*$  and  $R_n$  mainly differ in two quantities, i.e.  $\sup_{x,x' \in \mathcal{X}} \|x-x'\|_F$  and  $\sup_{x,x' \in \mathcal{X}} \|x-x'\|_\infty$ . We argue that  $\sup_{x,x' \in \mathcal{X}} \|x-x'\|_\infty$  can be much less than  $\sup_{x,x' \in \mathcal{X}} \|x-x'\|_F$ . For instance, consider the input space  $\mathcal{X} = [0, 1]^d$ . It is easy to see that  $\sup_{x,x' \in \mathcal{X}} \|x-x'\|_F = \sqrt{d}$  while  $\sup_{x,x' \in \mathcal{X}} \|x-x'\|_\infty \equiv 1$ . Consequently, we can summarise the estimations as follows:

- **Frobenius-norm:**  $X_* = d$ , and  $R_n \leq \frac{2d}{\sqrt{n}}$ .
- **Sparse  $L^1$ -norm:**  $X_* = 1$ , and  $R_n \leq \frac{4\sqrt{e \log d}}{\sqrt{n}}$ .
- **Mixed (2,1)-norm:**  $X_* = \sqrt{d}$ , and  $R_n \leq \frac{4\sqrt{ed \log d}}{\sqrt{n}}$ .

Therefore, when  $d$  is large, the generalisation bound with sparse  $L^1$ -norm regularisation is much better than that with Frobenius-norm regularisation while the bound with mixed (2,1)-norm are between the above two. These theoretical results are nicely consistent with the rationale that sparse methods are more effective in dealing with high-dimensional data.

We end this section with two remarks. Firstly, in the setting of trace-norm regularisation, it remains a question to us on how to establish more accurate estimation of  $R_n$  by using the Khinchin-Kahane inequality. Secondly, the bounds in the above examples are true for similarity learning with different matrix-norm regularisation. Indeed, the generalisation bound for similarity learning in Theorem 4 tells us that it suffices to estimate  $\tilde{X}_*$  and  $\tilde{R}_n$ . In analogy to the arguments in the above examples, we can get the following results. For similarity learning formulation (4) with Frobenius-norm regularisation, there holds

$$\tilde{X}_* = \sup_{x \in \mathcal{X}} \|x\|_F^2, \quad \tilde{R}_n \leq \frac{2 \sup_x \|x\|_F^2}{\sqrt{n}}.$$

For  $L^1$ -norm regularisation, we have

$$\tilde{X}_* = \sup_{x \in \mathcal{X}} \|x\|_\infty^2, \quad \tilde{R}_n \leq 4 \sup_{x \in \mathcal{X}} \|x\|_\infty^2 \sqrt{e \log d} / \sqrt{n}.$$

In the setting of (2,1)-norm, we obtain

$$\tilde{X}_* = \sup_{x \in \mathcal{X}} \|x\|_\infty \sup_{x \in \mathcal{X}} \|x\|_F, \quad \tilde{R}_n \leq 4 \left[ \sup_{x \in \mathcal{X}} \|x\|_F \sup_{x \in \mathcal{X}} \|x\|_\infty \right] \sqrt{e \log d} / \sqrt{n}.$$

Putting these estimations back into Theorem 4 yields generalisation bounds for similarity learning with different matrix norms. For simplicity, we omit the details here.

## 5 Conclusion and Discussion

In this paper we are mainly concerned with theoretical generalisation analysis of the regularized metric and similarity learning. In particular, we first showed that the generalisation analysis for metric/similarity learning reduces to the estimation of the Rademacher average over “sums-of-i.i.d.” sample-blocks. Then, we derived their generalisation bounds with different matrix regularisation terms. Our analysis indicates that sparse metric/similarity learning with  $L^1$ -norm regularisation could lead significantly better bounds than that with the Frobenius norm regularisation, especially when the dimension of the input data is high. Our novel generalisation analysis develops the techniques of U-statistics [20, 7] and Rademacher complexity analysis [2, 15]. Below we mention several questions remaining to be further studied.

Firstly, in Section 3, the derived bounds for metric and similarity learning with trace-norm regularisation were the same as those with Frobenius-norm regularisation. It would be very interesting to derive the bounds similar to those with sparse  $\ell^1$ -norm regularisation. The key issue is to estimate the Rademacher complexity term (12) related to the spectral norm using the Khinchin-Kahne inequality. However, we are not aware of such Khinchin-Kahne inequalities for general matrix spectral norms. Another alternative is to apply the advanced oracle inequalities in [16].

Secondly, this study only investigated the generalisation bounds for metric and similarity learning. We can further get the consistency estimation under strong assumptions on the loss function and underlying distribution. Specifically, we assume that the loss function is the least square loss, the matrix norm is the Frobenius norm and the bias term  $b$  is fixed to be zero. In addition, assume that the true minimizer  $M_* = \arg \min_{M \in \mathbb{S}^d} \mathcal{E}(M, 0)$  exists and let  $M_{\mathbf{z}} = \arg \min_{M \in \mathbb{S}^d} [\mathcal{E}_{\mathbf{z}}(M, 0) + \lambda \|M\|_F^2]$ . Observe that

$$\begin{aligned} \mathcal{E}(M_{\mathbf{z}}, 0) - \mathcal{E}(M_*, 0) &= \iint \langle M_{\mathbf{z}} - M_*, x(x')^T \rangle^2 d\rho(x)\rho(x') \\ &= \langle \mathcal{C}(M_{\mathbf{z}} - M_*), M_{\mathbf{z}} - M_* \rangle, \end{aligned} \quad (27)$$

where  $\mathcal{C} = \iint (x(x')^T) \otimes (x(x')^T) d\rho(x)\rho(x')$  and  $\otimes$  represents the tensor product of matrices. Equation (27) implies that  $\mathcal{E}(M_{\mathbf{z}}, 0) - \mathcal{E}(M_*, 0) = \iint \langle M_{\mathbf{z}} - M_*, x(x')^T \rangle^2 d\rho(x)\rho(x') \geq \lambda_{\min}(\mathcal{C}) \|M_{\mathbf{z}} - M_*\|_F^2$ , where  $\lambda_{\min}(\mathcal{C})$  is the minimum eigenvalue of the  $d^2 \times d^2$  matrix  $\mathcal{C}$ . Furthermore, observe that  $\mathcal{E}(M_{\mathbf{z}}, 0) - \mathcal{E}(M_*, 0)$  is further bounded by

$$\begin{aligned} & [\mathcal{E}(M_{\mathbf{z}}, 0) - \mathcal{E}_{\mathbf{z}}(M_{\mathbf{z}}, 0)] + [\mathcal{E}_{\mathbf{z}}(M_{\mathbf{z}}, 0) + \lambda \|M_{\mathbf{z}}\|_F^2] - \mathcal{E}(M_*, 0) \\ & \leq [\mathcal{E}(M_{\mathbf{z}}, 0) - \mathcal{E}_{\mathbf{z}}(M_{\mathbf{z}}, 0)] + [\mathcal{E}_{\mathbf{z}}(M_*, 0) + \lambda \|M_*\|_F^2] - \mathcal{E}(M_*, 0) \\ & = [\mathcal{E}(M_{\mathbf{z}}, 0) - \mathcal{E}_{\mathbf{z}}(M_{\mathbf{z}}, 0)] + [\mathcal{E}_{\mathbf{z}}(M_*, 0) - \mathcal{E}(M_*, 0)] + \lambda \|M_*\|_F^2, \end{aligned} \quad (28)$$

where the inequality follows from the definition of the minimizer  $M_{\mathbf{z}}$ . Combining

equation (27) with the above estimation together implies that

$$\lambda_{\min}(\mathcal{C})\|M_{\mathbf{z}} - M_*\|_F^2 \leq [\mathcal{E}(M_{\mathbf{z}}, 0) - \mathcal{E}_{\mathbf{z}}(M_{\mathbf{z}}, 0)] + [\mathcal{E}_{\mathbf{z}}(M_*, 0) - \mathcal{E}(M_*, 0)] + \lambda\|M_*\|_F^2. \quad (29)$$

Using a similar argument as that for proving Theorem 3 and Example 1, we can get that  $[\mathcal{E}(M_{\mathbf{z}}, 0) - \mathcal{E}_{\mathbf{z}}(M_{\mathbf{z}}, 0)] + [\mathcal{E}_{\mathbf{z}}(M_*, 0) - \mathcal{E}(M_*, 0)] \leq \frac{C \ln(\frac{2}{\delta})}{\lambda\sqrt{n}}$  with a high confidence  $1 - \delta$ , where the constant  $C$  does not depend on  $\mathbf{z}$ . Consequently, putting this estimation with inequality (29) together implies that  $\|M_{\mathbf{z}} - M_*\|_F^2 \leq \frac{1}{\lambda_{\min}(\mathcal{C})} \left[ C \frac{\ln(\frac{2}{\delta})}{\lambda\sqrt{n}} + \lambda\|M_*\|_F^2 \right]$ . Choosing  $\lambda = n^{-\frac{1}{4}}$  yields the consistency estimation:

$$\|M_{\mathbf{z}} - M_*\|_F^2 \leq \frac{C \ln(\frac{2}{\delta}) + \|M_*\|_F^2}{\lambda_{\min}(\mathcal{C}) n^{\frac{1}{4}}}.$$

For the hinge loss, equality (27) does not hold true any more. Hence, it remains a question on how to get the consistency estimation for metric and similarity learning with general loss functions.

Thirdly, in many applications involving multi-media data, different aspects of the data may lead to several different, and apparently equally valid notions of similarity. This leads to a natural question to combine multiple similarities and metrics for a unified data representation. An extension of multiple kernel learning approach was proposed in [3] to address this issue. It would be very interesting to investigate the theoretical generalisation analysis for this multi-modal similarity learning framework. A possible starting point would be the techniques established for learning the kernel problem [30, 31].

Finally, the target of supervised metric learning is to improve the generalisation performance of kNN classifiers. It remains a challenging question to investigate how the generalisation performance of kNN classifiers relates to the generalisation bounds of metric learning given here.

## Acknowledgement:

We are grateful to the referees for their constructive comments and suggestions. This work is supported by the EPSRC under grant EP/J001384/1. The corresponding author is Yiming Ying.

## References

- [1] A. Bar-Hillel, T. Hertz, N. Shental, and D. Weinshall. Learning a mahalanobis metric from equivalence constraints. *J. of Machine Learning Research*, **6**: 937–965, 2005.

- [2] P.L. Bartlett and S. Mendelson. Rademacher and Gaussian complexities: risk bounds and structural results. *J. of Machine Learning Research*, **3**: 463–482, 2002.
- [3] B. McFee and G. Lanckriet. Learning multi-modal similarity. *J. of Machine Learning Research*, **12**: 491–523, 2011.
- [4] O. Bousquet and A. Elisseeff. Stability and generalization. *J. of Machine Learning Research*, **2**: 499–526, 2002.
- [5] D.R. Chen, Q. Wu, Y. Ying and D.X. Zhou. Support vector machine soft margin classifiers: error analysis, *J. of Machine Learning Research*, **5**: 1143–1175, 2004.
- [6] G. Chechik, V. Sharma, U. Shalit, and S. Bengio. Large scale online learning of image similarity through ranking. *J. of Machine Learning Research*, **11**: 1109–1135, 2010.
- [7] S. Cléménçon, G. Lugosi, and N. Vayatis. Ranking and empirical minimization of U-statistics. *The Annals of Statistics*, **36**: 844–874, 2008.
- [8] J. Davis, B. Kulis, P. Jain, S. Sra, and I. Dhillon. Information-theoretic metric learning. *ICML*, 2007.
- [9] A. Globerson and S. Roweis. Metric learning by collapsing classes. *NIPS*, 2005.
- [10] J. Goldberger, S. Roweis, G. Hinton, and R. Salakhutdinov. Neighbourhood component analysis. *NIPS*, 2004.
- [11] M. Guillaumin, J. Verbeek and C. Schmid. Is that you? Metric learning approaches for face identification. *ICCV*, 2009.
- [12] S. C. H. Hoi, W. Liu, M. R. Lyu, and W.-Y. Ma. Learning distance metrics with contextual constraints for image retrieval. *CVPR*, 2006.
- [13] R. Jin, S. Wang and Y. Zhou. Regularized distance metric learning: theory and algorithm. *NIPS*, 2009.
- [14] P. Kar and P. Jain. Similarity-based learning via data-driven embeddings. *NIPS*, 2011.
- [15] V. Koltchinskii and V. Panchenko. Empirical margin distributions and bounding the generalization error of combined classifiers. *The Annals of Statistics*, **30**, 1–5, 2002.
- [16] V. Koltchinskii. Oracle Inequalities in Empirical Risk Minimization and Sparse Recovery Problems. Springer, 2011.

- [17] M. Ledoux and M. Talagrand. *Probability in Banach Spaces: Isoperimetry and Processes*. Springer Press, New York, 1991.
- [18] A. Maurer. Learning similarity with operator-valued large-margin classifiers, *J. of Machine Learning Research*. **9**: 1049-1082, 2008.
- [19] C. McDiarmid. Surveys in Combinatorics, Chapter On the methods of bounded differences, 148-188, 1989. Cambridge University Press, Cambridge (UK).
- [20] V.H. De La Peña and E. Giné. *Decoupling: from Dependence to Independence*. Springer, New York, 1999.
- [21] R. Rosales and G. Fung. Learning sparse metrics via linear programming, *KDD*, 2006.
- [22] O. Shalit, D. Weinshall and G. Chechik. Online learning in the manifold of low-rank matrices. *NIPS*, 2010.
- [23] C. Shen, J. Kim, L. Wang and A. Hengel. Positive semidefinite metric learning with boosting. *NIPS*, 2009.
- [24] L. Torresani and K. Lee. Large margin component analysis. *NIPS*, 2007.
- [25] K. Q. Weinberger and L. K. Saul. Fast solvers and efficient implementations for distance metric learning. *ICML*, 2008.
- [26] E. Xing, A. Ng, M. Jordan, and S. Russell. Distance metric learning with application to clustering with side information. *NIPS*, 2002.
- [27] L. Yang and R. Jin. Distance metric learning: A comprehensive survey. In *Technical report, Department of Computer Science and Engineering, Michigan State University*, 2007.
- [28] Y. Ying, K. Huang and C. Campbell. Sparse metric learning via smooth optimisation. *NIPS*, 2009.
- [29] Y. Ying and P. Li. Distance metric learning with eigenvalue optimisation. *J. of Machine Learning Research*, **13**: 1–26, 2012.
- [30] Y. Ying and Campbell. Generalization bounds for learning the kernel. *COLT*, 2009.
- [31] Y. Ying and C. Campbell. Rademacher chaos complexity for learning the kernel problem. *Neural Computation*, **22**: 2858–86, 2010.

## Appendix

In this appendix we assemble some facts, which were used to establish generalisation bounds for metric/similarity learning.

**Definition 5.** We say the function  $f : \prod_{k=1}^n \Omega_k \rightarrow \mathbb{R}$  with bounded differences  $\{c_k\}_{k=1}^n$  if, for all  $1 \leq k \leq n$ ,

$$\max_{z_1, \dots, z_k, z'_k, \dots, z_n} |f(z_1, \dots, z_{k-1}, z_k, z_{k+1}, \dots, z_n) - f(z_1, \dots, z_{k-1}, z'_k, z_{k+1}, \dots, z_n)| \leq c_k$$

**Lemma 6.** (McDiarmid's inequality [19]) Suppose  $f : \prod_{k=1}^n \Omega_k \rightarrow \mathbb{R}$  with bounded differences  $\{c_k\}_{k=1}^n$  then, for all  $\epsilon > 0$ , there holds

$$\Pr_{\mathbf{z}} \left\{ f(\mathbf{z}) - \mathbb{E}_{\mathbf{z}} f(\mathbf{z}) \geq \epsilon \right\} \leq e^{-\frac{2\epsilon^2}{\sum_{k=1}^n c_k^2}}.$$

Finally we list a useful property for U-statistics. Given the i.i.d. random variables  $z_1, z_2, \dots, z_n \in \mathcal{Z}$ , let  $q : \mathcal{Z} \times \mathcal{Z} \rightarrow \mathbb{R}$  be a symmetric real-valued function. Denote a U-statistic of order two by  $U_n = \frac{1}{n(n-1)} \sum_{i \neq j} q(x_i, x_j)$ . Then, the U-statistic  $U_n$  can be expressed as

$$U_n = \frac{1}{n!} \sum_{\pi} \frac{1}{\lfloor \frac{n}{2} \rfloor} \sum_{i=1}^{\lfloor \frac{n}{2} \rfloor} q(z_{\pi(i)}, z_{\pi(\lfloor \frac{n}{2} \rfloor + i)}) \quad (30)$$

where the sum is taken over all permutations  $\pi$  of  $\{1, 2, \dots, n\}$ . The main idea underlying this representation is to reduce the analysis to the ordinary case of i.i.d. random variable blocks. Based on the above representation, we can prove the following lemma which plays a critical role in deriving generalisation bounds for metric learning. For completeness, we include a proof here. For more details on U-statistics, one is referred to [7, 20].

**Lemma 7.** Let  $q_{\tau} : \mathcal{Z} \times \mathcal{Z} \rightarrow \mathbb{R}$  be real-valued functions indexed by  $\tau \in \mathcal{T}$  where  $\mathcal{T}$  is some index set. If  $z_1, \dots, z_n$  are i.i.d. then we have that

$$\mathbb{E} \left[ \sup_{\tau \in \mathcal{T}} \frac{1}{n(n-1)} \sum_{i \neq j} q_{\tau}(z_i, z_j) \right] \leq \mathbb{E} \left[ \sup_{\tau \in \mathcal{T}} \frac{1}{\lfloor \frac{n}{2} \rfloor} \sum_{i=1}^{\lfloor \frac{n}{2} \rfloor} q_{\tau}(z_i, z_{\lfloor \frac{n}{2} \rfloor + i}) \right].$$

*Proof.* From the representation of U-statistics (30), we observe that

$$\begin{aligned}
\mathbb{E} \left[ \sup_{\tau \in \mathcal{T}} \frac{1}{n(n-1)} \sum_{i \neq j} q_{\tau}(z_i, z_j) \right] &= \mathbb{E} \sup_{\tau} \frac{1}{n!} \sum_{\pi} \frac{1}{\lfloor \frac{n}{2} \rfloor} \sum_{i=1}^{\lfloor \frac{n}{2} \rfloor} q_{\tau}(z_{\pi(i)}, z_{\pi(\lfloor \frac{n}{2} \rfloor + i)}) \\
&\leq \frac{1}{n!} \mathbb{E} \sum_{\pi} \sup_{\tau} \frac{1}{\lfloor \frac{n}{2} \rfloor} \sum_{i=1}^{\lfloor \frac{n}{2} \rfloor} q_{\tau}(z_{\pi(i)}, z_{\pi(\lfloor \frac{n}{2} \rfloor + i)}) \\
&= \frac{1}{n!} \sum_{\pi} \mathbb{E} \sup_{\tau} \frac{1}{\lfloor \frac{n}{2} \rfloor} \sum_{i=1}^{\lfloor \frac{n}{2} \rfloor} q_{\tau}(z_{\pi(i)}, z_{\pi(\lfloor \frac{n}{2} \rfloor + i)}) \\
&= \mathbb{E} \left[ \sup_{\tau \in \mathcal{T}} \frac{1}{\lfloor \frac{n}{2} \rfloor} \sum_{i=1}^{\lfloor \frac{n}{2} \rfloor} q_{\tau}(z_i, z_{\lfloor \frac{n}{2} \rfloor + i}) \right].
\end{aligned}$$

This completes the proof of the lemma.  $\square$

We need the following contraction property of the Rademacher averages which is essentially implied by Theorem 4.12 in Ledoux and Talagrand [17], see also [2, 15].

**Lemma 8.** *Let  $F$  be a class of uniformly bounded real-valued functions on  $(\Omega, \mu)$  and  $m \in \mathbb{N}$ . If for each  $i \in \{1, \dots, m\}$ ,  $\Psi_i : \mathbb{R} \rightarrow \mathbb{R}$  is a function with  $\Psi_i(0) = 0$  having a Lipschitz constant  $c_i$ , then for any  $\{x_i\}_{i=1}^m$ ,*

$$\mathbb{E}_{\epsilon} \left( \sup_{f \in F} \left| \sum_{i=1}^m \epsilon_i \Psi_i(f(x_i)) \right| \right) \leq 2 \mathbb{E}_{\epsilon} \left( \sup_{f \in F} \left| \sum_{i=1}^m c_i \epsilon_i f(x_i) \right| \right). \quad (31)$$

The last property of Rademacher averages is the Khinchin-Kahne inequality (see e.g. [20, Theorem 1.3.1]).

**Lemma 9.** *For  $n \in \mathbb{N}$ , let  $\{f_i \in \mathbb{R} : i \in \mathbb{N}_n\}$ , and  $\{\sigma_i : i \in \mathbb{N}_n\}$  be a family of i.i.d. Rademacher variables. Then, for any  $1 < p < q < \infty$  we have*

$$\left( \mathbb{E}_{\sigma} \left| \sum_{i \in \mathbb{N}_n} \sigma_i f_i \right|^q \right)^{\frac{1}{q}} \leq \left( \frac{q-1}{p-1} \right)^{\frac{1}{2}} \left( \mathbb{E}_{\sigma} \left| \sum_{i \in \mathbb{N}_n} \sigma_i f_i \right|^p \right)^{\frac{1}{p}}$$