

About the Linear Complexity of Ding-Hellestth Generalized Cyclotomic Binary Sequences of Any Period

Vladimir Edemskiy

Novgorod State University,
B. St. Petersburgskaya, 41, 173003 Velikiy Novgorod, Russia
Vladimir.Edemskiy@novsu.ru

Abstract. We defined sufficient conditions for designing Ding-Hellestth sequences with arbitrary period and high linear complexity for generalized cyclotomies. Also we discuss the method of computing the linear complexity of Ding-Hellestth sequences in the general case.

Keywords: Generalized cyclotomic sequences, Linear complexity

1 Introduction

For cryptographic applications, the linear complexity (L) of a sequence is an important merit factor. It may be defined as the length of the shortest linear feedback shift register that is capable of generating the sequence. The feedback function of this shift register can be deduced from knowledge of just $2L$ consecutive digits of the sequence. Thus, it is reasonable to suggest that "good" sequences have $L > N/2$ (where N denotes the period of the sequence) [1].

Using classical cyclotomic classes and generalized cyclotomic classes to construct binary sequences, which are called classical cyclotomic sequences and generalized cyclotomic sequences respectively, is an important method for sequence design [1]. In their paper [2] C. Ding and T. Hellestth first introduced a new generalized cyclotomy of order 2 with respect to $p_1^{e_1} \cdots p_t^{e_t}$, which includes classical cyclotomy as a special case and they show how to construct binary sequences based on this new generalized cyclotomy. There are many works devoted to the investigation of the properties of the Ding-Hellestth sequences. In particular, the linear complexity of these sequences is studied in [3-11]. Most of the papers study a special case described in sections 1-5 of [2]. In addition, the linear complexity of the sequences based on cyclotomic classes of higher orders was considered for specific modules (p^n, pq) and for a special case [9, 12-14].

The purpose of this paper is to find Ding-Hellestth generalized cyclotomic sequences with arbitrary periods and high linear complexity. In particular, we generalize the result of Tongjiang Yan [15]. Also we discuss the computation of the linear complexity of these sequences in the general case.

2 Basic Definitions and Notations

In this section let us briefly recall the definition of Ding-Helleseth generalized cyclotomic sequences [2].

Let $n = p_1^{e_1} \cdots p_t^{e_t}$, when p_1, \dots, p_t be pairwise distinct odd primes satisfying

$$\gcd\left(p_i^{e_i-1}(p_i-1), p_j^{e_j-1}(p_j-1)\right) = 2 \text{ for all } i \neq j$$

and $e_1 \geq 1, \dots, e_t \geq 1$ be integers.

Let Z_n be the ring of residue classes modulo n . According to the Chinese Remainder Theorem

$$Z_n \cong Z_{p_1^{e_1}} \times \dots \times Z_{p_t^{e_t}} \quad (1)$$

relatively to isomorphism $\varphi(x) = (x \pmod{p_1^{e_1}}, \dots, x \pmod{p_t^{e_t}})$. Here and hereafter $x \pmod{n}$ denotes the least nonnegative integer that is congruent to x modulo n .

It is well known that exists a primitive root g_i modulo $p_i^{e_i}$. Let $D_0^{(p_i^{e_i})} = \{g_i^{2j} | j \in Z\}$ be the subgroup of $Z_{p_i^{e_i}}^*$, generated by g_i^2 , and $D_1^{(p_i^{e_i})} = g_i D_0^{(p_i^{e_i})}$, where the arithmetic is that of $Z_{p_i^{e_i}}$, $i = 1, 2, \dots, t$.

Let $\mathbf{a} = (a_1, \dots, a_t)$ be a nonzero vector from $(Z_2)^t$ and

$$I_0^{(\mathbf{a}, n)} = \left\{ (i_1, \dots, i_t) \in (Z_2)^t \mid \sum_{k=1}^t i_k a_k = 0 \right\}, \quad I_1^{(\mathbf{a}, n)} = (Z_2)^t \setminus I_0^{(\mathbf{a}, n)}.$$

By definition, put [2]

$$E_j^{(\mathbf{a}, n)} = \prod_{(i_1, \dots, i_t) \in I_j^{(\mathbf{a}, n)}} D_{i_1}^{(p_1^{e_1})} \times \dots \times D_{i_t}^{(p_t^{e_t})} \text{ and } D_j^{(\mathbf{a}, n)} = \varphi^{-1}\left(E_j^{(\mathbf{a}, n)}\right), j = 0, 1.$$

From our definition it follows that [2]

$$Z_n^* = D_0^{(\mathbf{a}, n)} \cup D_1^{(\mathbf{a}, n)}, \quad D_0^{(\mathbf{a}, n)} \cap D_1^{(\mathbf{a}, n)} = \emptyset. \quad (2)$$

Clearly there is an element $b \in Z_n^*$ such that $D_1^{(\mathbf{a}, n)} = bD_0^{(\mathbf{a}, n)}$. The $D_0^{(\mathbf{a}, n)}$ and $D_1^{(\mathbf{a}, n)}$ are called generalized cyclotomic classes of order 2 with respect to \mathbf{a} and n . In the following $D_j^{(\mathbf{a}, n)}$ will denote $D_j^{(\mathbf{a}, n)} \pmod{2}$.

Further, by [2] we have a partition

$$Z_n \setminus \{0\} = \bigcup_{d|n, d>1} \frac{n}{d} Z_d^* \quad (3)$$

Let $d > 1$ be a positive integer and $d|n$, and the nonzero vector $\mathbf{a}_d = (a_1^{(d)}, \dots, a_m^{(d)}) \in (Z_2)^m$, where m is a number of different prime numbers participating in the factorization d . By (2) and (3) we obtain

$$Z_n \setminus \{0\} = \bigcup_{d|n, d>1} \frac{n}{d} \left(D_0^{(\mathbf{a}_d, d)} \cup D_1^{(\mathbf{a}_d, d)} \right).$$

Let

$$C_0 = \bigcup_{d|n, d>1} \frac{n}{d} D_0^{(\mathbf{a}_d, d)} \quad \text{and} \quad C_1 = \bigcup_{d|n, d>1} \frac{n}{d} D_1^{(\mathbf{a}_d, d)} \cup \{0\}.$$

Then $\{C_0, C_1\}$ is a partition of Z_n , i.e. $Z_n = C_0 \cup C_1$ and $C_0 \cap C_1 = \emptyset$.

In accordance with [2], the binary sequence s^∞ is then defined by

$$s_i = j \text{ if and only if } j \pmod{n} \in C_j.$$

The s^∞ is called Ding-Helleseth sequence, with the most frequently discussed options when $\mathbf{a}_d = (0, \dots, 0, 1)$.

3 Evaluation of the Linear Complexity of Ding-Helleseth Sequences

In this section we find sufficient conditions for Ding-Helleseth sequences to have high linear complexity. Define g to be the unique solution of the following set of congruences

$$g \equiv g_i \pmod{p_i^{e_i}}, \quad i = 1, 2, \dots, t.$$

LEMMA 1. If $\sum_{k=1}^m a_k^{(d)}$ is an odd number, then

$$gD_j^{(\mathbf{a}_d, d)} = D_{j+1}^{(\mathbf{a}_d, d)}$$

for $j = 0, 1$, where the arithmetic is that of Z_d .

PROOF. Let $d = p_{j_1}^{l_1} \dots p_{j_m}^{l_m}$, where $j_k \in \{1, 2, \dots, t\}$, $k = 1, 2, \dots, m$ and integers l_k satisfy the set of inequalities $1 \leq l_k \leq e_{j_k}$, $k = 1, 2, \dots, m$. By the definitions of $D_j^{(\mathbf{a}_d, d)}$ and g we obtain

$$\phi\left(gD_0^{(\mathbf{a}_d, d)}\right) = \prod_{(j_1, \dots, j_m) \in I_0^{(\mathbf{a}_d, d)}} \left(g_{j_1} D_{j_1}^{(p_{j_1}^{l_1})}\right) \times \dots \times \left(g_{j_m} D_{j_m}^{(p_{j_m}^{l_m})}\right),$$

where $\phi(x) = (x \pmod{p_{j_1}^{l_1}}, \dots, x \pmod{p_{j_m}^{l_m}})$ or

$$\phi\left(gD_0^{(\mathbf{a}_d, d)}\right) = \prod_{(j_1, \dots, j_m) \in I_0^{(\mathbf{a}_d, d)}} D_{j_1+1}^{(p_{j_1}^{l_1})} \times \dots \times D_{j_m+1}^{(p_{j_m}^{l_m})}.$$

The sum $\sum_{k=1}^m (j_k + 1) a_k^{(d)} = \sum_{k=1}^m a_k^{(d)}$ for $(j_1, \dots, j_m) \in I_0^{(\mathbf{a}_d, d)}$, therefore, by the condition of Lemma $(j_1 + 1, \dots, j_m + 1) \in I_1^{(\mathbf{a}_d, d)}$ and by (2) we have

$$D_{j_1+1}^{(p_{j_1}^{l_1})} \times \dots \times D_{j_m+1}^{(p_{j_m}^{l_m})} \in E_1^{(\mathbf{a}_d, d)}.$$

Then $gD_0^{(a_d, d)} \subset D_1^{(a_d, d)}$, but as their orders are equal we obtain $gD_0^{(a_d, d)} = D_1^{(a_d, d)}$.

The assertion $gD_1^{(a_d, d)} = D_0^{(a_d, d)}$ can be proven similarly.

Let α be a primitive n -th root of unity in the extension of field $GF(2)$. Then by Blahut's theorem for the linear complexity L of the sequence s^∞ we have

$$L = n - |\{v \mid S(\alpha^v) = 0, v = 0, 1, \dots, n-1\}|, \quad (4)$$

where $S(x)$ is defined by $S(x) = \sum_{i \in C_1} x^i$.

In order to investigate the values of $S(\alpha^v)$, let us introduce subsidiary polynomials. Let $S_A(x) = \sum_{i \in A} x^i$, where A is a subset of Z_n . Then for any $v = 1, \dots, n-1$ we obtain

$$S_{C_0}(\alpha^v) + S_{C_1}(\alpha^v) = 0. \quad (5)$$

LEMMA 2. If $\sum_{k=1}^m a_k^{(d)}$ is an odd number, then for any $v = 1, 2, \dots, n-1$ we have

$$S_{\frac{n}{d}D_1^{(a_d, d)}}(\alpha^{vg}) = S_{\frac{n}{d}D_0^{(a_d, d)}}(\alpha^v).$$

PROOF. By Lemma 1 $gD_1^{(a_d, d)} = D_0^{(a_d, d)}$ in the ring Z_d , then $\frac{n}{d}D_1^{(a_d, d)} = \frac{n}{d}gD_0^{(a_d, d)}$ in the ring Z_n and the statement of Lemma follows from the definition of auxiliary polynomial.

Let $\delta = \begin{cases} 1, & \text{if } S(1) = 0, \\ 0, & \text{otherwise.} \end{cases}$, that is $\delta = \begin{cases} 1, & \text{if } n \equiv 3 \pmod{4}, \\ 0, & \text{otherwise.} \end{cases}$, because by definition of sequence $S(1) = (n+1)/2$.

THEOREM 1. Suppose that for any integer $d > 1$ with $d|n$ sum $\sum_{k=1}^m a_k^{(d)}$ is an odd number; then for the linear complexity L of the sequence s^∞ we have

$$L \geq (n+1)/2 - \delta.$$

PROOF. By definition of the sequence s^∞ for all $v = 1, \dots, n-1$ we obtain

$$S(\alpha^v) = \sum_{d|n, d>1} S_{\frac{n}{d}D_1^{(a_d, d)}}(\alpha^v) + 1.$$

Then, by Lemma 2

$$S(\alpha^{gv}) = \sum_{d|n, d>1} S_{\frac{n}{d}D_0^{(a_d, d)}}(\alpha^v) + 1,$$

therefore $S(\alpha^{gv}) = S_{C_0}(\alpha^v) + 1$. Hence, by (5) we get $S(\alpha^v) + S(\alpha^{gv}) = 1$ for all $v = 1, \dots, n-1$. So, the order of the set

$$|\{v \mid S(\alpha^v) = 0, v = 1, 2, \dots, n-1\}| \leq (n-1)/2,$$

then by (4) we have, $L \geq (n+1)/2 - \delta$, which was to be demonstrated.

COROLLARY. If $2 \equiv g \pmod{n}$ under conditions of Theorem 1, then $L = n - \delta$. Indeed, in this case $S(\alpha^v) + S^2(\alpha^v) = 1$ and $S(\alpha^v) \neq 0$ for all $v = 1, 2, \dots, n-1$.

Let us make some more remarks on Theorem 1. When $\mathbf{a}_d = (0, \dots, 0, 1)$, i.e. in the special case of Ding-Helleseth generalized cyclotomic sequence, the condition of the Theorem 1 is automatically satisfied, so for this kind of sequences the evaluation of the linear complexity $L \geq (n+1)/2 - \delta$ is always valid. It is easy to see that this is in accord with already known results about the linear complexity of the sequences of periods p^n, pq [3-12]. It should be noted that not all the sequences examined in [3-12] can be defined as s^∞ .

Further, if

$$\gcd(p_i^{e_i-1}(p_i-1), p_j^{e_j-1}(p_j-1)) = r \text{ for all } i \neq j,$$

then like in [9, 12-14] for $p_i^{e_i}$ we can define the generalized cyclotomic classes $H_k^{(p_i^{e_i})}$, $k = 0, 1, \dots, r-1$ of order r . Now, we suppose $D_0^{(p_i^{e_i})} = \bigcup_{k=0}^{r/2-1} H_k^{(p_i^{e_i})}$ and $D_1^{(p_i^{e_i})} = \bigcup_{k=r/2}^{d-1} H_k^{(p_i^{e_i})}$ [13]. Then, for $\mathbf{a}_d = (0, \dots, 0, 1)$ the evaluation of the linear complexity given in Theorem 1 is valid also for generalized cyclotomic sequences built on new classes. We can prove it by just replacing the element g with $g^{r/2}$ in Lemmas 1 and 2. This is consistent with the results from [9, 12-14].

So, Theorem 1 establishes sufficient conditions for existence of Ding-Helleseth sequences with high linear complexity.

In the next section we discuss how the linear complexity of the generalized cyclotomic sequences could be computed in the general case. In particular, we show that if amongst vectors \mathbf{a}_d there exists one with the even sum of coordinates, then there exists some n for which the statement of Theorem 1 is not true.

4 About Computing of the Linear Complexity of Ding-Helleseth Sequences

Now let us generalize the method of computing the linear complexity of the generalized cyclotomic sequences with period pq proposed in [16]. By construction we see that

$$S(\alpha^v) = \sum_{d|n, d>1} S_{\frac{n}{d}D_1^{(a_d, d)}}(\alpha^v) + 1, \quad (6)$$

First, we examine the items of this sum.

If $d = p_{j_1}^{l_1} \cdots p_{j_m}^{l_m}$, then, by (1) there exist integers $b_i, i = 1, \dots, m$ such that

$$b_1 \frac{n}{p_{j_1}^{l_1}} + \dots + b_m \frac{n}{p_{j_m}^{l_m}} = \frac{n}{d},$$

where each $b_i, i = 1, \dots, m$ is uniquely determined modulo $p_{j_i}^{l_i}$ and $b_i \not\equiv 0 \pmod{p_{j_i}}$.

Let $\beta_k = \alpha^{b_k n / p_{j_k}^{l_k}}$, $k = 1, \dots, m$, then β_k is a primitive $p_{j_k}^{l_k}$ -th root of unity in the extension of the field $GF(2)$ and

$$\alpha = \beta_1 \dots \beta_m.$$

Generalizing Theorem 1 from [16] we obtain the following.

LEMMA 3. If $d = p_{j_1}^{l_1} \dots p_{j_m}^{l_m}$, then for $v = 1, \dots, n - 1$ we have

$$S_{\frac{n}{d} D_1^{(\mathbf{a}_d, d)}}(\alpha^v) = \sum_{(i_1, \dots, i_m) \in I_1^{(\mathbf{a}_d, d)}} S_{i_1}^{(p_{j_1}^{l_1})}(\beta_{j_1}^{\frac{n}{d} v}) \dots S_{i_m}^{(p_{j_m}^{l_m})}(\beta_{j_m}^{\frac{n}{d} v}),$$

where $S_{i_k}^{(p_{j_k}^{l_k})}(x) = \sum_{i \in D_{i_k}^{(p_{j_k}^{l_k})}} x^i$.

The method of computation of $S_{i_k}^{(p_{j_k}^{l_k})}(\beta_{j_k}^v)$ over the values of the classical cyclotomic sequences polynomial was proposed in [7-9]. Thus, formula (6), Lemma 3 and the results from [10] allow us to compute the values of the polynomial $S(\alpha^v)$, and consequently the linear complexity of Ding-Helleseth sequence if we know the kind of factorization of n . By example let us look on the case when the conditions of Theorem 1 does not hold.

Let $n = p_1 p_2$ and $\mathbf{a}_{p_1 p_2} = (1, 1)$ then

$$I_1^{(\mathbf{a}_{p_1 p_2}, p_1 p_2)} = \{(0, 1), (1, 0)\} \quad \text{and} \quad I_1^{(\mathbf{a}_{p_1}, p_1)} = I_1^{(\mathbf{a}_{p_2}, p_2)} = \{(1)\},$$

because by definition $\mathbf{a}_{p_1} = \mathbf{a}_{p_2} = (1)$. If $d = p_1 p_2$, then $\beta_1 = \alpha^{b_1 p_2}$ and $\beta_2 = \alpha^{b_2 p_1}$, where $b_1 p_1 + b_2 p_2 = 1$. Therefore $\alpha^{p_1} = \beta_1^{p_2}$, $\alpha^{p_2} = \beta_2^{p_1}$.

Hence by Lemma 3 and (6) for $n = p_1 p_2$ and $\mathbf{a}_{p_1 p_2} = (1, 1)$ we obtain

$$S(\alpha^v) = S_0^{(p_1)}(\beta_1^v) S_1^{(p_2)}(\beta_2^v) + S_1^{(p_1)}(\beta_1^v) S_0^{(p_2)}(\beta_2^v) + S_1^{(p_1)}(\beta_1^{p_2 v}) + S_1^{(p_2)}(\beta_2^{p_1 v}) + 1. \quad (7)$$

The properties of the polynomials $S_j^{(p_i)}(x)$, $j, i = 0, 1$ were examined in [17].

LEMMA 4. If $v \in Z_n^*$ then

$$S(\alpha^v) = \begin{cases} 0, & \text{if } p_1 \equiv 3 \pmod{4} \text{ and } p_2 \equiv 3 \pmod{4}, \\ 1, & \text{otherwise.} \end{cases}$$

PROOF. If $v \in Z_n^*$, then from (7), we obtain

$$S(\alpha^v) = S_1^{(p_1)}(\beta_1^v) + S_0^{(p_2)}(\beta_2^v) + S_1^{(p_1)}(\beta_1^{p_2 v}) + S_1^{(p_2)}(\beta_2^{p_1 v}). \quad (8)$$

Let us consider two cases

1) If $p_1 \equiv p_2 \equiv 3 \pmod{4}$, then in accordance with the law of quadratic reciprocity the Legendre symbols $\left(\frac{p_1}{p_2}\right)$ and $\left(\frac{p_2}{p_1}\right)$ are different. Without loss of generality, we can assume that $\left(\frac{p_1}{p_2}\right) = 1$. Then

$$S_1^{(p_1)}(\beta_1^{p_2 v}) = S_0^{(p_1)}(\beta_1^v) \quad \text{and} \quad S_1^{(p_2)}(\beta_2^{p_1 v}) = S_1^{(p_2)}(\beta_2^v).$$

By means of these relations we replace the last two terms in (8) and now obtain $S(\alpha^v) = 0$ for all $v \in Z_n^*$.

2) If $p_1 \equiv 1 \pmod{4}$ or $p_2 \equiv 1 \pmod{4}$ then $\left(\frac{p_1}{p_2}\right) = \left(\frac{p_2}{p_1}\right)$. In this case without loss of generality, we can assume that

$$S_1^{(p_1)}(\beta_1^{p_2 v}) = S_1^{(p_1)}(\beta_1^v) \quad \text{and} \quad S_1^{(p_2)}(\beta_2^{p_1 v}) = S_1^{(p_2)}(\beta_2^v)$$

Then by [17] and (8) we get $S(\alpha^v) = 1$ for all $v \in Z_n^*$.

Lemma 4 makes it clear that the statement of Theorem 1 is not true for $n = p_1 p_2$ and $\mathbf{a}_{p_1 p_2} = (1, 1)$.

With Lemma 4 we can conclude the computation of the linear complexity. If $n = p_1 p_2$ and $\mathbf{a}_{p_1 p_2} = (1, 1)$, then for the sequence s^∞ we have

1. $L = p_1 + p_2 - 1$, if $p_1 \equiv 3 \pmod{8}$ and $p_2 \equiv 3 \pmod{8}$;
2. $L = p_1 + (p_2 - 1)/2$, if $p_1 \equiv 3 \pmod{8}$ and $p_2 \equiv 7 \pmod{8}$;
3. $L = p_2 + (p_1 - 1)/2$, if $p_1 \equiv 7 \pmod{8}$ and $p_2 \equiv 3 \pmod{8}$;
4. $L = (p_1 + p_2)/2$, if $p_1 \equiv 7 \pmod{8}$ and $p_2 \equiv 7 \pmod{8}$.

5 Conclusions

In the paper we defined sufficient conditions for designing Ding-Helleseth sequences with arbitrary period and high linear complexity for generalized cyclotomies. In particular, we have that the most frequently considered variant of Ding-Helleseth sequences possesses high linear complexity for any period. Also we discuss the method of computing the linear complexity of Ding-Helleseth sequences in the general case.

References

1. T. W. Cusick, C. Ding, A. Renvall, Stream Ciphers and Number Theory. Elsevier, Amsterdam, 1998.
2. C. Ding, T. Helleseht, New generalized cyclotomy and its applications. *Finite Fields and Their Applications*, 4(2), (1998), 140–166.
3. C. Ding, Linear complexity of some generalized cyclotomic sequences. *Int. J. Algebra Comput.* 8(4), (1998), 431–442.
4. Y. H. Park, D. Hong and E. Chun, On the linear complexity of some generalized cyclotomic sequences. *Int. J. Algebra Comput.* 14(4), (2004), 431–439
5. E. Bai, X. Liu, Some notes on prime-square sequences. *J. Comput. Sci. Technol.* 22(3), pp. 481–486 (2007)
6. Y. J. Kim, S.-Y. Jin and H.-Y. Song, Linear complexity and autocorrelation of prime cube sequences. In: *Lecture Notes in Computer Science*, vol. 4851, 188–197. Springer, Berlin (2007).
7. Y.J. Kim, H.Y. Song, Linear complexity of prime n-square sequences. In: *IEEE International Symposium on Information Theory*, Toronto, Canada, 2405–2408 (2008)
8. T. Yan, S. Li and G. Xiao, On the linear complexity of generalized cyclotomic sequences with the period p^m . *Appl. Math. Lett.* 21 (2008), 187–193.

9. V. Edemskiy, About computation of the linear complexity of generalized cyclotomic sequences with period p^{n+1} . *Designs, Codes and Cryptography*, vol. 61 (3), (2011), 251–260.
10. T. Yan, R. Sun and G. Xiao, Autocorrelation and linear complexity of the new generalized cyclotomic sequences. *IEICE Trans. Fundamentals of Electronics, Communications and Computer Sciences*, E90-A (4), (2007), 857–864.
11. T. Yan, Z. Chen and G. Xiao, Linear complexity of Ding generalized cyclotomic sequences. *Journal of Shanghai University*, 11(1), (2007), 22–26.
12. E. Bai, X. Liu and G. Xiao, Linear complexity of new generalized cyclotomic sequences of order two of length pq . *IEEE Transactions on Information Theory*, 51(5), (2005), 1849–1853.
13. T. Yan, L. Hong and G. Xiao, The linear complexity of new generalized cyclotomic binary sequences of order four. *Information Sciences*, 178(3), (2008), 807–815.
14. ZX. Chen, SQ. Li, Some notes on generalized cyclotomic sequences of length pq . *Journal of Computer Science and Technology*, 23(5), (2008), 843–850.
15. T. Yan, Linear Complexity of Ding-Helleseth Generalized Cyclotomic Binary Sequences of Any Order. Publication: eprint arXiv:1108.4450
16. V. Edemskiy, O. Antonova, About Computation of the Linear Complexity of Generalized Cyclotomic Sequences with Period pq . *Proc. Inter. Workshop on Signal Design and Its Applications in Communications (IWSDA'011)*. China. 2011, 9–12 (2011)
17. C. Ding, T. Hellesteth and W. Shan, On the linear complexity of Legendre sequences. *IEEE Trans. Inform. Theory* 44, (1998), 1276–1278.