

An isoperimetric inequality for extremal Sobolev functions

Tom Carroll ^{*} and Jesse Ratzkin [†]

April 23, 2018

Abstract

Let $D \subset \mathbf{R}^n$ be a bounded domain with a Lipschitz boundary, let $1 < p < \frac{2n}{n-2}$, and let ϕ minimize the ratio $\|\nabla u\|_{L^2}/\|u\|_{L^p}$. We prove a reverse-Hölder inequality, finding a lower bound for $\|\phi\|_{L^{p-1}}$ in terms of $\|\phi\|_{L^p}$, in which equality holds if and only if D is a ball. This result generalizes an inequality due to Payne and Rayner [6, 7] regarding eigenfunctions of the Laplacian.

1 Introduction and statement of results

Let $D \subset \mathbf{R}^n$ be a bounded domain with Lipschitz boundary, and let $1 < p < \frac{2n}{n-2}$ (or, $p > 1$ if $n = 2$). For this range of exponents, the Sobolev embedding $W_0^{1,2}(D) \hookrightarrow L^p(D)$ is compact, and so the infimum

$$C_p(D) = \inf \left\{ \frac{\int_D |\nabla u|^2 d\mu}{\left(\int_D |u|^p d\mu\right)^{2/p}} : u \in W_0^{1,2}(D), u \neq 0 \right\} \quad (1.1)$$

is finite and achieved by a nontrivial function $\phi = \phi_p$.

We take this opportunity to set notation for the remainder of the paper. We denote the volume element of the usual Lebesgue measure in \mathbf{R}^n by $d\mu$; when it will be necessary, we will denote the induced area element on a hypersurface $\Sigma \subset \mathbf{R}^n$ by $d\sigma$. We write the appropriate dimensional volume of a set as $|\Omega|$, *i.e.* if $\Omega \subset \mathbf{R}^n$ is an open set then $|\Omega| = \mu(\Omega)$ and if $\Sigma \subset \mathbf{R}^n$ is a hypersurface then $|\Sigma| = \sigma(\Sigma)$. If $\mathbf{B}_1 \subset \mathbf{R}^n$ is the unit ball, we denote $|\mathbf{B}_1| = \omega_n$, so that $|\mathbf{B}_r| = \omega_n r^n$ and $|\partial\mathbf{B}_r| = n\omega_n r^{n-1}$. The Sobolev space $W_0^{1,2}(D)$ is the closure of $C_0^\infty(D)$ under the norm $\|u\|_{W^{1,2}}^2 = \|u\|_{L^2}^2 + \|\nabla u\|_{L^2}^2$.

An extremal function ϕ for (1.1) will solve the boundary value problem in D :

$$\Delta\phi + \lambda\phi^{p-1} = 0, \quad \phi|_{\partial D} = 0. \quad (1.2)$$

Without loss of generality we can take $\phi > 0$ inside D . General regularity results imply that $\phi \in C_0^\infty(D)$, and a short integration by parts argument reveals that

$$\lambda = C_p(D) \left(\int_D \phi^p d\mu \right)^{\frac{2-p}{p}}. \quad (1.3)$$

^{*}School of Mathematical Sciences, University College Cork, t.carroll@ucc.ie

[†]Department of Mathematics and Applied Mathematics, University of Cape Town, jesse.ratzkin@uct.ac.za

This sharp Sobolev constant $C_p(D)$ and its associated extremal function ϕ_p are both the subject of a vast literature, and incorporate much information relating the function theory and the geometry of D . In particular, a long string of results (for example, [6, 4, 1, 3]) have uncovered isoperimetric-type inequalities of varying sorts. Our main theorem generalizes the reverse-Hölder inequalities of [6, 7, 2], and has the following form.

Theorem 1. *Let $D \subset \mathbf{R}^n$ be a bounded domain with Lipschitz boundary, let $C_p(D)$ be the sharp Sobolev constant defined by (1.1), and let ϕ be its associated extremal function. Let D^* be a ball with the same volume as D . Then*

$$\left(\int_D \phi^{p-1} d\mu \right)^2 \geq |D|^{\frac{n-2}{n}} \left(\int_D \phi^p d\mu \right)^{\frac{2(p-1)}{p}} \left[\frac{2n^2 \omega_n^{2/n}}{p C_p(D)} - (n-2) \frac{n \omega_n^{\frac{2}{n} + \frac{p^2-p+2}{p(p-1)}}}{C_p(D^*)} \right]. \quad (1.4)$$

Equality holds if and only if D is a ball.

Remark 1. • *The Hölder inequality implies that for any $u \in W_0^{1,2}$ we have*

$$\int_D |u|^{p-1} d\mu \leq |D|^{1/p} \left(\int_D |u|^p d\mu \right)^{\frac{p-1}{p}}.$$

For this reason, upper bounds of the form (1.4) are called reverse-Hölder inequalities.

- *Observe that we recover the main inequality of [7] in the case $p = 2$, and we recover the reverse-Hölder inequality of [2] in the case $n = 2$.*

ACKNOWLEDGEMENTS: T. C. is partially supported by the programme of the ESF Network ‘Harmonic and Complex Analysis and Applications’ (HCAA). J. R. is partially supported by a Carnegie Research grant from the University of Cape Town.

2 Proof of the main theorem

We begin by briefly outlining our strategy for proving (1.4), which we adapted from Payne and Rayner’s proof in [7]. Let $M = \sup_{x \in D} \phi(x)$, and for $0 \leq t \leq M$ we define

$$D_t = \{x \in D : \phi(x) > t\}, \quad \Sigma_t = \{x \in D : \phi(x) = t\}.$$

By Sard’s theorem, we have $\Sigma_t = \partial D_t$ for almost every value of t . To prove (1.4) we define the auxilliary function

$$H(t) = \int_{D_t} \phi^{p-1} d\mu = \int_t^M \tau^{p-1} \int_{\Sigma_\tau} \frac{d\sigma}{|\nabla \phi|} d\tau, \quad t \in [0, M].$$

In Section 2.1 we derive lower bounds for the second derivative of H , and in Section 2.2 we integrate these to obtain several integral inequalities for H and for powers of ϕ . In Section 2.3 we examine the one-dimensional eigenvalue problem which arises from the radially symmetric case, and in Section 2.4 complete the proof of (1.4).

2.1 Differential inequalities

We let $V(t) = |D_t|$. Then, by the co-area formula,

$$V'(t) = - \int_{\Sigma_t} \frac{d\sigma}{|\nabla\phi|} < 0.$$

Thus V is a monotone function of t , and we can invert it to obtain $t = t(V)$, with

$$\frac{dt}{dV} = \frac{1}{V'(t)} = - \frac{1}{\int_{\Sigma_t} \frac{d\sigma}{|\nabla\phi|}}.$$

This in turn implies that

$$\frac{dH}{dV} = \frac{dH}{dt} \frac{dt}{dV} = \left(-t^{p-1} \int_{\Sigma_t} \frac{d\sigma}{|\nabla\phi|} \right) \cdot \left(-\frac{1}{\int_{\Sigma_t} \frac{d\sigma}{|\nabla\phi|}} \right) = t^{p-1}, \quad (2.1)$$

a relation which will prove quite useful in our computations. Taking one more derivative shows that

$$\frac{d^2H}{dV^2} = \frac{d}{dV}(t^{p-1}) = -\frac{(p-1)t^{p-2}}{\int_{\Sigma_t} \frac{d\sigma}{|\nabla\phi|}}.$$

Lemma 1. *The function H satisfies*

$$\frac{d^2H}{dV^2} \geq -(p-1)(t(V))^{p-2} \frac{\Lambda H(V)}{n^2 \omega_n^{2/n} V^{\frac{2(n-1)}{n}}}, \quad V \in [0, |D|]. \quad (2.2)$$

with the boundary conditions $H(0) = 0$ and $H'(|D|) = 0$. Moreover, equality in (2.2) forces D to be a ball, and forces the function ϕ to be radially symmetric.

Proof. By the Cauchy-Schwarz inequality,

$$|\Sigma_t|^2 \leq \left(\int_{\Sigma_t} |\nabla\phi| d\sigma \right) \left(\int_{\Sigma_t} \frac{d\sigma}{|\nabla\phi|} \right),$$

which we can rearrange to read

$$\int_{\Sigma_t} \frac{d\sigma}{|\nabla\phi|} \geq \frac{|\Sigma_t|^2}{\int_{\Sigma_t} |\nabla\phi| d\sigma}. \quad (2.3)$$

Since Σ_t is a level-set of ϕ , we may use the divergence theorem and (1.2) to obtain

$$\begin{aligned} \int_{\Sigma_t} |\nabla\phi| d\sigma &= - \int_{\Sigma_t} \frac{\partial\phi}{\partial\eta} d\sigma = - \int_{D_t} \Delta\phi d\mu \\ &= \lambda \int_{D_t} \phi^{p-1} d\mu = \lambda H(t). \end{aligned}$$

Combining this with (2.3) we obtain

$$\int_{\Sigma_t} \frac{d\sigma}{|\nabla\phi|} \geq \frac{|\Sigma_t|^2}{\lambda H(t)}. \quad (2.4)$$

By the classical isoperimetric inequality,

$$|\Sigma_t|^2 \geq n^2 \omega_n^{2/n} |D_t|^{\frac{2(n-1)}{n}} = n^2 \omega_n^{2/n} V^{\frac{2(n-1)}{n}}. \quad (2.5)$$

Together with (2.4), this shows that

$$\begin{aligned} \frac{d^2 H}{dV^2} &= -(p-1) \frac{t^{p-2}}{\int_{\Sigma_t} \frac{d\sigma}{|\nabla\phi|}} \geq -(p-1) t^{p-2} \frac{\lambda H(V)}{|\Sigma_t|^2} \\ &\geq -(p-1) t^{p-2} \frac{\lambda H(V)}{n^2 \omega_n^{2/n} V^{\frac{2(n-1)}{n}}}. \end{aligned}$$

Notice that the boundary conditions for this differential inequality are

$$H(0) = 0, \quad H'(|D|) = t^{p-1}|_{t=0} = 0. \quad (2.6)$$

Moreover, we only have equality in (2.2) for each V in $[0, |D|]$ if we have equality in (2.5) for almost every t , which in turn implies that Σ_t is a round sphere for almost every $t \in [0, M]$. This is possible only if D is itself a ball. Also, equality in (2.2) forces equality in (2.3), which implies $\nabla\phi$ must be constant on each sphere Σ_t , and so ϕ must be radial. \square

We change variables by letting $\rho = (V/\omega_n)^{1/n}$ be the volume radius of D_t , so that $V = |D_t| = \omega_n \rho^n$. We also define $\rho_M = (|D|/\omega_n)^{1/n}$. As a function of ρ , the function H satisfies the boundary conditions

$$H(0) = H'(0) = \dots = H^{(n-1)}(0) = 0, \quad H'(\rho_M) = 0. \quad (2.7)$$

Lemma 2.

$$\frac{d}{d\rho} \left[\rho^{1-n} \left(\frac{dH}{d\rho} \right)^{\frac{1}{p-1}} \right] \geq - \frac{\lambda}{(n\omega_n)^{\frac{p-2}{p-1}}} \rho^{1-n} H(\rho), \quad 0 < \rho < \rho_M. \quad (2.8)$$

Proof. Taking derivatives, we see that

$$\frac{dH}{dV} = \frac{\rho^{1-n}}{n\omega_n} \frac{dH}{d\rho}, \quad \frac{d^2 H}{dV^2} = \frac{\rho^{1-n}}{n\omega_n} \frac{d}{d\rho} \left(\frac{\rho^{1-n}}{n\omega_n} \frac{dH}{d\rho} \right). \quad (2.9)$$

Substituting these expressions in (2.2) gives

$$\frac{d}{d\rho} \left(\rho^{1-n} \frac{dH}{d\rho} \right) \geq -(p-1) \lambda t^{p-2} \rho^{1-n} H(\rho). \quad (2.10)$$

However,

$$t^{p-1} = \frac{dH}{dV} = \frac{\rho^{1-n}}{n\omega_n} \frac{dH}{d\rho},$$

so that (2.10) becomes

$$\frac{d}{d\rho} \left(\rho^{1-n} \frac{dH}{d\rho} \right) \geq - \frac{(p-1) \lambda}{(n\omega_n)^{\frac{p-2}{p-1}}} \left(\rho^{1-n} \frac{dH}{d\rho} \right)^{\frac{p-2}{p-1}} \rho^{1-n} H(\rho).$$

This we can rewrite as

$$\frac{1}{p-1} \frac{\frac{d}{d\rho} \left(\rho^{1-n} \frac{dH}{d\rho} \right)}{\left(\rho^{1-n} \frac{dH}{d\rho} \right)^{\frac{p-2}{p-1}}} = \frac{d}{d\rho} \left[\rho^{1-n} \left(\frac{dH}{d\rho} \right)^{\frac{1}{p-1}} \right] \geq -\frac{\lambda}{(n\omega_n)^{\frac{p-2}{p-1}}} \rho^{1-n} H(\rho). \quad \square$$

Remark 2. Since (2.8) is really the same as (2.2) rewritten in different variables, equality holds in (2.8) for $0 < \rho < \rho_M$ if and only if D is a ball and ϕ is radial.

2.2 Integral inequalities

In this section we integrate (2.2) and (2.8) to obtain inequalities for the integral of H and the integral of powers of ϕ . As each of these inequalities is an integrated form of (2.2) and (2.8), equality holds if and only if D is a ball and ϕ is radial.

Lemma 3.

$$\begin{aligned} \left(\int_D \phi^{p-1} d\mu \right)^2 &\geq \frac{2n^2 \omega_n^{2/n} |D|^{\frac{n-2}{n}}}{p C_p(D)} \left(\int_D \phi^p d\mu \right)^{\frac{2(p-1)}{p}} \\ &\quad - \frac{n-2}{n} |D|^{\frac{n-2}{n}} \int_0^{|D|} V^{\frac{2(1-n)}{n}} H^2(V) dV. \end{aligned} \quad (2.11)$$

Proof. We multiply the inequality (2.2) by $\frac{p}{p-1} V \left(\frac{dH}{dV} \right)^{1/(p-1)}$ and integrate from 0 to $|D|$. Upon integration, the left hand side becomes

$$\begin{aligned} \int_0^{|D|} \frac{p}{p-1} V \left(\frac{dH}{dV} \right)^{1/(p-1)} \frac{d^2 H}{dV^2} dV &= \int_0^{|D|} V \frac{d}{dV} \left[\left(\frac{dH}{dV} \right)^{p/(p-1)} \right] dV \\ &= V \left(\frac{dH}{dV} \right)^{p/(p-1)} \Big|_0^{|D|} - \int_0^{|D|} \left(\frac{dH}{dV} \right)^{p/(p-1)} dV \\ &= - \int_0^{|D|} (t^{p-1}(V))^{p/(p-1)} dV \\ &= - \int_0^{|D|} t^p(V) dV = - \int_D \phi^p d\mu. \end{aligned}$$

The boundary terms in the integration by parts vanished since $H'(|D|) = 0$, while (2.1) was used at the third step. On the other hand, using (2.1) again, the right hand side becomes

$$\begin{aligned} -\frac{p\lambda}{n^2 \omega_n^{n/2}} \int_0^{|D|} V \left(\frac{dH}{dV} \right)^{1/(p-1)} t^{p-2} H(t) V^{\frac{2(1-n)}{n}} dV \\ &= -\frac{p\lambda}{n^2 \omega_n^{n/2}} \int_0^{|D|} t^{p-2} \left(\frac{dH}{dV} \right)^{\frac{2-p}{p-1}} V^{\frac{2-n}{n}} H(V) \frac{dH}{dV} dV \\ &= -\frac{p\lambda}{n^2 \omega_n^{n/2}} \int_0^{|D|} t^{p-2} (t^{p-1})^{\frac{2-p}{p-1}} V^{\frac{2-n}{n}} H(V) \frac{dH}{dV} dV \\ &= -\frac{p\lambda}{n^2 \omega_n^{n/2}} \int_0^{|D|} V^{\frac{2-n}{n}} H(V) \frac{dH}{dV} dV \end{aligned}$$

We combine these last two equations and replace λ by $\mathcal{C}_p(D) \left(\int_D \phi^p d\mu\right)^{(2-p)/p}$ to obtain

$$\begin{aligned} \left(\int_D \phi^p d\mu\right)^{\frac{2(p-1)}{p}} &\leq \frac{p\mathcal{C}_p(D)}{n^2\omega_n^{n/2}} \int_0^{|D|} V^{\frac{2-n}{n}} H(V) \frac{dH}{dV} dV \\ &= \frac{p\mathcal{C}_p(D)}{2n^2\omega_n^{2/n}} \int_0^{|D|} V^{\frac{2-n}{n}} \frac{d}{dV} (H^2(V)) dV \\ &= \frac{p\mathcal{C}_p(D)}{2n^2\omega_n^{2/n}} \left[|D|^{\frac{2-n}{n}} \left(\int_D \phi^{p-1} d\mu\right)^2 + \frac{n-2}{n} \int_0^{|D|} H^2(V) V^{\frac{2(1-n)}{n}} dV \right], \end{aligned}$$

which we can rearrange to give (2.11). \square

Lemma 4.

$$\int_0^{\rho_M} \rho^{\frac{1-n}{p-1}} \left(\frac{dH}{d\rho}\right)^{\frac{p}{p-1}} \leq \frac{\lambda}{(n\omega_n)^{\frac{p-2}{p-1}}} \int_0^{\rho_M} \rho^{1-n} H^2(\rho) d\rho. \quad (2.12)$$

Equality holds if and only if D is a ball.

Proof. We multiply (2.8) by H and integrate from 0 to ρ_M . The boundary conditions (2.7) imply that $\rho^{1-n} \frac{dH}{d\rho}$ is bounded at 0. Hence the boundary terms vanish in the integration parts below, and we obtain that

$$\int_0^{\rho_M} \rho^{\frac{1-n}{p-1}} \left(\frac{dH}{d\rho}\right)^{\frac{p}{p-1}} d\rho = \int_0^{\rho_M} \left[\rho^{1-n} \frac{dH}{d\rho}\right]^{\frac{1}{p-1}} \frac{dH}{d\rho} d\rho \leq \frac{\lambda}{(n\omega_n)^{\frac{p-2}{p-1}}} \int_0^{\rho_M} \rho^{1-n} H^2(\rho) d\rho.$$

\square

Lemma 5. *With ϕ , H , and ρ defined as above,*

$$\int_0^{\rho_M} \rho^{\frac{1-n}{p-1}} \left(\frac{dH}{d\rho}\right)^{\frac{p}{p-1}} d\rho = (n\omega_n)^{\frac{1}{p-1}} \int_D \phi^p d\mu. \quad (2.13)$$

Proof. We use (2.1) and (2.9) to conclude

$$\begin{aligned} \int_0^{\rho_M} \rho^{\frac{1-n}{p-1}} \left(\frac{dH}{d\rho}\right)^{\frac{p}{p-1}} &= \int_0^{\rho_M} (\rho^{1-n})^{\frac{p}{p-1}} \left(\frac{dH}{d\rho}\right)^{\frac{p}{p-1}} \rho^{n-1} d\rho \\ &= \int_0^{|D|} \left(\rho^{1-n} \frac{dH}{d\rho}\right)^{\frac{p}{p-1}} \frac{dV}{n\omega_n} \\ &= \int_0^{|D|} \left(n\omega_n \frac{dH}{dV}\right)^{\frac{p}{p-1}} \frac{dV}{n\omega_n} \\ &= (n\omega_n)^{\frac{1}{p-1}} \int_0^{|D|} \left(\frac{dH}{dV}\right)^{\frac{p}{p-1}} dV \\ &= (n\omega_n)^{\frac{1}{p-1}} \int_0^{|D|} (t^{p-1})^{\frac{p}{p-1}} dV \\ &= (n\omega_n)^{\frac{1}{p-1}} \int_0^{|D|} t^p dV = (n\omega_n)^{\frac{1}{p-1}} \int_D \phi^p d\mu. \quad \square \end{aligned}$$

Corollary 2.

$$n\omega_n \left(\int_D \phi^p d\mu \right)^{\frac{2(p-1)}{p}} \leq \mathcal{C}_p(D) \int_0^{\rho_M} \rho^{1-n} H^2(\rho) d\rho. \quad (2.14)$$

Moreover, we have equality if and only if D is a ball and ϕ is radial.

Proof. Combine (2.12), (2.13), and (1.3). \square

Lemma 6.

$$\begin{aligned} \left(\int_D \phi^{p-1} d\mu \right)^2 &\geq \frac{2n^2 \omega_n^{2/n} |D|^{\frac{n-2}{n}}}{p \mathcal{C}_p(D)} \left(\int_D \phi^p d\mu \right)^{\frac{2(p-1)}{p}} \\ &\quad - (n-2) \omega_n^{\frac{2-n}{n}} |D|^{\frac{n-2}{n}} \int_0^{\rho_M} \rho^{1-n} H^2(\rho) d\rho. \end{aligned} \quad (2.15)$$

Equality holds if and only if D is a ball.

Proof. Since $\rho(V) = (V/\omega_n)^{1/n}$, we have

$$n\omega_n^{1/n} \frac{d\rho}{dV} V^{\frac{n-1}{n}} = 1,$$

so that

$$\begin{aligned} \int_0^{|D|} V^{\frac{2(1-n)}{n}} H^2(V) dV &= \int_0^{|D|} H^2(\rho) \omega_n^{\frac{2(1-n)}{n}} \rho^{2(1-n)} n\omega_n^{1/n} (\omega_n \rho^n)^{\frac{n-1}{n}} \frac{d\rho}{dV} dV \\ &= n\omega_n^{\frac{2-n}{n}} \int_0^{\rho_M} \rho^{1-n} H^2(\rho) d\rho. \end{aligned}$$

Putting this into (2.11) gives (2.15). \square

2.3 The radially symmetric case

Motivated by (2.12) and (2.7), we define Λ_* by

$$\Lambda_* = \inf \left\{ \frac{\left(\int_0^{\rho_M} \rho^{\frac{1-n}{p-1}} f'(\rho)^{\frac{p}{p-1}} d\rho \right)^{\frac{2(p-1)}{p}}}{\int_0^{\rho_M} \rho^{1-n} f^2(\rho) d\rho} \right\} \quad (2.16)$$

where the infimum is over all functions on $[0, \rho_M]$ for which

$$f(0) = f'(0) = \dots = f^{(n-1)}(0) = 0 = f'(\rho_M), \quad f \not\equiv 0. \quad (2.17)$$

Remark 3. Notice that we have rescaled the numerator to make the quotient scale-invariant. This does not, however, affect the Euler-Lagrange equation involved.

Lemma 7. The Euler-Lagrange equation for the variational problem (2.16), with the boundary conditions (2.17), is

$$f''(\rho) - \frac{n-1}{\rho} f'(\rho) + \Lambda [\rho^{1-n} f'(\rho)]^{\frac{p-2}{p-1}} f(\rho) = 0. \quad (2.18)$$

Proof. Since the ratio defining Λ_* is scale-invariant, we may either restrict our attention to either of the constrained critical point problems:

$$\text{minimize } \int_0^{\rho_M} \rho^{\frac{1-n}{p-1}} f'(\rho)^{\frac{p}{p-1}} d\rho \text{ subject to } \int_0^{\rho_M} \rho^{1-n} f^2 d\rho = \text{constant}$$

or

$$\text{maximize } \int_0^{\rho_M} \rho^{1-n} f^2 d\rho \text{ subject to } \int_0^{\rho_M} \rho^{\frac{1-n}{p-1}} f'(\rho)^{\frac{p}{p-1}} d\rho = \text{constant.}$$

Regardless, the method of Lagrange multipliers implies that a constrained critical point f satisfies

$$\frac{d}{d\epsilon} \Big|_{\epsilon=0} \int_0^{\rho_M} \rho^{\frac{1-n}{p-1}} \left(\frac{df}{d\rho} + \epsilon \frac{dg}{d\rho} \right)^{\frac{p}{p-1}} d\rho = \Lambda \frac{d}{d\epsilon} \Big|_{\epsilon=0} \int_0^{\rho_M} \rho^{1-n} [f(\rho) + \epsilon g(\rho)]^2 d\rho,$$

for any admissible g . Having evaluated these derivatives, we use the boundary conditions (2.17) to see that $\rho^{1-n} f'(\rho)$ is bounded at 0 and that consequently the boundary terms arising from integration by parts vanish, and see that

$$\begin{aligned} & 2\lambda \int_0^{\rho_M} \rho^{1-n} f(\rho) g(\rho) d\rho \\ &= \frac{p}{p-1} \int_0^{\rho_M} \rho^{\frac{1-n}{p-1}} \left(\frac{df}{d\rho} \right)^{\frac{1}{p-1}} \frac{dg}{d\rho} d\rho \\ &= -\frac{p}{p-1} \int_0^{\rho_M} g(\rho) \frac{d}{d\rho} \left[\rho^{\frac{1-n}{p-1}} \left(\frac{df}{d\rho} \right)^{\frac{1}{p-1}} \right] d\rho \\ &= -\frac{p}{p-1} \int_0^{\rho_M} g(\rho) \left[\frac{1}{p-1} \rho^{\frac{1-n}{p-1}} \left(\frac{df}{d\rho} \right)^{\frac{2-p}{p-1}} \frac{d^2 f}{d\rho^2} + \frac{1-n}{p-1} \rho^{\frac{2-p-n}{p-1}} \left(\frac{df}{d\rho} \right)^{\frac{1}{p-1}} \right] d\rho. \end{aligned}$$

This must hold for all choices of g , hence (absorbing a factor of $2p/(p-1)^2$ into the Lagrange multiplier Λ) we must have

$$\begin{aligned} 0 &= \rho^{\frac{1-n}{p-1}} f'(\rho)^{\frac{2-p}{p-1}} f''(\rho) - (n-1) \rho^{\frac{2-p-n}{p-1}} f'(\rho)^{\frac{1}{p-1}} + \Lambda \rho^{1-n} f(\rho) \\ &= \rho^{\frac{1-n}{p-1}} f'(\rho)^{\frac{2-p}{p-1}} \left[f''(\rho) - (n-1) \rho^{-1} f'(\rho) + \Lambda [\rho^{1-n} f'(\rho)]^{\frac{p-2}{p-1}} f(\rho) \right], \end{aligned}$$

as claimed. \square

Lemma 8. *Let D^* be the ball \mathbf{B}_{ρ_M} of radius ρ_M . Then,*

$$\Lambda_* \leq (n\omega_n)^{\frac{2-p}{p}} \mathcal{C}_p(D^*). \quad (2.19)$$

Proof. We use the function $H(\rho)$ for the ball \mathbf{B}_{ρ_M} as a test function for the quotient

defining Λ_* and use the inequalities (2.12), (1.3), and (2.13):

$$\begin{aligned}
\Lambda_* &\leq \left(\int_0^{\rho_M} \rho^{\frac{1-n}{p-1}} H'(\rho)^{\frac{p}{p-1}} d\rho \right)^{\frac{2(p-1)}{p}} / \int_0^{\rho_M} \rho^{1-n} H^2(\rho) d\rho \\
&\leq \frac{\Lambda}{(n\omega_n)^{\frac{p-2}{p-1}}} \left(\int_0^{\rho_M} \rho^{\frac{1-n}{p-1}} H'(\rho)^{\frac{p}{p-1}} d\rho \right)^{\frac{p-2}{p}} \\
&= \frac{1}{(n\omega_n)^{\frac{p-2}{p-1}}} \mathcal{C}_p(D^*) \left(\int_{D^*} \phi^p d\mu \right)^{\frac{2-p}{p}} \left[\frac{1}{(n\omega_n)^{\frac{1}{p-1}}} \int_{D^*} \phi^p d\mu \right]^{\frac{p-2}{p}} \\
&= (n\omega_n)^{\frac{2-p}{p}} \mathcal{C}_p(D^*). \quad \square
\end{aligned}$$

In order to obtain a lower bound for Λ_* in terms of $\mathcal{C}_p(D^*)$, we first need to relate the particular Λ occurring in the Euler-Lagrange equation (2.18) to the eigenvalue Λ_* , just as (1.3) relates the number λ occurring in the Euler-Lagrange equation (1.2) to the eigenvalue $\mathcal{C}_p(D)$.

Lemma 9. *Let f be a minimizer for Λ_* given by (2.16) with the boundary conditions (2.17) and satisfy the Euler-Lagrange equation (2.18), written as*

$$\frac{d}{d\rho} \left[\rho^{\frac{1-n}{p-1}} f'(\rho)^{\frac{1}{p-1}} \right] + \Lambda \rho^{1-n} f(\rho) = 0. \quad (2.20)$$

Then

$$\Lambda = \Lambda_* \left(\int_0^{\rho_M} \rho^{\frac{1-n}{p-1}} f'(\rho)^{\frac{p}{p-1}} d\rho \right)^{\frac{2-p}{p}}. \quad (2.21)$$

Proof. Multiply the Euler-Lagrange equation(2.20) across by $f(\rho)$ and integrate from 0 to ρ_M to obtain

$$\int_0^{\rho_M} f(\rho) \frac{d}{d\rho} \left[\rho^{\frac{1-n}{p-1}} f'(\rho)^{\frac{1}{p-1}} \right] d\rho + \Lambda \int_0^{\rho_M} \rho^{1-n} f(\rho)^2 d\rho = 0.$$

Integrating by parts in the first term and using the boundary conditions (2.17) gives

$$\int_0^{\rho_M} f(\rho) \frac{d}{d\rho} \left[\rho^{\frac{1-n}{p-1}} f'(\rho)^{\frac{1}{p-1}} \right] d\rho = - \int_0^{\rho_M} \rho^{\frac{1-n}{p-1}} f'(\rho)^{\frac{p}{p-1}} d\rho,$$

from which it follows that

$$\Lambda = \int_0^{\rho_M} \rho^{\frac{1-n}{p-1}} f'(\rho)^{\frac{p}{p-1}} d\rho / \int_0^{\rho_M} \rho^{1-n} f(\rho)^2 d\rho.$$

We can use (2.16) to write $\int_0^{\rho_M} \rho^{1-n} f^2(\rho) d\rho$ in terms of Λ_* since f is a minimizer for this Rayleigh quotient, leading to

$$\Lambda = \Lambda_* \left(\int_0^{\rho_M} \rho^{\frac{1-n}{p-1}} f'(\rho)^{\frac{p}{p-1}} d\rho \right)^{1 - \frac{2(p-1)}{p}},$$

which is (2.21). □

Lemma 10.

$$\mathcal{C}_p(D^*) \leq (n\omega_n)^{\frac{p-2}{p}} \Lambda_*. \quad (2.22)$$

Proof. Let f be a minimizer for the generalized quotient (2.16) defining Λ_* . Set

$$\psi(\rho) = \int_{\rho}^{\rho_M} r^{1-n} f(r) dr, \quad 0 \leq \rho \leq \rho_M,$$

so that $\psi(\rho_M) = 0$. Then $\psi(\rho)$ (where $\rho = |x|$ for $x \in \mathcal{C}_p(D^*)$) is an admissible test function for the quotient defining $\mathcal{C}_p(D^*)$. Thus

$$\mathcal{C}_p(D^*) \leq (n\omega_n)^{\frac{p-2}{p}} \int_0^{\rho_M} \rho^{n-1} \psi'(\rho)^2 d\rho / \left(\int_0^{\rho_M} \rho^{n-1} \psi(\rho)^p d\rho \right)^{2/p}. \quad (2.23)$$

Now

$$\int_0^{\rho_M} \rho^{n-1} \psi'(\rho)^2 d\rho = \int_0^{\rho_M} \rho^{n-1} [\rho^{1-n} f(\rho)]^2 d\rho = \int_0^{\rho_M} \rho^{1-n} f(\rho)^2 d\rho. \quad (2.24)$$

Next, using the Euler-Lagrange equation (2.20),

$$\begin{aligned} \psi(\rho) &= \int_{\rho}^{\rho_M} r^{1-n} f(r) dr \\ &= -\frac{1}{\Lambda} \int_{\rho}^{\rho_M} \frac{d}{dr} \left[r^{\frac{1-n}{p-1}} f'(r)^{\frac{1}{p-1}} \right] dr \\ &= -\frac{1}{\Lambda} r^{\frac{1-n}{p-1}} f'(r)^{\frac{1}{p-1}} \Big|_{r=\rho}^{r=\rho_M} \\ &= \frac{1}{\Lambda} \rho^{\frac{1-n}{p-1}} f'(\rho)^{\frac{1}{p-1}}, \end{aligned}$$

where we used $f'(\rho_M) = 0$. From this we obtain that

$$\begin{aligned} \int_0^{\rho_M} \rho^{n-1} \psi(\rho)^p d\rho &= \int_0^{\rho_M} \rho^{n-1} \frac{1}{\Lambda^p} \rho^{\frac{p(1-n)}{p-1}} f'(\rho)^{\frac{p}{p-1}} d\rho \\ &= \frac{1}{\Lambda^p} \int_0^{\rho_M} \rho^{\frac{1-n}{p-1}} f'(\rho)^{\frac{p}{p-1}} d\rho. \end{aligned} \quad (2.25)$$

With the help of the identities (2.24) and (2.25), we can write the numerator and the denominator of (2.23) in terms of the minimizer f for Λ_* . We find, using that f minimizes the quotient for Λ_* at the second step and using (2.21) at the third step, that

$$\mathcal{C}_p(D^*) \leq (n\omega_n)^{\frac{p-2}{p}} \int_0^{\rho_M} \rho^{1-n} f(\rho)^2 d\rho / \left(\frac{1}{\Lambda^p} \int_0^{\rho_M} \rho^{\frac{1-n}{p-1}} f'(\rho)^{\frac{p}{p-1}} d\rho \right)^{\frac{2}{p}} \quad (2.26)$$

$$= (n\omega_n)^{\frac{p-2}{p}} \frac{\Lambda^2}{\Lambda_*^2} \left(\int_0^{\rho_M} \rho^{\frac{1-n}{p-1}} f'(\rho)^{\frac{p}{p-1}} d\rho \right)^{\frac{2(p-1)}{p} - \frac{2}{p}} \quad (2.27)$$

$$= (n\omega_n)^{\frac{p-2}{p}} \frac{1}{\Lambda_*} \left[\Lambda \left(\int_0^{\rho_M} \rho^{\frac{1-n}{p-1}} f'(\rho)^{\frac{p}{p-1}} d\rho \right)^{\frac{p-2}{p}} \right]^2 \quad (2.28)$$

$$= (n\omega_n)^{\frac{p-2}{p}} \frac{\Lambda_*^2}{\Lambda_*} = (n\omega_n)^{\frac{p-2}{p}} \Lambda_*. \quad \square$$

2.4 Completion of the proof of Theorem 1

We can now finally complete the proof of Theorem 1. Indeed, we have

$$\begin{aligned} \int_0^{\rho_M} \rho^{1-n} H^2(\rho) d\rho &\leq \frac{1}{\Lambda_*} \left(\int_0^{\rho_M} \rho^{\frac{1-n}{p-1}} \left(\frac{dH}{d\rho} \right)^{\frac{p}{p-1}} d\rho \right)^{\frac{2(p-1)}{p}} \\ &= \frac{1}{\Lambda_*} \left((n\omega_n)^{\frac{1}{p-1}} \int_D \phi^p d\mu \right)^{\frac{2(p-1)}{p}} \\ &= \frac{1}{\Lambda_*} (n\omega_n)^{2/p} \left(\int_D \phi^p d\mu \right)^{\frac{2(p-1)}{p}}, \end{aligned}$$

with equality if and only if D is a ball and ϕ is radial. Substituting this last inequality into (2.10), we have

$$\begin{aligned} \left(\int_D \phi^{p-1} d\mu \right)^2 &\leq \frac{2n^2 \omega_n^{n/2}}{p \mathcal{C}_p(D)} |D|^{\frac{n-2}{n}} \left(\int_D \phi^p d\mu \right)^{\frac{2(p-1)}{p}} \\ &\quad - (n-2) \omega_n^{\frac{2-n}{n}} |D|^{\frac{n-2}{n}} \frac{1}{\Lambda_*} (n\omega_n)^{2/p} \left(\int_D \phi^p d\mu \right)^{\frac{2(p-1)}{p}}. \end{aligned}$$

Since $\Lambda_* = (n\omega_n)^{\frac{2-p}{p}} \mathcal{C}_p(D^*)$ by (2.19) and (2.22), the main inequality (1.4) follows with equality if and only if D is a ball. \square

References

- [1] A. Alvino, V. Ferone and G. Trombetti, *On the properties of some nonlinear eigenvalues*. SIAM J. Math. Anal. **29** (1998), 437–451.
- [2] T. Carroll and J. Ratzkin, *Two isoperimetric inequalities for the Sobolev constant*. to appear in Z. Angew. Math. Phys.
- [3] A. Chianchi, N. Fusco, F. Maggi, and A. Pratelli, *The sharp Sobolev inequality in quantitative form*. J. Eur. Math. Soc. **11**(2009), 1105–1139.
- [4] G. Chiti, *A reverse Hölder inequality for the eigenfunctions of linear second order elliptic operators*. Z. Angew. Math. Phys. **33** (1982), 143–148.
- [5] B. Gidas, W.N. Ni, and L. Nirenberg, *Symmetry and related properties via the maximum principle*. Comm. Math. Phys. **68** (1979), 209–243.
- [6] L. Payne and M. Rayner, *An isoperimetric inequality for the first eigenfunction in the fixed membrane problem*. J. Angew. Math. Phys. **23** (1972), 13–15.
- [7] L. Payne and M. Rayner, *Some isoperimetric norm bounds for solutions of the Helmholtz equation*. Z. Angew. Math. Phys. **24** (1973), 105–110.
- [8] G. Pólya and G. Szegő, *Isoperimetric Inequalities in Mathematical Physics*. Princeton University Press (1951).