

**ON GENERALIZATION OF DIFFERENT TYPE INEQUALITIES
FOR SOME CONVEX FUNCTIONS VIA FRACTIONAL
INTEGRALS**

İMDAT İŞCAN

ABSTRACT. New identity for fractional integrals have been defined. By using of this identity, we obtained new estimates on generalization of Hadamard, Ostrowski and Simpson type inequalities for s -convex, quasi-convex, m -convex functions via Riemann Liouville fractional integral.

1. INTRODUCTION

Following inequalities are well known in the literature as Hermite-Hadamard inequality, Ostrowski inequality and Simpson inequality respectively:

Theorem 1. *Let $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be a convex function defined on the interval I of real numbers and $a, b \in I$ with $a < b$. The following double inequality holds*

$$(1.1) \quad f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x)dx \leq \frac{f(a)+f(b)}{2}.$$

Theorem 2. *Let $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be a mapping differentiable in I° , the interior of I , and let $a, b \in I^\circ$ with $a < b$. If $|f'(x)| \leq M$, $x \in [a, b]$, then we the following inequality holds*

$$\left| f(x) - \frac{1}{b-a} \int_a^b f(t)dt \right| \leq \frac{M}{b-a} \left[\frac{(x-a)^2 + (b-x)^2}{2} \right]$$

for all $x \in [a, b]$. The constant $\frac{1}{4}$ is the best possible in the sense that it cannot be replaced by a smaller one.

Theorem 3. *Let $f : [a, b] \rightarrow \mathbb{R}$ be a four times continuously differentiable mapping on (a, b) and $\|f^{(4)}\|_\infty = \sup_{x \in (a,b)} |f^{(4)}(x)| < \infty$. Then the following inequality holds:*

$$\left| \frac{1}{3} \left[\frac{f(a)+f(b)}{2} + 2f\left(\frac{a+b}{2}\right) \right] - \frac{1}{b-a} \int_a^b f(x)dx \right| \leq \frac{1}{2880} \|f^{(4)}\|_\infty (b-a)^2.$$

In [5], Hudzik and Maligranda considered among others the class of functions which are s -convex in the second sense.

2000 *Mathematics Subject Classification.* 26A51, 26A33, 26D10, 26D15.

Key words and phrases. Hermite-Hadamard inequality, Riemann-Liouville fractional integral, Ostrowski inequality, Simpson type inequalities, s -convex function, quasi-convex function, m -convex function.

Definition 1. A function $f : [0, \infty) \rightarrow \mathbb{R}$ is said to be s -convex in the second sense if

$$f(\alpha x + \beta y) \leq \alpha^s f(x) + \beta^s f(y)$$

for all $x, y \in [0, \infty)$, $\alpha, \beta \geq 0$ with $\alpha + \beta = 1$ and for some fixed $s \in (0, 1]$. This class of s -convex functions in the second sense is usually denoted by K_s^2 .

It can be easily seen that for $s = 1$, s -convexity reduces to ordinary convexity of functions defined on $[0, \infty)$.

In [13], G. Toader considered the class of m -convex functions: another intermediate between the usual convexity and starshaped convexity.

Definition 2. The function $f : [0, b] \rightarrow \mathbb{R}$, $b > 0$, is said to be m -convex, where $m \in [0, 1]$, if we have

$$f(tx + m(1-t)y) \leq tf(x) + m(1-t)f(y)$$

for all $x, y \in [0, b]$ and $t \in [0, 1]$. We say that f is m -concave if $(-f)$ is m -convex.

In [10], Pecaric et al. defined quasi-convex functions as following

Definition 3. A function $f : [a, b] \rightarrow \mathbb{R}$ is said quasi-convex on $[a, b]$ if

$$f(\lambda x + (1-\lambda)y) \leq \sup\{f(x), f(y)\},$$

for any $x, y \in [a, b]$ and $\lambda \in [0, 1]$.

The notion of quasi-convex functions generalizes the notion of convex functions. Clearly, any convex function is a quasi-convex function. Furthermore, there exist quasi-convex functions which are not convex (see [6]).

We give some necessary definitions and mathematical preliminaries of fractional calculus theory which are used throughout this paper.

Definition 4. Let $f \in L[a, b]$. The Riemann-Liouville integrals $J_{a+}^\alpha f$ and $J_{b-}^\alpha f$ of order $\alpha > 0$ with $a \geq 0$ are defined by

$$J_{a+}^\alpha f(x) = \frac{1}{\Gamma(\alpha)} \int_a^x (x-t)^{\alpha-1} f(t) dt, \quad x > a$$

and

$$J_{b-}^\alpha f(x) = \frac{1}{\Gamma(\alpha)} \int_x^b (t-x)^{\alpha-1} f(t) dt, \quad x < b$$

respectively, where $\Gamma(\alpha)$ is the Gamma function defined by $\Gamma(\alpha) = \int_0^\infty e^{-t} t^{\alpha-1} dt$

and $J_{a+}^0 f(x) = J_{b-}^0 f(x) = f(x)$.

In the case of $\alpha = 1$, the fractional integral reduces to the classical integral. Properties concerning this operator can be found [4, 7, 9].

For some recent result connected with fractional integral see [3, 8, 11, 12].

2. GENERALIZED INTEGRAL INEQUALITIES FOR SOME CONVEX FUNCTIONS VIA FRACTIONAL INTEGRALS

Let $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be a differentiable function on I° , the interior of I , throughout this section we will take

$$S_f(x, \lambda, \alpha, a, b) = (1 - \lambda) \left[\frac{(x-a)^\alpha + (b-x)^\alpha}{b-a} \right] f(x) + \lambda \left[\frac{(x-a)^\alpha f(a) + (b-x)^\alpha f(b)}{b-a} \right] - \frac{\Gamma(\alpha+1)}{b-a} [J_{x^-}^\alpha f(a) + J_{x^+}^\alpha f(b)]$$

where $a, b \in I$ with $a < b$, $x \in [a, b]$, $\lambda \in [0, 1]$, $\alpha > 0$ and Γ is Euler Gamma function. In order to prove our main results we need the following identity.

Lemma 1. *Let $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be a differentiable function on I° such that $f' \in L[a, b]$, where $a, b \in I$ with $a < b$. Then for all $x \in [a, b]$, $\lambda \in [0, 1]$ and $\alpha > 0$ we have:*

$$(2.1) \quad S_f(x, \lambda, \alpha, a, b) = \frac{(x-a)^{\alpha+1}}{b-a} \int_0^1 (t^\alpha - \lambda) f'(tx + (1-t)a) dt + \frac{(b-x)^{\alpha+1}}{b-a} \int_0^1 (\lambda - t^\alpha) f'(tx + (1-t)b) dt.$$

Proof. By integration by parts and changing the variable, we can state

$$(2.2) \quad \int_0^1 (t^\alpha - \lambda) f'(tx + (1-t)a) dt = (t^\alpha - \lambda) \frac{f(tx + (1-t)a)}{x-a} \Big|_0^1 - \int_0^1 \alpha t^{\alpha-1} \frac{f(tx + (1-t)a)}{x-a} dt = (1-\lambda) \frac{f(x)}{x-a} + \lambda \frac{f(a)}{x-a} - \frac{\alpha}{x-a} \int_a^x \left(\frac{u-a}{x-a} \right)^{\alpha-1} \frac{f(u)}{x-a} du = (1-\lambda) \frac{f(x)}{x-a} + \lambda \frac{f(a)}{x-a} - \frac{\Gamma(\alpha+1)}{(x-a)^{\alpha+1}} J_{x^-}^\alpha f(a)$$

and similarly we get

$$(2.3) \quad \int_0^1 (\lambda - t^\alpha) f'(tx + (1-t)b) dt = (\lambda - t^\alpha) \frac{f(tx + (1-t)b)}{x-b} \Big|_0^1 - \int_0^1 \alpha t^{\alpha-1} \frac{f(tx + (1-t)b)}{x-b} dt = (1-\lambda) \frac{f(x)}{b-x} + \lambda \frac{f(b)}{b-x} - \frac{\alpha}{b-x} \int_x^b \left(\frac{b-u}{b-x} \right)^{\alpha-1} \frac{f(u)}{b-x} du = (1-\lambda) \frac{f(x)}{b-x} + \lambda \frac{f(b)}{b-x} - \frac{\Gamma(\alpha+1)}{(b-x)^{\alpha+1}} J_{x^+}^\alpha f(b)$$

Multiplying both sides of (2.2) and (2.3) by $\frac{(x-a)^{\alpha+1}}{b-a}$ and $\frac{(b-x)^{\alpha+1}}{b-a}$, respectively, and adding the resulting identities we obtain the desired result. \square

2.1. For s -convex functions.

Theorem 4. *Let $f : I \subset [0, \infty) \rightarrow \mathbb{R}$ be a differentiable function on I° such that $f' \in L[a, b]$, where $a, b \in I^\circ$ with $a < b$. If $|f'|^q$ is s -convex on $[a, b]$ for some fixed $q \geq 1$, $x \in [a, b]$, $\lambda \in [0, 1]$ then the following inequality for fractional integrals holds*

$$(2.4) \quad |S_f(x, \lambda, \alpha, a, b)| \leq A_1^{1-\frac{1}{q}}(\alpha, \lambda) \left\{ \frac{(x-a)^{\alpha+1}}{b-a} (|f'(x)|^q A_2(\alpha, \lambda) + |f'(a)|^q A_3(\alpha, \lambda))^{\frac{1}{q}} + \frac{(b-x)^{\alpha+1}}{b-a} (|f'(x)|^q A_2(\alpha, \lambda) + |f'(b)|^q A_3(\alpha, \lambda))^{\frac{1}{q}} \right\}$$

where

$$\begin{aligned} A_1(\alpha, \lambda) &= \frac{2\alpha\lambda^{1+\frac{1}{\alpha}} + 1}{\alpha + 1} - \lambda, \\ A_2(\alpha, \lambda, s) &= \frac{2\alpha\lambda^{1+\frac{s+1}{\alpha}} + s + 1}{(s+1)(\alpha+s+1)} - \frac{\lambda}{s+1}, \\ A_3(\alpha, \lambda, s) &= \lambda \left[\frac{1 - 2\left(1 - \lambda^{\frac{1}{\alpha}}\right)^{s+1}}{s+1} \right] + \beta(\alpha+1, s+1) - 2\beta\left(\lambda^{\frac{1}{\alpha}}; \alpha+1, s+1\right), \end{aligned}$$

and β is Euler Beta function defined by

$$\beta(x, y) = \frac{\Gamma(x)\Gamma(y)}{\Gamma(x+y)} = \int_0^1 t^{x-1} (1-t)^{y-1} dt, \quad x, y > 0.$$

Proof. From Lemma 1, property of the modulus and using the power-mean inequality we have

$$(2.5) \quad \begin{aligned} |S_f(x, \lambda, \alpha, a, b)| &\leq \frac{(x-a)^{\alpha+1}}{b-a} \int_0^1 |t^\alpha - \lambda| |f'(tx + (1-t)a)| dt \\ &\quad + \frac{(b-x)^{\alpha+1}}{b-a} \int_0^1 |\lambda - t^\alpha| |f'(tx + (1-t)b)| dt \\ &\leq \frac{(x-a)^{\alpha+1}}{b-a} \left(\int_0^1 |t^\alpha - \lambda| dt \right)^{1-\frac{1}{q}} \left(\int_0^1 |t^\alpha - \lambda| |f'(tx + (1-t)a)|^q dt \right)^{\frac{1}{q}} \\ &\quad + \frac{(b-x)^{\alpha+1}}{b-a} \left(\int_0^1 |t^\alpha - \lambda| dt \right)^{1-\frac{1}{q}} \left(\int_0^1 |t^\alpha - \lambda| |f'(tx + (1-t)b)|^q dt \right)^{\frac{1}{q}} \end{aligned}$$

Since $|f'|^q$ is s -convex on $[a, b]$ we get

$$(2.6) \quad \int_0^1 |t^\alpha - \lambda| |f'(tx + (1-t)a)|^q dt \leq \int_0^1 |t^\alpha - \lambda| (t^s |f'(x)|^q + (1-t)^s |f'(a)|^q) dt \\ = |f'(x)|^q A_2(\alpha, \lambda, s) + |f'(a)|^q A_3(\alpha, \lambda, s),$$

$$(2.7) \quad \int_0^1 |t^\alpha - \lambda| |f'(tx + (1-t)b)|^q dt \leq \int_0^1 |t^\alpha - \lambda| (t^s |f'(x)|^q + (1-t)^s |f'(b)|^q) dt \\ = |f'(x)|^q A_2(\alpha, \lambda, s) + |f'(b)|^q A_3(\alpha, \lambda, s),$$

where we use the fact that

$$\int_0^1 |t^\alpha - \lambda| (1-t)^s dt = \int_0^{\lambda^{\frac{1}{\alpha}}} (\lambda - t^\alpha) (1-t)^s dt + \int_{\lambda^{\frac{1}{\alpha}}}^1 (t^\alpha - \lambda) (1-t)^s dt \\ = \lambda \int_0^{\lambda^{\frac{1}{\alpha}}} (1-t)^s dt - \int_0^{\lambda^{\frac{1}{\alpha}}} t^\alpha (1-t)^s dt + \int_{\lambda^{\frac{1}{\alpha}}}^1 t^\alpha (1-t)^s dt \\ - \lambda \int_{\lambda^{\frac{1}{\alpha}}}^1 (1-t)^s dt \\ = \lambda \left[\frac{1 - 2 \left(1 - \lambda^{\frac{1}{\alpha}}\right)^{s+1}}{s+1} \right] + \int_0^1 t^\alpha (1-t)^s dt - 2 \int_0^{\lambda^{\frac{1}{\alpha}}} t^\alpha (1-t)^s dt \\ = \lambda \left[\frac{1 - 2 \left(1 - \lambda^{\frac{1}{\alpha}}\right)^{s+1}}{s+1} \right] + \beta(\alpha + 1, s + 1) - 2\beta\left(\lambda^{\frac{1}{\alpha}}; \alpha + 1, s + 1\right),$$

$$\int_0^1 |t^\alpha - \lambda| t^s dt = \frac{2\alpha\lambda^{1+\frac{s+1}{\alpha}} + s + 1}{(s+1)(\alpha + s + 1)} - \frac{\lambda}{s+1},$$

and by simple computation

$$(2.8) \quad \int_0^1 |t^\alpha - \lambda| dt = \int_0^{\lambda^{\frac{1}{\alpha}}} (\lambda - t^\alpha) dt + \int_{\lambda^{\frac{1}{\alpha}}}^1 (t^\alpha - \lambda) dt \\ = \frac{2\alpha\lambda^{1+\frac{1}{\alpha}} + 1}{\alpha + 1} - \lambda.$$

Hence, If we use (2.6), (2.7) and (2.8) in (2.5), we obtain the desired result. This completes the proof. \square

Corollary 1. *Under the assumptions of Theorem 4 with $q = 1$, the inequality (2.4) reduced to the following inequality*

$$|S_f(x, \lambda, \alpha, a, b)| \leq \left\{ \frac{(x-a)^{\alpha+1}}{b-a} (|f'(x)| A_2(\alpha, \lambda, s) + |f'(a)| A_3(\alpha, \lambda, s)) + \frac{(b-x)^{\alpha+1}}{b-a} (|f'(x)| A_2(\alpha, \lambda, s) + |f'(b)| A_3(\alpha, \lambda, s)) \right\}.$$

Corollary 2. *Under the assumptions of Theorem 4 with $x = \frac{a+b}{2}$, $\lambda = \frac{1}{3}$, from the inequality (2.4) we get the following Simpson type inequality for fractional integrals*

$$\begin{aligned} & \left| \frac{2^{\alpha-1}}{(b-a)^{\alpha-1}} S_f \left(\frac{a+b}{2}, \frac{1}{3}, \alpha, a, b \right) \right| \\ &= \left| \frac{1}{6} \left[f(a) + 4f \left(\frac{a+b}{2} \right) + f(b) \right] - \frac{\Gamma(\alpha+1) 2^{\alpha-1}}{(b-a)^\alpha} \left[J_{\left(\frac{a+b}{2}\right)^-}^\alpha f(a) + J_{\left(\frac{a+b}{2}\right)^+}^\alpha f(b) \right] \right| \\ &\leq \frac{b-a}{4} A_1^{1-\frac{1}{q}} \left(\alpha, \frac{1}{3} \right) \left\{ \left(\left| f' \left(\frac{a+b}{2} \right) \right|^q A_2 \left(\alpha, \frac{1}{3}, s \right) + |f'(a)|^q A_3 \left(\alpha, \frac{1}{3}, s \right) \right)^{\frac{1}{q}} \right. \\ &\quad \left. + \left(\left| f' \left(\frac{a+b}{2} \right) \right|^q A_2 \left(\alpha, \frac{1}{3}, s \right) + |f'(b)|^q A_3 \left(\alpha, \frac{1}{3}, s \right) \right)^{\frac{1}{q}} \right\}. \end{aligned}$$

Corollary 3. *Under the assumptions of Theorem 4 with $x = \frac{a+b}{2}$, $\lambda = 0$, from the inequality (2.4) we get the following midpoint type inequality for fractional integrals*

$$\begin{aligned} & \left| \frac{2^{\alpha-1}}{(b-a)^{\alpha-1}} S_f \left(\frac{a+b}{2}, 0, \alpha, a, b \right) \right| \\ &= \left| f \left(\frac{a+b}{2} \right) - \frac{\Gamma(\alpha+1) 2^{\alpha-1}}{(b-a)^\alpha} \left[J_{\left(\frac{a+b}{2}\right)^-}^\alpha f(a) + J_{\left(\frac{a+b}{2}\right)^+}^\alpha f(b) \right] \right| \\ &\leq \frac{b-a}{4} \left(\frac{1}{\alpha+1} \right)^{1-\frac{1}{q}} \left\{ \left[\frac{|f' \left(\frac{a+b}{2} \right)|^q}{\alpha+s+1} + |f'(a)|^q \beta(\alpha+1, s+1) \right]^{\frac{1}{q}} \right. \\ &\quad \left. + \left[\frac{|f' \left(\frac{a+b}{2} \right)|^q}{\alpha+s+1} + |f'(b)|^q \beta(\alpha+1, s+1) \right]^{\frac{1}{q}} \right\}. \end{aligned}$$

Corollary 4. *Under the assumptions of Theorem 4 with $\lambda = 1$, from the inequality (2.4) we get the following trapezoid type inequality for fractional integrals*

$$\begin{aligned} & \left| S_f \left(\frac{a+b}{2}, 1, \alpha, a, b \right) \right| \\ &= \left| \frac{(x-a)^\alpha f(a) + (b-x)^\alpha f(b)}{b-a} - \frac{\Gamma(\alpha+1)}{b-a} [J_{x^-}^\alpha f(a) + J_{x^+}^\alpha f(b)] \right| \\ &\leq \left(\frac{\alpha}{\alpha+1} \right)^{1-\frac{1}{q}} \left\{ \frac{(x-a)^{\alpha+1}}{b-a} \left[\frac{\alpha |f'(x)|^q}{(s+1)(\alpha+s+1)} + |f'(a)|^q \left(\frac{1}{s+1} - \beta(\alpha+1, s+1) \right) \right]^{\frac{1}{q}} \right. \\ &\quad \left. + \frac{(b-x)^{\alpha+1}}{b-a} \left[\frac{\alpha |f'(x)|^q}{(s+1)(\alpha+s+1)} + |f'(b)|^q \left(\frac{1}{s+1} - \beta(\alpha+1, s+1) \right) \right]^{\frac{1}{q}} \right\} \end{aligned}$$

which is the same of the inequality in [8, Theorem 9].

Remark 1. In Corollary 4, if we choose $\alpha = 1$, we get the same inequality in [2, Theorem 7].

Corollary 5. Let the assumptions of Theorem 4 hold. If $|f'(x)| \leq M$ for all $x \in [a, b]$ and $\lambda = 0$, then from the inequality (2.4) we get the following Ostrowski type inequality for fractional integrals

$$\begin{aligned} & \left| \left[\frac{(x-a)^\alpha + (b-x)^\alpha}{b-a} \right] f(x) - \frac{\Gamma(\alpha+1)}{b-a} [J_{x^-}^\alpha f(a) + J_{x^+}^\alpha f(b)] \right| \\ & \leq M \left(\frac{1}{\alpha+1} \right)^{1-\frac{1}{q}} \left[\frac{1}{\alpha+s+1} + \beta(\alpha+1, s+1) \right]^{\frac{1}{q}} \left[\frac{(x-a)^{\alpha+1} + (b-x)^{\alpha+1}}{b-a} \right] \end{aligned}$$

for each $x \in [a, b]$.

Remark 2. In Corollary 5, if we choose $\alpha = 1$, we get the same inequality in [1, Theorem 4].

2.2. For quasi-convex functions.

Theorem 5. Let $f : I \subset [0, \infty) \rightarrow \mathbb{R}$ be a differentiable function on I° such that $f' \in L[a, b]$, where $a, b \in I^\circ$ with $a < b$. If $|f'|^q$ is quasi-convex on $[a, b]$ for some fixed $q \geq 1$, $x \in [a, b]$, $\lambda \in [0, 1]$ then the following inequality for fractional integrals holds

$$(2.9) |S_f(x, \lambda, \alpha, a, b)| \leq A_1(\alpha, \lambda) \left\{ \frac{(x-a)^{\alpha+1}}{b-a} (\sup \{|f'(x)|^q, |f'(a)|^q\})^{\frac{1}{q}} + \frac{(b-x)^{\alpha+1}}{b-a} (\sup \{|f'(x)|^q, |f'(b)|^q\})^{\frac{1}{q}} \right\}$$

where

$$A_1(\alpha, \lambda) = \frac{2\alpha\lambda^{1+\frac{1}{\alpha}} + 1}{\alpha+1} - \lambda.$$

Proof. We proceed similarly as in the proof Theorem 4. Since $|f'|^q$ is quasi-convex on $[a, b]$, for all $t \in [0, 1]$

$$|f'(tx + (1-t)a)|^q \leq \sup \{|f'(x)|^q, |f'(a)|^q\}$$

and

$$|f'(tx + (1-t)b)|^q \leq \sup \{|f'(x)|^q, |f'(b)|^q\}.$$

Hence, from the inequality (2.5) we get

$$\begin{aligned}
& |S_f(x, \lambda, \alpha, a, b)| \\
& \leq \left(\int_0^1 |t^\alpha - \lambda| dt \right) \left\{ \frac{(x-a)^{\alpha+1}}{b-a} (\sup \{|f'(x)|^q, |f'(a)|^q\})^{\frac{1}{q}} \right. \\
& \quad \left. + \frac{(b-x)^{\alpha+1}}{b-a} (\sup \{|f'(x)|^q, |f'(b)|^q\})^{\frac{1}{q}} \right\} \\
& \leq A_1(\alpha, \lambda) \left\{ \frac{(x-a)^{\alpha+1}}{b-a} (\sup \{|f'(x)|^q, |f'(a)|^q\})^{\frac{1}{q}} \right. \\
& \quad \left. + \frac{(b-x)^{\alpha+1}}{b-a} (\sup \{|f'(x)|^q, |f'(b)|^q\})^{\frac{1}{q}} \right\}
\end{aligned}$$

which completes the proof. \square

Corollary 6. *Under the assumptions of Theorem 5 with $q = 1$, the inequality (2.9) reduced to the following inequality*

$$\begin{aligned}
|S_f(x, \lambda, \alpha, a, b)| & \leq A_1(\alpha, \lambda) \left\{ \frac{(x-a)^{\alpha+1}}{b-a} (\sup \{|f'(x)|, |f'(a)|\}) \right. \\
& \quad \left. + \frac{(b-x)^{\alpha+1}}{b-a} (\sup \{|f'(x)|, |f'(b)|\}) \right\}.
\end{aligned}$$

Corollary 7. *Under the assumptions of Theorem 5 with $x = \frac{a+b}{2}$, $\lambda = \frac{1}{3}$, from the inequality (2.9) we get the following Simpson type inequality or fractional integrals*

$$\begin{aligned}
& \left| \frac{1}{6} \left[f(a) + 4f\left(\frac{a+b}{2}\right) + f(b) \right] - \frac{\Gamma(\alpha+1)2^{\alpha-1}}{(b-a)^\alpha} \left[J_{\left(\frac{a+b}{2}\right)^-}^\alpha f(a) + J_{\left(\frac{a+b}{2}\right)^+}^\alpha f(b) \right] \right| \\
& \leq \frac{b-a}{4} A_1\left(\alpha, \frac{1}{3}\right) \left[\sup \left\{ \left| f'\left(\frac{a+b}{2}\right) \right|, |f'(a)| \right\} + \sup \left\{ \left| f'\left(\frac{a+b}{2}\right) \right|, |f'(b)| \right\} \right].
\end{aligned}$$

Corollary 8. *Under the assumptions of Theorem 5 with $x = \frac{a+b}{2}$, $\lambda = 0$, from the inequality (2.9) we get the following midpoint type inequality or fractional integrals*

$$\begin{aligned}
& \left| f\left(\frac{a+b}{2}\right) - \frac{\Gamma(\alpha+1)2^{\alpha-1}}{(b-a)^\alpha} \left[J_{\left(\frac{a+b}{2}\right)^-}^\alpha f(a) + J_{\left(\frac{a+b}{2}\right)^+}^\alpha f(b) \right] \right| \\
& \leq \frac{b-a}{4} \left(\frac{1}{\alpha+1}\right) \left\{ \left[\sup \left\{ \left| f'\left(\frac{a+b}{2}\right) \right|^q, |f'(a)|^q \right\} \right]^{\frac{1}{q}} \right. \\
& \quad \left. + \left[\sup \left\{ \left| f'\left(\frac{a+b}{2}\right) \right|^q, |f'(b)|^q \right\} \right]^{\frac{1}{q}} \right\}.
\end{aligned}$$

Corollary 9. *Under the assumptions of Theorem 5 with $\lambda = 1$, from the inequality (2.9) we get the following trapezoid type inequality or fractional integrals*

$$\begin{aligned} & \left| \frac{(x-a)^\alpha f(a) + (b-x)^\alpha f(b)}{b-a} - \frac{\Gamma(\alpha+1)}{b-a} [J_{x^-}^\alpha f(a) + J_{x^+}^\alpha f(b)] \right| \\ & \leq \left(\frac{\alpha}{\alpha+1} \right) \left\{ \frac{(x-a)^{\alpha+1}}{b-a} [\sup \{|f'(x)|^q, |f'(a)|^q\}]^{\frac{1}{q}} \right. \\ & \quad \left. + \frac{(b-x)^{\alpha+1}}{b-a} [\sup \{|f'(x)|^q, |f'(b)|^q\}]^{\frac{1}{q}} \right\}. \end{aligned}$$

Corollary 10. *Let the assumptions of Theorem 5 hold. If $|f'(x)| \leq M$ for all $x \in [a, b]$ and $\lambda = 0$, then from the inequality (2.9) we get the following Ostrowski type inequality or fractional integrals*

$$\begin{aligned} & \left| \left[\frac{(x-a)^\alpha + (b-x)^\alpha}{b-a} \right] f(x) - \frac{\Gamma(\alpha+1)}{b-a} [J_{x^-}^\alpha f(a) + J_{x^+}^\alpha f(b)] \right| \\ & \leq \frac{M}{\alpha+1} \left[\frac{(x-a)^{\alpha+1} + (b-x)^{\alpha+1}}{b-a} \right], \end{aligned}$$

2.3. For m -convex functions. Similarly lemma 1, we can proved the following lemma

Lemma 2. *Let $f : I \subset \mathbb{R} \rightarrow \mathbb{R}$ be a differentiable mapping on I° such that $f' \in L[ma, mb]$, where $m \in (0, 1]$, $ma, mb \in I$ with $a < b$. Then for all $x \in [a, b]$, $\lambda \in [0, 1]$ and $\alpha > 0$ we have:*

$$\begin{aligned} S_f(mx, \lambda, \alpha, ma, mb) &= \frac{m^\alpha (x-a)^{\alpha+1}}{b-a} \int_0^1 (t^\alpha - \lambda) f'(tmx + m(1-t)a) dt \\ &+ \frac{m^\alpha (b-x)^{\alpha+1}}{b-a} \int_0^1 (\lambda - t^\alpha) f'(tmx + m(1-t)b) dt. \end{aligned}$$

Theorem 6. *Let $f : I \subset [0, \infty) \rightarrow \mathbb{R}$ be a differentiable function on I° such that $f' \in L[ma, mb]$, where $m \in (0, 1]$, $a, b \in I^\circ$ with $a < b$. If $|f'|^q$ is m -convex on $[ma, mb]$ for some fixed $q \geq 1$, $x \in [a, b]$, $\lambda \in [0, 1]$ and $\alpha > 0$ then the following inequality for fractional integrals holds*

$$\begin{aligned} (2.10) \quad & |S_f(mx, \lambda, \alpha, ma, mb)| \\ & \leq A_1(\alpha, \lambda)^{1-\frac{1}{q}} \left\{ \frac{m^\alpha (x-a)^{\alpha+1}}{b-a} (|f'(mx)|^q A_2(\alpha, \lambda) + m |f'(a)|^q A_3(\alpha, \lambda))^{\frac{1}{q}} \right. \\ & \quad \left. + \frac{m^\alpha (b-x)^{\alpha+1}}{b-a} (|f'(mx)|^q A_2(\alpha, \lambda) + m |f'(b)|^q A_3(\alpha, \lambda))^{\frac{1}{q}} \right\} \end{aligned}$$

where

$$\begin{aligned} A_1(\alpha, \lambda) &= \frac{2\alpha\lambda^{1+\frac{1}{\alpha}} + 1}{\alpha + 1} - \lambda, \\ A_2(\alpha, \lambda) &= \frac{\alpha\lambda^{1+\frac{2}{\alpha}} + 1}{\alpha + 2} - \frac{\lambda}{2}, \\ A_3(\alpha, \lambda) &= \frac{2\alpha\lambda^{1+\frac{1}{\alpha}} + 1}{\alpha + 1} - \frac{\alpha\lambda^{1+\frac{2}{\alpha}} + 1}{\alpha + 2} - \frac{\lambda}{2}. \end{aligned}$$

Proof. We proceed similarly as in the proof Theorem 4. From Lemma 2, property of the modulus and using the power-mean inequality we have

$$\begin{aligned} & |S_f(mx, \lambda, \alpha, ma, mb)| \\ & \leq \frac{m^\alpha (x-a)^{\alpha+1}}{b-a} \left(\int_0^1 |t^\alpha - \lambda| dt \right)^{1-\frac{1}{q}} \left(\int_0^1 |t^\alpha - \lambda| |f'(tmx + m(1-t)a)|^q dt \right)^{\frac{1}{q}} \\ & \quad (2.11) \quad \frac{m^\alpha (b-x)^{\alpha+1}}{b-a} \left(\int_0^1 |t^\alpha - \lambda| dt \right)^{1-\frac{1}{q}} \left(\int_0^1 |t^\alpha - \lambda| |f'(tmx + m(1-t)b)|^q dt \right)^{\frac{1}{q}}. \end{aligned}$$

Since $|f'|^q$ is m -convex on $[ma, mb]$, for all $t \in [0, 1]$

$$|f'(tmx + m(1-t)a)|^q \leq t|f'(mx)|^q + m(1-t)|f'(a)|^q$$

and

$$|f'(tmx + m(1-t)b)|^q \leq t|f'(mx)|^q + m(1-t)|f'(b)|^q.$$

Hence by simple computation we get

$$\begin{aligned} \int_0^1 |t^\alpha - \lambda| |f'(tmx + m(1-t)a)|^q dt & \leq \int_0^1 |t^\alpha - \lambda| t |f'(mx)|^q + m(1-t) |f'(a)|^q dt \\ & = |f'(mx)|^q \left(\frac{\alpha\lambda^{1+\frac{2}{\alpha}} + 1}{\alpha + 2} - \frac{\lambda}{2} \right) \\ (2.12) \quad & + m |f'(a)|^q \left(\frac{2\alpha\lambda^{1+\frac{1}{\alpha}} + 1}{\alpha + 1} - \frac{\alpha\lambda^{1+\frac{2}{\alpha}} + 1}{\alpha + 2} - \frac{\lambda}{2} \right), \end{aligned}$$

and similarly

$$\begin{aligned} \int_0^1 |t^\alpha - \lambda| |f'(tmx + m(1-t)b)|^q dt & \leq \int_0^1 |t^\alpha - \lambda| t |f'(mx)|^q + m(1-t) |f'(b)|^q dt \\ & = |f'(mx)|^q \left(\frac{\alpha\lambda^{1+\frac{2}{\alpha}} + 1}{\alpha + 2} - \frac{\lambda}{2} \right) \\ (2.13) \quad & + m |f'(b)|^q \left(\frac{2\alpha\lambda^{1+\frac{1}{\alpha}} + 1}{\alpha + 1} - \frac{\alpha\lambda^{1+\frac{2}{\alpha}} + 1}{\alpha + 2} - \frac{\lambda}{2} \right). \end{aligned}$$

If we use (2.12), (2.13) and (2.8) in (2.11), we obtain the desired result. This completes the proof. \square

Corollary 11. *Under the assumptions of Theorem 6 with $q = 1$, the inequality (2.10) reduced to the following inequality*

$$|S_f(mx, \lambda, \alpha, ma, mb)| \leq \left\{ \frac{m^\alpha (x-a)^{\alpha+1}}{b-a} (|f'(mx)| A_2(\alpha, \lambda) + m |f'(a)| A_3(\alpha, \lambda)) + \frac{m^\alpha (b-x)^{\alpha+1}}{b-a} (|f'(mx)| A_2(\alpha, \lambda) + m |f'(b)| A_3(\alpha, \lambda)) \right\}.$$

Corollary 12. *Under the assumptions of Theorem 6 with $x = \frac{a+b}{2}$, $\lambda = \frac{1}{3}$, from the inequality (2.10) we get the following Simpson type inequality or fractional integrals*

$$\begin{aligned} & \left| \frac{2^{\alpha-1}}{m^\alpha (b-a)^{\alpha-1}} S_f \left(m \left(\frac{a+b}{2} \right), \frac{1}{3}, \alpha, ma, mb \right) \right| \\ &= \left| \frac{1}{6} \left[f(ma) + 4f \left(\frac{m(a+b)}{2} \right) + f(mb) \right] - \frac{\Gamma(\alpha+1) 2^{\alpha-1}}{m^\alpha (b-a)^\alpha} \left[J_{\left(\frac{m(a+b)}{2} \right)^-}^\alpha f(ma) + J_{\left(\frac{m(a+b)}{2} \right)^+}^\alpha f(mb) \right] \right| \\ &\leq \frac{m(b-a)}{4} A_1^{1-\frac{1}{q}} \left(\alpha, \frac{1}{3} \right) \left\{ \left(\left| f' \left(\frac{m(a+b)}{2} \right) \right|^q A_2 \left(\alpha, \frac{1}{3} \right) + m |f'(a)|^q A_3 \left(\alpha, \frac{1}{3} \right) \right)^{\frac{1}{q}} \right. \\ &\quad \left. + \left(\left| f' \left(\frac{m(a+b)}{2} \right) \right|^q A_2 \left(\alpha, \frac{1}{3} \right) + m |f'(b)|^q A_3 \left(\alpha, \frac{1}{3} \right) \right)^{\frac{1}{q}} \right\}. \end{aligned}$$

Corollary 13. *Under the assumptions of Theorem 6 with $x = \frac{a+b}{2}$, $\lambda = 0$, from the inequality (2.10) we get the following midpoint type inequality or fractional integrals*

$$\begin{aligned} & \frac{2^{\alpha-1}}{m^\alpha (b-a)^{\alpha-1}} \left| S_f \left(m \left(\frac{a+b}{2} \right), 0, \alpha, ma, mb \right) \right| \\ &= \left| f \left(\frac{m(a+b)}{2} \right) - \frac{\Gamma(\alpha+1) 2^{\alpha-1}}{m^\alpha (b-a)^\alpha} \left[J_{\left(\frac{m(a+b)}{2} \right)^-}^\alpha f(ma) + J_{\left(\frac{m(a+b)}{2} \right)^+}^\alpha f(mb) \right] \right| \\ &\leq \frac{m(b-a)}{4} \left(\frac{1}{\alpha+1} \right) \left(\frac{1}{\alpha+2} \right)^{\frac{1}{q}} \left\{ \left[(\alpha+1) \left| f' \left(\frac{m(a+b)}{2} \right) \right|^q + m |f'(a)|^q \right]^{\frac{1}{q}} \right. \\ &\quad \left. + \left[(\alpha+1) \left| f' \left(\frac{m(a+b)}{2} \right) \right|^q + m |f'(b)|^q \right]^{\frac{1}{q}} \right\}. \end{aligned}$$

Corollary 14. *Under the assumptions of Theorem 6 with $\lambda = 1$, from the inequality (2.10) we get the following trapezoid type inequality or fractional integrals*

$$\begin{aligned} & |S_f(mx, 1, \alpha, ma, mb)| \\ &= \left| \frac{(x-a)^\alpha f(ma) + (b-x)^\alpha f(mb)}{b-a} - \frac{\Gamma(\alpha+1)}{m^{\alpha+1} (b-a)} [J_{m^-}^\alpha f(ma) + J_{m^+}^\alpha f(mb)] \right| \\ &\leq \left(\frac{\alpha}{\alpha+1} \right) \left(\frac{1}{2(\alpha+2)} \right)^{\frac{1}{q}} \left\{ \frac{(x-a)^{\alpha+1}}{b-a} [(\alpha+1) |f'(mx)|^q + (\alpha+3) m |f'(a)|^q]^{\frac{1}{q}} \right. \\ &\quad \left. + \frac{(b-x)^{\alpha+1}}{b-a} [(\alpha+1) |f'(mx)|^q + (\alpha+3) m |f'(b)|^q]^{\frac{1}{q}} \right\}. \end{aligned}$$

Corollary 15. *Let the assumptions of Theorem 6 hold. If $|f'(u)| \leq M$ for all $u \in [ma, mb]$ and $\lambda = 0$, then from the inequality (2.10) we get the following*

Ostrowski type inequality or fractional integrals

$$\left| \left[\frac{(x-a)^\alpha + (b-x)^\alpha}{b-a} \right] f(mx) - \frac{\Gamma(\alpha+1)}{m^{\alpha+1}(b-a)} [J_{mx^-}^\alpha f(ma) + J_{mx^+}^\alpha f(mb)] \right|$$

$$\leq mM \left(\frac{1}{\alpha+1} \right) \left(\frac{\alpha+m+1}{\alpha+2} \right)^{\frac{1}{q}} \left[\frac{(x-a)^{\alpha+1} + (b-x)^{\alpha+1}}{b-a} \right],$$

for each $x \in [a, b]$.

REFERENCES

- [1] M. Alomaria, M. Darus, S.S. Dragomir, P. Cerone, Ostrowski type inequalities for functions whose derivatives are s -convex in the second sense, Applied Mathematics Letters 23 (2010) 1071–1076.
- [2] M. Avci, H. Kavurmaci and M.E. Ozdemir, New inequalities of Hermite-Hadamard type via s -convex functions in the second sense with applications, Appl. Math. Comput., 217 (2011) 5171–5176.
- [3] Z. Dahmani, On Minkowski and Hermite-Hadamard integral inequalities via fractional via fractional integration, Ann. Funct. Anal. 1 (1) (2010), 51-58
- [4] R. Gorenflo, F. Mainardi, Fractional calculus; integral and differential equations of fractional order, Springer Verlag, Wien (1997), 223-276.
- [5] H. Hudzik and L. Maligranda, Some remarks on s -convex functions, Aequationes Math. 48 (1994), 100-111.
- [6] D.A. Ion, Some estimates on the Hermite-Hadamard inequality through quasi-convex functions, Annals of University of Craiova, Math. Comp. Sci. Ser. 34 (2007) 82-87.
- [7] S. Miller and B. Ross, An introduction to the Fractional Calculus and Fractional Differential Equations, John Wiley & Sons, USA (1993), 2.
- [8] M.E. Ozdemir, M. Avci, H. Kavurmaci, Hermite-Hadamard type inequalities for s -convex and s -concave functions via fractional integrals, arXiv:1202.0380v1.
- [9] I. Podlubni, Fractional Differential Equations, Academic Press, San Diego, 1999.
- [10] J. Pečarić, F. Proschan, and Y.L. Tong, Convex Functions, Partial Orderings and Statistical Applications, Academic Press Inc. 1992.
- [11] E. Set, New inequalities of Ostrowski type for mapping whose derivatives are s -convex in the second sense via fractional integrals, Computers and Math. with Appl. 63 (2012) 1147-1154.
- [12] M.Z. Sarıkaya, E. Set, H. Yıldız and N. Başak, Hermite-Hadamard's inequalities for fractional integrals and related fractional inequalities, Mathematical and Computer Modelling, DOI:10.1016/j.mcm.2011.12.048.
- [13] G. Toader, Some generalizations of the convexity, Proceedings of The Colloquium On Approximation And Optimization, Univ. Cluj-Napoca, Cluj-Napoca, 1985, 329-338.

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCES AND ARTS, GİRESUN UNIVERSITY, GİRESUN, TURKEY

E-mail address: imdat.iscan@giresun.edu.tr