### OPTIMUM BOUNDS FOR THE DISTRIBUTIONS OF MARTINGALES IN BANACH SPACES<sup>1</sup>

BY IOSIF PINELIS

Michigan Technological University

A general device is proposed, which provides for extension of exponential inequalities for sums of independent real-valued random variables to those for martingales in the 2-smooth Banach spaces. This is used to obtain optimum bounds of the Rosenthal-Burkholder and Chung types on moments of the martingales in the 2-smooth Banach spaces. In turn, it leads to best-order bounds on moments of the sums of independent random vectors in any separable Banach spaces. Although the emphasis is put on the infinite-dimensional martingales, most of the results seem to be new even for the onedimensional ones. Moreover, the bounds on the Rosenthal-Burkholder type of moments seem to be to certain extent new even for the sums of independent real-valued random variables. Analogous inequalities for (one-dimensional) supermartingales are given.

**1. Introduction.** For a separable Banach space  $(\mathcal{X}, \|\cdot\|)$ , let  $\mathcal{S}(\mathcal{X})$  denote the class of all sequences  $f = (f_j) = (f_0, f_1, ...)$  of Bochner-integrable random vectors in  $\mathcal{X}$ , with  $f_0 \equiv 0$ , defined on a probability space  $(\Omega, F, P)$  and adapted to a non-decreasing sequence  $(F_j) = (F_0, F_1, ...)$  of sub- $\sigma$ -fields of F; here,  $n, (\Omega, F, P)$  and  $(F_j)$  are considered attributes of f and may be different for different  $f \in \mathcal{S}(\mathcal{X})$ .

For  $f \in \mathcal{S}(\mathcal{X})$ , put  $f^* = \sup\{||f_j||: j = 0, 1, ...\}, d_0 = d_0(f) \equiv 0, d_j = d_j(f) = f_j - f_{j-1}, j = 1, 2, ..., S_p = S_p(f) = \left(\sum_{j=1}^{\infty} ||d_j||^p\right)^{1/p}, p > 0, s_2 = s_2(f)$ 

<sup>&</sup>lt;sup>1</sup>Research partially supported by NSF grant DMS-8921369.

AMS 1980 subject classifications. 60B12, 60E15, 60G42, 60G50, 60F10.

Key words and phrases. Distribution inequalities, exponential inequalities, bounds on moments, martingales in Banach spaces, 2-smooth Banach spaces, sums of independent random variables.

=  $\left(\sum_{j=1}^{\infty} \mathbf{E}_{j-1} \|d_j\|^2\right)^{1/2}$ , where  $\mathbf{E}_{j-1}$  stands for the conditional expectation given  $F_{j-1}$ .

Let  $\mathcal{M}(\mathcal{X})$  denote the class of all sequences  $(f_j) \in \mathcal{S}(\mathcal{X})$  that are martingales and  $\mathcal{M}_{ind}(\mathcal{X})$ , the class of all sequences  $(f_j) \in \mathcal{S}(\mathcal{X})$  having independent increments  $d_j$ 's.

For any two nonnegative expressions  $\mathcal{E}_1$  and  $\mathcal{E}_2$ , let us write  $\mathcal{E}_1 \leq \mathcal{E}_2$  (or, equivalently,  $\mathcal{E}_2 \succeq \mathcal{E}_1$ ) if  $\mathcal{E}_1 \leq A\mathcal{E}_2$ , and  $\mathcal{E}_1 \asymp \mathcal{E}_2$  if  $\mathcal{E}_1 \leq \mathcal{E}_2 \leq \mathcal{E}_1$ . Here, A denotes a positive absolute constant.

We assume that  $\inf \emptyset = \infty$ ,  $\sup \emptyset = 0$ ,  $\sum_{j \in \emptyset} u_j = 0$ ,  $\prod_{j \in \emptyset} u_j = 1$ .

Symbol  $\blacksquare$  indicates the end of a proof or a remark.

In section 2, some preliminary results on the 2-smooth Banach spaces and on the martingales in such spaces are given.

In section 3, a device is suggested, which provides for the extension of exponential inequalities for the sums of independent real-valued random variables to those for the martingales in the 2-smooth Banach spaces. In particular, by that means an exponential inequality for the martingales in the 2-smooth spaces optimal in terms of  $||d^*||_{\infty}$ ,  $||s_2||_{\infty}$  is obtained, which is a generalization of an inequality of Bennett (1962) and Hoeffding (1963).

In section 4, using methods of Burkholder (1973) and results of section 3, we obtain optimal (to the above defined relation  $\approx$ ) upper bounds of the Rosenthal (1970)-Burkholder (1973) type on moments of the martingales in the 2-smooth Banach spaces, i.e., optimal in terms of  $||d^*||_p$ ,  $||s_2||_p$ , for  $p \ge 2$ .

In section 5, via a modification of the martingale decomposition method by Yurinskii (1974), we apply the inequalities of section 4 to obtain bounds of the Rosenthal-Burkholder type on  $|| ||f_n|| - \mathbf{E} ||f_n|| ||_p$ ,  $p \ge 2$ , for an arbitrary separable Banach space  $\mathcal{X}$ , but only for  $f \in \mathcal{M}_{ind}(\mathcal{X})$ .

In section 6, we show that the inequalities derived in sections 4, 5 are optimal

in the terms used. The key roles in the proof of the optimality are played by the results and methods of Pinelis and Utev (1984).

In section 7, we obtain bounds on  $|| || f_n || ||_p$ ,  $p \ge 2$ ,  $f \in \mathcal{M}(\mathcal{X})$ , which are optimal in terms of n and  $|| S_p ||_p$ . We refer to them as to bounds of Chung type.

In section 8, inequalities for super-martingales (of course, in  $\mathcal{X} = \mathbf{R}$ ) similar to those in sections 3, 4 and certain refinements for real-valued martingales are presented.

In section 9, some details are given. For instance, it is shown that, at least for the sums of independent random vectors, the bounds in Pinelis (1980) are equivalent to those in section 4.

Substantial part of the results was announced in Pinelis (1992).

I am grateful to Donald Burkholder and Walter Philipp for their support and stimulating conversations and to Pawel Hitczenko, who has acquainted me with his latest results here referred to. I would like to thank the organizers of the work-shop "Linear Analysis and Probability" in the Department of Mathematics, Texas A&M University for the great opportunity to discuss the results of this paper.

2. Preliminaries: 2-smooth Banach spaces and a reduction of martingales. Let us call a Banach space  $(\mathcal{X}, \|\cdot\|)$  2-smooth [or, more exactly, (2, D)-smooth, for some  $D = D(\mathcal{X}) > 0$ ] if  $\forall x \in \mathcal{X} \ \forall y \in \mathcal{X}$ 

(2.1) 
$$\|x+y\|^2 + \|x-y\|^2 \le 2\|x\|^2 + 2D^2\|y\|^2$$

The importance of the 2-smooth spaces was elucidated in the paper of Pisier (1975): they play the same role with respect to the vector martingales as the spaces of type 2 do with respect to the sums of independent random vectors.

The definition assumed in this paper is slightly different from that given by Pisier [which required only that (2.1) hold for an equivalent norm], because, in the subsequent account, we would like to follow the dependence of certain constants on D, the constant of the 2-smoothness.

Substituting  $\lambda x$  for y in (2.1), where  $\lambda \in \mathbf{R}$ , one observes that except for  $\mathcal{X} = \{0\}, D \geq 1.$ 

It is easily seen that the condition

(2.2) 
$$(\|x\|^2)''(v,v) \le 2D^2 \|v\|^2 \quad \forall x \in \mathcal{X} \ \forall v \in \mathcal{X}$$

is sufficient for the (2, D)-smoothness, where  $(||x||^2)''(v, v)$  stands for the second directional derivative of the function  $x \mapsto ||x||^2$  in the direction v.

By way of illustration, we give

PROPOSITION 2.1. For any  $p \geq 2$  and any measure space  $(T, \mathcal{A}, \nu)$ ,  $L^p := L^p(T, \mathcal{A}, \nu)$  is  $(2, \sqrt{p-1})$ -smooth.

PROOF. For  $||x|| := (\int_T |x|^p d\nu)^{1/p}$ , one has

(2.3)  
$$\frac{1}{2}(\|x\|^2)''(v,v) = (p-1)\|x\|^{2-p} \int_T |x|^{p-2} v^2 d\nu - (p-2)\|x\|^{2-2p} \left(\int_T |x|^{p-2} xv \, d\nu\right)^2 \leq (p-1)\|v\|^2 - 0$$

if  $x \in \mathcal{X} \setminus \{0\}, v \in \mathcal{X}$  (in view of Hölder's inequality), and  $\frac{1}{2}(||x||^2)''(v,v) = ||v||^2$ if x = 0. Thus, (2.2) is checked.

In particular, what is obvious and well-known, if  $\mathcal{X}$  is a Hilbert space, then it is (2,1)-smooth.

REMARK. If  $L^p = L^p(T, \mathcal{A}, \nu)$  is at least two-dimensional, i.e., if there exist  $T_1 \in \mathcal{A}, T_2 \in \mathcal{A}$  such that  $T_1 \cap T_2 = \emptyset, 0 < \nu(T_1) < \infty, 0 < \nu(T_2) < \infty$ , then  $L^p$  is not (2, D)-smooth if  $0 < D < \sqrt{p-1}$ , so that Proposition 2.1 gives the best bound. Indeed, put  $x \equiv \nu(T_2)^{1/p}$  on  $T_1, x \equiv \nu(T_1)^{1/p}$  on  $T_2, v \equiv x$  on  $T_1, v \equiv -x$  on  $T_2, x \equiv v \equiv 0$  on  $T \setminus T_1 \setminus T_2$ . Then  $x \neq 0$ , and (2.3) turns into an equality.

Condition (2.2) is not only sufficient but also necessary for the (2, D)-smoothness if the derivatives are understood in a generalized sense. To state this remark rigorously, let us give a definition somewhat extending the notion of the 2smoothness.

For any Banach space  $(\mathcal{X}, \|\cdot\|)$ , we call a function  $\Psi: \mathcal{X} \to [0, \infty)$  (2, D)-smooth, D > 0, if it satisfies the conditions:  $\Psi(0) = 0$ ,

$$|\Psi(x+v) - \Psi(x)| \le ||v||,$$
$$\Psi^2(x+v) - 2\Psi^2(x) + \Psi^2(x-v) \le 2D^2 ||v||^2$$

for all  $x \in \mathcal{X}, v \in \mathcal{X}$ .

Evidently, a Banach space  $(\mathcal{X}, \|\cdot\|)$  is (2, D)-smooth if and only if its norm function is (2, D)-smooth.

The results stated in the subsequent sections for the norms of the martingales in the 2-smooth spaces can be extended to those for the (2, D)-smooth functions of martingales in any Banach spaces.

For any (2, D)-smooth function  $\Psi$  on a *finite-dimensional* Banach space  $(\mathcal{X}, \|\cdot\|)$  and  $\varepsilon > 0$ , define

$$\Psi_{\varepsilon}(x) = \sqrt{\int_{\mathcal{X}} \Psi^2(x - \varepsilon y) \gamma(\mathrm{d}y)},$$

where  $\gamma$  is, say, a zero-mean Gaussian measure on  $\mathcal{X}$  with support $(\gamma) = \mathcal{X}$ .

LEMMA 2.2. If  $\Psi$  is a (2, D)-smooth function on a finite-dimensional Banach space

 $(\mathcal{X}, \|\cdot\|)$ , then for all  $\varepsilon > 0$ ,  $\Psi_{\varepsilon}$  has the Fréchet derivatives  $\Psi'_{\varepsilon}(x)$ ,  $\Psi''_{\varepsilon}(x)$ , ... of any order, and the directional derivatives in any direction  $v \in \mathcal{X}$  satisfy the inequalities

(2.4) 
$$|\Psi_{\varepsilon}'(x)(v)| \le ||v||, \qquad (\Psi_{\varepsilon}^2)''(x)(v,v) \le 2D^2 ||v||^2$$

for all  $x \in \mathcal{X}$ . Besides, for each  $x \in \mathcal{X}$ ,  $\Psi_{\varepsilon}(x) \to \Psi(x)$  as  $\varepsilon \downarrow 0$ . [In this generalized sense, sufficient condition (2.2) is also necessary for a Banach space

 $(\mathcal{X}, \|\cdot\|)$  to be (2, D)-smooth; note that (2.1) may be considered locally – for any two-dimensional subspace containing x and y.]

PROOF. Among the statements of the lemma, only the first of the inequalities (2.4) is comparatively non-trivial. Observe that

$$\begin{split} |(\Psi_{\varepsilon}^{2})'(x)(v)| &\leq \limsup_{t \to 0} \frac{1}{t} \int_{\mathcal{X}} |\Psi(x + tv - \varepsilon y) - \Psi(x - \varepsilon y)| \left(\Psi(x + tv - \varepsilon y) + \Psi(x - \varepsilon y)\right) \gamma(\mathrm{d}y) \\ &\leq \|v\| \int_{\mathcal{X}} 2\Psi(x - \varepsilon y) \gamma(\mathrm{d}y). \end{split}$$

One can assume that  $\Psi_{\varepsilon}(x) \neq 0$  for all  $x \in \mathcal{X}$ . Now,

$$\begin{aligned} |(\Psi_{\varepsilon})'(x)(v)| &= \left| \frac{(\Psi_{\varepsilon}^{2})'(x)(v)}{2\Psi_{\varepsilon}(x)} \right| \\ &\leq \frac{\|v\| \int_{\mathcal{X}} 2\Psi(x - \varepsilon y)\gamma(\mathrm{d}y)}{2\sqrt{\int_{\mathcal{X}} \Psi^{2}(x - \varepsilon y)\gamma(\mathrm{d}y)}} \\ &\leq \|v\| \end{aligned}$$

by Schwartz inequality.

We also need the following folklorish lemma.

LEMMA 2.3. Let  $(f_j)_{j=0}^{\infty} \in \mathcal{M}(\mathcal{X})$  be a martingale in a separable Banach space  $(\mathcal{X}, \|\cdot\|)$  relative to a filtration  $(F_j)_{j=0}^{\infty}$ . Then for any  $\varepsilon > 0$ , there exists a martingale  $(f_{j,\varepsilon})_{j=0}^{\infty} \in \mathcal{M}(\mathcal{X})$  relative to a filtration  $(F_{j,\varepsilon})_{j=0}^{\infty}$  such that  $\forall j = 1, 2, \ldots,$ 

- (2.5)  $f_{j,\varepsilon}$  is a random variable having only a finite number of values,
- (2.6)  $f_{j,\varepsilon} \to f_j$  in probability as  $\varepsilon \downarrow 0$ ,
- (2.7)  $\mathbf{E}g(f_{j,\varepsilon}) \leq \mathbf{E}g(f_j),$
- (2.8)  $\|\mathbf{E}(g(f_{j,\varepsilon})|F_{j-1,\varepsilon})\|_{\infty} \leq \mathbf{E}(g(f_j)|F_{j-1})\|_{\infty},$

where g is any nonnegative convex real function on  $\mathcal{X}$ .

PROOF. Consider the approximation  $f_{j,\varepsilon} := \mathbf{E}(f_j|F_{j,\varepsilon})$ , where  $F_{j,\varepsilon}$  is the  $\sigma$ -field generated by all the events of the form  $\{f_i \in B_{k,\varepsilon}\}, i = 0, 1, \ldots, j$ ,

 $k = 1, \ldots, k(j, \varepsilon)$ , where  $(\{B_{k,\varepsilon} : k = 1, \ldots, k(j, \varepsilon)\})_{j=0}^{\infty}$  is an increasing sequence of sets of balls in  $\mathcal{X}$  of the radius  $\varepsilon$  such that  $k(0, \varepsilon) < k(1, \varepsilon) < k(2, \varepsilon) < \ldots$ and  $\mathbf{Pr}(\bigcap_{i=1}^{j} \bigcup_{k=1}^{k(j,\varepsilon)} \{f_i \in B_{k,\varepsilon}\}) \ge 1 - \varepsilon, j = 1, 2, \ldots$  [the existence of such a sequence of sets is guaranteed by the tightness of any probability measure on a separable Banach space].

Then, (2.5) and (2.6) are satisfied. The Jensen's type inequality  $g(f_{j,\varepsilon}) \leq \mathbf{E}(g(f_j)|F_{j,\varepsilon})$  implies (2.7) and  $\|\mathbf{E}(g(f_{j,\varepsilon})|F_{j-1,\varepsilon})\|_{\infty} \leq \|\mathbf{E}(g(f_j)|F_{j-1,\varepsilon})\|_{\infty} \leq \|\mathbf{E}(g(f_j)|F_{j-1})\|_{\infty}$ , so that (2.8) is also true.

REMARK 2.4. Using the standard construction  $(f_{j\wedge n\wedge \tau})_{j=0}^{\infty}$ , where n = 1, 2, ..., $\tau = \inf\{j: ||f_j|| \ge M\}$  with large n and M, we will need only to consider the bounded stopped martingales. By Lemma 2.3 and Lemma 2.2, when proving subsequent results for the martingales in the 2-smooth spaces, we may and shall, without loss of generality, restrict the consideration only to the bounded stopped martingales  $(0, f_1, \ldots, f_n, f_n, f_n, \ldots)$  with each of  $f_j$ 's having only a finite number of values in a finite-dimensional Banach space  $(\mathcal{X}, \|\cdot\|)$  satisfying condition (2.2); for such martingales, we put  $f_{\infty} = f_n$ .

PROPOSITION 2.5. If  $f \in \mathcal{M}(\mathcal{X})$  and  $\mathcal{X}$  is a (2, D)-smooth separable Banach space, then

$$||f_n||_2 \le D||s_2||_2 = D||S_2||_2.$$

PROOF. For j = 1, 2, ..., put

$$g(t) = \mathbf{E} \| f_{j-1} + t d_j \|^2.$$

In view of Remark 2.4, we may assume that  $g'(0) = 0, g''(t) \le 2D^2 ||d_j||_2^2$ , and so,

$$\mathbf{E} \|f_j\|^2 - \mathbf{E} \|f_{j-1}\|^2 = g(1) - g(0) = \int_0^1 g''(t)(1-t)dt$$
  
$$\leq D^2 \|d_j\|_2^2, \quad j = 1, 2, \dots$$

It remains to sum up these inequalities.

The upper bounds provided by Theorems 15.1 and 21.1 of Burkholder (1973)

can be immediately extended to the martingales in the 2-smooth separable Banach spaces. Let us state this as

THEOREM 2.6. If  $f \in \mathcal{M}(\mathcal{X})$ ,  $\mathcal{X}$  is a 2-smooth separable Banach space, a function  $g: [0, \infty) \to [0, \infty)$  is non-decreasing, g(0) = 0, and  $g(2u) \leq c_g g(u)$ ,  $u \geq 0$ , for some  $c_g$ , then

(2.9) 
$$\mathbf{E}g(f^*) \le c_{2.9}[\mathbf{E}g(D\,s_2) + \mathbf{E}g(d^*)];$$

if, moreover, g is convex, then

(2.10) 
$$\mathbf{E}g(f^*) \le c_{2.10}\mathbf{E}g(D\,S_2);$$

here,  $c_{2.9}$  and  $c_{2.10}$  depend only on  $c_g$ .

PROOF repeats that in Burkholder (1973) with the following exceptions: 1) use  $\|\cdot\|$  instead of  $|\cdot|$ , and 2) use Proposition 2.5 instead of the identities (in the notation therein):  $\|h\|_2 = \|S(h)\|_2$  and  $\|h\|_2 = \|s(h)\|_2$ .

It is well-known [see (11.1) in Burkholder (1991)] that  $(Ap)^p$  is optimum for  $c_{2.10}$  at least if  $\mathcal{X}$  is a Hilbert space. As to optimum bounds like (2.10) for conditionally symmetric martingales, see section 4 below.

# 3. Exponential bounds on tail probabilities for the martingales in the 2-smooth spaces.

THEOREM 3.1. Suppose that  $f \in \mathcal{M}(\mathcal{X})$ ,  $\mathcal{X}$  is a (2, D)-smooth separable Banach space, and  $\lambda > 0$  is such that  $\mathbf{E}e^{\lambda \|d_j\|} < \infty$  for j = 1, 2, ... Then for all  $r \geq 0$ ,

$$\begin{aligned} \mathbf{Pr}\left(f^* \geq r\right) &\leq 2\mathrm{e}^{-\lambda r} \left\| \left\| \prod_{j=1}^{\infty} [1 + D^2 \mathbf{E}_{j-1}(\mathrm{e}^{\lambda \|d_j\|} - 1 - \lambda \|d_j\|)] \right\|_{\infty} \\ &\leq 2\exp\left\{ -\lambda r + D^2 \left\| \left| \sum_{j=1}^{\infty} \mathbf{E}_{j-1}(\mathrm{e}^{\lambda \|d_j\|} - 1 - \lambda \|d_j\|) \right\|_{\infty} \right\}. \end{aligned}$$

We shall obtain this theorem as a particular case of the following result for the

(2, D)-smooth functions defined in section 2.

THEOREM 3.2. Suppose that  $f \in \mathcal{M}(\mathcal{X})$ ,  $\mathcal{X}$  is any separable Banach space,  $\lambda > 0$  is such that  $\mathbf{E}e^{\lambda \|d_j\|} < \infty$  for j = 1, 2, ..., and a function  $\Psi$  is (2, D)-smooth. Then for all  $r \geq 0$ ,

(3.1) 
$$\mathbf{Pr}\left(\sup_{j}\Psi(f_{j}) \ge r\right) \le 2\mathrm{e}^{-\lambda r} \left\| \prod_{j=1}^{\infty} (1+e_{j}) \right\|_{\infty},$$

(3.2) 
$$\leq 2 \exp\left\{-\lambda r + \left\|\sum_{j=1}^{\infty} e_j\right\|_{\infty}\right\},$$

where

$$e_j = D_*^2 \mathbf{E}_{j-1}(e^{\lambda ||d_j||} - 1 - \lambda ||d_j||), \quad D_* = 1 \lor D.$$

PROOF. Put  $u(t) = u_{x,v}(t) = \Psi(x + tv)$  for any x, v in  $\mathcal{X}$ . By Lemma 2.2 and Remark 2.4, we may assume that u is Fréchet-differentiable,  $|u'(t)| \leq ||v||$ ,  $(u^2)''(t) \leq 2D^2 ||v||^2$ . Hence, when u''u > 0, one has  $(\cosh u)'' = u'^2 \cosh u + u'' \sinh u$ 

 $\leq (u'^2 + u''u)\cosh u = \frac{1}{2}(u^2)''\cosh u \leq D^2 ||v||^2\cosh u, \text{ and, otherwise, } (\cosh u)'' \leq u'^2\cosh u \leq ||v||^2\cosh u; \text{ so, in any case,}$ 

(3.3) 
$$(\cosh u)'' \le D_*^2 ||v||^2 \cosh u$$

Consider now  $\varphi(t) := \mathbf{E}_{j-1} \cosh(\lambda \Psi(f_{j-1} + td_j)), \quad |t| \le 1$ . In view of (3.3) and Remark 2.4,

$$\varphi''(t) \leq D_*^2 \lambda^2 \mathbf{E}_{j-1} \|d_j\|^2 \cosh(\lambda \Psi(f_{j-1} + td_j))$$
$$\leq D_*^2 \lambda^2 \mathbf{E}_{j-1} \|d_j\|^2 e^{\lambda t \|d_j\|} \cosh(\lambda \Psi(f_{j-1})),$$

|t| < 1. But  $\varphi'(0) = 0$  since  $(f_j)$  is a martingale, and therefore,

$$\mathbf{E}_{j-1}\cosh(\lambda\Psi(f_j)) = \varphi(1) = \varphi(0) + \int_0^1 (1-t)\varphi''(t)dt \le (1+e_j)\cosh(\lambda\Psi(f_{j-1})).$$

Thus, putting  $G_0 = 1$ ,  $G_j = \cosh(\lambda \Psi(f_j)) / \prod_{i=1}^j (1 + e_i)$ , j = 1, 2, ...,one has a positive supermartingale. Hence, if  $\tau := \inf\{j: \Psi(f_j) \ge r\}$ , then  $\mathbf{E}G_{\tau} \le \mathbf{E}G_0 = 1$ , and so,

$$\mathbf{Pr}\left(\sup_{j}\Psi(f_{j}) \geq r\right) \leq \mathbf{Pr}\left(G_{\tau} \geq \cosh(\lambda r) \left| \left| \left| \prod_{j=1}^{\infty} (1+e_{j}) \right| \right|_{\infty} \right).$$

Now, (3.1) follows from Chebyshev's inequality and  $\cosh u > e^u/2$ ; (3.2) is elementary.

REMARK. For  $f \in \mathcal{M}_{ind}(\mathbf{R})$ , i.e., for the sums of independent zero-mean realvalued random variables, the following is used as a starting point when proving exponential inequalities:

$$\mathbf{Pr}(f_n \ge r) \le e^{-\lambda r} \prod_{j=1}^n [1 + \mathbf{E}(e^{\lambda d_j} - 1 - \lambda d_j)]$$
$$\le \exp\left\{-\lambda r + \sum_{j=1}^n \mathbf{E}(e^{\lambda d_j} - 1 - \lambda d_j)\right\}.$$

Thus, Theorem 3.1 provides a similar starting point for  $f \in \mathcal{M}(\mathcal{X})$ ,  $\mathcal{X}$  being 2-smooth. [In this sense, it is analogous to the results of Pinelis and Sakhanenko (1985) for the sums of independent random vectors.] A general method of obtaining exact exponential inequalities for the sums of independent real-valued random variables is proposed in Pinelis and Utev (1989). So, for the martingales in the 2-smooth spaces, these two devices taken together produce analogues of exact "independent real-valued" exponential bounds. For instance, this remark easily leads to the following analogues of classical results of Bernstein and Bennett (1962)-Hoeffding (1963) [cf. Theorem 9 and 3, resp., in Pinelis and Utev (1989)].

THEOREM 3.3. Suppose that  $f \in \mathcal{M}(\mathcal{X}), \mathcal{X}$  is a (2, D)-smooth separable Banach space, and

$$\left\| \sum_{j=1}^{\infty} \mathbf{E}_{j-1} \| d_j \|^m \right\|_{\infty} \le m! \Gamma^{m-2} B^2 / (2D^2)$$

for some  $\Gamma > 0$ , B > 0 and  $m = 2, 3, \ldots$ . Then for all  $r \ge 0$ ,

$$\mathbf{Pr}\left(f^* \ge r\right) \le 2\exp\left(-\frac{r^2}{B^2 + B\sqrt{B^2 + 2\Gamma r}}\right).$$

PROOF. Under the conditions given,

$$D^{2} \left\| \sum_{j=1}^{\infty} \mathbf{E}_{j-1}(\mathrm{e}^{\lambda \| d_{j} \|} - 1 - \lambda \| d_{j} \|) \right\|_{\infty} \leq \frac{1}{2} \sum_{m=2}^{\infty} \lambda^{m} \Gamma^{m-2} B^{2} = \frac{B^{2} \lambda^{2}}{2(1 - \lambda \Gamma)}, 0 \leq \lambda < \frac{1}{\Gamma}.$$

Now Theorem 3.1 yields

$$\mathbf{Pr}\left(f^* \ge r\right) \le 2\exp\left\{-\lambda r + \frac{B^2\lambda^2}{2(1-\lambda\Gamma)}\right\}.$$
  
mize the B.H.S. in  $\lambda$ .

It remains to minimize the R.H.S. in  $\lambda$ .

THEOREM 3.4. Suppose that  $f \in \mathcal{M}(\mathcal{X})$ ,  $\mathcal{X}$  is a (2, D)-smooth separable Banach space, and  $||d^*||_{\infty} \leq a$ ,  $||s_2||_{\infty} \leq b/D$  for some a > 0, b > 0. Then for all  $r \geq 0$ ,

(3.4) 
$$\mathbf{Pr}\left(f^* \ge r\right) \le 2 \exp\left[\frac{r}{a} - \left(\frac{r}{a} + \frac{b^2}{a^2}\right) \ln\left(1 + \frac{ra}{b^2}\right)\right]$$
  
(3.5) 
$$\le 2 \left(\frac{eb^2}{ra}\right)^{r/a}.$$

PROOF. Since the function  $g(u) := u^{-2}(e^u - 1 - u)$  for  $u \neq 0$ , g(0) := 1/2 is increasing in  $u \in \mathbf{R}$ ,

$$\mathbf{E}_{j-1}(e^{\lambda \|d_j\|} - 1 - \lambda \|d_j\|) \le \frac{e^{\lambda a} - 1 - \lambda a}{a^2} \mathbf{E}_{j-1} \|d_j\|^2.$$

Now Theorem 3.1 yields

$$\mathbf{Pr}\left(f^* \ge r\right) \le 2\exp\left\{-\lambda r + \frac{\mathrm{e}^{\lambda a} - 1 - \lambda a}{a^2}b^2\right\},\,$$

and the minimization in  $\lambda$  gives (3.4). Inequality (3.5) is trivial.

In the special case  $\mathcal{X} = L^2$ , a bound similar to (3.4), but somewhat weaker, was proved by Kallenberg and Sztencel (1991); their method seems to be confined only to Hilbert spaces.

Theorem 3.4 was proved in Pinelis (1992) for  $\mathcal{X} = L^p$ ,  $p \ge 2$ ; a version for the general 2-smooth spaces was given therein too, but with another, greater constant in place of D. THEOREM 3.5. Suppose that  $f \in \mathcal{M}(\mathcal{X}), \mathcal{X}$  is a (2, D)-smooth separable Banach space, and  $\sum_{j=1}^{\infty} ||d_j||_{\infty}^2 \leq b_*^2$  for some  $b_* > 0$ . Then for all  $r \geq 0$ ,

$$\mathbf{Pr}\left(f^* \geq r\right) \leq 2\exp\left\{-\frac{r^2}{2D^2b_*^2}\right\}$$

PROOF is the same as that of Theorem 3 in Pinelis (1992) except that, in view of (3.3), one can use  $D^2$  instead of B therein.

Theorem 3.5 can be improved in the special case of conditionally symmetric martingales.

THEOREM 3.6. Suppose that  $\mathcal{X}$  is a (2, D)-smooth separable Banach space,  $f \in \mathcal{M}(\mathcal{X}), \|S_2(f)\|_{\infty} \leq b$  for some b > 0, and the increments  $d_j$ 's are  $F_{j-1}$ conditionally symmetrically distributed,  $j = 1, 2, \ldots$ . Then for all  $r \geq 0$ ,

$$\mathbf{Pr}\left(f^* \geq r\right) \leq 2\exp\left\{-\frac{r^2}{2D^2b^2}\right\}.$$

PROOF. Being conditionally symmetric,  $(f_j)$  is a martingale also relative to the sequence  $(G_j)$ , where  $G_j$  is the  $\sigma$ -field generated by  $F_j$  and  $||d_{j+1}||$ ; see, e.g. Lemma 10.2 in Burkholder (1991). Now, the proof can be concluded as that of Theorem 3 in Pinelis (1992): only the conditional expectations given  $G_j$ 's are taken instead of those given  $F_j$ 's, and  $D^2$ ,  $||d_j||^2$  are used in place of B,  $b_n^2$ therein, resp.

In the case when  $\mathcal{X} = \mathbf{R}$  and  $d_j$ 's are simple functions, Theorem 3.6 was given in Hitczenko (1990a).

An analogous result for sums of independent random vectors in arbitrary separable Banach spaces is given in Pinelis (1990).

### 4. A spectrum of Rosenthal-Burkholder type bounds on moments of the martingales in the 2-smooth spaces.

THEOREM 4.1. If  $f \in \mathcal{M}(\mathcal{X})$ ,  $\mathcal{X}$  is a (2, D)-smooth separable Banach space,

 $p \geq 2, 1 \leq c \leq p$ , then

(4.1) 
$$\|f^*\|_p \leq c \|d^*\|_p + \sqrt{c} e^{p/c} D \|s_2\|_p;$$

in particular,

(4.2) 
$$||f^*||_p \leq p ||d^*||_p + \sqrt{p} D ||s_2||_p$$

(4.3) 
$$||f^*||_p \leq \frac{p}{\ln p} (||d^*||_p + D||s_2||_p)$$

(4.4) 
$$||f^*||_p \preceq \alpha ||d^*||_p + e^{p/\alpha} D ||s_2||_p, \quad 1 \le \alpha \le \frac{p}{\ln(ep)}.$$

We need

LEMMA 4.2. Suppose that  $f \in \mathcal{M}(\mathcal{X})$ , the increments  $d_j$ 's are  $F_{j-1}$ -conditionally symmetrically distributed,  $j = 1, 2, ..., \lambda > 0, \delta_1 > 0, \delta_2 > 0, \beta - 1 - \delta_2 > 0$ . Then

$$\mathbf{Pr}\left(f^* > \beta\lambda, w^* \le \lambda\right) \le \varepsilon \mathbf{Pr}\left(f^* > \lambda\right),$$

where

$$w^* = \left(\frac{d^*}{\delta_2}\right) \lor \left(\frac{D s_2}{\delta_1}\right),$$

(4.5) 
$$\varepsilon = 2\left(\frac{\mathrm{e}}{N}\frac{\delta_1^2}{\delta_2^2}\right)^N, \quad N = \frac{\beta - 1 - \delta_2}{\delta_2}$$

PROOF. Put  $\bar{d}_j = d_j I\{||d_j|| \leq \delta_2 \lambda\}$ ,  $\bar{f}_j = \sum_{i=0}^j \bar{d}_i$ ,  $h_j = \bar{f}_{(j \wedge \tau \wedge \nu) \lor \mu} - \bar{f}_{\mu}$ , where  $\mu = \inf\{j: ||\bar{f}_j|| > \lambda\}$ ,  $\nu = \inf\{j: ||\bar{f}_j|| > \beta\lambda\}$ ,  $\tau = \inf\{j: \bar{s}_{j+1} > \lambda\}$ ,  $\bar{s}_{j+1} = \sqrt{\sum_{i=1}^{j+1} \mathbf{E}_{i-1} ||\bar{d}_i||^2}$ ,  $j = 0, 1, \dots$  Then  $\mathbf{E}_{j-1} \bar{d}_j = 0$ ,  $h_j - h_{j-1} = \bar{d}_j I\{\mu < j \leq \tau \land \nu\}$ ,  $j = 1, 2, \dots$ , and so,  $(h_j)$  is a martingale in  $\mathcal{X}$  conditionally on  $F_{\mu}$ [the  $\sigma$ -field consisting of all  $\Omega_0 \in F$  such that  $\Omega_0 \cap \{\mu = j\} \in F_j$  for all j]. Besides,

$$\begin{aligned} \mathbf{Pr}\left(f^* > \beta\lambda, w^* \leq \lambda\right) &= \mathbf{Pr}\left(\bar{f}^* > \beta\lambda, w^* \leq \lambda\right) \leq \mathbf{Pr}\left(h^* > (\beta - 1 - \delta_2)\lambda\right) \\ &= \mathbf{EPr}\left(h^* > (\beta - 1 - \delta_2)\lambda \mid F_{\mu}\right)I\{\mu < \infty\} \\ &\leq \varepsilon \mathbf{Pr}\left(\mu < \infty\right) = \varepsilon \mathbf{Pr}\left(f^* > \lambda\right); \end{aligned}$$

here, we have applied (3.5) with  $r = (\beta - 1 - \delta_2)\lambda$ ,  $a = \delta_2\lambda$ ,  $b = \delta_1\lambda$ .

PROOF OF THEOREM 4.1. Argument in Hitczenko (1990) shows that we need to consider only the following two cases: 1) the increments  $d_j$ 's are  $F_{j-1}$ conditionally symmetrically distributed, and 2)  $f \in \mathcal{M}_{ind}(\mathcal{X})$ . But via the standard symmetrization formula  $\widetilde{X} = X - X'$ , where X, X' are independent copies, one can easily reduce case 2) to 1). Thus, one can use Lemma 4.2.

Applying now Lemma 7.1 in Burkholder (1973) with  $\Phi(\lambda) = \lambda^p$ ,  $\gamma = \beta^p$ ,  $\delta = \eta = 1$ , one has

(4.6) 
$$\|f^*\|_p \le 2\beta \left(\frac{\|d^*\|_p}{\delta_2} + \frac{D\|s_2\|_p}{\delta_1}\right) \quad \text{if} \quad \beta^p \varepsilon \le 1/2,$$

where  $\varepsilon$  is given by (4.5). Choose now, for any  $c \in [1, p]$ ,

$$\beta = 1 + e^{-p/c} + \frac{1}{c}, \quad \delta_2 = \frac{1}{10c}, \quad \delta_1 = \frac{1}{10\sqrt{c}e^{p/c}}.$$

Then  $\beta < 3$ ,

$$\begin{split} N\left(=\frac{\beta-1-\delta_2}{\delta_2}\right) &= 9+10c\mathrm{e}^{-p/c} > 9,\\ \beta^p &= [1+(N+1)\delta_2]^p < \mathrm{e}^{2pN\delta_2},\\ (\beta^p\varepsilon)^{1/N} &< 2^{1/9}\mathrm{e}^{2p\delta_2}\frac{\mathrm{e}}{10c\mathrm{e}^{-p/c}}\frac{c}{\mathrm{e}^{2p/c}} < \frac{1}{2}, \end{split}$$

so that (4.6) implies

$$||f^*||_p \le 60(c||d^*||_p + \sqrt{c} e^{p/c} D||s_2||_p);$$

thus, (4.1) is proved.

Let  $c_p$  stand for the unique solution to the equation  $\sqrt{c_p} = e^{p/c_p}$ . Then,  $c_p \sim 2p/\ln p$  as  $p \to \infty$ . Hence, putting c = p,  $c = c_p$ , in (4.1), one comes to (4.2), (4.3), resp.

The function  $g(c) := \sqrt{c} e^{p/c}$  decreases on  $[1, p], g(1) = e^p, g(p) = e\sqrt{p} < ep$ . Hence, for each  $\alpha \in [1, p/\ln(ep)]$ , there exists  $z_{\alpha} \in [1, p]$ , the unique solution to the equation  $g(z_{\alpha}) = e^{p/\alpha}$ ; besides,  $g(2\alpha) = e^{p/\alpha}\sqrt{2\alpha} e^{-p/(2\alpha)} \leq e^{p/\alpha}\sqrt{2\alpha_p} e^{-p/(2\alpha_p)} < g(z_{\alpha})$ , where  $\alpha_p := p/\ln(ep)$ . Thus,  $z_{\alpha} < 2\alpha$ . Now we see that (4.1) with  $c = z_{\alpha}$  yields (4.4).

For  $\mathcal{X} = \mathbf{R}$ , inequalities (4.3) and (4.2) were proved in Hitczenko (1990) and Hitczenko (1991), resp.

A spectrum of bounds on moments of martingales in Hilbert spaces with bounded second conditional moments was found in Pinelis (1980). It is essentially equivalent to (4.1) at least in the case of independent increments  $d_j$ 's (see Proposition 9.2 below) but has a much more cumbersome expression.

The infimum in c of the R.H.S. of (4.1), evaluated in section 6 below, turns out to be an upper bound on  $||f^*||_p$  optimal in terms of  $||d^*||_p$ ,  $||s_2||_p$ , the optimum choice of c depending, obviously, on  $||d^*||_p/||s_2||_p$ . Besides, for each  $c \in [1, p]$ , the "individual" bound  $c||d^*||_p + \sqrt{c} e^{p/c} ||s_2||_p$  is optimal for a certain corresponding value of  $||d^*||_p/||s_2||_p$ ; in particular, all the bounds in (4.2)-(4.4) are optimal. The issue of optimality is treated rigorously in section 6.

THEOREM 4.3. If  $\mathcal{X}$  is a (2, D)-smooth separable Banach space,  $f \in \mathcal{M}(\mathcal{X})$ , and the increments  $d_j$ 's are  $F_{j-1}$ -conditionally symmetrically distributed, then

$$||f^*||_p \leq \sqrt{p} D||S_2||_p, \quad p \ge 1.$$

PROOF. Consider f as a martingale relative to the sequence of  $\sigma$ -fields  $(G_j)$  defined in the proof of Theorem 3.6. Then  $s_2 = S_2$ . Reasoning as in the proof of Lemma 4.2 but using Theorem 3.6 instead of Theorem 3.4, we see that

(4.7) 
$$\mathbf{Pr}\left(f^* > \beta\lambda, D S_2 \le \delta\lambda\right) \le \varepsilon \mathbf{Pr}\left(f^* > \lambda\right),$$

 $\lambda > 0, \delta > 0, \beta - 1 - \delta > 0$ , where  $\varepsilon = \exp[-\delta^{-2}(\beta - 1 - \delta)^2/2]$ . It remains to choose, say,  $\beta = 2, \ \delta = (0.1)p^{-1/2}$  and apply Lemma 7.1 of Burkholder (1973).

REMARK. For conditionally symmetric martingales in Hilbert spaces, the exact constant  $A_p$  in the inequality

(4.8) 
$$||f_n||_p \le A_p ||S_2||_p$$

was found [see Burkholder (1991, chapter 11), Davis (1976), Wang (1991)]. For any real martingale with independent symmetrically distributed increments, it follows from the result of Whittle (1960) [proved correctly only for p = 2 and  $p \geq 3$ ] and of Haagerup (1982) that (4.8) takes place with the exact, in this "independent increments" case, constant  $A_p = \|\xi\|_p$ , where  $\xi \sim N(0,1), p \geq 2$ . Since  $\|\xi\|_p \asymp \sqrt{p}$ , the bound in Theorem 4.3 is optimal (to  $\asymp$ ).

REMARK. Bounds given in Theorems 3.1, 4.1, 4.3 are only possible in the 2smooth Banach spaces, even if we need a bound like those in Theorem 4.1 for just one particular p. Indeed, all the mentioned here results imply  $||f^*||_p \leq CD||S_2||_p$ for at least one particular  $p \geq 2$ , some C > 0 and all Walsch-Paley martingales in  $\mathcal{X}$  since, for those martingales,  $s_2 = S_2 \geq d^*$ . Thus, one has (4.7) with  $\varepsilon = (CD\delta/(\beta - 1 - \delta))^p$ , and so,  $||f^*||_2 \leq C_1D||S_2||_2$  for some  $C_1 > 0$ . It remains now to recall the characterization of the 2-smooth Banach spaces given by Pisier (1975).

## 5. Applications: bounds on central moments of the norm of the sum of independent random vectors in arbitrary Banach spaces.

THEOREM 5.1. If  $f \in \mathcal{M}_{ind}(\mathcal{X})$ ,  $(\mathcal{X}, \|\cdot\|)$  is any separable Banach space,  $p \geq 2, 1 \leq c \leq p$ , and x is any non-random vector in  $\mathcal{X}$ , then for all n = 1, 2, ...,

(5.1) 
$$\| \|f_n + x\| - \mathbf{E} \|f_n + x\| \|_p \leq c \|d^*\|_p + \sqrt{c} e^{p/c} \|S_2\|_2;$$

in particular,

(5.2)  $\|\|f_n + x\| - \mathbf{E}\|f_n + x\|\|_p \leq p\|d^*\|_p + \sqrt{p}\|S_2\|_2,$ 

(5.3) 
$$\|\|f_n + x\| - \mathbf{E}\|f_n + x\|\|_p \preceq \frac{p}{\ln p} (\|d^*\|_p + \|S_2\|_2),$$

(5.4) 
$$\| \| f_n + x \| - \mathbf{E} \| f_n + x \| \|_p \leq \alpha \| d^* \|_p + e^{p/\alpha} \| S_2 \|_2.$$

 $1 \le \alpha \le p/\ln(\mathrm{e}p).$ 

PROOF. We use the following modification of the method of Yurinskii (1974), being stated here as

LEMMA 5.2. For  $f \in \mathcal{M}_{ind}(\mathcal{X})$ , let  $F_j$  be the  $\sigma$ -field generated by  $f_0, f_1, \ldots, f_j$ ; put  $\zeta_j = \mathbf{E}_j ||f_n + x|| - \mathbf{E} ||f_n + x||, \ \xi_{j+1} = \zeta_{j+1} - \zeta_j, \ j = 0, 1, \ldots, n$ . Then  $(\zeta_j)$  is a martingale,

$$|\xi_j| \le ||d_j|| + \mathbf{E} ||d_j||, \quad \mathbf{E}_{j-1}\xi_j^2 \le ||d_j||^2,$$

j = 1, ..., n, and

$$||f_n + x|| - \mathbf{E}||f_n + x|| = \zeta_n.$$

PROOF can be found in each of the papers of Pinelis (1981, 1986), Pinelis and Sakhanenko (1985). Since it is very short, we repeat it here for the reader's convenience. Put  $\eta_j = \mathbf{E}_j(||f_n + x|| - ||f_n + x - d_j||)$ . Then  $|\eta_j| \le ||d_j||$ . Besides,  $\xi_j = \eta_j - \mathbf{E}_{j-1}\eta_j, \ j = 1, \dots, n$ , in view of the independence of  $d_j$ 's; hence,

$$\begin{aligned} |\xi_j| &\le |\eta_j| + |\mathbf{E}_{j-1}\eta_j| \le ||d_j|| + \mathbf{E}||d_j||, \\ \mathbf{E}_{j-1}\xi_j^2 &= \mathbf{E}_{j-1}\eta_j^2 - (\mathbf{E}_{j-1}\eta_j)^2 \le \mathbf{E}_{j-1}\eta_j^2 \le \mathbf{E}||d_j||^2, \quad j = 1, \dots, n. \quad \blacksquare \end{aligned}$$

To complete now the proof of Theorem 5.1, apply Theorem 4.1 to the martingale  $(0, \zeta_1, \zeta_2, \ldots, \zeta_n, \zeta_n, \zeta_n, \ldots), \zeta_j$ 's defined in Lemma 5.2. Using the lemma and the inequalities of Minkowsky and Hölder, we see that

$$\sum_{j=1}^{n} \mathbf{E}_{j-1} \xi_{j}^{2} \leq \sum_{j=1}^{n} \mathbf{E} \|d_{j}\|^{2} = \|S_{2}\|_{2}^{2},$$
$$\|\max_{j\leq n} |\xi_{j}|\|_{p} \leq \|\max_{j} (\|d_{j}\| + \mathbf{E} \|d_{j}\|)\|_{p}$$
$$\leq \|d^{*}\|_{p} + \max_{j} \mathbf{E} \|d_{j}\|$$
$$\leq 2\|d^{*}\|_{p}.$$

Finally, note that the constant of smoothness  $D(\mathbf{R}) = 1$ . Now the statement of Theorem 5.1 is evident.

Hoffmann-Jørgensen (1974) found the following extension of the Rosenthal's inequality for  $f \in \mathcal{M}_{ind}(\mathcal{X}), p \geq 1$ :

(5.5) 
$$|| ||f_n|| ||_p \le c(p)(||d^*||_p + || ||f_n|| ||_1)$$

with c(p) depending only on p [it can be seen that the best choice of parameters in the method of Hoffmann-Jørgensen gives (5.5) with  $c(p) \approx p$ ].

For  $f \in \mathcal{M}_{ind}(\mathcal{X})$ , Pinelis (1978) proved that

$$|| || f_n || ||_p \leq c_1(p) || || f_n || ||_1 + c_2(p) ||S_p||_p + c_3(p) ||S_2||_2,$$

which is also a generalization of the Rosenthal's (1970) inequality; the method can actually yield  $c_1(p) = 1$ ,  $c_2(p) = p$ ,  $c_3(p) = \sqrt{p}$ .

An inequality, implying (5.1), was obtained in Pinelis (1980) [see Proposition 9.2 below].

De Acosta (1981) proved a version of (5.3) but with x = 0 and with an implicit factor c(p) instead of  $p/\ln p$ . Since he used inequality (21.1) of Burkholder (1973), Proposition 9.1 below shows that in fact his method leads to c(p) = p.

Using isoperimetric technique, Talagrand (1989) proved the following version of (5.5):

(5.6) 
$$\| \|f_n\| \|_p \leq \frac{p}{\ln(2p)} (\|d^*\|_p + \| \|f_n\| \|_1)$$

for  $f \in \mathcal{M}_{ind}(\mathcal{X})$ ,  $p \geq 1$ , which was also proved in Kwapień and Szulga (1991) by a different method. The Talagrand's inequality may be compared with (5.1) and, in particular, with (5.3). If  $||S_2||_2$  is much greater than  $||d^*||_p + || ||f_n|| ||_1$ , then (5.1) may lose to (5.6) in certain cases. If however  $\mathcal{X}$  is e.g. of cotype 2, then (5.3) is often no worse then (5.6); moreover, say, in the typical case of the increments with the same or almost the same distribution, (5.2) is significantly better. On the other hand, if there are heavy distribution tails, i.e., if  $||d^*||_p$  is much greater than both  $||S_2||_2$  and  $|| ||f_n|| ||_1$ , then (5.4) with  $\alpha \approx 1$  does better than (5.6). Other advantages of bounds (5.1)-(5.4) are that they are applicable to the sums of non-zero-mean random vectors (owing to the presence of x) and better reflect the concentration phenomenon of the distribution of the sum of independent random vectors.

Modifications of the method of Yurinskii (1974) for  $f \in \mathcal{M}_{ind}(\mathcal{X})$ , allowing reduction of the problem of upper bounds on the L.H.S. of (5.1) for any separable Banach space  $\mathcal{X}$  to that of upper bounds on  $||f_n||_p$  for  $f \in \mathcal{M}_{ind}(\mathbf{R})$  were proposed in Berger (1991) [for x = 0] and in Pinelis (1992a) [with the best constant, for any  $x \in \mathcal{X}$ ]; actually, instead of the power moment function  $u \mapsto$  $|u|^p$ , one can use any convex function there. A straightforward application of Theorem 4.1 yields the following bounds in the case of sums of independent zero-mean random variables.

THEOREM 5.3. Suppose that  $f \in \mathcal{M}_{ind}(\mathcal{X})$ ,  $\mathcal{X}$  is a Hilbert space,  $p \geq 2$ ,  $1 \leq c \leq p$ . Then

(5.7) 
$$\| \|f_n\| \|_p \leq c \|S_p\|_p + \sqrt{c} e^{p/c} \|S_2\|_2;$$

in particular,

(5.8) 
$$|| || f_n || ||_p \preceq p ||S_p||_p + \sqrt{p} ||S_2||_2$$

(5.9) 
$$||| ||f_n||||_p \leq \frac{p}{\ln p} (||S_p||_p + ||S_2||_2),$$

(5.10) 
$$||| \|f_n\|||_p \leq \alpha \|S_p\|_p + e^{p/\alpha} \|S_2\|_2, \quad 1 \leq \alpha \leq \frac{p}{\ln(ep)}.$$

PROOF, in view of Theorem 4.1 and the equality  $D(\mathcal{X}) = 1$  for any Hilbert space  $\mathcal{X}$ , consists in the following trivial remarks:  $|| \|f_n\| \|_p \leq \|f^*\|_p, \|d^*\|_p \leq \|S_p\|_p$ , and, for  $f \in \mathcal{M}_{ind}(\mathcal{X}), s_2 = S_2$ .

As it was said in section 4, the results of Pinelis (1980) imply (5.1), (5.7); it was also explained in that paper how to elicit bounds like (5.8) and (5.10) (for  $\alpha = 4$ ). Nevertheless, it is not obvious how to deduce a general inequality like (5.7) from the spectrum of bounds in Pinelis (1980) (again, we refer to Proposition 9.2). From this point of view, even in the classical case of sums of independent real-valued zero-mean random variables, (5.7) is apparently new.

An inequality similar to, say, (5.9), but with  $2^p$  instead of  $p/\ln p$ , was probably first found by Rosenthal (1970) for  $f \in \mathcal{M}_{ind}(\mathbf{R})$ , who also obtained a lower bound, which differs from the upper one by at most a factor depending only on p.

Rosén (1970) proved a result for  $f \in \mathcal{M}_{ind}(\mathbf{R})$ , which implies the upper Rosenthal's bound for  $p = 2, 4, 6, \ldots$  [this implication was unnoticed; to demonstrate it, one can put, in the notation of Rosén,  $\lambda_{\nu}(p) = (\mathbf{E}X_{\nu}^{2p}/\mathbf{E}X_{\nu}^{2})^{1/(2p-2)}$ ,  $\rho_{\nu}(p) = \mathbf{E}X_{\nu}^{2}/(\lambda_{\nu}(p)^{2})$ ]. Moreover, using some ideas of Dharmadhikari and Jogdeo (1969), it is possible to deduce the upper bound by Rosenthal for all real  $p \ge 2$  from the Rosén's result.

A method like just described was used in the student diploma work of Pinelis (1974) to prove an upper bound of the Rosenthal type for  $f \in \mathcal{M}_{ind}(\mathbf{R})$  [via the Marcienkiewicz-Zygmund (1937) inequality, the lower Rosenthal bound was also obtained therein]. While the constants in Pinelis (1974) were implicit, the method could yield (5.9); regrettably, the results of Rosenthal (1970) and Burkholder (1973) had long remained unknown to the author, and so, the problem of the constants was not among the ones considered then by him as most urgent.

Inequality (5.9) for  $f \in \mathcal{M}_{ind}(\mathbf{R})$  and for sums of exchangeable random variables, with the proof that  $p/\ln p$  is optimal in (5.9), was first given by Johnson, Schechtman and Zinn (1985).

A.I. Sakhanenko, a referee of the mentioned diploma work, upon having reviewed it, suggested another approach, giving in effect (5.8), again for  $f \in \mathcal{M}_{ind}(\mathbf{R})$  [see Nagaev and Pinelis (1977)].

An inequality, similar to (5.10) with  $\alpha = 4$ , was found by Sazonov (1974).

All bounds in Theorems 4.1, 5.1, 5.3 are optimum. We shall prove the optimality in the next section, using results and ideas from Pinelis and Utev (1984), where, in particular, for any  $p \ge 2$ ,

$$\sup\{\|f_n\|_p: f \in \mathcal{M}_{ind}(\mathbf{R}), \|S_2\|_2 \text{ and } \|S_p\|_p \text{ fixed}\}\$$

was computed up to  $\approx$ , which, for instance, implies all inequalities (5.7)-(5.10) for  $f \in \mathcal{M}_{ind}(\mathbf{R})$ ; also, it was noted in Pinelis and Utev (1984) that bounds like (5.8) and (5.10) [for  $\alpha \approx 1$ ] represent in a certain sense the two extreme bounds in the spectrum of all optimal bounds on moments, the optimum value of a "spectrum parameter" depending on  $\|S_p\|_p/\|S_2\|_2$ .

#### 6. Optimality of the bounds on moments. Let us consider the following

upper bounds for any  $a_2 > 0$ ,  $a_p > 0$ ,  $p \ge 2$ :

$$B_{p} := B_{p}(a_{p}, a_{2}) := \sup\{\|f^{*}\|_{p}: f \in \mathcal{M}(\mathcal{X}), \ \mathcal{X} \text{ is } (2, D)\text{-smooth}, \\ \|d^{*}\|_{p} = a_{p}, \ D\|s_{2}\|_{p} = a_{2}\}, \\ B_{p,ind}^{\mathcal{X}} := B_{p,ind}^{\mathcal{X}}(a_{p}, a_{2}) := \sup\{\|\|f_{n} + x\| - \mathbf{E}\|f_{n} + x\|\|_{p}: \ f \in \mathcal{M}_{ind}(\mathcal{X}), \\ x \in \mathcal{X}, \ \|d^{*}\|_{p} = a_{p}, \ \|S_{2}\|_{2} = a_{2}, \ n = 1, 2, \ldots\}, \\ B_{p,ind}^{\mathcal{X}} := B_{p,ind}^{\mathcal{X}}(a_{p}, a_{2}) := \sup\{\|f_{n}\| : \ f \in \mathcal{M}_{ind}(\mathbf{R})\}$$

 $B_{p,ind,0} := B_{p,ind,0}(a_p, a_2) := \sup\{\|f_n\|_p: f \in \mathcal{M}_{ind}(\mathbf{R}), \\ \|d^*\|_p = a_p, \|S_2\|_2 = a_2, n = 1, 2, \ldots\},\$ 

and their analogues  $B_{p,S} = B_{p,S}(a_p, a_2)$ ,  $B_{p,ind,S}^{\mathcal{X}} = B_{p,ind,S}^{\mathcal{X}}(a_p, a_2)$ ,  $B_{p,ind,0,S}(a_p, a_2)$ ,  $B_{p,ind,0,S}(a_p, a_2)$  obtained by replacing the equality  $||d^*||_p = a_p$  in the above definitions by  $||S_p||_p = a_p$ .

We shall show that all the introduced bounds are  $\approx$ -equivalent to each of the following:

$$B_p^* := B_p^*(a_p, a_2) := a_p + \sqrt{p} a_2 + \frac{p a_p}{\ln(2 + \frac{a_p}{a_2}\sqrt{p})},$$
$$\hat{B}_p := \hat{B}_p(a_p, a_2) := \min\{ca_p + \sqrt{c} e^{p/c} a_2 \colon 1 \le c \le p\},$$
$$\check{B}_p := \check{B}_p(a_p, a_2) := \max\{(p\alpha + 1)^{1 - \alpha/2} a_p^{1 - \alpha} a_2^{\alpha} \colon 0 \le \alpha \le 1\}$$

Theorem 6.1 below principally means that for any pair  $(a_p, a_2)$  of the values of the characteristics used in the bounds (4.1), (5.1) and (5.7) there exists a value of the "spectrum parameter" c providing an optimum bound. Roughly speaking, it means that spectra (4.1), (5.1) and (5.7) are rich enough, so Theorem 6.1 may be called the "spectrum completeness theorem". It also means that it is not essential in this context values of which of the two pairs are fixed:  $||d^*||_p$  and  $||s_2||_p$ , or  $||S_p||_p$  and  $||s_2||_p$ .

THEOREM 6.1. For all p > 2,  $a_2 > 0$ ,  $a_p > 0$  and for all separable Banach spaces  $(\mathcal{X}, \|\cdot\|)$ ,

$$B_p \asymp B_{p,ind}^{\mathcal{X}} \asymp B_{p,ind,0} \asymp B_{p,S} \asymp B_{p,ind,S}^{\mathcal{X}} \asymp B_{p,ind,0,S}$$
$$\approx \check{B}_p \asymp \hat{B}_p \asymp B_p^*.$$

Proof is comparatively long and will be given later in this section.

The "spectrum"  $ca_p + \sqrt{c} e^{p/c}a_2$ ,  $1 \leq c \leq p$ , turns out to be not only "complete" but also "minimal" in the sense that for each  $c \in [1, p]$ , there exist  $a_p > 0$ ,  $a_2 > 0$  such that the "individual" bound  $ca_p + \sqrt{c} e^{p/c}a_2$  is the best possible. Let us now give the rigorous statement.

THEOREM 6.2. For any  $c \in [1, p]$ , p > 2,

$$\sup\left\{\frac{B_p(a_p, a_2)}{ca_p + \sqrt{c} e^{p/c} a_2}: a_p > 0, a_2 > 0\right\} \asymp 1;$$

here any of the other 5 bounds  $B_{p,ind}^{\mathcal{X}}$ ,  $B_{p,ind,0}$ ,  $B_{p,S}$ ,  $B_{p,ind,S}^{\mathcal{X}}$ ,  $B_{p,ind,0,S}$  may be used in place of  $B_p$ . In particular, all bounds (4.2)-(4.4), (5.2)-(5.4), (5.8)-(5.10) are optimal.

Proof will be given after that of Theorem 6.1.

The following proposition might seem analogous to Theorem 6.2, but it is less important since, for an "individual" value of  $\alpha$ ,  $(p\alpha + 1)^{1-\alpha/2}a_p^{1-\alpha}a_2^{\alpha}$  does not represent an upper bound on the moments – actually, by Theorem 6.1,  $B_p(a_p, a_2) \succeq (p\alpha + 1)^{1-\alpha/2}a_p^{1-\alpha}a_2^{\alpha}$ .

PROPOSITION 6.3. For any  $\alpha \in [0, 1], p > 2$ ,

$$\inf\left\{\frac{B_p(a_p, a_2)}{(p\alpha + 1)^{1-\alpha/2}a_p^{1-\alpha}a_2^{\alpha}}: a_p > 0, a_2 > 0\right\} \asymp 1;$$

here, any of the other 5 bounds  $B_{p,ind}^{\mathcal{X}}$ ,  $B_{p,ind,0}$ ,  $B_{p,S}$ ,  $B_{p,ind,S}^{\mathcal{X}}$ ,  $B_{p,ind,0,S}$  may be used in place of  $B_p$ .

Proof will be given after that of Theorem 6.2.

REMARK. It is easy to see that Doob's inequality

$$||f^*||_p \le \frac{p}{p-1} \sup_n || ||f_n|| ||_p, \quad p > 1, f \in \mathcal{M}(\mathcal{X}),$$

(see also (1.4) in Burkholder (1973)) remains true for all separable Banach spaces  $\mathcal{X}$ . Therefore, one could replace  $||f^*||_p$  in the definition of  $B_p$  by  $\sup_n || ||f_n|| ||_p$ , and statements

6.1–6.3 would hold. This remark can be also deduced from the proof of these statements.  $\blacksquare$ 

Proof of Theorem 6.1 is based on lemmas 6.4 - 6.9 below.

LEMMA 6.4. If p > 2, m is a positive integer,  $m \le p/2$ ,  $a_p > 0$ ,  $a_2 > 0$ , then

$$B_{p,ind,0}(a_p, a_2) \succeq a_p + B_{2m,ind,sym,S}((a_p^{2m-2}a_2^{p-2m})^{\frac{1}{p-2}}, a_2),$$

where  $B_{2m,ind,sym,S}$  may be defined by (6.6), (6.7) below.

PROOF. Put  $g_n(t) = t^{-p/2}[1 - (1 - t/n)^n]$ ,  $g_{\infty}(t) = t^{-p/2}(1 - e^{-t})$ , t > 0, n > 0. Then  $g_n(t)$  decreases in n to  $g_{\infty}(t)$ ,  $g_n(t)$  decreases in  $t \in (0, n]$  from  $g_n(0+) = \infty$  to  $g_n(n) = n^{-p/2}$ ,  $g_{\infty}(t)$  decreases in  $t \in (0, \infty)$  from  $g(0+) = \infty$ to  $g(\infty) = 0$ . Hence, for any  $n \in (a_2^2/a_p^2, \infty]$ , there exists the unique solution  $t_n \in (0, n)$  to the equation  $g_n(t_n) = (a_p/a_2)^p$ , and  $t_n \downarrow t_{\infty}$  as  $n \uparrow \infty$ .

For any  $p > 0, a_p > 0, a_2 > 0$ , and any integer  $n > a_2^2/a_p^2$ , consider  $f \in \mathcal{M}_{ind}(\mathbf{R})$  such that

$$\mathbf{Pr}(d_j = \pm u_n) = q_n/2, \quad \mathbf{Pr}(d_j = 0) = 1 - q_n \text{ for } j = 1, \dots, n,$$
  
 $\mathbf{Pr}(d_j = 0) = 1 \text{ for } j = n + 1, n + 2, \dots,$ 

where

$$q_n := \frac{t_n}{n}, \quad u_n := \frac{a_2}{\sqrt{t_n}}.$$

Then, for the so constructed f,

(6.1) 
$$||d^*||_p = a_p, \quad ||S_2||_2 = a_2,$$

(6.2) 
$$\|S_{2m}\|_{2m} = t_n^{(1-m)/(2m)} a_2 \to t_\infty^{1/(2m)} u_\infty,$$

$$\mathbf{E}e^{i\lambda f_n} = [1 + q_n(\cos\lambda u_n - 1)]^n \to \mathbf{E}e^{i\lambda Z}, \quad \lambda \in \mathbf{R},$$

as  $n \to \infty$ , where

(6.3) 
$$u_{\infty} = \lim_{n \to \infty} u_n = \frac{a_2}{\sqrt{t_{\infty}}},$$

Z is a (symmetrized Poisson) random variable with the characteristic function

(6.4) 
$$\mathbf{E}e^{i\lambda Z} = \exp[(\cos\lambda u_{\infty} - 1)t_{\infty}].$$

Hence, by the analogue of the Fatou lemma for the convergence in distribution [see, e.g., Theorem 5.3 in Billingsley (1968)],

(6.5) 
$$\liminf_{n \to \infty} \|f_n\|_{2m} \ge \|Z\|_{2m}.$$

Consider now, for  $b_{2m} > 0, b_2 > 0$ ,

 $B_{2m,ind,sym,S}(b_{2m},b_2) := \sup\{\|f_n\|_{2m}: f \in \mathcal{M}_{ind}(\mathbf{R}), \|S_{2m}\|_{2m} = b_{2m}, \|S_2\|_2 = b_2, \\ d_i$ 's symmetrically distributed  $\}.$ 

The theorem in Pinelis and Utev (1984) states that

(6.6) 
$$B_{2m,ind,sym,S}(b_{2m},b_2) = \left(\sum_{j=1}^m \Gamma_{j,m}(b_{2m}^{m-j}b_2^{j-1})^{\frac{2m}{m-1}}\right)^{\frac{1}{2m}},$$

where

(6.7) 
$$\Gamma_{j,m} = (2m)! \sum_{r=1}^{j} \sum_{k=1}^{r} \prod_{k=1}^{r} \frac{1}{j_k! ((2m_k)!)^{j_k}};$$

the inner summation is over all positive integers  $m_1 > \cdots > m_r$  and  $j_1, \ldots, j_r$ such that  $m_1 j_1 + \cdots + m_r j_r = m$ ,  $j_1 + \cdots + j_r = j$ . It follows from its proof that

(6.8) 
$$B_{2m,ind,sym,S}(t_{\infty}^{\frac{1}{2m}}u_{\infty}, t_{\infty}^{\frac{1}{2}}u_{\infty}) = ||Z||_{2m},$$

where Z is defined by (6.4) [note the misprint in formula (7) of Pinelis and Utev (1984): there must be  $A_{2m_k,n}^{j_k}$  instead of  $A_{2m_k,n}$ ]. Observe that for the f constructed above in this proof,

$$||S_{2m}||_{2m} = (||S_p||_p^{2m-2}||S_2||_2^{p-2m})^{\frac{1}{p-2}}.$$

Now, (6.2), (6.1) and the inequality  $||S_p||_p \ge ||d^*||_p$  imply

(6.9) 
$$t_{\infty}^{\frac{1}{2m}} u_{\infty} \ge (a_p^{2m-2} a_p^{p-2m})^{\frac{1}{p-2}}.$$

But (6.6), (6.7) show that  $B_{2m,ind,sym,S}$  is an increasing function in either of its arguments. Thus, using (6.5), (6.8), (6.9) and (6.3), we deduce

$$\liminf_{n \to \infty} \|f_n\|_{2m} \ge B_{2m,ind,sym,S}\left( (a_p^{2m-2} a_2^{p-2m})^{\frac{1}{p-2}}, a_2 \right).$$

The lower Rosenthal's bound and Hölder's inequality give

$$||f_n||_p \succeq ||S_p||_p \lor ||f_n||_{2m} \ge a_p \lor ||f_n||_{2m}.$$

Now the statement of the lemma is obvious.  $\hfill\blacksquare$ 

LEMMA 6.5. Under the conditions of Lemma 6.4,

$$B_{p,ind,0,S}(a_1,a_2) \succeq a_p + B_{2m,ind,sym,S}((a_p^{2m-2}a_2^{p-2m})^{\frac{1}{p-2}},a_2).$$

PROOF is similar to that of Lemma 6.4 but easier. In fact, Lemma 6.5 was proved in Pinelis and Utev (1984).  $\blacksquare$ 

LEMMA 6.6. For all  $p \ge 2$ ,  $a_p > 0$ ,  $a_2 > 0$ , one has  $\hat{B}_p(a_p, a_2) \preceq \check{B}_p(a_p, a_2)$ .

PROOF. Consider the functions

(6.10) 
$$g_p(\alpha) := (p\alpha + 1)^{1-\alpha/2} q^{\alpha}, \qquad q := a_2/a_p, \qquad 0 \le \alpha \le 1,$$

(6.11) 
$$q_p(\alpha) := \sqrt{p\alpha + 1} \exp\left\{\frac{1}{2}\frac{p\alpha - 2p}{p\alpha + 1}\right\}, \qquad 0 \le \alpha \le 1.$$

Then, by the definition of  $\check{B}_p$ ,

(6.12) 
$$\check{B}_p(a_p, a_2) = \max\{a_p g_p(\alpha) \colon 0 \le \alpha \le 1\}.$$

Besides,  $q_p$  is continuous and increasing and maps  $\left[0,1\right]$  onto the segment

(6.13) 
$$I_p := \left[ e^{-p}, \sqrt{p+1} \exp\left\{ -\frac{1}{2} \frac{p}{p+1} \right\} \right].$$

An essential relation between  $g_p$  and  $q_p$  is

(6.14) 
$$g'_p(\alpha) = g_p(\alpha) \ln \frac{q}{q_p(\alpha)}.$$

If we have  $q \in I_p$ , take  $\alpha_p = q_p^{-1}(q)$ ; then  $\alpha_p \in [0, 1]$ ,  $q = q_p(\alpha_p)$ , and, in view of (6.10), (6.11),

(6.15) 
$$(p\alpha_p + 1)a_p = a_p g_p(\alpha_p) \exp\left\{-\frac{\alpha_p}{2} \frac{p\alpha_p - 2p}{p\alpha_p + 1}\right\} \asymp a_p g_p(\alpha_p),$$
$$\sqrt{p\alpha_p + 1} e^{p/(p\alpha_p + 1)}a_2 = a_p g_p(\alpha_p) \exp\left\{\frac{p\alpha_p(3 - \alpha_p)}{2(p\alpha_p + 1)}\right\} \asymp a_p g_p(\alpha_p).$$

This implies, with  $c_p := p\alpha_p + 1$ , that

(6.16) 
$$c_p a_p + \sqrt{c_p} e^{p/c_p} a_2 \asymp a_p g_p(\alpha_p)$$

(if  $q \in I_p$ ). Obviously,  $c_p \in [1, p + 1]$ . Hence, in view of (6.12), if  $q \in I_p$  and  $c_p \leq p$ , the lemma is true; if however  $c_p \in (p, p + 1]$ , then

$$pa_p + \sqrt{p} e^{p/p} a_2 \le c_p a_p + \sqrt{c_p} e^{p/c_p} ea_2 \asymp a_p g_p(\alpha_p)$$

by (6.16). Thus, the lemma is true whenever  $q \in I_p$ .

Consider now the two cases when  $q \notin I_p$ .

First, suppose that

$$(6.17) \qquad \qquad q < \min I_p = \mathrm{e}^{-p}.$$

Then  $q < q_p(\alpha)$  for all  $\alpha \in [0, 1]$ , and (6.14) implies  $g'_p < 0$  on [0, 1]. Hence,  $\check{B}_p(a_p, a_2) = a_p g_p(0) = a_p$ ; on the other hand,  $\hat{B}_p(a_p, a_2) \leq a_p + e^p a_2 = (1 + e^p q) a_p \leq 2a_p$  by (6.17). Thus, the lemma is true in the case (6.17), too.

Finally, let

(6.18) 
$$q > \max I_p(>\sqrt{p}/e)$$

Then  $q > q_p(\alpha)$  for all  $\alpha \in [0,1]$ , and (6.11) implies  $\check{B}_p(a_p, a_2) = a_p g_p(1) \ge \sqrt{p} a_2$ , whereas  $\hat{B}_p(a_p, a_2) \le p a_p + \sqrt{p} a_2 = (p/q + \sqrt{p} e) a_2 \le 2e\sqrt{p} a_2$ , in view of (6.18).

LEMMA 6.7. For all p > 2,  $a_p > 0$ ,  $a_2 > 0$ , one has  $B_{p,ind,0}(a_p, a_2) \succeq \check{B}_p(a_p, a_2)$ .

PROOF. By Lemma 6.4 and the estimate

(6.19) 
$$\Gamma_{j,m}^{1/(2m)} \asymp j^{1-j/(2m)}$$

[see (9) in Pinelis and Utev (1984)],

(6.20) 
$$B_{p,ind,0}(a_p, a_2) \succeq a_p + B_p(a_p, a_2),$$

where

$$\widetilde{B}_p(a_p, a_2) = \max\{j^{1-j/2m}a_p^{1-\alpha_j}a_2^{\alpha_j}: j = 1, \dots, m\},\$$

 $\alpha_j := 1 - \frac{2(m-j)}{p-2}$ , *m* is the integer part of p/2. Since  $j \succeq p\alpha_j + 1$  and  $j/(2m) \le \alpha_j/2 + 1/p$ , one has  $j^{1-j/2m} \succeq (p\alpha_j + 1)^{1-\alpha_j/2}$ ,  $j = 1, \ldots, m$ . Therefore,

(6.21) 
$$\widetilde{B}_p(a_p, a_2) \succeq \widetilde{\widetilde{B}}_p(a_p, a_2) := \max\{a_p g_p(\alpha_j): j = 1, \dots, m\},$$

where  $g_p$  is defined by (6.10).

Consider first the case  $q \in I_p$ , where q and  $I_p$  are given by (6.10) and (6.13), resp. We can assume that p > 3, because for  $p \in [2,3]$ ,  $\check{B}_p(a_p, a_2) \preceq a_p + a_2 \preceq B_{p,ind,0}(a_p, a_2)$  in view of the Rosenthal's lower bound. Then,

$$0 \le \alpha_1 < \frac{2}{p-2} \le \frac{1}{p}, \qquad 0 < \alpha_{j+1} - \alpha_j = \frac{2}{p-2} \le \frac{1}{p};$$

besides,  $\alpha_m = 1$ . Note also that  $q \in I_p$  implies  $|\ln q| \leq p$ . This, along with (6.14), (6.11), leads to  $|g'_p(\alpha)| \leq pg_p(\alpha)$ , and hence to  $g_p(\alpha + \delta) \approx g_p(\alpha)$  if  $0 \leq \alpha \leq \alpha + \delta \leq 1$  and  $\delta \leq 1/p$ . Thus,

$$\check{B}_p(a_p, a_2) = \max\{a_p g_p(\alpha): 0 \le \alpha \le 1\} \asymp \max\{a_p g_p(\alpha_j): j = 1, \dots, m\},\$$

and, in virtue of (6.20), (6.21), the lemma is true if  $q \in I_p$ .

Let us now consider the case (6.17). Then, as at the end of the proof Lemma 6.6, we see that  $\check{B}_p(a_p, q_2) = a_p g_p(0) = a_p$ , and (6.20) implies the lemma.

Finally, suppose that (6.18) takes place. Then  $\widetilde{\widetilde{B}}_p(a_p, a_2) = a_p g_p(1) = \check{B}_p(a_p, a_2)$ , and now (6.20) and (6.21) imply the lemma.

LEMMA 6.8. For all  $p > 2, a_p > 0, a_2 > 0$ , one has  $B_{p,ind,0,S}(a_p, a_2) \succeq \check{B}_p(a_p, a_2)$ .

PROOF differs from that of Lemma 6.7 only in that we have to refer to Lemma 6.5 instead of Lemma 6.4.  $\hfill\blacksquare$ 

LEMMA 6.9. For all  $p \ge 2, a_p > 0, a_2 > 0$ , one has  $B_p^*(a_p, a_2) \asymp \check{B}_p(a_p, a_2)$ .

PROOF. Consider first the case  $q \in I_p$ , where q and  $I_p$  are defined in (6.10) and (6.13), resp. Then (6.15) implies

(6.22) 
$$\check{B}_p(a_p, a_2) = a_p g_p(\alpha_p) \asymp c_p a_p$$

where  $g_p$  is given by (6.10),  $\alpha_p = q_p^{-1}(q)$ ,  $c_p = p\alpha_p + 1$ . Putting  $z = (p + 1/2)/c_p$ , one has z > 1/2. We can rewrite the equation  $\alpha_p = q_p^{-1}(q)$  as  $\sqrt{z} e^z = \sqrt{e(p+1/2)} a_p/a_2$ . Since  $e^z/2 < \sqrt{z} e^z < e^{2z}$  when z > 1/2, we deduce  $z \approx \ln(2 + a_p\sqrt{p}/a_2)$ ,

(6.23) 
$$c_p \asymp \frac{p}{\ln(2 + \frac{a_p}{a_2}\sqrt{p})}$$

This implies  $\check{B}_p \preceq B_p^*$  if  $q \in I_p$ .

In the cases (6.17) and (6.18),  $\check{B}_p(a_p, a_2)$  equals  $a_p g_p(0) = a_p$  and  $a_p g_p(1) \approx \sqrt{p} a_2$ , resp. Taking into account (6.22), (6.23), (6.12), we see that in any case,  $\check{B}_p \preceq B_p^*$ .

It remains to prove that  $\check{B}_p \succeq B_p^*$ . By Lemma 6.6,  $a_p + \sqrt{p}a_2 \preceq \check{B}_p(a_p, a_2)$ since  $\sqrt{c}e^{p/c} \ge \sqrt{p}$  if  $1 \le c \le p$ . Thus [in view of (6.22), (6.23) for  $q \in I_p$ ], we only need to show that in the cases (6.17) and (6.18),

(6.24) 
$$\frac{pa_p}{\ln\left(2+\sqrt{p}\frac{a_p}{a_2}\right)} \preceq \check{B}_p(a_p, a_2).$$

If (6.17) is true, then, as it was said above in this proof,  $\check{B}_p(a_p, a_2) = a_p$ ; on the other hand,  $\frac{a_2}{a_p} = q < e^{-p}$ , and so,  $\ln\left(2 + \sqrt{p}\frac{a_p}{a_2}\right) > \ln(2 + \sqrt{p}e^p) > p$ ; now, (6.24) follows.

If (6.18) is true, then  $\check{B}_p(a_p, a_2) \approx \sqrt{p}a_2$ ; on the other hand,  $\frac{a_2}{a_p} = q > \sqrt{p}/e$ ,

$$\frac{pa_p}{\ln\left(2+\sqrt{p}\frac{a_p}{a_2}\right)} < \frac{e}{\ln 2} a_2 \sqrt{p},$$

and, again, (6.24) follows.

PROOF OF THEOREM 6.1. Observe that

$$B_p \succeq B_{p,ind,0} \succeq \check{B}_p \succeq \hat{B}_p \succeq B_p;$$

the first inequality in this chain is trivial; then we successively apply Lemma 6.7, Lemma 6.6 and (4.1). Hence,

(6.25) 
$$B_p \asymp B_{p,ind,0} \asymp \check{B}_p \asymp \hat{B}_p.$$

Note that for any separable Banach space  $\mathcal{X}$ ,

$$(6.26) B_{p,ind,0} \preceq B_{p,ind}^{\mathcal{X}}$$

since **R** may be isometrically embedded into  $\mathcal{X}$  and, for  $f \in \mathcal{M}_{ind}(\mathbf{R})$ ,

$$|| |f_n + x| - \mathbf{E} |f_n + x| ||_p \to ||f_n||_p \text{ as } x \to \infty.$$

It follows from Theorem 5.1 that

$$(6.27) B_{p,ind}^{\mathcal{X}} \preceq \hat{B}_p.$$

Now (6.25)-(6.27) yield

$$B_p \asymp B_{p,ind}^{\mathcal{X}} \asymp B_{p,ind,0} \asymp \check{B}_p \asymp \check{B}_p.$$

Analogously,

$$B_{p,S} \asymp B_{p,ind,S}^{\mathcal{X}} \asymp B_{p,ind,0,S} \asymp \check{B}_p \asymp \hat{B}_p$$

here, we need to refer to Lemma 6.8 instead of Lemma 6.7 and also to  $||d^*||_p \le ||S_p||_p$ . It remains to recall Lemma 6.9.

PROOF OF THEOREM 6.2. For any given  $c \in [1, p]$ , choose  $a_p > 0$ ,  $a_2 > 0$ so that  $a_2/a_p = \sqrt{c} e^{-p/c}$ . Then  $ca_p = \sqrt{c} e^{p/c}a_2$ . Put  $\alpha = (c-1)/p$ , so that  $c=p\alpha+1,\,0\leq\alpha\leq1.$  Then

$$\check{B}_p(a_p, a_2) \ge a_p g_p(\alpha) = a_p c e^{-p\alpha/c} \ge e^{-1} c a_p$$
  
=  $(2e)^{-1} (c a_p + \sqrt{c} e^{p/c} a_2),$ 

where  $g_p$  is defined by (6.10). It remains to apply Theorem 6.1 and also to recall that inequalities (4.2)-(4.4), (5.2)-(5.4), (5.8)-(5.10) were obtained by choosing particular values of c.

PROOF OF PROPOSITION 6.3. For any given  $\alpha \in [0,1]$ , choose  $a_p > 0$ ,  $a_2 > 0$  so that

(6.28) 
$$\frac{a_2}{a_p} = q_p(\alpha),$$

where  $q_p(\alpha)$  is defined by (6.11). Then, putting  $c = p\alpha + 1$ , we see that  $1 \le c \le p + 1$  and, in view of (6.10), (6.28),

$$a_p g_p(\alpha) = a_p c \exp\left\{\frac{p\alpha}{2} \frac{\alpha - 2}{p\alpha + 1}\right\} \approx a_p c \approx c a_p + \sqrt{c} e^{p/c} a_2$$

since, in consequence of (6.28), (6.11),

$$\sqrt{c} e^{p/c} a_2 = c a_p \exp\left\{\frac{p\alpha}{2(p\alpha+1)}\right\} \asymp c a_p.$$

Thus, if the defined above  $c \leq p$ , we see that

$$(6.29) \qquad \qquad \hat{B}_p \preceq a_p g_p(\alpha)$$

If however  $c \in (p, p+1]$ , then

$$a_p g_p(\alpha) \asymp c a_p + \sqrt{c} \operatorname{e}^{p/c} a_2 \ge p a_p + \sqrt{p} \operatorname{e}^{p/(p+1)} a_2 \succeq \hat{B}_p,$$

so that (6.29) holds. Now it remains to apply Theorem 6.1.

#### 7. Chung type bounds on moments. Consider

$$B_p^{Ch}(a_p, n) := \sup\{ \| \|f_n\| \|_p: (0, f_1, \dots, f_n, f_n, f_n, \dots) \in \mathcal{M}(\mathcal{X}),$$
  
*n* is fixed,  $D \|S_p\|_p = a_p, \quad \mathcal{X} \text{ is } (2, D) \text{-smooth} \}.$ 

THEOREM 7.1. For all  $p > 2, a_p > 0, n = 1, 2, ...,$  one has

$$B_p^{\operatorname{Ch}}(a_p, n) \asymp \sqrt{p \wedge n} \, n^{\frac{p-2}{2p}} a_p.$$

PROOF. Let  $(0, f_1, \ldots, f_n, f_n, f_n, f_n, \ldots) \in \mathcal{M}(\mathcal{X})$ . Using Hölder's inequality two times, we see that  $\|s_2\|_p \leq n^{(p-2)/(2p)} \|S_p\|_p$ . If  $p \leq 3n^{(p-2)/p}$ , then  $p\|d^*\|_p \leq$  $p\|S_p\|_p \leq p^{1/2} n^{(p-2)/2p} \|S_p\|_p$ , and hence, (4.2) and  $p \leq 3n$  yield

(7.1) 
$$B_p^{\mathrm{Ch}}(a_p, n) \preceq \sqrt{p \wedge n} \, n^{\frac{p-2}{2p}} a_p$$

If now  $p > 3n^{(p-2)/p}$ , then p > 3,  $n < p^{p/(p-2)} \preceq p$ . The inequalities of Minkowsky and Hölder give  $|| ||f_n|| ||_p \leq \sum_{j=1}^n || ||d_j|| ||_p \leq n^{(p-1)/p} ||S_p||_p$ ; this and  $p \succeq n$  show that (7.1) holds. It remains to prove that

(7.2) 
$$B_p^{\mathrm{Ch}}(a_p, n) \succeq \sqrt{p \wedge n} n^{\frac{p-2}{2p}} a_p.$$

Let  $d_j$ 's be independent,  $\mathbf{Pr}(d_j = \pm u) = 1/2, u > 0, j = 1, \dots, n$ . Then the multinomial formula yields

(7.3) 
$$\|f_n\|_{2m} = \left(\sum_{j=1}^{m\wedge n} \binom{n}{j} j! \Gamma_{j,m}\right)^{\frac{1}{2m}} u \ \asymp \left(\sum_{j=1}^{m\wedge n} \binom{n}{j} j^{2m}\right)^{\frac{1}{2m}} u,$$

where  $\Gamma_{j,m}$  is defined by (6.7), m = 1, 2, ..., and (6.19) is used. Let m be the integer part of p/2. If  $m \leq n$ , then (7.3) shows that

$$\|f_n\|_p \ge \|f_n\|_{2m} \ge \binom{n}{m}^{\frac{1}{2m}} mu \ge \left(\frac{n}{m}\right)^{\frac{1}{2}} mu = \sqrt{m} n^{(p-2)/2p} \|S_p\|_p$$

which implies (7.2) in the case  $m \leq n$ . If, finally, m > n, then it follows from (7.3) that

$$||f_n||_p \ge ||f_n||_{2m} \succeq {\binom{n}{n}}^{\frac{1}{2m}} nu = n^{(p-1)/p} ||S_p||_p,$$

so that (7.2) holds.

Chung (1951), pp.348-349, showed that in the case  $f \in \mathcal{M}_{ind}(\mathbf{R})$ , the inequality of Marcinkiewicz and Zygmund (1937), p. 87, implies an estimate like (7.1) but with some  $C_0(p)$  depending only on p, instead of  $\sqrt{p \wedge n}$ . As it was pointed out in Dharmadhikari, Fabian and Jogdeo (1968), an analogous result for  $f \in \mathcal{M}(\mathbf{R})$  is implied by the generalization of the Marcienkiewicz-Zygmund inequality obtained by Burkholder (1966), Theorem 9 [see also Burkholder (1973), Theorems 3.2, 15.1, or (9.2) below]; using Proposition 9.1, one can see that Theorem 15.1 in Burkholder (1973) in fact gives  $C_0(p) = p$ ; a constant of the same order p is given in Theorem 3.2 of Burkholder (1973).

The direct proof due to Dharmadhihari, Fabian and Jogdeo (1968) yields  $C_0(p) = p2^p$ . For  $f \in \mathcal{M}_{ind}(\mathbf{R})$ , Dharmadhikari and Jogdeo (1969) obtained  $C_0(p) = p/\ln p$ .

For  $f \in \mathcal{M}_{ind}(\mathbf{R})$ , the result of Whittle (1960) implies the Marcinkiewicz-Zygmund inequality with the best constant and, along with the mentioned above Chung's remark (see also Rosén (1970)), leads to (7.1) but with  $\sqrt{p}$  instead of  $\sqrt{p \wedge n}$ ; so, for n > p, it gives the optimum.

What has been said is a reason for the referring to (7.1) as to an optimum bound on moments of the Chung type.

REMARK. Bounds of the Chung type on central moments of the norm of the sum of independent random vectors in any separable Banach space can be easily derived from Theorem 7.1 (cf. Theorem 5.1).  $\blacksquare$ 

8. One-sided bounds for the distributions of real-valued (super)martingales. Let  $\mathcal{M}_{-}$  stand for the set of all real-valued supermartingales  $f \in \mathcal{S}(\mathbf{R})$ . For  $f \in \mathcal{M}_{-}$ , put  $f_{+}^{*} = \sup_{j} f_{j}, d_{+}^{*} = \sup_{j} d_{j}$ .

THEOREM 8.1. If  $f \in \mathcal{M}_{-}$  and  $\lambda > 0$  is such that  $\mathbf{E}e^{\lambda d_{j}} < \infty$  for each j,

then for all  $r \geq 0$ ,

$$\begin{aligned} \mathbf{Pr}\left(f_{+}^{*} \geq r\right) &\leq \mathrm{e}^{-\lambda r} \left\| \prod_{j=1}^{\infty} (1+e_{j}) \right\|_{\infty} \\ &\leq \exp\left\{ -\lambda r + \left\| \sum_{j=1}^{\infty} e_{j} \right\|_{\infty} \right\}, \end{aligned}$$

where  $e_j := \mathbf{E}_{j-1}(e^{\lambda d_j} - 1 - \lambda d_j).$ 

PROOF follows from the trivial remark that the sequence:  $G_0 := 1$ ,  $G_j := e^{\lambda f_j} \prod_{i=1}^j (1+e_i)^{-1}, \ j = 1, 2, \ldots$ , is a positive supermartingale [cf. the end of the proof of Theorem 3.2].

THEOREM 8.2. Suppose that  $f \in \mathcal{M}_-$ ,  $||d_+^*||_{\infty} \leq a$ ,  $||s_2||_{\infty} \leq b$  for some a > 0, b > 0. Then for all  $r \geq 0$ ,

$$\mathbf{Pr}\left(f_{+}^{*} \geq r\right) \leq \exp\left[\frac{r}{a} - \left(\frac{r}{a} + \frac{b^{2}}{a^{2}}\right)\ln\left(1 + \frac{ra}{b^{2}}\right)\right] \leq \left(\frac{\mathrm{e}b^{2}}{ra}\right)^{r/a}.$$

PROOF is quite similar to that of Theorem 3.4.

THEOREM 8.3. If  $f \in \mathcal{M}_{-}$ ,  $p \geq 2$ ,  $1 \leq c \leq p$ , then

$$||f_+^*||_p \leq c ||d_+^*||_p + \sqrt{c} e^{p/c} ||s_2||_p.$$

We use

LEMMA 8.4. (cf. Lemma 4.2.) If  $\lambda > 0$ ,  $\delta_1 > 0$ ,  $\delta_2 > 0$ ,  $\beta - 1 - \delta_2 > 0$ ,  $f \in \mathcal{M}_-$ , then

$$\mathbf{Pr}\left(f_{+}^{*} > \beta\lambda, w_{+}^{*} \leq \lambda\right) \leq \varepsilon \mathbf{Pr}\left(f_{+}^{*} > \lambda\right),$$

where

$$w_{+}^{*} = \left(\frac{d_{+}^{*}}{\delta_{2}}\right) \lor \left(\frac{s_{2}}{\delta_{1}}\right),$$
$$\varepsilon = \left(\frac{e}{N}\frac{\delta_{1}^{2}}{\delta_{2}^{2}}\right)^{N}, \quad N = \frac{\beta - 1 - \delta_{2}}{\delta_{2}}.$$

PROOF. Put  $\bar{d}_j = d_j I\{d_j \le \delta_2 \lambda\}, \ \bar{f}_j = \bar{d}_0 + \dots + \bar{d}_j, \ \bar{s}_{2,j+1} = (\sum_{i=1}^{j+1} \mathbf{E}_{i-1} \bar{d}_i^2)^{1/2},$ 

 $\bar{h}_j = \bar{f}_{(j \wedge \tau \wedge \nu) \vee \mu} - \bar{f}_{\mu}, \ j = 0, 1, \dots,$ where  $\mu = \inf\{j: \ \bar{f}_j > \lambda\}, \quad \nu = \inf\{j: \ \bar{f}_j > \beta\lambda\},$  $\sigma = \inf\{j: \ \bar{s}_{2,j+1} > \delta_1\lambda\}.$  Then  $(\bar{h}_j)$  is a supermartingale conditionally on  $F_{\mu}$ , and

$$\begin{aligned} \mathbf{Pr}\left(f_{+}^{*} > \beta\lambda, w_{+}^{*} \leq \lambda\right) &= \mathbf{Pr}\left(\bar{f}_{+}^{*} > \beta\lambda, w_{+}^{*} \leq \lambda\right) \\ &\leq \mathbf{Pr}\left(\bar{h}_{+}^{*} > (\beta - 1 - \delta_{2})\lambda\right) \\ &= \mathbf{E}\,\mathbf{Pr}\left(h_{+}^{*} > (\beta - 1 - \delta_{2})\lambda \mid F_{\mu}\right)I\{\mu < \infty\} \\ &= \varepsilon\mathbf{Pr}\left(\bar{f}_{+}^{*} > \lambda\right) \leq \varepsilon\mathbf{Pr}\left(f_{+}^{*} > \lambda\right); \end{aligned}$$

here, we put  $\bar{f}^*_+ = \sup_j \bar{f}_j$ ,  $\bar{h}^*_+ = \sup_j \bar{h}_j$  and took into account Theorem 8.2.

PROOF OF THEOREM 8.3 is similar to that of Theorem 4.1, but simpler. Here, we do not need to symmetrize; instead, we can apply Lemma 7.1 of Burkholder (1973) directly to  $f_{+}^{*}$ .

The following is a refinement of Theorem 8.1.

THEOREM 8.5. Let  $f \in \mathcal{M}(\mathbf{R}), d_j = u_j - \mathbf{E}_{j-1}u_j, \lambda > 0, \mathbf{E}e^{\lambda u_j} < \infty,$  $j = 1, 2, \dots$ . Then for all  $r \ge 0$ ,

$$\mathbf{Pr}\left(f_{+}^{*} \geq r\right) \leq \exp\left\{-\lambda r + \left\|\sum_{j=1}^{\infty} e_{j}\right\|_{\infty}\right\},\,$$

where  $e_j = \mathbf{E}_{j-1}(e^{\lambda u_j} - 1 - \lambda u_j)$ . If, besides,  $\mathbf{E}_{j-1}u_j \ge 0$  for all  $j = 1, 2, \dots$ , then, moreover,

$$\mathbf{Pr}\left(f_{+}^{*} \geq r\right) \leq e^{-\lambda r} \left\| \prod_{j=1}^{\infty} (1+e_{j}) \right\|_{\infty}$$

PROOF is analogous to that in Pinelis and Sakhanenko (1985) but simpler. The elementary inequalities

$$(1 + a + b)e^{-b} \le e^a$$
  $(a \in \mathbf{R}, b \in \mathbf{R}),$   
 $(1 + a + b)e^{-b} \le 1 + a$   $(a \ge 0, b \ge 0)$ 

imply, resp., that

$$\mathbf{E}_{j-1} \mathrm{e}^{\lambda d_j} \le \mathrm{e}^{e_j},$$

and, if  $\mathbf{E}_{j-1}u_j \geq 0$ ,

$$\mathbf{E}_{j-1}\mathrm{e}^{\lambda d_j} \le 1 + e_j,$$

 $j = 1, 2, \ldots$ , if one chooses  $a = e_j, b = \lambda \mathbf{E}_{j-1} u_j$ . Hence,

$$\exp\left\{\lambda f_j - \sum_{i=1}^j e_j\right\}, \qquad j = 0, 1, 2, \dots,$$

is a supermartingale and, if  $\mathbf{E}_{j-1}u_j \ge 0 \quad \forall j$ , so is

$$e^{\lambda f_j} \prod_{i=1}^j (1+e_i)^{-1}, \qquad j=0,1,2,\dots$$

It remains to use reasoning like that at the end of the proof of Theorem 3.2.

REMARK. Martingales like those in Theorem 8.5 may arise, e.g., as a result of truncating and subsequent centering of the increments of other martingales. The aim of Theorem 8.5 is to provide for the best constants in exponential inequalities for martingales, which cannot be reached via, e.g., the straightforward estimate  $|d_j| \leq |u_j| + |\mathbf{E}_{j-1}u_j|$ .

As illustration, let us give the following corollaries to Theorem 8.5, which are refinements of Theorems 3.3, 3.4, resp., for the particular case  $\mathcal{X} = \mathbf{R}$ .

THEOREM 8.6. Suppose that  $f \in \mathcal{M}(\mathbf{R}), d_j = u_j - \mathbf{E}_{j-1}u_j, j = 1, 2, \dots,$ and

$$\left\|\sum_{j=1}^{\infty} \mathbf{E}_{j-1} |u_j|^m\right\|_{\infty} \le m! \Gamma^{m-2} B^2/2$$

for some  $\Gamma > 0$ , B > 0 and  $m = 2, 3, \ldots$ . Then for all  $r \ge 0$ ,

$$\mathbf{Pr}\left(f_{+}^{*} \geq r\right) \leq \exp\left(-\frac{r^{2}}{B^{2} + B\sqrt{B^{2} + 2\Gamma r}}\right).$$

PROOF is almost literally the same as that of Theorem 3.3, but rests upon Theorem 8.5, instead of Theorem 3.1.  $\blacksquare$ 

THEOREM 8.7. Suppose that  $f \in \mathcal{M}(\mathbf{R}), d_j = u_j - \mathbf{E}_{j-1}u_j, j = 1, 2, ..., \text{ and}$  $\|u^*\|_{\infty} \leq a, \left\|\sum_{j=1}^{\infty} \mathbf{E}_{j-1}u_j^2\right\|_{\infty} \leq b^2 \text{ for some } a > 0, b > 0. \text{ Then for all } r \geq 0,$  $\mathbf{Pr}\left(f_+^* \geq r\right) \leq \exp\left[\frac{r}{a} - \left(\frac{r}{a} + \frac{b^2}{a^2}\right)\ln\left(1 + \frac{ra}{b^2}\right)\right]$  $\leq \left(\frac{eb^2}{ra}\right)^{r/a}.$ 

PROOF differs from that of Theorem 3.4 only in that we use Theorem 8.5 in place of Theorem 3.1.  $\blacksquare$ 

If, instead of Theorem 8.5, we had used the "naive" estimate  $|d_j| \leq |u_j| + |\mathbf{E}_{j-1}u_j|$  and, say, Theorem 3.1, we would have hardly been able to obtain inequalities better than ones like those in Theorems 8.6, 8.7 but with  $2\Gamma$ , 2B, 2a, 2b in place of  $\Gamma$ , B, a, b, resp. The gain provided, e.g., by Theorem 8.6 is quite significant. If, say,

$$\exp\left(-\frac{r^2}{(2B)^2 + 2B\sqrt{(2B)^2 + 2(2\Gamma)r}}\right) = 10^{-1},$$

then the bound given in Theorem 8.6 varies from  $10^{-2\sqrt{2}} < (1.5)10^{-3}$  (when  $\Gamma r$  is much greater than  $B^2$ ) to  $10^{-4}$  (when  $B^2$  is much greater than  $\Gamma r$ ).

**9. Appendix.** Here, we find explicitly the best constants in the mentioned in the preceding sections bounds due to Burkholder (1973) obtainable via the methods used therein. Besides, we prove that, at least for the sums of independent random vectors in Hilbert spaces, the bound on moments found in Pinelis (1980) is equivalent to (4.1).

It follows from Theorems 15.1 and 21.1 of Burkholder (1973) that for  $f \in \mathcal{M}(\mathbf{R}), p \geq 1$ ,

(9.1) 
$$||f^*||_p \leq C_1(||d^*||_p + ||s_2||_p),$$

(9.2) 
$$||f^*||_p \leq (C_2 + p) ||S_2||_p;$$

analysis of the proof in Burkholder (1973) shows that one can put

$$C_{i} = \inf \left\{ \frac{\beta/\delta}{[1 - i\beta^{p}\delta^{2}/(\beta - 1 - \delta)^{2}]^{1/p}}; \ \beta - 1 - \delta > 0, \ \delta > 0, \\ \frac{i\beta^{p}\delta^{2}}{(\beta - 1 - \delta)^{2}} < 1 \right\}, \quad i = 1, 2$$

PROPOSITION 9.1. For  $i = 1, 2, p \ge 1$ , one has  $C_i \asymp p$ .

PROOF. Suppose that  $\beta - 1 - \delta > 0$ ,  $\delta > 0$ ,  $i\beta^p \delta^2/(\beta - 1 - \delta)^2 < 1$ , i = 1or 2. Then  $\beta^{p-2}\delta^2 < 1$ , and we see that  $C_1 > \beta/\delta > \beta^{p/2}$ . If  $\beta \ge 2$ , then  $C_1 > 2^{p/2} \succeq p$ . Assume now that  $\beta < 2$ . Put  $c = p\delta$ ,  $k = (\beta - 1 - \delta)/\delta$ . Then  $\beta = 1 + (k + 1)c/p \ge e^{(k+1)c/2p}$  since  $1 + \alpha > e^{\alpha/2}$  for  $0 < \alpha < 1$ . Hence,  $1 > \beta^p \delta^2/(\beta - 1 - \delta)^2 = \beta^p/k^2 \ge k^{-2}e^{(k+1)c/2}$ ,  $c < 4(k+1)^{-1}\ln k < 4$ ,  $\delta = c/p < 4/p$ ; therefore,

$$(9.3) C_1 > 1/\delta > p/4$$

On the other hand,  $\beta \leq e^{(k+1)c/p}$  since  $1 + \alpha \leq e^{\alpha}$ , and so, for k = 3, c = 1/4, one has  $\beta \leq e$ ,  $2\beta^p \delta^2/(\beta - 1 - \delta)^2 = 2\beta^p/k^2 \leq 2e^{(k+1)c}/k^2 < 2/3$ . Hence,

(9.4) 
$$C_2 \le 3\beta/\delta < 3e/\delta = 3ep/c = 12ep.$$

Note that  $C_1 \leq C_2$ . Now (9.3), (9.4) imply the proposition.

Finally, we give an interpretation of results of Pinelis (1980). Let  $\mathcal{F}(p)$ , p > 0 stand for the set of all the pairs  $(c_1, c_2)$  that may be represented in the form

(9.5) 
$$c_1 = \sum_{i=0}^{m-1} \frac{c_i(p)b_i^{-2i}}{i!},$$

(9.6) 
$$c_2 = \frac{\tilde{c}_m(p)}{m!} + \sum_{i=0}^{m-1} \frac{c_i(p)b_i^{p-2i-2}}{i!},$$

where m is the integer part of p/2,  $b_i > 0$ ,

(9.7) 
$$c_i(p) = v(p-2i)2^{-i}(2i)! \binom{p}{2i} \prod_{k=0}^{i-1} u(p-2k),$$

(9.8) 
$$\tilde{c}_m(p) = 2^{-m} (2m)! {p \choose 2m} \prod_{i=0}^{m-1} u(p-2i),$$
$${p \choose k} = \frac{p(p-1)\dots(p-(k-1))}{k!},$$

and  $u(\cdot), v(\cdot)$  are any functions such that

$$(9.9) u(2) + v(2) \ge 1, \quad u(2) > 0, \quad v(2) > 0,$$

(9.10) 
$$u(s) \ge 1, \quad v(s) \ge 1 \quad \text{for} \quad s > 2,$$

(9.11)  $u(s)^{\frac{1}{3-s}} + v(s)^{\frac{1}{3-s}} \le 1 \text{ for } s > 3.$ 

For  $a_p > 0, a_2 > 0, p > 0$ , put

$$B_{p,1980}(a_p, a_2) = \inf\{(c_1 a_p^p + c_2 a_2^p)^{1/p} \colon (c_1, c_2) \in \mathcal{F}(p)\}.$$

In Pinelis (1980), bounds on moments of martingales in Hilbert spaces with bounded conditional second moments were found. If applied, in particular, to any  $f \in \mathcal{M}_{ind}(\mathcal{X})$ , where  $\mathcal{X}$  is any separable Hilbert space, the Corollary in Pinelis (1980) states that

$$|| || f_n || ||_p \le B_{p,1980}(||S_p||_p, ||S_2||_2), \quad p > 0.$$

In turn, this implies (cf. Theorems 5.1, 6.1)

(9.12) 
$$B_{p,ind,S}(a_p, a_2) \le 2 B_{p,1980}(a_p, a_2)$$

for any  $a_p > 0, a_2 > 0, p > 0$  and any separable Banach space  $\mathcal{X}$ .

PROPOSITION 9.2. For  $a_p > 0, a_2 > 0, p > 2$ ,

$$B_{p,1980}(a_p, a_2) \simeq B_p(a_p, a_2).$$

In view of Theorem 6.1, any of the following functions:  $B_p^*$ ,  $B_{p,ind}^{\mathcal{X}}$ ,  $B_{p,ind,0,S}$ ,  $B_{p,ind,0,S}$ ,  $\check{B}_p$ ,  $\hat{B}_p$  may be used here in place of  $B_p$ .

PROOF. For  $s \ge 0$ , put

$$(9.13) y_s = g_s(\alpha_s) = \max\{g_s(\alpha): \ 0 \le \alpha \le 1\}, 0 \le \alpha_s \le 1,$$

and, for  $s \geq 2$ ,

(9.14) 
$$u(s) = \frac{\mathrm{e}^3 y_s^s}{sq^2 y_{s-2}^{s-2}}, \quad v(s) = 10^s y_s^{s-2},$$

where (cf. (6.10))

(9.15) 
$$g_s(\alpha) = (s\alpha + 1)^{1-\alpha/2}q^{\alpha}, \quad q = a_2/a_p.$$

Note that for  $s \ge 0$ ,

(9.16) 
$$y_s \ge g_s(0) \lor g_s(1) = 1 \lor (s+1)^{1/2} q.$$

Besides,  $y_s$  is increasing in s because so is  $g_s(\alpha)$ . Hence, for  $s \ge 2$ , one has  $u(s) \ge e^3 y_s^2/(sq^2) \ge e^3 > 1$ ,  $v(s) \le 2 \cdot 10^s y_s^s/(1+q^2)$ ,  $v(s) \ge (10^{s-3} y_s^{s-3}) \lor 1$ , and so, for s > 3,

(9.17) 
$$u(s)^{\frac{1}{3-s}} + v(s)^{\frac{1}{3-s}} \le \left[\frac{sq^2}{e^3y_s^2}\right]^{\frac{1}{s}} + \frac{1}{10y_s}.$$

Consider first the case  $q \in I_s$ , where  $I_s$  is defined by (6.13). Then,  $y_s = g_s(\alpha_s) > e^{-1}(s\alpha_s + 1)$  [cf. (6.15)], and, in view of (6.11),

$$q = q_s(\alpha_s) < \sqrt{s\alpha_s + 1} \exp\left\{\frac{1}{2} - \frac{s}{s\alpha_s + 1}\right\} < \sqrt{y_s} e^{1 - s/(y_s e)};$$

hence,

$$\left(\frac{sq^2}{e^3y_s^2}\right)^{\frac{1}{s}} < \left(\frac{s}{ey_s}\right)^{\frac{1}{s}} e^{-1/(y_s e)} \le \exp\left\{\frac{1}{e^2y_s} - \frac{1}{ey_s}\right\} < 1 - \frac{1}{10y_s}$$

since  $(s/a)^{1/s} \leq e^{1/(ae)}$  for a > 0 and  $e^{-\alpha} < 1 - \alpha/2$  for  $0 < \alpha < 1$ . Thus, (9.17) yields (9.11) if  $q \in I_s$ .

If  $q < \min I_s = e^{-s}$ , then  $y_s = g_s(0) = 1$ , and

$$\left(\frac{sq^2}{e^3y_s^2}\right)^{\frac{1}{s}} < \left(\frac{s}{e^3}\right)^{\frac{1}{s}} e^{-2} \le e^{e^{-4}-2} < e^{-1},$$

so that (9.17) again yields (9.11).

Now suppose that  $q > \max I_s(> [(s+1)/e]^{1/2})$ . Then  $y_s = g_s(1) = (s+1)^{1/2}q > s^{1/2}q$ , and for s > 3,  $[(sq^2)/(e^3y_s^2)]^{\frac{1}{s}} < e^{-3/s} < 1 - 1/s$ ,

$$\frac{1}{10y_s} = \frac{1}{10\sqrt{s+1}\,q} < \frac{\sqrt{e}}{10(s+1)} < \frac{1}{s},$$

so that (9.17) once more yields (9.11).

Thus, u(s) and v(s) defined by (9.14) satisfy (9.9)-(9.11).

Let us write  $\mathcal{E}_1 \ll \mathcal{E}_2$  if  $\mathcal{E}_1^{1/p} \preceq \mathcal{E}_2^{1/p}$ , p > 2.

Using (9.7), (9.14), (9.16) and the Stirling's formula, we see that for p>2, $i=1,\ldots,m-1,$ 

(9.18) 
$$\frac{c_i(p)}{i!} \ll y_p^p y_{p-2i}^{-2} q^{-2i} \ll y_p^p (1+q)^{-2} q^{-2i}$$
$$\leq y_p^p q^{-2pi/(p-2)};$$

similarly, (9.8), (9.14), (9.16) imply

(9.19) 
$$\frac{\tilde{c}_m(p)}{m!} \ll y_p^p q^{-2m} y_{p-2m}^{-(p-2m)} \le y_p^p q^{-p}.$$

Choose now  $b_i \equiv q^{-p/(p-2)}$  so that  $b_i^{-2i}a_p^p = b_i^{p-2i-2}a_2^p$ , i = 1, ..., m, in accordance with (9.15); then (9.5), (9.6), (9.18), (9.19), (9.13), (6.12) yield  $(c_1a_p^p + c_2a_2^p)^{1/p} \preceq y_pa_p = \check{B}_p$ . Thus,  $B_{p,1980}(a_p, a_2) \preceq \check{B}_p(a_p, a_2)$ . It remains to recall (9.12) and Theorem 6.1.

### REFERENCES

- Bennett, G. (1962). Probability inequalities for the sum of independent random variables. J. Amer. Statist. Assoc. 57 33-45.
- Berger, E. (1991). Marjorization, exponential inequalities and almost sure behavior of vector-valued random variables. Ann. Probab. 19 1206-1226.
- Billingsley, P. (1968). Convergence of probability measures. John Wiley, New York.
- Burkholder, D. (1966). Martingale transforms. Ann. Math. Statist. 37 1494-1504.
- Burkholder, D. (1973). Distribution function inequalities for martingales. Ann. Probab. 1 19-42.
- Burkholder, D. (1991). Explorations in martingale theory and its applications. Saint-Flour Lectures (1989). Lecture Notes in Mathematics 1464 1-66.
- Chung, K.L. (1951). The strong law of large numbers. Proc. Second Berkeley Symp. Math. Statist. Prob. 341-352. Univ. of California Press.
- Davis, B. (1970). On the integrability of the martingale square function. Israel J. Math. 8 187-190.
- De Acosta, A. (1981). Inequalities for *B*-valued random vectors with applications to the strong law of large numbers. *Ann. Probab.* **9**, 157-161.
- Dharmadhikari, S.W., Fabian, V. and Jogdeo, K. (1968). Bounds on the moments of martingales. Ann. Math. Statist. 39 1719-1723.
- Dharmadhikari, S.W. and Jogdeo, K. (1969). Bounds on moments of certain random variables. *Ann. Math. Statist.* **40** 1506-1508.
- Haagerup, U. (1982). The best constants in the Khintchine inequality. Stu-

dia Math. 70 232-283.

- Hall, P. and Marron, J.S. (1987). Choice of kernel order in density estimation. Ann. Statist. 16 161-173.
- Hitczenko, P. (1990). Best constants in martingale version of Rosenthal's inequality. Ann. Probab. 18 1656-1668.
- Hitczenko, P. (1990a). Upper bounds for the  $L_p$ -norms of martingales. *Probab. Th. Rel. Fields* 86 225-237.
- Hitczenko, P. (1991). Private communication.
- Hoeffding, W. (1963). Probability inequalities for sums of bounded random variables. J. Amer. Statist. Assoc. 58 13-30.
- Hoffmann-Jørgensen, J. (1974). Sums of independent Banach space valued random variables. *Studia Math.* **52** 159-186.
- Johnson, W.B., Schechtman, G. and Zinn, J. (1985). Best constants in moment inequalities for linear combination of independent and exchangeable random variables. Ann. Probab. 13 234-253.
- Kallenberg, O. and Sztencel, R. (1991). Some dimension-free features of vector-valued martingales. Probab. Th. Rel. Fields 88 215-247.
- Kwapień, S. and Szulga, J. (1991). Hypercontraction methods in moment inequalities for series of independent random variables in normed spaces. Ann. Probab. 19 369-379.
- Kwapień, S. and Woyczyński, W.A. (1989). Tangent sequences of random variables: Basic inequalities and their applications. Proc. Conf. on Almost Everywhere Convergence in Probab. and Ergodic Theory, 237-265. Academic, New York.
- Marcinkiewicz, J. and Zygmund, A. (1937), Sur les fonctions independantes. Fund. Math. **89** 60-90.
- Nagaev, S.V. and Pinelis, I.F. (1977). Some inequalities for the distributions

of sums of independent random variables. *Theory Probab. Appl.* **22** 248-256.

- Pinelis, I.F. (1974). On some inequalities for the distributions of sums of independent random variables. *Diploma work*. Novosibirsk State University. (Russian).
- Pinelis, I.F. (1978). Distributions of sums of independent random variables with values in a Banach space. *Theory Probab. Appl.* 23 608-615.
- Pinelis, I.F. (1980) Estimates of moments of infinite-dimensional martingales. Math. Notes 27 459-462.
- Pinelis, I.F. (1981). Limit theorems on large deviations for sums of infinitedimensional random variables when the Cramér's condition is violated. Manuscript deposited at All-Union Institute for Scientific and Technical Information (VINITI), no. 1674-81. (Russian).
- Pinelis, I.F. (1986). Probability inequalities for sums of independent random variables with values in a Banach space. Math. Notes 39 241-244.
- Pinelis, I.F. (1990). Inequalities for the distributions of sums of independent random vectors and their applications to the density estimation. *Theory Probab. Appl.* 35 500-512.
- Pinelis, I. (1992). An approach to inequalities for the distributions of infinite-dimensional martingales. Probability in Banach Spaces,
  8: Proceedings of the Eighth International Conference. 128-134. Birkhäuser, Boston.
- Pinelis, I. (1992a). On a distribution inequality for sums of independent random vectors. *Submitted.*
- Pinelis, I.F. and Sakhanenko, A.I. (1985). Remarks on inequalities for large deviation probabilities. *Theory Probab. Appl.* **30** 143-148.

- Pinelis, I.F. and Utev, S.A. (1984). Estimates of the moments of sums of independent random variables. *Theory Probab. Appl.* 29 574-577.
- Pinelis, I.F. and Utev, S.A. (1989). Exact exponential bounds for sums of independent random variables. *Theory Probab. Appl.* 34 340-346.
- Pisier, G. (1975). Martingales with values in uniformly convex spaces. Israel J. Math. 20 326-350.
- Rosén, B. (1970). On bounds on the central moments of even order of a sum of independent random variables. Ann. Math. Statist. 41 1074-1077.
- Rosenthal, H.P. (1970). On the subspaces of  $L^p(p > 2)$  spanned by sequences of independent random variables. *Israel J. Math.* 8 273-303.
- Sazonov, V.V. (1974). On the estimation of moments of sums of independent random variables. *Theory Probab. Appl.* **19** 371-374.
- Talagrand, M. (1989). Isoperimetric and integrability of the sum of independent Banach-space valued random variables. Ann. Probab. 17 1546-1570.
- Wang, M. (1991). Sharp square function inequlities for conditionally symmetric martngales. Trans. AMS 328 393-419.
- Whittle, P. (1960). Bounds for the moments of linear and quadratic forms in independent random variables. *Theory Probab. Appl.* 5 302-305.
- Yurinskii, V.V. (1974). Exponential bounds for large deviations. Theory Probab. Appl. 19 154-155.

DEPARTMENT OF MATHEMATICAL

SCIENCES

MICHIGAN TECHNOLOGICAL UNIVERSITY HOUGHTON, MICHIGAN 49931