

PHASE TRANSITIONS FOR p -ADIC POTTS MODEL ON THE CAYLEY TREE OF ORDER THREE

FARRUKH MUKHAMEDOV AND HASAN AKIN

ABSTRACT. In the present paper, we study a phase transition problem for the q -state p -adic Potts model over the Cayley tree of order three. We consider a more general notion of p -adic Gibbs measure which depends on parameter $\rho \in \mathbb{Q}_p$. Such a measure is called *generalized p -adic quasi Gibbs measure*. When ρ equals to p -adic exponent, then it coincides with the p -adic Gibbs measure. When $\rho = p$, then it coincides with p -adic quasi Gibbs measure. Therefore, we investigate two regimes with respect to the value of $|\rho|_p$. Namely, in the first regime, one takes $\rho = \exp_p(J)$ for some $J \in \mathbb{Q}_p$, in the second one $|\rho|_p < 1$. In each regime, we first find conditions for the existence of generalized p -adic quasi Gibbs measures. Furthermore, in the first regime, we establish the existence of the phase transition under some conditions. In the second regime, when $|\rho|_p, |q|_p \leq p^{-2}$ we prove the existence of a quasi phase transition. It turns out that if $|\rho|_p < |q - 1|_p^2 < 1$ and $\sqrt{-3} \in \mathbb{Q}_p$, then one finds the existence of the strong phase transition.

Mathematics Subject Classification: 46S10, 82B26, 12J12, 39A70, 47H10, 60K35.

Key words: p -adic numbers, Potts model; p -adic quasi Gibbs measure, phase transition.

1. INTRODUCTION

It is known that (see [75]) the q -state Potts model is one of the most studied models in statistical mechanics. It has wide theoretical interest and practical applications. Originally, the Potts model was introduced as a generalization of the Ising model to more than two spin components. The model has enough rich structure to illustrate almost every conceivable nuance of statistical mechanics. In [60, 61, 22] the phase diagrams of the model on the Bethe lattices (Cayley tree in our terminology) were studied and the pure phases of the ferromagnetic Potts model were found. Note that the Bethe lattices were fruitfully used, providing a deeper insight into the behavior of the Potts models.

We have to stress that one of the central problems in the theory of Gibbs measures of lattice systems is to describe infinite-volume (or limiting) Gibbs measures corresponding to a given Hamiltonian. A complete analysis of this set is often a difficult problem. Many papers have been devoted to these studies when the underlying lattice is a Cayley tree (see for example, [22, 26, 25, 65, 74]).

On the other hand, since the 1980s various models in physics described in the language of p -adic analysis (see [7, 8, 21, 50, 71, 72]), and numerous applications of such an analysis to mathematical physics have been studied in [11, 35, 36, 46, 50, 69, 70]. In those studies, it appeared unconventional probability measures. This means that a number of p -adic models in physics cannot be described using ordinary Kolmogorov's probability theory. New probability models, namely p -adic ones were investigated in [13, 34, 45, 42, 49, 63, 20]. Note that such kind of probabilities measures have many applications in algebraic number theory (see for example, [14, 15, 16, 28, 69]).

In [24, 58, 59] we have developed a p -adic probability theory approaches to study p -adic Potts models on Cayley tree of order two. We especially interested in the construction of p -adic Gibbs measures for the mentioned model. It was proved that if q is divisible by p then a phase transitions occurs for the q -state Potts model. Further, in [55] we have introduced a new kind of p -adic measures, associated with $q + 1$ -state Potts model, called *p -adic quasi Gibbs measure*, which is totally different from the p -adic Gibbs measure. Note that such measures present more natural concrete examples of p -adic Markov processes (see [76]). We established the existence p -adic quasi Gibbs measures for the mentioned model on a Cayley tree of order two. Moreover, if q is divisible by p , then we prove the occurrence of a strong phase transition. If q and p are relatively prime, then there is a quasi phase transition. These results totally different from the results of [58, 59]. It is natural to ask: do we have the similar phase transitions if one decreases the order of the tree?

In the present paper, our aim is to investigate a phase transition problem for the q -state p -adic Potts model over the Cayley tree of order three. In this paper, we consider a more general notion of p -adic Gibbs measure which depends on parameter $\rho \in \mathbb{Q}_p$ (see [56]). Such a measure is called *generalized p -adic quasi Gibbs measure*. When ρ equals to p -adic exponent, then it coincides with the p -adic Gibbs measure. When $\rho = p$, then it coincides with p -adic quasi Gibbs measure. Therefore, in the sequel, we will consider two regimes with respect to the values of $|\rho|_p$. Namely, in the first regime, one takes $\rho = \exp_p(J)$ for some $J \in \mathbb{Q}_p$, in the second one $|\rho|_p < 1$. In each regime, we first find conditions for the existence of generalized p -adic quasi Gibbs measures. Furthermore, we will study the occurrence of phase transitions. In the first regime, in [58] it was predicted that a phase transition may occur if $|q|_p < 1$ (i.e. q is divisible by p). In this paper, we able to establish the existence of the phase transition under a stronger condition, i.e. when $|q|_p \leq p^{-3}$. If $|q|_p = 1$, such a case was not investigated neither in [58] nor [55]. So, in that case, we proved the existence of a phase transition under some conditions. In the second regime, when $|\rho|_p, |q|_p \leq p^{-2}$ we will prove the existence of a quasi phase transition. It turns out that if $|\rho|_p < |q - 1|_p^2 < 1$, then one finds the existence of the strong phase transition.

2. PRELIMINARIES

2.1. p -adic numbers. In what follows p will be a fixed prime number, and by \mathbb{Q}_p it is denoted the field of p -adic numbers, which is a completion of the rational numbers \mathbb{Q} with respect to the norm $|\cdot|_p : \mathbb{Q} \rightarrow \mathbb{R}$ given by

$$(2.1) \quad |x|_p = \begin{cases} p^{-r} & x \neq 0, \\ 0, & x = 0, \end{cases}$$

here, $x = p^r \frac{m}{n}$ with $r, m \in \mathbb{Z}$, $n \in \mathbb{N}$, $(m, p) = (n, p) = 1$. A number r is called a p -order of x and it is denoted by $ord_p(x) = r$. The absolute value $|\cdot|_p$, is non-Archimedean, meaning that it satisfies the ultrametric triangle inequality $|x + y|_p \leq \max\{|x|_p, |y|_p\}$.

Any p -adic number $x \in \mathbb{Q}_p$, $x \neq 0$ can be uniquely represented in the form

$$(2.2) \quad x = p^{ord_p(x)}(x_0 + x_1p + x_2p^2 + \dots),$$

where x_j are integers, $0 \leq x_j \leq p - 1$, $x_0 > 0$, $j = 0, 1, 2, \dots$. In this case $|x|_p = p^{-ord_p(x)}$.

We respectively denote the set of all p -adic integers and units of \mathbb{Q}_p by

$$\mathbb{Z}_p = \{x \in \mathbb{Q}_p : |x|_p \leq 1\}, \quad \mathbb{Z}_p^* = \{x \in \mathbb{Q}_p : |x|_p = 1\}.$$

Any nonzero p -adic number $x \in \mathbb{Q}_p$ has a unique representation of the form $x = \frac{x^*}{|x|_p}$, where $x^* \in \mathbb{Z}_p^*$.

Lemma 2.1 (Hensel's Lemma, [70]). *Let $f(x)$ be polynomial whose the coefficients are p -adic integers. Let θ be a p -adic integer such that for some $i \geq 0$ we have*

$$f(\theta) \equiv 0 \pmod{p^{2i+1}},$$

$$f'(\theta) \equiv 0 \pmod{p^i}, \quad f'(\theta) \not\equiv 0 \pmod{p^{i+1}}.$$

Then $f(x)$ has a unique p -adic integer root x_0 which satisfies $x_0 \equiv \theta \pmod{p^{i+1}}$.

Let p be a prime number, $q \in \mathbb{N}$, $a \in \mathbb{F}_p$ with $a \neq \bar{0}$. The number a is called a q -th power residue modulo p if the the following equation

$$(2.3) \quad x^q = a$$

has a solution in \mathbb{F}_p .

Proposition 2.2 ([64]). *Let p be an odd prime number, $q \in \mathbb{N}$, $d = (q, p-1)$, and $a \in \mathbb{F}_p$ with $a \neq \bar{0}$. Then the following statements hold true:*

- (i) *a is the q -th power residue modulo p if and only if one has $a^{\frac{p-1}{d}} = \bar{1}$;*
- (ii) *If $a^{\frac{p-1}{d}} = \bar{1}$ then the equation (2.3) has d number of solutions in \mathbb{F}_p .*

Let $B(a, r) = \{x \in \mathbb{Q}_p : |x - a|_p < r\}$, where $a \in \mathbb{Q}_p$, $r > 0$. The p -adic logarithm is defined by series

$$\log_p(x) = \log_p(1 + (x - 1)) = \sum_{n=1}^{\infty} (-1)^{n+1} \frac{(x - 1)^n}{n},$$

which converges for every $x \in B(1, 1)$. And p -adic exponential is defined by

$$\exp_p(x) = \sum_{n=1}^{\infty} \frac{x^n}{n!},$$

which converges for every $x \in B(0, p^{-1/(p-1)})$.

Lemma 2.3. [46],[70] *Let $x \in B(0, p^{-1/(p-1)})$ then we have*

$$|\exp_p(x)|_p = 1, \quad |\exp_p(x) - 1|_p = |x|_p < 1, \quad |\log_p(1 + x)|_p = |x|_p < p^{-1/(p-1)}$$

and

$$\log_p(\exp_p(x)) = x, \quad \exp_p(\log_p(1 + x)) = 1 + x.$$

In what follows we will use the following

Lemma 2.4 ([43]). *If $|a_i|_p \leq 1$, $|b_i|_p \leq 1$, $i = 1, \dots, n$, then*

$$\left| \prod_{i=1}^n a_i - \prod_{i=1}^n b_i \right|_p \leq \max_{i \leq i \leq n} \{|a_i - b_i|_p\}$$

Denote

$$(2.4) \quad \mathcal{E}_p = \{x \in \mathbb{Q}_p : |x|_p = 1, \quad |x - 1|_p < p^{-1/(p-1)}\}.$$

So, from Lemma 2.3 one concludes that if $x \in \mathcal{E}_p$, then there is an element $h \in B(0, p^{-1/(p-1)})$ such that $x = \exp_p(h)$.

Note that the basics of p -adic analysis, p -adic mathematical physics are explained in [46, 51, 66, 63, 70].

2.2. p -adic measure. Let (X, \mathcal{B}) be a measurable space, where \mathcal{B} is an algebra of subsets X . A function $\mu : \mathcal{B} \rightarrow \mathbb{Q}_p$ is said to be a p -adic measure if for any $A_1, \dots, A_n \subset \mathcal{B}$ such that $A_i \cap A_j = \emptyset$ ($i \neq j$) the equality holds

$$\mu\left(\bigcup_{j=1}^n A_j\right) = \sum_{j=1}^n \mu(A_j).$$

A p -adic measure is called a *probability measure* if $\mu(X) = 1$. One of the important condition (which was already invented in the first Monna–Springer theory of non-Archimedean integration [52]) is boundedness, namely a p -adic probability measure μ is called *bounded* if $\sup\{|\mu(A)|_p : A \in \mathcal{B}\} < \infty$. We should stress that the boundedness condition by itself provides a fruitful integration theory (see for example [38]). Note that, in general, a p -adic probability measure need not be bounded [34, 42, 46]. For more detail information about p -adic measures we refer to [6],[34],[44],[63].

2.3. Cayley tree. Let $\Gamma_+^k = (V, L)$ be a semi-infinite Cayley tree of order $k \geq 1$ with the root x^0 (whose each vertex has exactly $k + 1$ edges, except for the root x^0 , which has k edges). Here V is the set of vertices and L is the set of edges. The vertices x and y are called *nearest neighbors* and they are denoted by $l = \langle x, y \rangle$ if there exists an edge connecting them. A collection of the pairs $\langle x, x_1 \rangle, \dots, \langle x_{d-1}, y \rangle$ is called a *path* from the point x to the point y . The distance $d(x, y)$, $x, y \in V$, on the Cayley tree, is the length of the shortest path from x to y .

Recall a coordinate structure in Γ_+^k : every vertex x (except for x^0) of Γ_+^k has coordinates (i_1, \dots, i_n) , here $i_m \in \{1, \dots, k\}$, $1 \leq m \leq n$ and for the vertex x^0 we put (0) . Namely, the symbol (0) constitutes level 0, and the sites (i_1, \dots, i_n) form level n (i.e. $d(x^0, x) = n$) of the lattice.

Let us set

$$W_n = \{x \in V | d(x, x^0) = n\}, \quad V_n = \bigcup_{m=1}^n W_m, \quad L_n = \{l = \langle x, y \rangle \in L | x, y \in V_n\}.$$

For $x \in \Gamma_+^k$, let us denote

$$(2.5) \quad S(x) = \{y \in W_{n+1} : d(x, y) = 1\} \quad x \in W_n.$$

here This set is called a set of *direct successors* of x .

Using the coordinate system one can define translations of Γ_+^k by

$$(2.6) \quad \tau_{(i_1, \dots, i_n)}(j_1, \dots, j_m) = (i_1, \dots, i_n, j_1, \dots, j_m).$$

2.4. Generalized p -adic quasi Gibbs measure. In this section we define a notion of generalized p -adic quasi Gibbs measure in a general setting, i.e. for arbitrary nearest-neighbor models (see [56]).

Let $\Phi = \{0, 1, 2, \dots, q-1\}$, here $q \geq 1$, (Φ is called a *state space*) and is assigned to the vertices of the tree $\Gamma_+^k = (V, \Lambda)$. A configuration σ on V is then defined as a function $x \in V \rightarrow \sigma(x) \in \Phi$; in a similar manner one defines configurations σ_n and ω on V_n and W_n , respectively. The set of all configurations on V (resp. V_n, W_n) coincides with $\Omega = \Phi^V$ (resp. $\Omega_{V_n} = \Phi^{V_n}, \Omega_{W_n} = \Phi^{W_n}$). One can see that $\Omega_{V_n} = \Omega_{V_{n-1}} \times \Omega_{W_n}$. Using this, for given configurations $\sigma_{n-1} \in \Omega_{V_{n-1}}$ and $\omega \in \Omega_{W_n}$ we define their concatenations by

$$(\sigma_{n-1} \vee \omega)(x) = \begin{cases} \sigma_{n-1}(x), & \text{if } x \in V_{n-1}, \\ \omega(x), & \text{if } x \in W_n. \end{cases}$$

It is clear that $\sigma_{n-1} \vee \omega \in \Omega_{V_n}$.

Let H_n be a Hamiltonian given by

$$(2.7) \quad H_n(\sigma) = \sum_{\langle x,y \rangle \in L_n} \Psi_{x,y}(\sigma), \quad \sigma \in \Omega_{V_n}, \quad n \in \mathbb{N},$$

where $\Psi_{x,y}(\sigma) : \Omega_{V_n} \rightarrow \mathbb{Z}$ is a given function for every x, y .

A construct of a generalized p -adic quasi Gibbs measure corresponding to the model is given below.

Assume that $\mathbf{h} : V \setminus \{x^{(0)}\} \rightarrow \mathbb{Q}_p^\Phi$ is a mapping, i.e. $\mathbf{h}_x = (h_{0,x}, h_{1,x}, \dots, h_{q-1,x})$, where $h_{i,x} \in \mathbb{Q}_p$ ($i \in \Phi$) and $x \in V \setminus \{x^{(0)}\}$. Given $n \in \mathbb{N}$, we consider a p -adic probability measure $\mu_{\mathbf{h},\rho}^{(n)}$ on Ω_{V_n} defined by

$$(2.8) \quad \mu_{\mathbf{h},\rho}^{(n)}(\sigma) = \frac{1}{Z_{n,\rho}^{(\mathbf{h})}} \rho^{H_n(\sigma)} \prod_{x \in W_n} h_{\sigma(x),x}$$

Here, $\rho \in \mathbb{Q}_p$, $\sigma \in \Omega_{V_n}$, and $Z_{n,\rho}^{(\mathbf{h})}$ is the corresponding normalizing factor called a *partition function* given by

$$(2.9) \quad Z_{n,\rho}^{(\mathbf{h})} = \sum_{\sigma \in \Omega_{V_n}} \rho^{H_n(\sigma)} \prod_{x \in W_n} h_{\sigma(x),x}.$$

Note that, in general, $Z_{n,\rho}^{(\mathbf{h})}$ could be zero, but we will show that, in a Potts model case, the partition function is not zero. In this paper, we are interested in a construction of an infinite volume distribution with given finite-dimensional distributions. More exactly, we would like to find a p -adic probability measure μ on Ω which is compatible with given ones $\mu_{\mathbf{h},\rho}^{(n)}$, i.e.

$$(2.10) \quad \mu(\sigma \in \Omega : \sigma|_{V_n} = \sigma_n) = \mu_{\mathbf{h},\rho}^{(n)}(\sigma_n), \quad \text{for all } \sigma_n \in \Omega_{V_n}, \quad n \in \mathbb{N}.$$

In general, à priori the existence such a kind of measure μ is not known, since there is not much information on topological properties, such as compactness, of the set of all p -adic measures defined even on compact spaces [77]. Note that certain properties of the set of p -adic measures has been studied in [32, 33], but those properties are not enough to prove the existence of the limiting measure. Therefore, at a moment, we are going to employ so called the p -adic Kolmogorov's extension Theorem (see [23],[42]) which is based on *compatibility condition* for

the measures $\mu_{\mathbf{h},\rho}^{(n)}$, $n \geq 1$, i.e.

$$(2.11) \quad \sum_{\omega \in \Omega_{W_n}} \mu_{\mathbf{h},\rho}^{(n)}(\sigma_{n-1} \vee \omega) = \mu_{\mathbf{h},\rho}^{(n-1)}(\sigma_{n-1}),$$

for any $\sigma_{n-1} \in \Omega_{V_{n-1}}$. This condition according to the theorem implies the existence of a unique p -adic measure $\mu_{\mathbf{h},\rho}$ defined on Ω with a required condition (2.10). Such a measure $\mu_{\mathbf{h},\rho}$ is said to be a *generalized p -adic quasi Gibbs measure* corresponding to the model. Note that more general theory of p -adic measures has been developed in [31].

By $GQG_\rho(H)$ we denote the set of all generalized p -adic quasi Gibbs measures associated with functions $\mathbf{h} = \{\mathbf{h}_x, x \in V\}$. If there are at least two distinct generalized p -adic quasi Gibbs measures such that at least one of them is unbounded, then we say that a *phase transition* occurs. If there are two different functions \mathbf{s} and \mathbf{h} defined on \mathbb{N} such that there exist the corresponding measures $\mu_{\mathbf{s},\rho}$, $\mu_{\mathbf{h},\rho}$, and they are bounded, then we say there is a *quasi phase transition*. If one finds two different functions \mathbf{s} and \mathbf{h} defined on \mathbb{N} such that there exist the corresponding measures $\mu_{\rho,\mathbf{s}}$ and $\mu_{\rho,\mathbf{h}}$, for which one is bounded, another one is unbounded, and there is a sequence of sets $\{A_n\}$ such that $A_n \in \Omega_{V_n}$ with $|\mu_{\rho,\mathbf{s}}(A_n)|_p \rightarrow 0$, $|\mu_{\rho,\mathbf{h}}(A_n)|_p \rightarrow \infty$ as $n \rightarrow \infty$, then we say that there occurs a *strong phase transition* (see [56]).

3. p -ADIC POTTS MODEL AND ITS p -ADIC QUASI GIBBS MEASURES

In this section we consider the p -adic Potts model where spin takes values in the set $\Phi = \{0, 1, 2, \dots, q-1\}$. In what follows, we will always assume that $q \geq 3$. In this case, for every $\sigma \in \Omega_{V_n}$ one has $\Psi_{x,y}(\sigma) = \delta_{\sigma(x),\sigma(y)}$, i.e. the Hamiltonian of q -state Potts model has the following form

$$(3.1) \quad H_n(\sigma) = \sum_{\langle x,y \rangle \in L_n} \delta_{\sigma(x),\sigma(y)}, \quad \sigma \in \Omega_{V_n}, \quad n \in \mathbb{N},$$

where δ is the Kronecker symbol. Note that when $q = 1$, then the corresponding model reduces to the p -adic Ising model. Such a model was investigated in [23, 30].

Given $\rho \in \mathbb{Q}_p$ one can construct generalized p -adic quasi Gibbs measures corresponding to the q -state Potts model. In what follows, we are interested in the existence of phase transition for the Potts model.

Remark 3.1. Note that if one takes $\rho = p$, then the generalized p -adic quasi Gibbs measure reduces to the *p -adic quasi Gibbs measure* (see [55]). If one takes $\rho \in \mathcal{E}_p$ with $h_x \in \mathcal{E}_p$, then the defined measure reduces to *p -adic Gibbs measure* (see [58]).

Remark 3.2. Note that in [23, 30, 55, 58, 59] phase transitions in class of p -adic quasi Gibbs measures, p -adic Gibbs measures for Ising and Potts models on Cayley tree have been studied over the Cayley tree of order two. When a state space Φ is countable, the corresponding p -adic Gibbs measures (resp. generalized p -adic quasi Gibbs measures) have been investigated in [43, 53] (resp. [54]).

Using the same argument as [58, 43] we can prove the following

Theorem 3.1. *The measures $\mu_{\mathbf{h},\rho}^{(n)}$, $n = 1, 2, \dots$ (see (2.8)) associated with q -state Potts model (3.1) satisfy the compatibility condition (2.11) if and only if for any $n \in \mathbb{N}$ the following equation*

holds:

$$(3.2) \quad \hat{h}_x = \prod_{y \in \mathcal{S}(x)} \mathbf{F}(\hat{\mathbf{h}}_y; \rho),$$

here and below a vector $\hat{\mathbf{h}} = (\hat{h}_1, \dots, \hat{h}_{q-1}) \in \mathbb{Q}_p^{q-1}$ is defined by a vector $\mathbf{h} = (h_0, h_1, \dots, h_{q-1}) \in \mathbb{Q}_p^q$ as follows

$$(3.3) \quad \hat{h}_i = \frac{h_i}{h_0}, \quad i = 1, 2, \dots, q-1$$

and mapping $\mathbf{F} : \mathbb{Q}_p^{q-1} \times \mathbb{Q}_p \rightarrow \mathbb{Q}_p^{q-1}$ is defined by $\mathbf{F}(\mathbf{x}; \rho) = (F_1(\mathbf{x}; \rho), \dots, F_{q-1}(\mathbf{x}; \rho))$ with

$$(3.4) \quad F_i(\mathbf{x}; \rho) = \frac{(\rho-1)x_i + \sum_{j=1}^{q-1} x_j + 1}{\sum_{j=1}^{q-1} x_j + \rho}, \quad \mathbf{x} = \{x_i\} \in \mathbb{Q}_p^{q-1}, \quad i = 1, 2, \dots, q-1.$$

Remark 3.3. In what follows, without loss of generality, we may assume that $h_0 = 1$. Otherwise, in (2.8) we multiply and divide the expression on the right hand side by $\prod_{x \in W_n} h_{0,x}$, and after replacing h_i by h_i/h_0 , we get the desired equality.

4. EXISTENCE OF GENERALIZED p -ADIC QUASI GIBBS MEASURES

In this section we are going to establish the existence of generalized p -adic quasi Gibbs measures for the Potts model (3.1) on a Cayley tree of order 3, i.e. $k = 3$. In the sequel, for the sake of simplicity we will always assume that $p > 3$.

Recall that a function $\mathbf{h} = \{\mathbf{h}_x\}_{x \in V \setminus \{x^0\}}$ is called *translation-invariant* if $\mathbf{h}_{\tau_x(y)} = \mathbf{h}_y$ for all $x, y \in V \setminus \{x^0\}$. A p -adic measure $\mu_{\mathbf{h}}$, corresponding to a translation-invariant function \mathbf{h} , is called *translation-invariant generalized p -adic quasi Gibbs measure*. Note that the translational-invariance of $\mathbf{h} = \{\mathbf{h}_x\}_{x \in V \setminus \{x^0\}}$ implies that $\mathbf{h}_x = \mathbf{h}_y$ for all $x, y \in V \setminus \{x^0\}$.

Let us first restrict ourselves to the description of translation-invariant solutions of (3.2), namely $\mathbf{h}_x = \mathbf{h} (= (h_0, h_1, \dots, h_q))$ for all $x \in V$. Then (3.2) can be rewritten as follows

$$(4.1) \quad \hat{h}_i = \left(\frac{(\theta-1)\hat{h}_i + \sum_{j=1}^q \hat{h}_j + 1}{\sum_{j=1}^q \hat{h}_j + \theta} \right)^3, \quad i = 1, 2, \dots, q.$$

One can see that $(\underbrace{1, \dots, 1}_m, h, 1, \dots, 1)$ is an invariant line for (4.1) ($m = 1, \dots, q-1$). Over such a line equation (4.1) reduces to the following fixed point problem

$$(4.2) \quad x = f_\rho(x),$$

where

$$(4.3) \quad f_\rho(x) = \left(\frac{\rho x + q - 1}{x + \rho + q - 2} \right)^3.$$

A simple calculation shows that (4.2) has a form

$$(x-1)(x^3 - Ax^2 - Bx + C) = 0,$$

where

$$(4.4) \quad \begin{cases} A = \rho^3 - 3\rho - 3q + 5, \\ B = \rho(\rho - 3)^2 + 9q + 3\rho q(\rho - 2) - 3q^2 - 7, \\ C = (q - 1)^3. \end{cases}$$

Hence, $x_0 = 1$ solution defines a generalized p -adic quasi Gibbs measure μ_0 .

Now we are interested in finding other solutions of (4.2), which means we need to solve the following one

$$(4.5) \quad x^3 - Ax^2 - Bx + C = 0.$$

By performing standard substitution $x = z + A/3$, one can reduce the equation (4.5) to

$$(4.6) \quad z^3 + \alpha z = \beta,$$

where

$$(4.7) \quad \alpha = -\left(B + \frac{A^2}{3}\right), \quad \beta = \frac{2}{3^3}A^3 + \frac{AB}{3} - C.$$

Now we are going to consider two regimes with respect to ρ . Namely, (i) $\rho \in \mathcal{E}_p$, and (ii) $|\rho|_p < 1$.

4.1. Regime $\rho \in \mathcal{E}_p$. In this case, due to Lemma 2.3 one has $|\rho|_p = 1$ and $|\rho - 1|_p < 1$.

Note that in this regime, the measure corresponding to the solution $x = 1$ is always p -adic Gibbs measure.

Direct calculations show that one has

$$(4.8) \quad \begin{aligned} A &= 3 + 3(1 - \rho) + (\rho^3 - 1) - 3q, \\ \alpha &= \frac{1}{3}(\rho - 1)^2 \left(-9 + 9q - 9(\rho - 1) + 6q(\rho - 1) - 9(\rho - 1)^2 \right. \end{aligned}$$

$$(4.9) \quad \begin{aligned} &\left. -6(\rho - 1)^3 - (\rho - 1)^4 \right) \\ \beta &= \frac{1}{27}(\rho - 1)^3 \left(54 - 81q + 27q^2 + 81(\rho - 1) - 81q(\rho - 1) \right. \\ &\quad \left. + 108(\rho - 1)^2 - 81q(\rho - 1)^2 + 81(\rho - 1)^3 - 18q(\rho - 1)^3 \right. \\ (4.10) \quad &\left. + 54(\rho - 1)^4 + 18(\rho - 1)^5 + 2(\rho - 1)^6 \right) \end{aligned}$$

In [58] we have proved the following

Theorem 4.1 ([58]). *Let $|q|_p = 1$ and $\rho \in \mathcal{E}_p$. Then there is a unique p -adic Gibbs measure for the Potts model over the Cayley tree of order k ($k \geq 1$).*

This theorem implies that if $|q|_p = 1$ then there is no p -adic Gibbs measure except for μ_0 . But further, we will show that under the mentioned condition, one can find generalized p -adic quasi Gibbs measures.

Now we first assume that $|q|_p < 1$ to find nontrivial p -adic Gibbs measures. Under this condition, from the strong triangle inequality from (4.9),(4.10) one finds $|\alpha|_p = |\rho - 1|_p^2$, and $|\beta|_p = |\rho - 1|_p^3$. We emphasize that the existence of nontrivial p -adic Gibbs measure means that a solution of (4.5) belongs to \mathcal{E}_p , which implies that one can find $h \in \mathbb{Q}_p$ such that $x = \exp_p(h)$.

Lemma 4.2. *Let $\rho \in \mathcal{E}_p$ and $|q|_p < 1$. Then a solution x of (4.5) belongs to \mathcal{E}_p iff the corresponding solution of (4.6) belongs to $\mathbb{Z}_p \setminus \mathbb{Z}_p^*$.*

Proof. First denote $\gamma = 3(1 - \rho) + (\rho^3 - 1) - 3q$. It is clear that $|\gamma|_p < 1$. Assume that $x \in \mathcal{E}_p$ is a solution of (4.5). Then for the solution of (4.6) with (4.8) we get

$$|z|_p = \left| x - \frac{A}{3} \right|_p = \left| x - 1 + 1 - \frac{3 + \gamma}{3} \right|_p = |x - 1 - \gamma|_p < 1.$$

This means that a solution of (4.6) must be in $\mathbb{Z}_p \setminus \mathbb{Z}_p^*$.

Now assume that a solution z of (4.6) belongs to $\mathbb{Z}_p \setminus \mathbb{Z}_p^*$. Then $x = z + A/3$ is a solution of (4.5). From (4.8) we find that $|A|_p = 1$, and one gets $|x|_p = |z + A/3|_p = 1$. Moreover, we have

$$|x - 1|_p = \left| z + \frac{A}{3} - 1 \right|_p = |z + \gamma|_p < 1,$$

hence $x \in \mathcal{E}_p$. □

So, to find nontrivial p -adic Gibbs measures, we need to solve (4.6) over the set $\mathbb{Z}_p \setminus \mathbb{Z}_p^*$. Hence, assume that a solution z of (4.6) has the following form $z = p^k y$, where $y \in \mathbb{Z}_p^*$ and $k \geq 1$. Let us substitute the last form to (4.6) and one finds

$$(4.11) \quad y^3 + ay = b$$

where $a = p^{-2k}\alpha$, $b = p^{-3k}\beta$. It is clear that $|a|_p = (p^k|\rho - 1|_p)^2$, $|b|_p = (p^k|\rho - 1|_p)^3$. Now we chose k such a way that $p^k|\rho - 1|_p = 1$. Note that such a number does exist. So, $|a|_p = |b|_p = 1$.

Now we want to solve (4.11) in \mathbb{Z}_p^* . To do it we are going to apply Theorem A.4.

Let us consider the canonical forms of a and b , i.e.

$$a = a_0 + a_1p + \cdots, \quad b = b_0 + b_1p + \cdots,$$

where $a_0, b_0 \in \{1, \dots, p - 1\}$. It then follows from (4.9) and (4.10) that

$$(4.12) \quad a = p - 3 + (p - 1)p + 3q - 3(\rho - 1) + \eta p^n, \quad b = 2 - 3q + 3(\rho - 1) + \eta_1 p^m, \quad n, m \geq 2,$$

where $\eta, \eta_1 \in \mathbb{Z}_p$. So, $a_0 = p - 3$ and $b_0 = 2$. Hence, $D_0 = -4p(p^2 - 9p + 27)$, where $D_0 = -4a_0^3 - 27b_0^2$. This means $D_0 \equiv 0 \pmod{p}$. In this case, due to Theorem A.5 we need to compute $D = -4a^3 - 27b^2$. Let us assume that $|q|_p \leq p^{-3}, |\rho - 1|_p \leq p^{-3}$. Then from (4.12) one finds that

$$a^3 \equiv (p - 3)^3 + 3(p - 3)^2(p - 1)p + 3(p - 3)(p - 1)^2p^2 \pmod{p^3}, \quad b^2 \equiv 4 \pmod{p^3}.$$

hence, we have

$$D = -\left(4p(p^2 - 9p + 27) - 4 \cdot 3^3p + 45p^2 + \varepsilon p^3\right) = -p^2(9 + \varepsilon p)$$

where $\varepsilon \in \mathbb{Z}_p$. This means $d_0 = 9$ and $\text{ord}_p(D)$ is divisible by 2. Moreover, $d_0^{p-2/2} \equiv 1 \pmod{p}$ holds. Then according to Theorem A.5 we conclude that the equation (4.11) has three distinct solutions in \mathbb{Z}_p^* .

Thus, the equation (4.6) has three distinct solutions belonging to $\mathbb{Z}_p \setminus \mathbb{Z}_p^*$. Hence, by Lemma 4.2 the equation (4.5) has three solutions. Therefore, by μ_i ($i = 1, 2, 3$) we denote the corresponding p -adic Gibbs measures. Due to Theorem 3.1 one can formulate the following

Theorem 4.3. *Let $p > 3$, $|q|_p \leq p^{-3}$, $\rho \in \mathcal{E}_p$ with $|\rho - 1|_p \leq p^{-3}$. Then for the p -adic Potts model (3.1) on the Cayley tree of order three, there exist four translation-invariant p -adic Gibbs measures.*

Note that in [58] if $|q|_p < 1$ we did not have other extra conditions for the existence of the p -adic Gibbs measures. From Theorem 4.3 we see that for the existence of the Gibbs measures one needs $|q|_p < p^{-3}$ extra condition.

Now assume that $|q|_p = 1$. In this case we are going to establish the existence of a generalized p -adic quasi Gibbs measure. In the sequel, for the sake of simplicity we impose on q the following constrains:

$$(4.13) \quad |q - 1|_p = 1, \quad |2q - 1|_p = 1.$$

Then from (4.8),(4.9) and (4.10) we immediately find that $|A|_p = 1$, $|\alpha|_p = |\rho - 1|_p^2$ and $|\beta|_p = |\rho - 1|_p^3$. In this case, the equation (4.6) has no solution belonging to \mathbb{Z}_p^* . Indeed, assume that $z \in \mathbb{Z}_p^*$ is a solution, then we have $|\alpha|_p = |z\alpha|_p = |\beta - z^3|_p = 1$, but this contradicts to $|\alpha|_p < 1$. Consequently, any solution of (4.6) belongs to $\mathbb{Z}_p \setminus \mathbb{Z}_p^*$. Now using the same argument as above we reduce the equation (4.6) to (4.11) with $y \in \mathbb{Z}_p^*$, $a = p^{-2k}\alpha$, $b = p^{-3k}\beta$. By the same argument, we choose k such a way that $p^k|\rho - 1|_p = 1$. So, $|a|_p = |b|_p = 1$.

Assume that $|\rho - 1|_p \leq p^{-3}$. Then from (4.9), (4.10) one gets

$$(4.14) \quad a \equiv 3(q - 1) \pmod{p^3}, \quad b \equiv 2(q - 1)(2q - 1) \pmod{p^3}.$$

Let us compute $D = -4a^3 - 27b^2$. Then from (4.14) one finds that

$$\begin{aligned} D &= -\left(4 \cdot 27(q - 1)^3 + 27 \cdot 4(q - 1)^2(2q - 1)^2 + \eta|\rho - 1|\right) \\ &= -108q(q - 1)^2(4q - 3) - \eta|\rho - 1|_p, \quad \text{for some } \eta \in \mathbb{Z}_p. \end{aligned}$$

Hence, if $|\rho - 1|_p < |4q - 3|_p$ we conclude that $|D|_p = |4q - 3|_p$.

Now taking into account Theorem A.5 we can formulate the following

Theorem 4.4. *Let $p > 3$, $\rho \in \mathcal{E}_p$ with $|\rho - 1|_p \leq p^{-3}$, $|\rho - 1|_p < |4q - 3|_p$ and $|q|_p = |q - 1|_p = |2q - 1|_p = 1$. Let $u_{n+3} = b_0u_n - a_0u_{n+1}$ with $u_1 = 0$, $u_2 = -a_0$, $u_3 = -b_0$, where $a_0 \equiv 3(q - 1) \pmod{p}$, $b_0 \equiv 2(q - 1)(2q - 1) \pmod{p}$ with $a_0, b_0 \in \{1, 2, \dots, p - 1\}$. Then for the q -state p -adic Potts model (3.1) on the Cayley tree of order three, there exist one translation-invariant p -adic Gibbs measure and*

- (i) *three translation-invariant generalized p -adic quasi Gibbs measures if one of the following conditions are satisfied:*
 - (a) $|4q - 3|_p = 1$ and $u_{p-2} \equiv 0 \pmod{p}$;
 - (b) $|4q - 3|_p = p^{-2k}$ for some $k \geq 1$, and $-108q(q - 1)^2(4q - 3)p^{-2k}$ is a quadratic residue modulo p .
- (ii) *one translation-invariant generalized p -adic quasi Gibbs measure if one of the following conditions are satisfied:*

- (c) $D_0 u_{p-2}^2 \not\equiv 0$ and $9a_0^2 \pmod{p}$;
- (d) $|4q - 3|_p = p^{-(2k+1)}$, $k \geq 1$;
- (e) $|4q - 3|_p = p^{-2k}$, and $-108q(q-1)^2(4q-3)p^{-2k}$ is not a quadratic residue modulo p .

(iii) otherwise, there is not translation-invariant generalized p -adic quasi Gibbs measure.

We should stress that in [58] it was not investigated the case $|q|_p = 1$. From the theorem we conclude in the mentioned setting one can find new kind of Gibbs measures.

Let us consider some more concrete examples.

EXAMPLE 1. Assume that $p = 5$ and $q = 4$. It is clear that $|q|_5 = 1$ and (4.13) is satisfied. From (4.9) and (4.10) we get that $a_0 = 4$, $b_0 = 2$. One can check that $D_0 = -4 \cdot 91$, $u_3 = -2$. Hence, we have $D_0 u_3^2 \equiv 9 \cdot a_0^2 \pmod{5}$, this with Theorem A.5 implies (4.6) has no solution. In this case, for the model there is only p -adic Gibbs measure.

EXAMPLE 2. Assume that $p = 5$ and $q = 7$. It is clear that $|q|_5 = 1$ and (4.13) with $|4q - 3|_5 = 5^{-2}$ is satisfied. One can see that $108q(q-1)^2(4q-3)5^{-2} \equiv 1 \pmod{5}$. Since 1 is a quadratic residue modulo 5, this yields that $108q(q-1)^2(4q-3)5^{-2}$ is so. Hence, due to Theorem 4.4 there are three translation-invariant generalized p -adic quasi Gibbs measures.

4.2. **Regime** $|\rho|_p < 1$. This regime is totally different from the previous case. Now we are interested in the existence of generalized p -adic quasi Gibbs measures.

Direct calculations show that one has

$$(4.15) \quad \alpha = \frac{1}{3}(\rho - 1)^2 \left(-4 + 3q - 5\rho + 6q\rho + 3\rho^2 - 2\rho^3 - \rho^4 \right)$$

$$(4.16) \quad \beta = \frac{1}{27}(\rho - 1)^3 \left(38 - 63q + 27q^2 - 30\rho + 27q\rho + \right. \\ \left. + 39\rho^2 - 27q\rho^2 + 5\rho^3 - 18q\rho^3 - 6\rho^4 + 6\rho^5 + 2\rho^6 \right)$$

Assume that $|q|_p < 1$. Then from (4.15),(4.16) one finds that $|\alpha|_p = 1$, and $|\beta|_p = 1$ if $p \neq 19$, and $|\beta|_p < 1$ if $p = 19$.

Let us first consider $p \neq 19$. Furthermore, for the sake of simplicity, we will assume that $|\rho|_p, |q|_p \leq p^{-2}$. Then one can see that

$$(4.17) \quad \alpha = -\frac{4}{3} + \epsilon p^2, \quad \beta = -\frac{38}{27} + \epsilon_1 p^2, \quad \epsilon, \epsilon_1 \in \mathbb{Z}_p.$$

So, $a_0 \equiv -4/3 \pmod{p}$, and $b_0 \equiv -38/27 \pmod{p}$. Moreover, one can find that $D = -(4\alpha^3 + 27\beta^2) = -44 + \eta p^2$, where $\eta \in \mathbb{Z}_p$. Hence,

$$|D|_p = \begin{cases} 1, & \text{if } p \neq 11, \\ p^{-1}, & \text{if } p = 11. \end{cases}$$

Therefore, due to Theorem A.5 we conclude that the equation (4.6) has a unique solution either $p = 11$ or $p \neq 11$ and $D_0 u_{p-2}^2 \not\equiv 0; 9 \pmod{p}$.

Now assume that $p = 19$. Then $|\beta|_p = p^{-1}$, $|\alpha|_p = 1$. One can see that $a_0 = 5$. We know that -5 is not a quadratic residue modulo 19, therefore, due to Theorem A.4 I.2 the equation (4.6) has no solution in \mathbb{Z}_p^* . Hence, the equation may have a solution in \mathbb{Z}_p . In this case, we have $z = p^k y$, where $k \geq 1$ and $y \in \mathbb{Z}_p^*$. Then the equation is reduced to (4.11) with $a = p^{-2k} \alpha$

and $b = p^{-3k}\beta$. It is clear that $|a|_p = p^{2k}$, $|b|_p = p^{2k}$. Take $k = 1$, then again Theorem A.4 and Theorem A.5 I.4 imply that the equation (4.11) has a unique solution. Hence, the equation (4.6) has a unique solution belonging to $\mathbb{Z}_p \setminus \mathbb{Z}_p^*$.

So, one can formulate the following

Theorem 4.5. *Let $p > 3$. Assume that $|\rho|_p, |q|_p \leq p^{-2}$. Then for the q -state p -adic Potts model (3.1) on the Cayley tree of order three, there exist two translation-invariant generalized p -adic quasi Gibbs measures*

- (i) if $p = 11$;
- (ii) if $p = 19$;
- (iii) if $p \neq 11, 19$ and $D_0 u_{p-2}^2 \not\equiv 0; 9 \pmod{p}$, where $a_0 \equiv -4/3 \pmod{p}$, and $b_0 \equiv -38/27 \pmod{p}$, where $a_0, b_0 \in \{1, 2, \dots, p-1\}$.

Now we assume that $|q-1|_p < 1$. In what follows, we will assume that $|\rho|_p < |q-1|_p^2$. Then from (4.15),(4.16) one gets

$$(4.18) \quad \alpha = \frac{1}{3}(\rho-1)^2 \left(-1 + 3(q-1) + \epsilon\rho \right)$$

$$(4.19) \quad \beta = \frac{1}{27}(\rho-1)^3 \left(2 - 9(q-1) + 27(q-1)^2 + \epsilon_1\rho \right)$$

where $\epsilon, \epsilon_1 \in \mathbb{Z}_p$. Hence, we rewrite (4.18),(4.19) as follows

$$(4.20) \quad \alpha = -\frac{1}{3} + (q-1) + \eta(q-1)^3, \quad \beta = -\frac{2}{27} + \frac{q-1}{3} - (q-1)^2 + \eta_1(q-1)^3$$

where $\eta, \eta_1 \in \mathbb{Z}_p$. It is clear that $|\alpha|_p = |\beta|_p = 1$. Then one finds that

$$(4.21) \quad \begin{aligned} -D &= 4\alpha^3 + 27\beta^3 = -\frac{4}{27} + \frac{4}{3}(q-1) - 4(q-1)^2 \\ &\quad + \frac{4}{27} + 3(q-1)^2 - \frac{4}{3}(q-1) + 4(q-1)^2 + \tilde{\eta}(q-1)^3 \\ &= (q-1)^2(3 + \tilde{\eta}(q-1)) \end{aligned}$$

where $\tilde{\eta} \in \mathbb{Z}_p$. This means $|D|_p = |q-1|_p^2$ and $d_0 = -3$. Hence, Theorem A.5 implies that the equation (4.6) has three solutions if $\sqrt{-3}$ does exist. Otherwise, it has a unique solution. Therefore, we have the following

Theorem 4.6. *Let $p > 3$. Assume that $|\rho|_p < |q-1|_p^2 < 1$. Then for the q -state p -adic Potts model (3.1) on the Cayley tree of order three, there exist*

- (i) four translation-invariant generalized p -adic quasi Gibbs measures if $\sqrt{-3}$ exists;
- (ii) two translation-invariant generalized p -adic quasi Gibbs measures if $\sqrt{-3}$ does not exist.

Remark 4.1. Let $|\rho|_p < |q-1|_p^2 < 1$ be satisfied, and we assume that the equation (4.5) has three solutions x_1, x_2, x_3 , which means $\sqrt{-3} \in \mathbb{Q}_p$. Due to Vieta's Theorem one has

$$(4.22) \quad |x_1x_2 + x_1x_3 + x_2x_3|_p = 1, \quad |x_1x_2x_3|_p = |q-1|_p^3 < 1.$$

Since from (4.4) we have $|A|_p = |B|_p = 1$. Now from $x_i = z_i + A/3$, where z_i solutions of (4.6), one gets that $|x_i|_p \leq 1$. It then follows from (4.22) that at least one of the solutions' norm should be strictly less than 1. Suppose that the norm of two of solutions is strictly

less 1, say $|x_1|_p < 1$ and $|x_2|_p < 1$. Then from (4.22) one finds that $|B|_p < 1$, which is impossible. So, only one solution's norm is less than strictly 1, say x_1 , i.e. $|x_1|_p < 1$, $|x_2|_p = |x_3|_p = 1$.

Now we are going to investigate solutions of (3.2) over the invariant line $(1, 1, \dots, h, 1, \dots, 1)$. Let us introduce some notations. If $x \in W_n$, then instead of h_x we use the symbol $h_x^{(n)}$.

Denote

$$(4.23) \quad g_\rho(x) = \frac{\rho x + q - 1}{x + \rho + q - 2}.$$

Note that $f_\rho(x) = (g_\rho(x))^3$. Then one can see that

$$(4.24) \quad |g_\rho(x) - g_\rho(y)|_p = \frac{|x - y|_p |\rho - 1|_p |\rho + q - 1|_p}{|x + \rho + q - 2|_p |y + \rho + q - 2|_p},$$

$$(4.25) \quad g_\rho^{-1}(x) = \frac{(\rho + q - 2)x - q + 1}{\rho - x}$$

Moreover, one has the following

Lemma 4.7. *Let $|q - 1|_p < 1$, and $|\rho|_p \leq |q - 1|_p^2$. The following assertions hold true:*

- (i) *If $|x|_p \neq 1$, then $|g_\rho(x)|_p \leq \max\{|q - 1|_p, |\rho|_p\}$;*
- (ii) *If $|g_\rho(x)|_p > 1$, then $|x|_p = 1$.*

Proof. (i). Let $|x|_p < 1$, then from (4.23) we get

$$|g_\rho(x)|_p = \left| \frac{\rho x + q - 1}{x + \rho + q - 2} \right|_p = |q - 1|_p < 1.$$

Now assume $|x|_p > 1$, then analogously one finds

$$|g_\rho(x)|_p \begin{cases} = |\rho|_p, & \text{if } |x|_p > \frac{|q-1|_p}{|\rho|_p}, \\ \leq |q - 1|_p, & \text{if } 1 < |x|_p \leq \frac{|q-1|_p}{|\rho|_p}. \end{cases}$$

(ii) Denoting $y = g_\rho(x)$, from (4.25) one finds

$$|x|_p = |g_\rho^{-1}(y)|_p = \left| \frac{(\rho + q - 2)y - q + 1}{\rho - y} \right|_p = \frac{|y|_p}{|y|_p} = 1.$$

□

Theorem 4.8. *Let $|\rho|_p < |q - 1|_p^2 < 1$ and $\sqrt{-3} \in \mathbb{Q}_p$. Assume that $\{h_x\}_{x \in V \setminus \{(0)\}}$ is a solution of (3.2) such that $|h_x|_p \neq 1$ for all $x \in V \setminus \{(0)\}$. Then $h_x = x_1$ for every x .*

Proof. Let us first show that $|h_x|_p < 1$ for all x . Suppose that $|h_x^{(n_0)}|_p > 1$ for some $n_0 \in \mathbb{N}$ and $x \in W_{n_0}$. Since $\{h_x\}$ is a solution of (3.2), therefore, we have

$$(4.26) \quad h_x^{(n_0)} = g_\rho(h_{(x,1)}^{(n_0+1)}) g_\rho(h_{(x,2)}^{(n_0+1)}) g_\rho(h_{(x,3)}^{(n_0+1)}),$$

here we have used coordinate structure of the tree.

Now according to $|h_{(x,i)}^{(n_0+1)}|_p \neq 1$, ($i = 1, 2, 3$), then Lemma 4.7 (i) implies that $|g_\rho(h_{(x,i)}^{(n_0+1)})|_p < 1$, which with (4.26) means $|h_x^{(n_0)}|_p < 1$. It is a contradiction.

Hence, $|h_x|_p < 1$ for all x . Then from (4.24) we obtain

$$(4.27) \quad |g_\rho(h_x) - g_\rho(x_1)|_p = |q - 1|_p |h_x - x_1|_p$$

for any $x \in V \setminus \{(0)\}$.

Now denote

$$\|h^{(n)}\|_p = \max\{|h_x^{(n)}|_p : x \in W_n\}.$$

Let $\epsilon > 0$ be an arbitrary number. Then from the proof of Lemma 4.7 (i) with (4.27), Lemma 2.4 one finds

$$\begin{aligned} |h_x^{(n)} - x_1|_p &= \left| \prod_{i=1}^3 g_\rho(h_{(x,i)}^{(n+1)}) - (g_\rho(x_1))^3 \right|_p \\ &\leq |q-1|_p^2 \max_{i=1,2,3} \left\{ |h_{(x,i)}^{(n+1)} - x_1|_p \right\}. \end{aligned}$$

Thus, we derive

$$\|h^{(n)} - x_1\|_p \leq |q-1|_p^2 \|h^{(n+1)} - x_1\|_p.$$

So, iterating the last inequality N times one gets

$$(4.28) \quad \|h^{(n)} - x_1\|_p \leq |q-1|_p^{2N} \|h^{(n+N)} - x_1\|_p.$$

Choosing N such that $|q-1|_p^{2N} < \epsilon$, from (4.28) we find $\|h^{(n)} - x_1\|_p < \epsilon$. Arbitrariness of ϵ yields that $h_x = x_1$. This completes the proof. \square

From this theorem we conclude that other solutions of (3.2) may exist under condition $|h_x|_p = 1$ for all $x \in V \setminus \{(0)\}$. Note that using arguments of [59] one can show the existence of periodic solutions of (3.2).

5. BOUNDEDNESS AND PHASE TRANSITIONS

In the previous section we have established the existence of generalized p -adic quasi Gibbs measures. In this section we are going to investigate boundedness of the measures, and moreover, establish the occurrence of phase transitions.

First we need an auxiliary result. Assume that \mathbf{h} is a solution of (3.2), then one finds a constant $a_{\mathbf{h}}(x) \in \mathbb{Q}_p$ such that

$$(5.1) \quad \prod_{y \in S(x)} \sum_{j=0}^{q-1} \rho^{\delta_{ij}} h_{j,y} = a_{\mathbf{h}}(x) h_{i,x}$$

for any $i \in \{0, \dots, q-1\}$. This implies

$$(5.2) \quad \prod_{x \in W_n} \prod_{y \in S(x)} \sum_{j=0}^{q-1} \rho^{\delta_{ij}} h_{j,y} = \prod_{x \in W_n} a_{\mathbf{h}}(x) h_{i,x} = A_{\mathbf{h},n} \prod_{x \in W_n} h_{i,x},$$

where

$$(5.3) \quad A_{\mathbf{h},n} = \prod_{x \in W_n} a_{\mathbf{h}}(x).$$

Given $j \in \Phi$, by $\eta^{(j)} \in \Omega_{W_n}$ we denote a configuration on W_n defined as follows: $\eta^{(j)}(x) = j$ for all $x \in W_n$.

Hence, by (2.8),(5.2) we have

$$\begin{aligned}
1 &= \sum_{\sigma \in \Omega_n} \sum_{\omega \in \Omega_{W_n}} \mu_{\mathbf{h}}^{(n+1)}(\sigma \vee \omega) \\
&= \sum_{\sigma \in \Omega_n} \sum_{\omega \in \Omega_{W_n}} \frac{1}{Z_{n+1,\rho}^{(\mathbf{h})}} \rho^{H(\sigma \vee \omega)} \prod_{x \in W_{n+1}} h_{\omega(x),x} \\
&= \frac{1}{Z_{n+1,\rho}^{(\mathbf{h})}} \sum_{\sigma \in \Omega_n} \rho^{H(\sigma)} \prod_{x \in W_n} \prod_{y \in S(x)} \sum_{j=0}^q \rho^{\delta_{\sigma(x),j}} h_{j,y} \\
&= \frac{A_{\mathbf{h},n}}{Z_{n+1,\rho}^{(\mathbf{h})}} \sum_{\sigma \in \Omega_n} \rho^{H(\sigma)} \prod_{x \in W_n} h_{\sigma(x),x} \\
&= \frac{A_{\mathbf{h},n}}{Z_{n+1,\rho}^{(\mathbf{h})}} Z_{n,\rho}^{(\mathbf{h})}
\end{aligned}$$

which implies

$$(5.4) \quad Z_{n+1,\rho}^{(\mathbf{h})} = A_{\mathbf{h},n} Z_{n,\rho}^{(\mathbf{h})}.$$

In this section we basically investigate measures corresponding to solutions of (4.2). Let \tilde{x} be a solution of (4.2), and $\tilde{\mu}$ be the corresponding p -adic measure to such a solution. Then according to (5.4) the partition function $Z_{\rho,n}^{(\tilde{x})}$ corresponding to the measure $\tilde{\mu}$ has the following form

$$(5.5) \quad Z_{\rho,n}^{(\tilde{x})} = a_i^{|V_{n-1}|}$$

where $a = (\tilde{x} + \rho + q - 2)^3$. Note that when $q \geq 3$, one can check that $a \neq 0$.

For a given configuration $\sigma \in \Omega_{V_n}$ denote

$$\#\sigma = \{x \in W_n : \sigma(x) = 1\}.$$

From (2.8),(3.3) and (5.5) we find

$$\begin{aligned}
|\tilde{\mu}(\sigma)|_p &= \frac{|\rho|_p^{H(\sigma)}}{|Z_{\rho,n}^{(\tilde{x})}|_p} \cdot \prod_{x \in W_n} |h_{\sigma(x),x}|_p \\
(5.6) \quad &= \frac{|\tilde{x}|_p^{\#\sigma} |\rho|_p^{H(\sigma)}}{|\tilde{x} + \rho + q - 2|_p^{3|V_{n-1}|}}.
\end{aligned}$$

5.1. Regime $\rho \in \mathcal{E}_p$. In this subsection we consider two modes: $|q|_p < 1$ and $|q|_p = 1$.

In this section we shall prove the existence of a phase transition. Namely one has the following

Theorem 5.1. *Let $p > 3$, $|q|_p \leq p^{-3}$, $\rho \in \mathcal{E}_p$ with $|\rho - 1|_p \leq p^{-3}$. Then for the p -adic q -state Potts model (3.1) on the Cayley tree of order three the p -adic Gibbs measures μ_k , $k = 0, 1, 2, 3$ are unbounded. Moreover, there is a phase transition.*

Proof. Note that under the condition of the theorem, there are four p -adic Gibbs measures μ_k , $k = 0, 1, 2, 3$ (see Theorem 4.3). In this case, for the solutions x_k ($k = 0, 1, 2, 3$) of (4.2) we have $x_k \in \mathcal{E}_p$. Therefore, one gets

$$|x_k + \rho + q - 2|_p = |x_k - 1 + \rho - 1 + q|_p \leq \frac{1}{p}.$$

Hence, from (5.6) one finds

$$|\mu_k(\sigma)|_p = \frac{1}{|x_k + \rho + q - 2|_p^{3|V_{n-1}|}} \geq p^{3|V_{n-1}|}$$

which implies the unboundedness of μ_k . Thus, there is a phase transition. \square

Now consider the case $|q|_p = 1$. In this setting one has

Theorem 5.2. *Let $p > 3$, $\rho \in \mathcal{E}_p$ with $|\rho - 1|_p \leq |4q - 3|_p$, and $|q|_p = |q - 1|_p = |2q - 1|_p = 1$. Let μ_0 be the p -adic Gibbs measure and assume that the conditions of Theorem 4.4 are satisfied, i.e. there is at least one translation-invariant generalized p -adic quasi Gibbs measure $\tilde{\mu}$ for the p -adic q -state Potts model (3.1) on the Cayley tree of order three. Then the measure μ_0 is bounded, and $\tilde{\mu}$ is unbounded. This means that there is a phase transition.*

Proof. We recall that the measure μ_0 corresponds to the solution $x_0 = 1$. It is clear that

$$|x_0 + \rho + q - 2|_p = |\rho - 1 + q|_p = 1.$$

Hence, from (5.6) one finds

$$|\mu_0(\sigma)|_p = \frac{1}{|x_0 + \rho + q - 2|_p^{3|V_{n-1}|}} = 1$$

which yields the boundedness of μ_0 .

Now assume that $\tilde{\mu}$ corresponds to a solution \tilde{x} of (4.5). Note that one has $\tilde{x} = \tilde{z} + A/3$, where \tilde{z} is a solution of (4.6) such that $|\tilde{z}|_p < 1$.

So, from (4.8) we have

$$\begin{aligned} |\tilde{x} + \rho + q - 2|_p &= \left| \tilde{z} + \frac{A}{3} + \rho + q - 2 \right|_p \\ &= \left| \tilde{z} + 1 - (\rho - 1) + \frac{\rho^3 - 1}{3} - q + \rho + q - 2 \right|_p \\ (5.7) \qquad &= |3\tilde{z} + (\rho^3 - 1)|_p \leq \frac{1}{p} \end{aligned}$$

Hence, from (5.6) with (5.7) one gets $|\tilde{\mu}(\sigma)|_p \geq p^{3|V_{n-1}|}$, so $\tilde{\mu}$ is unbounded. This implies that the existence of a phase transition. \square

5.2. Regime $|\rho|_p < 1$. In this subsection we are going to establish the existence of quasi phase transition.

Theorem 5.3. *Let $p > 3$, $|\rho|_p, |q|_p \leq p^{-2}$. Assume that the conditions of Theorem 4.5 are satisfied. In this case, there two translation-invariant generalized p -adic Gibbs measures μ_0 and $\tilde{\mu}$. Then the measures $\mu_0, \tilde{\mu}$ are bounded. This means that there is a quasi phase transition.*

Proof. Note that the measure μ_0 corresponds to the solution $x_0 = 1$. So, we have

$$|x_0 + \rho + q - 2|_p = |\rho + q - 1|_p = 1.$$

Hence, from (5.6) one finds

$$|\mu_0(\sigma)|_p = \frac{|\rho|_p^{H(\sigma)}}{|x_0 + \rho + q - 2|_p^{3|V_{n-1}|}} \leq |\rho|_p.$$

Therefore, μ_0 is bounded.

Now assume that $\tilde{\mu}$ corresponds to a solution \tilde{x} of (4.5). Note that one has $\tilde{x} = \tilde{z} + A/3$, where \tilde{z} is a solution of (4.6) such that

$$|\tilde{z}|_p = \begin{cases} 1, & \text{if } p \neq 19, \\ p^{-1}, & \text{if } p = 19. \end{cases}$$

for the detail we refer to the proof of Theorem 4.5.

So, from (4.8) we have

$$(5.8) \quad \begin{aligned} |\tilde{x} + \rho + q - 2|_p &= \left| \tilde{z} + \frac{A}{3} + q - 1 \right|_p \\ &= \left| \tilde{z} - \frac{1}{3} + \frac{\rho^3}{3} \right|_p. \end{aligned}$$

We want to show that $|\tilde{z} - 1/3|_p = 1$. If $|\tilde{z}|_p < 1$, then it is evident.

Let $|\tilde{z}|_p = 1$, and assume that $|\tilde{z} - 1/3|_p < 1$. Then it yields that $\tilde{z} = 1/3 + \theta\eta_1 p$ for some $\eta_1 \in \mathbb{Z}_p$. Since, \tilde{z} is a solution of (4.6), then from (4.17) one gets

$$\frac{1}{3^3} + \tilde{\eta}_1 p - \left(-\frac{4}{3} + \epsilon p^2 \right) \left(\frac{1}{3} + \eta_1 p \right) = -\frac{38}{27} + \epsilon_1 p^2.$$

This implies that

$$\frac{1}{27} - \frac{4}{9} + \frac{38}{27} \equiv 0 \pmod{p}.$$

The last one is equivalent to $1 \equiv 0 \pmod{p}$ which is impossible. Therefore, $|\tilde{z} - 1/3|_p = 1$, this with (5.8) yields that $|\tilde{x} + \rho + q - 2|_p = 1$. Consequently, from (5.6) one gets that the measure $\tilde{\mu}$ is bounded. Thus, a quasi phase transition occurs. \square

Note that when $k = 2$ the similar phase transition has occurred for the model (see [55]).

Theorem 5.4. *Let $p > 3$, $|\rho|_p < |q - 1|_p^2 < 1$, and assume that $\sqrt{-3}$ exists. Then for the translation-invariant generalized p -adic quasi Gibbs measures μ_k ($k = 0, 1, 2, 3$) one has: the measures μ_0 is unbounded, but the measures μ_k ($k = 1, 2, 3$) are bounded. Moreover, there is a strong phase transition.*

Proof. We again recall that the measure μ_0 corresponds to the solution $x_0 = 1$. So, one has

$$|x_0 + \rho + q - 2|_p = |\rho + q - 1|_p = |q - 1|_p.$$

Hence, from (5.6) one finds

$$(5.9) \quad |\mu_0(\sigma)|_p = \frac{|\rho|_p^{H(\sigma)}}{|x_0 + \rho + q - 2|_p^{3|V_{n-1}|}} = \frac{|\rho|_p^{H(\sigma)}}{|q - 1|_p^{3|V_{n-1}|}}.$$

Now let us choose $\sigma_{0,n} \in \Omega_{V_{2n}}$ as follows

$$\sigma_{0,n}(x) = \begin{cases} 1, & x \in W_{2k}, \\ 0, & x \in W_{2k-1}, \end{cases} \quad 1 \leq k \leq n.$$

Then one can see that $H(\sigma_{0,n}) = 0$, therefore it follows from (5.9) that

$$|\mu_0(\sigma_{0,n})|_p \geq p^{2|V_{2n-1}|} \rightarrow \infty \quad \text{as } n \rightarrow \infty.$$

This yields that the measure μ_0 is not bounded.

From Remark 4.1 we know that the equation (4.5) has three solutions x_1, x_2 and x_3 such that $|x_1|_p < 1$ and $|x_2|_p = |x_3|_p = 1$. Let μ_i be the corresponding generalized p -adic quasi Gibbs measures ($k = 1, 2, 3$). Note that the solutions x_i are defined by z_i solutions of (4.6) via $x_i = z_i + A/3$.

Now consider the measure μ_1 . From $|x_1|_p < 1$ it follows that $|x_1 + \rho + q - 2|_p = 1$, which immediately yields that μ_1 is bounded.

To study $\mu_{2,3}$ we need some analysis. From the proof of Theorem 4.6 we know that $|D|_p < 1$. Due to (A.5) one finds that

$$\left| -3 \left(\frac{z_1}{2} \right)^2 + \alpha \right|_p = |D|_p < 1$$

Now from (A.6) we obtain that

$$(5.10) \quad \left| z_{2,3} + \frac{z_1}{2} \right|_p < 1.$$

On the other hand, from the proof of Theorem A.5 one finds that $|3z_1^2 + \alpha|_p < 1$. The last inequality with (4.20) yields that

$$\left| 3z_1^2 - \frac{1}{3} + \eta p \right|_p < 1,$$

where $\eta \in \mathbb{Z}_p$. Then using the last one we have

$$(5.11) \quad \begin{aligned} \left| z_1 - \frac{1}{3} \right|_p \left| z_1 + \frac{1}{3} \right|_p &= \left| z_1^2 - \frac{1}{9} \right|_p \\ &= \left| 3z_1^2 - \frac{1}{3} \right|_p \\ &= \left| \left(3z_1^2 - \frac{1}{3} + \eta p \right) - \eta p \right|_p < 1 \end{aligned}$$

Hence, (5.11) implies that

$$\left| z_1 - \frac{1}{3} \right|_p < 1 \quad \text{or} \quad \left| z_1 + \frac{1}{3} \right|_p < 1.$$

In both cases, we infer that

$$(5.12) \quad \left| \frac{z_1}{2} + \frac{1}{3} \right|_p = \left| z_1 + \frac{2}{3} \right|_p = 1.$$

Now using (5.10) and (5.12) one gets

$$(5.13) \quad |3z_{2,3} - 1|_p = \left| z_{2,3} - \frac{1}{3} \right|_p = \left| \left(z_{2,3} + \frac{z_1}{2} \right) - \left(\frac{z_1}{2} + \frac{1}{3} \right) \right|_p = 1.$$

By means of (5.13) we find

$$|x_{2,3} + \rho + q - 2|_p = \left| z_{2,3} + A/3 + \rho + q - 2 \right|_p = |3z_{2,3} - 1 + \rho^3|_p = 1.$$

Consequently, the last equality with (5.6) yields that

$$(5.14) \quad \mu_{2,3}(\sigma) = |\rho|_p^{H(\sigma)} \leq 1.$$

So, the measures $\mu_{2,3}$ are bounded.

Now consider the following multiplication (see (5.9),(5.14))

$$(5.15) \quad |\mu_0(\sigma)\mu_2(\sigma)|_p = \frac{|\rho|_p^{2H(\sigma)}}{|q-1|_p^{3|V_{n-1}|}}$$

Let $|\rho|_p = p^{-k}$ and $|q-1|_p = p^{-m}$. Then from $|\rho|_p < |q-1|_p^2 < 1$ one finds that $0 < 2m < k$. Hence, (5.15) can be rewritten as follows

$$(5.16) \quad |\mu_0(\sigma)\mu_2(\sigma)|_p = p^{3m|V_{n-1}|-2kH(\sigma)}.$$

Now we choose $\tilde{\sigma}_n \in \Omega_{V_n}$ as follows

$$\tilde{\sigma}_n(x) = \begin{cases} 1, & x \in V_{n-1}, \\ 0, & x \in W_n, \end{cases}$$

Then one can see that $H(\tilde{\sigma}_n) = |V_{n-1}|$.

From $3m < 4m < 2k$ we conclude that

$$3m|V_{n-1}| < 2k|V_{n-1}| = 2kH(\tilde{\sigma}_n)$$

so, the last inequality with (5.16) yields that

$$(5.17) \quad |\mu_0(\tilde{\sigma}_n)\mu_2(\tilde{\sigma}_n)|_p \leq 1 \quad \text{for all } n \geq 1.$$

For the sake of simplicity we assume that $3m > k$. Then one finds

$$3m|V_{n-1}| - kH(\tilde{\sigma}_n) = (3m - k)|V_{n-1}| > 0$$

which with (5.9) implies that $|\mu_0(\tilde{\sigma}_n)|_p \rightarrow \infty$ as $n \rightarrow \infty$. This with (5.17) yields that $|\mu_2(\tilde{\sigma}_n)|_p \rightarrow 0$ as $n \rightarrow \infty$. Consequently, there exists a strong phase transition.

Note that if $3m \leq k$, the same phase transition occurs, but it is technically involved. Therefore, we leave this case. \square

6. CONCLUSIONS

In the present paper, we have studied a phase transition problem for the q -state p -adic Potts model over the Cayley tree of order three. We consider a more general notion of p -adic Gibbs measure which depends on parameter $\rho \in \mathbb{Q}_p$. Such a measure is called *generalized p -adic quasi Gibbs measure*. When ρ equals to p -adic exponent, then it coincides with the usual p -adic Gibbs measure (see [58]). When $\rho = p$, then it coincides with p -adic quasi Gibbs measure (see [55]). In the present paper we have considered two regimes with respect to the values of $|\rho|_p$. Namely, in the first regime, one takes $\rho = \exp_p(J)$ for some $J \in \mathbb{Q}_p$, in the second one we let $|\rho|_p < 1$. In each regime, we first find conditions for the existence of generalized p -adic quasi Gibbs measures. Furthermore, in the first regime, we established the existence of the phase transition under the condition $|q|_p \leq p^{-3}$. If $|q|_p = 1$, we proved the existence of a phase transition under some conditions. In the second regime, when $|\rho|_p, |q|_p \leq p^{-2}$ we proved the existence of a quasi phase transition. It turns out that if $|\rho|_p < |q-1|_p^2 < 1$, then one finds the existence of the strong phase transition.

ACKNOWLEDGEMENT

The first named author (F.M.) acknowledges the Scientific and Technological Research Council of Turkey (TUBITAK) for support, and Zirve University for kind hospitality.

APPENDIX A. CUBIC EQUATION $x^3 + ax = b$ OVER \mathbb{Z}_p^*

In this section we are going to provide a criterion of the existence of a solution of the cubic equation

$$(A.1) \quad x^3 + ax = b,$$

where $a, b \in \mathbb{Q}_p$.¹

Let us consider the following depressed cubic equation in the field $\mathbb{F}_p = \mathbb{Z}/p\mathbb{Z}$

$$(A.2) \quad x^3 + \bar{a}x = \bar{b},$$

where $\bar{a}, \bar{b} \in \mathbb{F}_p$. We assume that $\bar{a} \neq \bar{0}$ and $\bar{b} \neq \bar{0}$. The number of solutions $\mathbf{N}_{\mathbb{F}_p}(x^3 + \bar{a}x - \bar{b})$ of this equation was described in [68].

Proposition A.1 ([68]). *Let $p > 3$ be a prime number and $\bar{a}, \bar{b} \in \mathbb{F}_p$ with $\bar{a}\bar{b} \neq \bar{0}$. Let $\bar{D} = -4\bar{a}^3 - 27\bar{b}^2$ and $u_{n+3} = \bar{b}u_n - \bar{a}u_{n+1}$ for $n \in \mathbb{N}$ with $u_1 = \bar{0}$, $u_2 = -\bar{a}$, $u_3 = \bar{b}$. Then the following holds true:*

$$\mathbf{N}_{\mathbb{F}_p}(x^3 + \bar{a}x - \bar{b}) = \begin{cases} 3 & \text{if } \bar{D}u_{p-2}^2 = \bar{0} \\ 0 & \text{if } \bar{D}u_{p-2}^2 = 9\bar{a}^2 \\ 1 & \text{if } \bar{D}u_{p-2}^2 \neq \bar{0}, 9\bar{a}^2 \end{cases}$$

Proposition A.2 ([68]). *Let $p > 3$ be a prime number and $\bar{a}, \bar{b} \in \mathbb{F}_p$, $\bar{a}\bar{b} \neq \bar{0}$. Let $\bar{D} = -4\bar{a}^3 - 27\bar{b}^2$ and $u_{n+3} = \bar{b}u_n - \bar{a}u_{n+1}$ with $\bar{D}u_{p-2}^2 \neq 9\bar{a}^2$, $u_1 = \bar{0}$, $u_2 = -\bar{a}$, $u_3 = \bar{b}$.*

I Let $\bar{D}u_{p-2}^2 = \bar{0}$. Then the following statements holds true:

I.1 The equation (A.2) has 3 distinct solutions in \mathbb{F}_p if and only if $\bar{D} \neq \bar{0}$. Moreover, one has that $3\bar{x}^2 + \bar{a} \neq 0$ for any root \bar{x} ;

I.2 The equation (A.2) has 2 distinct solutions in \mathbb{F}_p while one of them of multiplicity 2 if and only if $\bar{D} = \bar{0}$. If \bar{x}_1, \bar{x}_2 are 2 distinct solutions while \bar{x}_1 is a multiple solution then $\bar{x}_1 = \frac{3\bar{b}}{2\bar{a}}$, $\bar{x}_2 = -\frac{3\bar{b}}{\bar{a}}$, and $3\bar{x}_2^2 + \bar{a} \neq \bar{0}$;

I.3 The equation (A.2) does not have any solution of multiplicity 3.

II Let $\bar{D}u_{p-2}^2 \neq \bar{0}, 9\bar{a}^2$. If \bar{x} is a solution of the equation (A.2) then $3\bar{x}^2 + \bar{a} \neq \bar{0}$.

Remark A.1. Due to Proposition A.2, one may conclude that under the assumption of Proposition A.2, there always exists at least one solution \bar{x} of the equation (A.2) such that $3\bar{x}^2 + \bar{a} \neq \bar{0}$.

In what follows we need the following auxiliary result.

Proposition A.3. *Let p be any prime. Suppose that the equation (A.1) has a solution in \mathbb{Z}_p^* , where $a, b \in \mathbb{Q}_p$ with $ab \neq 0$. Then one of the following conditions holds true:*

- (i) $|a|_p < |b|_p = 1$;
- (ii) $|b|_p < |a|_p = 1$;
- (iii) $|a|_p = |b|_p \geq 1$.

¹Material of this section is taken from [57]

The proof is straightforward.

We are going to state the solvability criterion for the depressed cubic equation (A.1) in domain \mathbb{Z}_p^* at $p > 3$.

Let $a, b \in \mathbb{Q}_p$ be two nonzero p -adic numbers with $a = \frac{a^*}{|a|_p}$, $b = \frac{b^*}{|b|_p}$ where $a^*, b^* \in \mathbb{Z}_p^*$ with $a^* = a_0 + a_1 \cdot p + a_2 \cdot p^2 + \dots$ and $b^* = b_0 + b_1 \cdot p + b_2 \cdot p^2 + \dots$.

We set $D_0 = -4a_0^3 - 27b_0^2$ and $u_{n+3} = b_0u_n - a_0u_{n+1}$ with $u_1 = 0$, $u_2 = -a_0$, and $u_3 = b_0$ for $n = \overline{1, p-3}$

Theorem A.4. *Let $p > 3$ be a prime. Then the equation (A.1) has a solution in \mathbb{Z}_p^* if and only if one of the following conditions hold true:*

- I.1 $|a|_p < |b|_p = 1$ and $b_0^{\frac{p-1}{(3,p-1)}} \equiv 1 \pmod{p}$;
- I.2 $|b|_p < |a|_p = 1$ and $(-a_0)^{\frac{p-1}{2}} \equiv 1 \pmod{p}$;
- I.3 $|a|_p = |b|_p = 1$ and $D_0u_{p-2}^2 \not\equiv 9a_0^2 \pmod{p}$;
- I.4 $|a|_p = |b|_p > 1$.

Proof. Note that if the equation (A.1) has a solution in \mathbb{Z}_p^* , then due to Proposition A.3 there are three possible cases. Let us consider every case.

I.1. Assume that the equation (A.1) has a solution $x \in \mathbb{Z}_p^*$, when $|a|_p < 1$, $|b|_p = 1$. Due to $|a|_p < 1$ and $|b|_p = 1$, we get

$$x_0^3 + ax_0 \equiv x_0^3 \equiv b_0 \pmod{p}.$$

Hence Proposition 2.2 yields that $b_0^{\frac{p-1}{(3,p-1)}} \equiv 1 \pmod{p}$.

Now assume that $|a|_p < 1$, $|b|_p = 1$, and $b_0^{\frac{p-1}{(3,p-1)}} \equiv 1 \pmod{p}$ are satisfied. From $b_0^{\frac{p-1}{(3,p-1)}} \equiv 1 \pmod{p}$ due to Proposition 2.2 one can find $x_0 \in \mathbb{Z}$ such that $x_0^3 \equiv b_0 \pmod{p}$ and $(x_0, p) = 1$. Let us consider the following polynomial $f_{a,b}(x) = x^3 + ax - b$. By $|a|_p < 1$ one gets

$$f_{a,b}(x_0) = x_0^3 + ax_0 - b \equiv x_0^3 - b_0 \equiv 0 \pmod{p}, \quad f'_{a,b}(x_0) = 3x_0^2 + a \not\equiv 0 \pmod{p}.$$

Hence, Hensel's Lemma 2.1 implies the existence of a solution $x \in \mathbb{Z}_p$ of (A.1) with $|x - x_0|_p \leq \frac{1}{p}$, where $|x_0|_p = 1$. This means that $|x|_p = 1$, i.e., $x \in \mathbb{Z}_p^*$.

I.2. Assume that the equation (A.1) has a solution $x \in \mathbb{Z}_p^*$ when $|a|_p = 1$, $|b|_p < 1$. Then it follows that

$$x_0^3 + a_0x_0 \equiv x_0(x_0^2 + a_0) \equiv b \equiv 0 \pmod{p}.$$

This means that $x_0^2 \equiv -a_0 \pmod{p}$, and again Proposition 2.2 implies $(-a_0)^{\frac{p-1}{2}} \equiv 1 \pmod{p}$.

Now we suppose that $|a|_p = 1$, $|b|_p < 1$, and $(-a_0)^{\frac{p-1}{2}} \equiv 1 \pmod{p}$. The congruence $(-a_0)^{\frac{p-1}{2}} \equiv 1 \pmod{p}$ yields the existence $x_0 \in \mathbb{Z}$ such that $x_0^2 + a_0 \equiv 0 \pmod{p}$ and $(x_0, p) = 1$. Again consider the polynomial $f_{a,b}(x)$. Due to $|b|_p < 1$ one finds

$$\begin{aligned} f_{a,b}(x_0) &= x_0^3 + ax_0 - b \equiv x_0(x_0^2 + a_0) \equiv 0 \pmod{p}, \\ f'_{a,b}(x_0) &= 3x_0^2 + a \equiv 3(x_0^2 + a_0) - 2a_0 \not\equiv 0 \pmod{p}. \end{aligned}$$

Hence, Hensel's Lemma 2.1 implies that the equation (A.1) has a solution $x \in \mathbb{Z}_p$ with $|x - x_0|_p \leq \frac{1}{p}$ where $|x_0|_p = 1$. This means $x \in \mathbb{Z}_p^*$.

I.3. Assume that the equation (A.1) has a solution $x \in \mathbb{Z}_p^*$ when $|a|_p = 1$, $|b|_p = 1$. Then we have

$$x_0^3 + a_0x_0 \equiv b_0 \pmod{p}.$$

This means that the depressed cubic equation $x^3 + a_0x = b_0$ has at least one solution in the finite field \mathbb{F}_p . According to Proposition A.1 one has $D_0u_{p-2}^2 \not\equiv 9a_0^2 \pmod{p}$.

Now we assume that $|a|_p = 1$, $|b|_p = 1$, and $D_0u_{p-2}^2 \not\equiv 9a_0^2 \pmod{p}$ are satisfied. The congruence $D_0u_{p-2}^2 \not\equiv 9a_0^2 \pmod{p}$ yields the equation $x^3 + a_0x \equiv b_0 \pmod{p}$ has at least one solution. Proposition A.2 implies that among all solutions of $x^3 + a_0x \equiv b_0 \pmod{p}$, there always exists at least one solution x_0 such that $3x_0^2 + a_0 \not\equiv 0 \pmod{p}$ with $(x_0, p) = 1$.

Again, if we apply Hensel's Lemma 2.1 to $f_{a,b}(x)$ at the point $x = x_0$ then we get that (A.1) has a solution $x \in \mathbb{Z}_p$ with $|x - x_0|_p \leq \frac{1}{p}$ where $|x_0|_p = 1$. So, $x \in \mathbb{Z}_p^*$.

I.4. In this case, it is enough to establish the existence of a solution of (A.1) under the condition $|a|_p = |b|_p > 1$. We have $a = p^{-m}a^*$, $b = p^{-m}b^*$ with $a^*, b^* \in \mathbb{Z}_p^*$, for some $m \geq 1$. It is clear that the depressed equation $x^3 + p^{-m}a^*x = p^{-m}b^*$ has a solution in \mathbb{Z}_p^* if and only if the equation $p^m x^3 + a^*x = b^*$ has a solution in \mathbb{Z}_p^* , where $a^*, b^* \in \mathbb{Z}_p^*$. To this end, we consider the polynomial $g(x) = p^m x^3 + a^*x - b^*$. If a_0, b_0 are the first digits of $a^*, b^* \in \mathbb{Z}_p^*$ then one finds is $x_0 \in \mathbb{Z}$ such that $a_0x_0 \equiv b_0 \pmod{p}$ with $(x_0, p) = 1$. Then, we obtain

$$g(x_0) \equiv p^m x_0^3 + a_0x_0 - b_0 \equiv 0 \pmod{p}, \quad g'(x_0) \equiv 3p^m x_0^2 + a_0 \not\equiv 0 \pmod{p}.$$

Again, due to Hensel's Lemma 2.1, the equation $p^m x^3 + a^*x = b^*$ has a solution $x \in \mathbb{Z}_p^*$ with $|x - x_0|_p \leq \frac{1}{p}$. This means that $x \in \mathbb{Z}_p^*$. \square

Let $D = -4(a|a|_p)^3 - 27(b|b|_p)^2$. We have $D = \frac{D^*}{|D|_p}$ whenever $D \neq 0$, where $D^* \in \mathbb{Z}_p^*$ and $D^* = d_0 + d_1 \cdot p + d_2 \cdot p^2 + \dots$

Theorem A.5. *Let $p > 3$ be a prime. Then one has*

$$\mathbf{N}_{\mathbb{Z}_p^*}(x^3 + ax - b) = \begin{cases} 3, & |a|_p < |b|_p = 1, p \equiv 1 \pmod{3}, b_0^{\frac{p-1}{3}} \equiv 1 \pmod{p} \\ 3, & |a|_p = |b|_p = 1, D = 0 \\ 3, & |a|_p = |b|_p = 1, 0 < |D|_p < 1, 2 \mid \log_p |D|_p, d_0^{\frac{p-1}{2}} \equiv 1 \pmod{p} \\ 3, & |a|_p = |b|_p = 1, |D|_p = 1, u_{p-2} \equiv 0 \pmod{p} \\ 2, & |b|_p < |a|_p = 1, (-a_0)^{\frac{p-1}{2}} \equiv 1 \pmod{p} \\ 1, & |a|_p < |b|_p = 1, p \equiv 2 \pmod{3} \\ 1, & |a|_p = |b|_p = 1, 0 < |D|_p < 1, 2 \mid \text{ord}_p(D), d_0^{\frac{p-1}{2}} \not\equiv 1 \pmod{p} \\ 1, & |a|_p = |b|_p = 1, 0 < |D|_p < 1, 2 \nmid \text{ord}_p(D), \\ 1, & |a|_p = |b|_p = 1, D_0u_{p-2}^2 \not\equiv 0, 9a_0^2 \pmod{p} \\ 1, & |a|_p = |b|_p > 1 \\ 0, & \text{otherwise} \end{cases}$$

Proof. Let $p > 3$ be a prime number and $a, b \in \mathbb{Q}_p$ be two nonzero p -adic numbers.

We want to describe the number $\mathbf{N}_{\mathbb{Z}_p^*}(x^3 + ax - b)$ of solutions of the depressed cubic equation (A.1) in the domain \mathbb{Z}_p^* .

According to Theorem A.4, the equation (A.1) has a solution in \mathbb{Z}_p^* if and only if one of the conditions I.1-I.4 holds true. We want to find a number of solutions in every case.

First assume that $|a|_p < |b|_p = 1$ and $b_0^{\frac{p-1}{(3,p-1)}} \equiv 1 \pmod{p}$. In this case, we showed in the proof of I.1, the number of solutions of (A.1) in \mathbb{Z}_p^* is the same as the number of solutions of the equation $x_0^3 = b_0$ in \mathbb{F}_p . Hence, Proposition 2.2 implies that the last equation has 3 distinct solutions if $p \equiv 1 \pmod{3}$ and it has a unique solution if $p \equiv 2 \pmod{3}$.

Therefore, if $|a|_p < |b|_p = 1$, $p \equiv 1 \pmod{3}$, and $b_0^{\frac{p-1}{3}} \equiv 1 \pmod{p}$ then $\mathbf{N}_{\mathbb{Z}_p^*}(x^3 + ax - b) = 3$, and if $|a|_p < |b|_p = 1$, $p \equiv 2 \pmod{3}$ and $b_0^{p-1} \equiv 1 \pmod{p}$ then $\mathbf{N}_{\mathbb{Z}_p^*}(x^3 + ax - b) = 1$. It is worth mentioning that one always has $b_0^{p-1} \equiv 1 \pmod{p}$ since $(b_0, p) = 1$.

Now consider $|b|_p < |a|_p = 1$ and $(-a_0)^{\frac{p-1}{2}} \equiv 1 \pmod{p}$. We showed that the number of solutions of (A.1) in \mathbb{Z}_p^* is the same as the number of solutions of the equation $x_0^2 = -a_0$ in \mathbb{F}_p . From $(-a_0)^{\frac{p-1}{2}} \equiv 1 \pmod{p}$, one concludes that the last equation has 2 distinct solutions in \mathbb{F}_p . Hence, $\mathbf{N}_{\mathbb{Z}_p^*}(x^3 + ax - b) = 2$.

Now assume that $|a|_p = |b|_p = 1$ and $D_0 u_{p-2}^2 \not\equiv 9a_0^2 \pmod{p}$. In this case, we know that the equation (A.1) has a solution \bar{x} such that $\bar{x} \equiv \bar{x}_0 \pmod{p}$, where \bar{x}_0 is a solution of

$$(A.3) \quad x_0^3 + a_0 x_0 \equiv b_0 \pmod{p}$$

such that $3\bar{x}_0^2 + a_0 \not\equiv 0 \pmod{p}$.

Due to Proposition A.1, if $D_0 u_{p-2}^2 \not\equiv 0, 9a_0^2 \pmod{p}$ then (A.3) does not have any solution except \bar{x}_0 . Therefore, due to Hensel's Lemma 2.1, if $|a|_p = |b|_p = 1$ and $D_0 u_{p-2}^2 \not\equiv 0, 9a_0^2 \pmod{p}$ then $\mathbf{N}_{\mathbb{Z}_p^*}(x^3 + ax - b) = 1$.

Now, assume that $|a|_p = |b|_p = 1$ and $D_0 u_{p-2}^2 \equiv 0 \pmod{p}$. Then Proposition A.1 yields that the equation (A.3) has 2 more solutions besides \bar{x}_0 . We denote them by x_0 and y_0 . There is no loss of generality in assuming that $0 < x_0, y_0 < p$.

If $D_0 \not\equiv 0 \pmod{p}$ then Proposition A.2 implies that all solutions x_0, y_0, \bar{x}_0 of (A.3) are distinct from each other and one has $3x_0^2 + a_0 \not\equiv 0 \pmod{p}$, $3y_0^2 + a_0 \not\equiv 0 \pmod{p}$. Therefore, by applying Hensel's Lemma 2.1 to (A.1) one finds 2 more solutions x, y . Moreover, we have \bar{x} such that $x \equiv x_0 \pmod{p}$, $y \equiv y_0 \pmod{p}$. Hence, $|a|_p = |b|_p = 1$, $D_0 \not\equiv 0 \pmod{p}$ (equivalently $|D|_p = 1$) and $u_{p-2} \equiv 0 \pmod{p}$ then $\mathbf{N}_{\mathbb{Z}_p^*}(x^3 + ax - b) = 3$.

Now we suppose that $D_0 \equiv 0 \pmod{p}$. Then Proposition A.2 implies that two solutions x_0, y_0 of (A.3) are equal to each other, and one has $\bar{x}_0 = -2x_0$, $3x_0^2 + a_0 \equiv 0 \pmod{p}$. Let us study this case in a detail.

Since \bar{x} is a solution of the equation (A.1), then one finds

$$x^3 + ax - b = (x - \bar{x})(x^2 + \bar{x}x + \bar{x}^2 + a).$$

We are going to study the quadratic equation

$$(A.4) \quad x^2 + \bar{x}x + \bar{x}^2 + a = 0.$$

It is clear that

$$\left(x + \frac{\bar{x}}{2}\right)^2 = -\left(3\left(\frac{\bar{x}}{2}\right)^2 + a\right).$$

Since $a, \bar{x} \in \mathbb{Z}_p^*$ and $p > 3$, we have that $\left|3\left(\frac{\bar{x}}{2}\right)^2 + a\right|_p \leq 1$. Since \bar{x} is a solution of (A.1), one gets $4\bar{x}\left(3\left(\frac{\bar{x}}{2}\right)^2 + a\right) = a\bar{x} + 3b$. We then have that

$$\begin{aligned} D &= -4a^3 - 27b^2 = 3(a^2\bar{x}^2 - 9b^2) - a^2(3\bar{x}^2 + 4a) \\ &= 12\bar{x}\left(3\left(\frac{\bar{x}}{2}\right)^2 + a\right)(a\bar{x} - 3b) - 4a^2\left(3\left(\frac{\bar{x}}{2}\right)^2 + a\right) \\ &= 4\left(3\left(\frac{\bar{x}}{2}\right)^2 + a\right)(3a\bar{x}^2 - 9b\bar{x} - a^2). \end{aligned}$$

Due to $\bar{x} \equiv \bar{x}_0 = -2x_0 \pmod{p}$, $3x_0^2 + a_0 \equiv 0 \pmod{p}$, and $2x_0a_0 \equiv 3b_0 \pmod{p}$ we find that $3a\bar{x}^2 - 9b\bar{x} - a^2 \equiv -9a_0^2 \pmod{p}$. This means that $|3a\bar{x}^2 - 9b\bar{x} - a^2|_p = 1$, and $-(3a\bar{x}^2 - 9b\bar{x} - a^2)$ is a complete square of some p -adic integer number. Therefore, we obtain that

$$(A.5) \quad -\left(3\left(\frac{\bar{x}}{2}\right)^2 + a\right) = \frac{D}{-4(3a\bar{x}^2 - 9b\bar{x} - a^2)}$$

Let us analyze the quadratic equation (A.4).

If $D = 0$ then (A.4) has solutions $x_1 = x_2 = \frac{3b}{2a}$ in \mathbb{Z}_p^* . Therefore, $|a|_p = |b|_p = 1$ and $D = 0$, then (A.1) has 3 solutions such that $x_1 = x_2 = \frac{3b}{2a}$ and $x_3 = -\frac{3b}{a}$.

Let $D \neq 0$. Since $D \equiv D_0 \equiv 0 \pmod{p}$, there exists $k \in \mathbb{N}$ such that $|D|_p = p^{-k}$, i.e., $D = \frac{D^*}{|D|_p}$, where $D^* \in \mathbb{Z}_p^*$ with $D^* = d_0 + d_1 \cdot p + d_2 \cdot p^2 + \dots$.

The quadratic equation (A.4) has a solution if and only if $\text{ord}_p(D) = -k$ is even number and $d_0^{\frac{p-1}{2}} \equiv 1 \pmod{p}$. In this case $-\left(3\left(\frac{\bar{x}}{2}\right)^2 + a\right)$ is a complete square and (A.4) has two distinct solutions in \mathbb{Z}_p^* , which are given by

$$(A.6) \quad x_{\pm} = -\frac{\bar{x}}{2} \pm \sqrt{-\left(3\left(\frac{\bar{x}}{2}\right)^2 + a\right)}$$

So, if $|a|_p = |b|_p = 1$, $0 < |D|_p < 1$, $2 \mid \text{ord}_p(D)$, and $d_0^{\frac{p-1}{2}} \equiv 1 \pmod{p}$ then $\mathbf{N}_{\mathbb{Z}_p^*}(x^3 + ax - b) = 3$. If $|a|_p = |b|_p = 1$, $0 < |D|_p < 1$, $2 \mid \text{ord}_p(D)$, and $d_0^{\frac{p-1}{2}} \not\equiv 1 \pmod{p}$ or $|a|_p = |b|_p = 1$, $0 < |D|_p < 1$, and $2 \nmid \text{ord}_p(D)$ then $\mathbf{N}_{\mathbb{Z}_p^*}(x^3 + ax - b) = 1$.

Now consider the case $|a|_p = |b|_p > 1$. We know that $\mathbf{N}_{\mathbb{Z}_p^*}(x^3 + ax - b)$ is the same as the number of solutions of the linear equation $a_0x_0 = b_0$ in \mathbb{F}_p . Since $a_0 \neq 0$, the last equation has a unique solution. Therefore, if $|a|_p = |b|_p > 1$ then $\mathbf{N}_{\mathbb{Z}_p^*}(x^3 + ax - b) = 1$. □

REFERENCES

- [1] Albeverio S., Karwowski W. A random walk on p -adics, the generator and its spectrum, *Stochastic. Process. Appl.* **53** (1994) 1-22.
- [2] Albeverio S., Zhao X. On the relation between different constructions of random walks on p -adics, *Markov Process. Related Fields* **6** (2000) 239-256.
- [3] Albeverio S., Zhao X. Measure-valued branching processes associated with random walks on p -adics, *Ann. Probab.* **28**(2000) 1680-1710.
- [4] Albeverio S., Khrennikov A., Kloeden P.E., Memory retrieval as a p -adic dynamical system, *BioSys.* **49**(1999), 105-115.
- [5] Albeverio S., Khrennikov A. Yu., Shelkovich, V.M. *Theory of p -adic Distributions. Linear and Nonlinear Models*, Cambridge Univ. Press, Cambridge, 2010.
- [6] Anashin V., Khrennikov A., *Applied Algebraic Dynamics*, Walter de Gruyter, Berlin, New York, 2009.
- [7] Areféva I. Ya., Dragovic B., Volovich I.V. p -adic summability of the anharmonic oscillator, *Phys. Lett. B* **200**(1988) 512-514.
- [8] Areféva I. Ya., Dragovic B., Frampton P.H., Volovich I.V. The wave function of the Universe and p -adic gravity, *Int. J. Modern Phys. A* **6**(1991) 4341-4358.
- [9] Arrowsmith D.K., Vivaldi F., Some p -adic representations of the Smale horseshoe, *Phys. Lett. A* **176**(1993), 292-294.
- [10] Arrowsmith D.K., Vivaldi F., Geometry of p -adic Siegel discs. *Physica D*, **71**(1994), 222-236.

- [11] Avetisov V.A., Bikulov A.H., Kozyrev S.V. Application of p -adic analysis to models of spontaneous breaking of the replica symmetry, *J. Phys. A: Math. Gen.* **32**(1999) 8785–8791.
- [12] Beltrametti E., Cassinelli G. Quantum mechanics and p -adic numbers, *Found. Phys.* **2**(1972) 1–7.
- [13] Besser A., Deninger C., p -adic Mahler measures, *J. Reine Angew. Math.* **517** (1999), 1950.
- [14] Böcherer, S., Panchishkin, A. A. Admissible p -adic measures attached to triple products of elliptic cusp forms. *Doc. Math.* 2006, Extra Vol., 77-132.
- [15] Böcherer, S., Schmidt, C.-G. p -adic measures attached to Siegel modular forms. *Ann. Inst. Fourier (Grenoble)* **50** (2000), 1375-1443
- [16] Childress, N., The coefficients of a p -adic measure and its Γ -transform. *Manuscripta Math.* **116** (2005), 249-263.
- [17] Del Muto M., Figà-Talamanca A. Diffusion on locally compact ultrametric spaces, *Expo. Math.* **22**(2004) 197-211.
- [18] Dobrushin R.L. The problem of uniqueness of a Gibbsian random field and the problem of phase transitions, *Funct. Anal. Appl.* **2** (1968) 302–312.
- [19] Dobrushin R.L. Prescribing a system of random variables by conditional distributions, *Theor. Probab. Appl.* **15**(1970) 458–486.
- [20] We point out that stochastic processes on the field \mathbb{Q}_p of p -adic numbers with values of real numbers have been studied by many authors, for example, [1, 2, 3, 17, 47, 73]. In those investigations wide classes of Markov processes on \mathbb{Q}_p were constructed and studied. In our case the situation is different, since probability measures take their values in \mathbb{Q}_p . This leads our investigation to some difficulties. For example, there is no information about the compactness of p -adic values probability measures.
- [21] Freund P.G.O., Olson M. Non-Archimedean strings, *Phys. Lett. B* **199**(1987) 186–190.
- [22] Ganikhodjaev N.N., On pure phases of the three-state ferromagnetic Potts model on the Bethe lattice order two, *Theor. Math. Phys.* **85** (1990) 163–175.
- [23] Ganikhodjaev N.N., Mukhamedov F.M., Rozikov U.A. Phase transitions of the Ising model on \mathbb{Z} in the p -adic number field, *Uzbek. Math. Jour.* **4** (1998) 23–29 (Russian).
- [24] Ganikhodjaev N.N., Mukhamedov F.M., Rozikov U.A. Existence of a phase transition for the Potts p -adic model on the set \mathbb{Z} , *Theor. Math. Phys.* **130** (2002) 425–431.
- [25] Ganikhodjaev N.N., Rozikov U.A. Description of periodic extreme Gibbs measures of some lattice model on the Cayley tree. *Theor. Math. Phys.* **111** (1997), 480–486.
- [26] Georgii H.O. *Gibbs measures and phase transitions*, Walter de Gruyter, Berlin, 1988.
- [27] Gandolfo D., Rozikov U.A., Ruiz J. On p -adic Gibbs Measures for Hard Core Model on a Cayley Tree, **ArXiv: arXiv:1107.4884**.
- [28] Harris, M., Tilouine, J., p -adic measures and square roots of special values of triple product L-functions. *Math. Ann.* **320** (2001), 127-147.
- [29] Kaneko H., Kochubei A.N., Weak solutions of stochastic differential equations over the field of p -adic numbers, *Tohoku Math. J.* **59**(2007), 547–564.
- [30] Khamraev M., Mukhamedov F.M. On p -adic λ -model on the Cayley tree, *Jour. Math. Phys.* **45**(2004) 4025–4034.
- [31] Katsaras A.K. Extensions of p -adic vector measures, *Indag. Math.N.S.* **19** (2008) 579–600.
- [32] Katsaras A.K. On spaces of p -adic vector measures, *P-Adic Numbers, Ultrametric Analysis, Appl.* **1** (2009) 190–203.
- [33] Katsaras A.K. On p -adic vector measures, *Jour. Math. Anal. Appl.* **365** (2010), 342-357.
- [34] Khrennikov A.Yu. p -adic valued probability measures, *Indag. Mathem. N.S.* **7**(1996) 311–330.
- [35] Khrennikov A.Yu. *p -adic Valued Distributions in Mathematical Physics*, Kluwer Academic Publisher, Dordrecht, 1994.
- [36] Khrennikov A.Yu. *Non-Archimedean analysis: quantum paradoxes, dynamical systems and biological models*, Kluwer Academic Publisher, Dordrecht, 1997.
- [37] Khrennikov A.Yu., The Bernoulli theorem for probabilities that take p -adic values. *Doklady Mathematics*, **55** (1998), N. 3, 402–405.

- [38] Khrennikov A.Yu., Generalized Probabilities Taking Values in Non-Archimedean Fields and in Topological Groups, *Russian J. Math. Phys.* **14** (2007), 142-159.
- [39] Khrennikov A.Yu., Kozyrev S.V., Wavelets on ultrametric spaces, *Appl. Comput. Harmonic Anal.* **19**(2005), 61–76.
- [40] Khrennikov A.Yu., Kozyrev S.V., Ultrametric random field. *Infin. Dimens. Anal. Quantum Probab. Relat. Top.* **9**(2006), 199-213.
- [41] Khrennikov A.Yu., Kozyrev S.V., Replica symmetry breaking related to a general ultrametric space I,II,III, *Physica A*, **359**(2006), 222-240; 241-266; **378**(2007), 283-298.
- [42] Khrennikov A.Yu., Ludkovsky S. Stochastic processes on non-Archimedean spaces with values in non-Archimedean fields, *Markov Process. Related Fields* **9**(2003) 131–162.
- [43] Khrennikov A., Mukhamedov F., Mendes J.F.F. On p -adic Gibbs measures of countable state Potts model on the Cayley tree, *Nonlinearity* **20**(2007) 2923-2937.
- [44] Khrennikov A.Yu., Nilsson M. *p -adic deterministic and random dynamical systems*, Kluwer, Dordrecht, 2004.
- [45] Khrennikov A.Yu., Yamada S., van Rooij A., Measure-theoretical approach to p -adic probability theory, *Annals Math. Blaise Pascal* **6** (1999) 21–32.
- [46] Koblitz N. *p -adic numbers, p -adic analysis and zeta-function*, Berlin, Springer, 1977.
- [47] Kochubei A.N. *Pseudo-differential equations and stochastics over non-Archimedean fields*, Mongr. Text-books Pure Appl. Math. 244 Marcel Dekker, New York, 2001.
- [48] Kozyrev S.V., Wavelets and spectral analysis of ultrametric pseudodifferential operators *Sbornik Math.* **198**(2007), 97–116.
- [49] Ludkovsky S.V. Non-Archimedean valued quasi-invariant descending at infinity measures, *Int. J. Math. Math. Sci.* **2005**(2005) N. 23, 3799–3817.
- [50] Marinary E., Parisi G. On the p -adic five point function, *Phys. Lett. B* **203**(1988) 52–56.
- [51] Mahler K., *p -adic numbers and their functions*. Cambridge Tracts in Mathematics, **76**, Cambridge Univ. Press, Cambridge-New York, 1981.
- [52] Monna A., Springer T., Integration non-Archimedienne 1, 2. *Indag. Math.* **25** (1963) 634-653.
- [53] Mukhamedov F.M., On the existence of generalized Gibbs measures for the one-dimensional p -adic countable state Potts model, *Proc. Steklov Inst. Math.* **265** (2009) 165-176.
- [54] Mukhamedov F.M., Existence of P -adic quasi Gibbs measure for countable state Potts model on the Cayley tree, *J. Inequal. Appl.* 2012, **2012**:104.
- [55] Mukhamedov F., On p -adic quasi Gibbs measures for $q + 1$ -state Potts model on the Cayley tree, *P -adic Numbers, Ultrametric Anal. Appl.* **2**(2010), 241–251.
- [56] Mukhamedov F., A dynamical system approach to phase transitions for p -adic Potts model on the Cayley tree of order two, *Rep. Math. Phys.* (accepted).
- [57] Mukhamedov F., Omirov B., Saburov M. On cubic equations over P -adic field, *ArXiv: 1204.1743*.
- [58] Mukhamedov F.M., Rozikov U.A. On Gibbs measures of p -adic Potts model on the Cayley tree, *Indag. Math. N.S.* **15** (2004) 85–100.
- [59] Mukhamedov F.M., Rozikov U.A. On inhomogeneous p -adic Potts model on a Cayley tree, *Infin. Dimens. Anal. Quantum Probab. Relat. Top.* **8**(2005) 277–290.
- [60] Peruggi, F., di Liberto F., Monroy G., Phase diagrams of the q -state Potts model on Bethe lattices. *Phys. A* **141** (1987), 151-186.
- [61] Peruggi, F., di Liberto F., Monroy G., The Potts model on Bethe lattices. I. General results. *J. Phys. A* **16** (1983), 811-827.
- [62] A.M.Robert, *A course of p -adic analysis*, Springer, New York, 2000.
- [63] van Rooij A., *Non-archimedean functional analysis*, Marcel Dekker, New York, 1978.
- [64] Rosen K.H., *Elementary number theory and its applications*, Pearson 2011.
- [65] Rozikov U.A. A contour method on Cayley trees. *Jour. Stat. Phys.* **130** (2008), 801–813.
- [66] Schikhof W.H. *Ultrametric Calculus*, Cambridge University Press, Cambridge, 1984.
- [67] Shiryaev A.N. *Probability*, Nauka, Moscow, 1980.
- [68] Z.H. Sun, Cubic and quartic congruences modulo a prime, *J. Num. Theory*, **102** (2003) 41–89.

- [69] Vishek M.M. Nonarchimedean measures connected with Dirichlet series. *Mat. Sb., N. Ser.* **99**(1976), 248–260.
- [70] Vladimirov V.S., Volovich I.V., Zelenov E.I. *p -adic Analysis and Mathematical Physics*, World Scientific, Singapore, 1994.
- [71] Volovich I.V. Number theory as the ultimate physical theory, *p -Adic Numbers, Ultrametric Analysis Appl.* **2**(2010), 77-87; // Preprint TH.4781/87, 1987.
- [72] Volovich I.V. p -adic string, *Classical Quantum Gravity* **4** (1987) L83-L87.
- [73] Yasuda K., Extension of measures to infinite-dimensional spaces over p -adic field, *Osaka J. Math.* **37**(2000) 967-985.
- [74] Zachary S. Countable state space Markov random fields and Markov chains on trees, *Ann. Prob.* **11** (1983), 894-903.
- [75] Wu F.Y., The Potts model, *Rev. Mod. Phys.* **54** (1982) 235–268.
- [76] In [39, 40] a notion of ultrametric Markovianity, which describes independence of contributions to random field from different ultrametric balls, has been introduced, and showed that Gaussian random fields on general ultrametric spaces (which were related with hierarchical trees), which were defined as a solution of pseudodifferential stochastic equation [5, 29], satisfies the Markovianity. In addition, covariation of the defined random field was computed with the help of wavelet analysis on ultrametric spaces (see also [5, 48]). Some applications of the results to replica matrices, related to general ultrametric spaces have been investigated in [41].
- [77] In the real case, when the state space is compact, then the existence follows from the compactness of the set of all probability measures (i.e. Prohorov's Theorem). When the state space is non-compact, then there is a Dobrushin's Theorem [18, 19] which gives a sufficient condition for the existence of the Gibbs measure for a large class of Hamiltonians.

FARRUKH MUKHAMEDOV, DEPARTMENT OF COMPUTATIONAL & THEORETICAL SCIENCES, FACULTY OF SCIENCE, INTERNATIONAL ISLAMIC UNIVERSITY MALAYSIA, P.O. BOX, 141, 25710, KUANTAN, PAHANG, MALAYSIA

E-mail address: far75m@yandex.ru farrukh_m@iiu.edu.my

HASAN AKIN, DEPARTMENT OF MATHEMATICS, FACULTY OF EDUCATION, ZIRVE UNIVERSITY, KIZILHISAR CAMPUS, GAZIANTEP, 27260, TURKEY

E-mail address: hasanakin69@gmail.com