Dynamical System with Boundary Control Associated with Symmetric Semi-Bounded Operator

M.I.Belishev *

Abstract

Let L_0 be a closed densely defined symmetric semi-bounded operator with nonzero defect indexes in a separable Hilbert space \mathcal{H} . It determines a *Green system* $\{\mathcal{H}, \mathcal{B}; L_0, \Gamma_1, \Gamma_2\}$, where \mathcal{B} is a Hilbert space, and $\Gamma_i : \mathcal{H} \to \mathcal{B}$ are the operators related through the Green formula

 $(L_0^*u, v)_{\mathcal{H}} - (u, L_0^*v)_{\mathcal{H}} = (\Gamma_1 u, \Gamma_2 v)_{\mathcal{B}} - (\Gamma_2 u, \Gamma_1 v)_{\mathcal{B}}.$

The boundary operators Γ_i are chosen canonically in the framework of the Vishik theory.

With the Green system one associates a *dynamical system with* boundary control (DSBC)

$u_{tt} + L_0^* u = 0$	in	$\mathcal{H},$	t > 0
$u _{t=0} = u_t _{t=0} = 0$	in	${\mathcal H}$	
$\Gamma_1 u = f$	in	$\mathcal{B},$	$t \ge 0.$

We show that this system is *controllable* if and only if the operator L_0 is completely non-self-adjoint.

A version of the notion of a *wave spectrum* of L_0 is introduced. It is a topological space determined by L_0 and constructed from reachable sets of the DSBC.

^{*}Saint-Petersburg Department of the Steklov Mathematical Institute (POMI), 27 Fontanka, St. Petersburg 191023, Russia; belishev@pdmi.ras.ru. Supported by RFBR grants 11-01-00407A and NSh-4210.2010.1.

0 Introduction

0.1 About the paper

We develop ideas and results of the papers [2] and [5]. The future prospect and goal is a *functional model* of a symmetric semi-bounded operator outlined in [5]. Our paper is a step towards this model, which prepares two of the model basic elements: *DSBC* and *wave spectrum*.

Motivation comes from inverse problems. Namely, the inspiring role is played by the problem of reconstruction of a Riemannian manifold via its boundary inverse data [1],[3],[4]. In accordance with the program, which we promote, to solve the latter problem (and a class of closely related problems) is to construct a certain functional model of a relevant symmetric operator. The operator codes information about the manifold and is determined by inverse data. Its wave spectrum turns out to be isometric to the manifold. By the latter, to decode the information one can find the wave spectrum [5].

The main subjects of the paper are the following.

0.2 Operator L_0

Let \mathcal{H} be a (separable) Hilbert space, L_0 a closed operator in \mathcal{H} such that clos Dom $L_0 = \mathcal{H}$, $L_0 \subset L_0^*$, $(L_0 y, y) \ge \varkappa ||y||^2$ ($\varkappa = \text{const}$). In what follows, without loss of generality, we deal with $\varkappa > 0$.

Also, we assume that L_0 has nonzero defect indexes: $n_+ = n_- = \dim \operatorname{Ker} L_0^* \geq 1$. Note that such an operator is necessarily unbounded.

No more assumptions on L_0 are imposed. It is a class of operators, for which we plan to construct the above-mentioned functional model.

0.3 Green system

Let L be the Friedrichs extension of L_0 , so that $L_0 \subset L \subset L_0^*$, $L^* = L$, and $(L_0y, y) \ge \varkappa ||y||^2$ holds (see, e.g., [7]). The inverse L^{-1} is bounded and defined on \mathcal{H} .

The well-known decomposition by Vishik [13] is

$$\operatorname{Dom} L_0^* = \operatorname{Dom} L_0 + L^{-1} \operatorname{Ker} L_0^* + \operatorname{Ker} L_0^*$$

(direct sums). By this, for a $y \in \text{Dom } L_0^*$ one has

$$y = y_0 + L^{-1}g + h$$

with $y_0 \in \text{Dom } L_0$ and $g, h \in \text{Ker } L_0^*$. Denote $h = \Gamma_1 y$ and $g = \Gamma_2 y$. We derive the *Green formula*

$$(L_0^*u, v)_{\mathcal{H}} - (u, L_0^*v)_{\mathcal{H}} = (\Gamma_1 u, \Gamma_2 v)_{\operatorname{Ker} L_0^*} - (\Gamma_2 u, \Gamma_1 v)_{\operatorname{Ker} L_0^*},$$

which is in fact a partial case of more general relation established in [13]. Hence, a collection $\{\mathcal{H}, \text{Ker } L_0^*; L_0, \Gamma_1, \Gamma_2\}$ constitutes a *Green system* determined by the operator L_0 .

0.4 DSBC

The Green system, in turn, determines a dynamical system with boundary control

$$\begin{split} u_{tt} + L_0^* u &= 0 & \text{in } \mathcal{H}, \ t > 0 \\ u|_{t=0} &= u_t|_{t=0} &= 0 & \text{in } \mathcal{H} \\ \Gamma_1 u &= f & \text{in } \operatorname{Ker} L_0^*, \ t \ge 0, \end{split}$$

where f = f(t) is a Ker L_0^* -valued function of time (boundary control). By $u = u^f(t)$ we denote the (generalized) solution, which is well defined for a class \mathcal{M} of smooth enough controls f.

A set

$$\mathcal{U}^t := \{ u^f(t) \mid f \in \mathcal{M} \}$$

is called *reachable* (at the moment t), whereas

$$\mathcal{U} := \bigvee_{t \geqslant 0} \mathcal{U}^t$$

(algebraic sum) is a *total reachable set*. The DSBC is said to be *controllable* if

$$\operatorname{clos} \mathcal{U} = \mathcal{H}$$

We prove that this relation holds if and only if L_0 is completely non-selfadjoint operator. The latter means that there is no nonzero subspace in \mathcal{H} , in which L_0 has a self-adjoint part.

0.5 Wave spectrum

This notion is introduced in a few steps.

First, we define a so-called *inflation* I_{L_0} , which is an operation on the lattice $\mathfrak{L}(\mathcal{H})$ of subspaces in \mathcal{H} . Inflation extends subspaces and is determined by the operator L_0 (more precisely, by its Friedrichs extension L).

Second, we define \mathfrak{L}_{L_0} as the *minimal sublattice* of $\mathfrak{L}(\mathcal{H})$, which contains all reachable subspaces $\operatorname{clos} \mathcal{U}^t$, $t \ge 0$ and is invariant with respect to the inflation I_{L_0} .

Next, we introduce a family $I_{L_0} \mathfrak{L}_{L_0}$ of monotone (growing) \mathfrak{L}_{L_0} -valued functions of $t \ge 0$ provided with standard lattice topology. This family is a partially ordered set. As such, it may content a set At $I_{L_0} \mathfrak{L}_{L_0}$ of minimal nonzero elements (*atoms*).

At last, At $I_{L_0} \mathfrak{L}_{L_0}$ is endowed with a relevant (*ball*-) topology β , and we arrive at a topological space (Ω_{L_0}, β) . It is the space, which we call a *wave spectrum* of the operator L_0 .

0.6 Reconstruction of manifolds

As was noted in 0.1, our program is motivated by inverse problems. As is shown in [5], to recover a Riemannian manifold Ω from the boundary inverse data one can

- determine a unitary copy \hat{L}_0 of the minimal Laplacian $L_0 = -\Delta$ in Ω from the data
- find the wave spectrum $\Omega_{\tilde{L}_0}$.

For a generic class of manifolds, the space $\Omega_{\tilde{L}_0}$ turns out to be isometric to Ω . By this, the wave spectrum provides a representative of the class of manifolds, which possess the given inverse data. Thus, $\Omega_{\tilde{L}_0}$ solves the reconstruction problem.

1 DSBC

1.1 Operator L_0

Let us specify the class of operators, which we deal with.

Let \mathcal{H} be a (separable) Hilbert space, L_0 an operator in \mathcal{H} . We assume that

- 1. L_0 is closed and densely defined: clos Dom $L_0 = \mathcal{H}$
- 2. L_0 is positive definite: there is a constant $\varkappa > 0$ such that $(L_0y, y) \ge \varkappa ||y||^2$ holds as $y \in \text{Dom } L_0$
- 3. L_0 has the nonzero defect indexes $n_+ = n_- = \dim \operatorname{Ker} L_0^*$: $1 \leq \dim \operatorname{Ker} L_0^* \leq \infty$.

Note that by 3. such an operator is necessarily unbounded.

By L we denote the Friedrichs extension of L, so that $L_0 \subset L \subset L_0^*$, $L^* = L$, and $(Ly, y) \ge \varkappa ||y||^2$ hold for all $y \in \text{Dom } L$ (see, e.g., [7]). Its inverse L^{-1} is a bounded operator defined on \mathcal{H} .

1.2 Green system

We begin with basic definitions, which go back to the pioneer paper by A.N.Kochubei [10] (see also [11]).

Let \mathcal{H} and \mathcal{B} be the Hilbert spaces, $A : \mathcal{H} \to \mathcal{H}$ and $\Gamma_i : \mathcal{H} \to \mathcal{B}$ (i = 1, 2)the operators such that clos Dom $A = \mathcal{H}$, Dom $\Gamma_i \supset$ Dom A, clos $\bigvee_{i=1,2}$ Ran $\Gamma_i = \mathcal{B}$.

A collection $\mathfrak{Gr} = \{\mathcal{H}, \mathcal{B}; A, \Gamma_1, \Gamma_2\}$ is said to be a *Green system*, if its elements are related via the *Green formula*

$$(Au, v)_{\mathcal{H}} - (u, Av)_{\mathcal{H}} = (\Gamma_1 u, \Gamma_2 v)_{\mathcal{B}} - (\Gamma_2 u, \Gamma_1 v)_{\mathcal{B}}$$
(1.1)

for all $u, v \in \text{Dom } A$. The space \mathcal{H} is called *inner*, \mathcal{B} is a space of boundary values, A is a basic operator, $\Gamma_{0,1}$ are the boundary operators.

In a Green system with the given $\mathcal{H}, \mathcal{B}, \mathcal{A}$, there is a freedom of choice of the boundary operators. For instance, taking an $S = S^*$, $\text{Dom } S \supset \text{Ran } \Gamma_1$ and putting $\widetilde{\Gamma}_2 := \Gamma_2 + S\Gamma_1$, one gets a collection $\widetilde{\mathfrak{Sr}} = \{\mathcal{H}, \mathcal{B}; \mathcal{A}, \Gamma_1, \widetilde{\Gamma}_2\}$, which is also a Green system.

1.3 System \mathfrak{Gr}_{L_0}

Here we associate with L_0 a Green system with the canonically chosen boundary operators. Denote

$$\mathcal{K} := \operatorname{Ker} L_0^*$$

and recall that $\dim \mathcal{K} \ge 1$. Let *P* be the (orthogonal) projection in \mathcal{H} onto \mathcal{K} , \mathbb{O} and \mathbb{I} the zero and unit operators. Also, introduce the operators

$$\Gamma_1 := L^{-1}L_0^* - \mathbb{I}, \quad \Gamma_2 := PL_0^*.$$

Lemma 1 The collection $\mathfrak{Gr}_{L_0} := \{\mathcal{H}, \mathcal{K}; L_0^*, \Gamma_1, \Gamma_2\}$ is a Green system.

Proof

1. Since $L_0^*\Gamma_1 = L_0^*L^{-1}L_0^* - L_0^* = LL^{-1}L_0^* - L_0^* = \mathbb{O}$, we have $\operatorname{Ran}\Gamma_i \subset \mathcal{K}$. The density of $\operatorname{Ran}L_0^* \supset \operatorname{Ran}L$ in \mathcal{H} implies $\operatorname{clos}\operatorname{Ran}\Gamma_2 = \mathcal{K}$. Thus, $\operatorname{clos} \vee_{i=1,2}\operatorname{Ran}\Gamma_i = \mathcal{K}$ does hold.

2. Recall the Vishik decomposition [13]

$$\operatorname{Dom} L_0^* = \operatorname{Dom} L_0 + L^{-1} \mathcal{K} + \mathcal{K}$$
(1.2)

(direct sums). By this, for a $u, v \in \text{Dom } L_0^*$ one can represent

$$u = u_0 + L^{-1}g_u + h_u, \quad v = v_0 + L^{-1}g_v + h_v$$
 (1.3)

with $u_0, v_0 \in \text{Dom } L_0$ and $g_u, h_u, g_v, h_v \in \mathcal{K}$. Therefore, with regard to $L_0 \subset L_0^*$ and $L^{-1}L_0 = \mathbb{I}$ one has

$$\begin{split} (L_0^*u, v) &- (u, L_0^*v) = \\ (L_0u_0 + g_u, v_0 + L^{-1}g_v + h_v) - (u_0 + L^{-1}g_u + h_u, L_0v + g_v) = \\ (L_0u_0, v_0) &+ (L_0u_0, L^{-1}g_v) + (L_0u_0, h_v) + (g_u, v_0) + (g_u, L^{-1}g_v) + (g_u, h_v) \\ - (u_0, L_0v_0) - (u_0, g_v) - (L^{-1}g_u, L_0v_0) - (L^{-1}g_u, g_v) - (h_u, L_0v_0) - (h_u, g_v) \,. \end{split}$$

Numbering by [...] the terms in the r.h.s. of the latter equality from the beginning, we have

$$[1] + [7] = 0, \quad [3] = (u_0, L_0^* h_v) = 0, \quad [11] = (L_0^* h_u, v_0) = 0, \\ [2] = (L^{-1} L_0 u_0, g_v) = (u_0, g_v), \quad [9] = (g_u, L^{-1} L_0 v_0) = (g_u, v_0),$$

and arrive at

$$(L_0^*u, v) - (u, L_0^*v) = (u_0, g_v) + (g_u, v_0) + (g_u, L^{-1}g_v) + (g_u, h_v) -(u_0, g_v) - (g_u, v_0) - (L^{-1}g_u, g_v) - (h_u, g_v) = (g_u, h_v) - (h_u, g_v).$$
(1.4)

3. Take a $y \in \text{Dom } L_0^*$ and represent by (1.3)

$$y = y_0 + L^{-1}g_y + h_y. (1.5)$$

Let us check that

$$y_0 = L^{-1} P_\perp L_0^* y, \quad g_y = P L_0^* y, \quad h_y = y - L^{-1} L_0^* y, \quad (1.6)$$

where $P_{\perp} = \mathbb{I} - P$ is the projection onto $\mathcal{H} \ominus \mathcal{K}$.

Indeed, since L_0^{-1} is bounded on Ran L_0 , we have Ran $L_0 = \operatorname{clos} \operatorname{Ran} L_0$. Hence, $\mathcal{H} = \operatorname{Ran} L_0 \oplus \mathcal{K}$. Therefore, $P_{\perp} L_0^* y \in \operatorname{Ran} L_0$ and $y = L^{-1} P_{\perp} L_0^* y = L_0^{-1} P_{\perp} L_0^* y$, so that $y_0 \in \operatorname{Dom} L_0$. The inclusion $g_y = P L_0^* y \in \mathcal{K}$ is evident. The relations $L_0^* h_y = L_0^* y - L_0^* L^{-1} L_0^* y = L_0^* y - \mathbb{I} L_0^* y = 0$ show that $h_y \in \mathcal{K}$. Thus, the summands in the r.h.s. of (1.5) belong to $\operatorname{Dom} L_0$, $L^{-1} \mathcal{K}$, and \mathcal{K} respectively.

In the mean time, we have

$$y = y_0 + L^{-1}g_y + h_y = \langle \sec(1.6) \rangle = L^{-1}P_{\perp}L_0^*y + L^{-1}PL_0^*y + y - L^{-1}L_0^*y = L^{-1}L_0^*y + y - L^{-1}L_0^*y = y,$$

so that (1.5) is valid.

4. Return to (1.4). By (1.5) and (1.6), we have

$$(L_0^*u, v) - (u, L_0^*v) = \left(\left[L^{-1}L_0^* - \mathbb{I} \right] u, PL_0^*v \right) - \left(PL_0^*u, \left[L^{-1}L_0^* - \mathbb{I} \right] v \right) = (\Gamma_1 u, \Gamma_2 v)_{\mathcal{K}} - (\Gamma_2 u, \Gamma_1 v)_{\mathcal{K}}$$

that proves the lemma. \blacksquare

Thus, the operator L_0 determines the Green system \mathfrak{Gr}_{L_0} in a canonical way.

1.4 System α_{L_0}

In its turn, \mathfrak{Gr}_{L_0} determines an evolutionary dynamical system of the form

$$u_{tt} + L_0^* u = 0$$
 in $\mathcal{H}, t > 0$ (1.7)

$$u|_{t=0} = u_t|_{t=0} = 0 \qquad \text{in } \mathcal{H} \tag{1.8}$$

$$\Gamma_1 u = h \qquad \qquad \text{in } \mathcal{K}, \ t \ge 0, \tag{1.9}$$

where h = h(t) is a \mathcal{K} -valued function of time, $u = u^{h}(t)$ is a solution.

In control theory, problem (1.7)–(1.9) is referred to as a dynamical system with boundary control (DSBC), h is a boundary control, $u^{h}(\cdot)$ is a trajectory, $u^{h}(t)$ is a state at the moment t. As is clear, the system (1.7)–(1.9) is determined by the operator L_0 and we denote it by α_{L_0} .

Recall that L is the Friedrichs extension of L_0 . Let $L^{\frac{1}{2}}$ be the positive square root of L. Denote $(\cdot)' := \frac{d}{dt}$. For a control $h \in C^2_{\text{loc}}([0,\infty);\mathcal{K})$ provided h(0) = h'(0) = 0, an \mathcal{H} -valued

For a control $h \in C^2_{\text{loc}}([0,\infty);\mathcal{K})$ provided h(0) = h'(0) = 0, an \mathcal{H} -valued function

$$u^{h}(t) := -h(t) + \int_{0}^{t} L^{-\frac{1}{2}} \sin\left[(t-s)L^{\frac{1}{2}}\right] h''(s) \, ds \,, \qquad t \ge 0 \qquad (1.10)$$

is said to be a *weak solution* to (1.7)–(1.9). This definition is motivated by the following fact. Introduce a class of smooth controls

$$\mathcal{M} := \{ h \in C^3_{\text{loc}}([0,\infty);\mathcal{K}) \mid h(0) = h'(0) = h''(0) = 0 \}$$

Lemma 2 If $h \in \mathcal{M}$ then u^h is a classical solution to (1.7) - (1.9).

Proof

1. Assuming $h \in \mathcal{M}$, let us derive a relevant representation for the weak solution.

Take a $y \in \mathcal{H}$. Representing $L = \int_0^\infty \lambda \, dE_\lambda$ via the spectral measure E_λ of L and integrating by parts, one has

$$\frac{d}{ds} \int_0^\infty \frac{\cos\sqrt{\lambda}(t-s)}{\lambda} d\left(E_{\lambda}h''(s), y\right) = \frac{d}{ds} \left(h''(s), L^{-1}\cos\left[(t-s)L^{\frac{1}{2}}\right]y\right) = \left(h'''(s), L^{-1}\cos\left[(t-s)L^{\frac{1}{2}}\right]y\right) + \left(h''(s), L^{-\frac{1}{2}}\sin\left[(t-s)L^{\frac{1}{2}}\right]y\right) = \left(L^{-1}\cos\left[(t-s)L^{\frac{1}{2}}\right]h'''(s), y\right) + \left(L^{-\frac{1}{2}}\sin\left[(t-s)L^{\frac{1}{2}}\right]h''(s), y\right).$$

Applying $\int_0^t (\dots) ds$, we get

$$\int_0^\infty \frac{1}{\lambda} d\left(E_\lambda h'''(t), y\right) = \left(L^{-1} h'''(t), y\right) = \left(\int_0^t L^{-1} \cos\left[(t-s)L^{\frac{1}{2}}\right] h'''(s), y\right) + \left(\int_0^t L^{-\frac{1}{2}} \sin\left[(t-s)L^{\frac{1}{2}}\right] h''(s), y\right).$$

By arbitrariness of y, we get

$$\int_0^t L^{-\frac{1}{2}} \sin\left[(t-s)L^{\frac{1}{2}}\right] h''(s) \, ds = L^{-1} \int_0^t \left\{ \mathbb{I} - \cos\left[(t-s)L^{\frac{1}{2}}\right] \right\} h'''(s) \, ds.$$

Therefore, for $h \in \mathcal{M}$ one can write (1.10) in the form

$$u^{h}(t) = -h(t) + L^{-1} \int_{0}^{t} \left\{ \mathbb{I} - \cos\left[(t-s)L^{\frac{1}{2}}\right] \right\} h'''(s) \, ds \tag{1.11}$$

with $h \in \mathcal{K}$ and $L^{-1} \int_0^t (\dots) \in \text{Dom } \subset \text{Dom } L_0^*$. Hence, we have $u^h(t) \in \text{Dom } L_0^*$ for all $t \ge 0$.

2. Show that (1.11) provides a classical solution to (1.7)–(1.9). Differentiation in (1.11) implies

$$u_t^h(t) = -h'(t) + L^{-\frac{1}{2}} \int_0^t \sin\left[(t-s)L^{\frac{1}{2}}\right] h'''(s) \, ds, \qquad (1.12)$$

$$u_{tt}^{h}(t) = -h''(t) + \int_{0}^{t} \cos\left[(t-s)L^{\frac{1}{2}}\right]h'''(s)\,ds.$$
(1.13)

Therefore,

$$u_{tt}^{h}(t) + L_{0}^{*}u^{h}(t) = -h''(t) + \int_{0}^{t} \cos\left[(t-s)L^{\frac{1}{2}}\right]h'''(s)\,ds + \int_{0}^{t} \left\{\mathbb{I} - \cos\left[(t-s)L^{\frac{1}{2}}\right]\right\}h'''(s)\,ds = -h''(t) + \int_{0}^{t}h'''(s)\,ds = 0,$$

so that (1.7) does hold.

As is seen from (1.11),(1.12), $u^h(0) = u_t^h(0) = 0$, i.e., the initial conditions (1.8) are fulfilled.

Applying Γ_1 in (1.11), we have

$$\Gamma_{1}u^{h}(t) = \left(L^{-1}L_{0}^{*} - \mathbb{I}\right)\left(-h(t) + L^{-1}\int_{0}^{t} \left\{\mathbb{I} - \cos\left[(t-s)L^{\frac{1}{2}}\right]\right\}h'''(s)\,ds\right) = L^{-1}\int_{0}^{t} \left\{\mathbb{I} - \cos\left[(t-s)L^{\frac{1}{2}}\right]\right\}h'''(s)\,ds + h(t) - L^{-1}\int_{0}^{t} \left\{\mathbb{I} - \cos\left[(t-s)L^{\frac{1}{2}}\right]\right\}h'''(s)\,ds = h(t).$$

Hence, the 'boundary condition' (1.9) is fulfilled.

Note in addition, that one can prove a uniqueness of the classical u^h .

1.5 Controllability

A set of all possible states of the system α_{L_0}

 $\mathcal{U}^t := \{ u^h(t) \mid u^h \text{ is a week solution to } (1.7) - (1.9) \}$

is said to be *reachable* (at the moment $t \ge 0$). Representing (1.10) in the convolution form

$$u^{h}(t) = \int_{0}^{t} \left\{ -(t-s)\mathbb{I} + L^{-\frac{1}{2}} \sin\left[(t-s)L^{\frac{1}{2}}\right] \right\} h''(s) \, ds \,, \tag{1.14}$$

one can easily see that \mathcal{U}^t extends as t grows.

Also, define a *total* reachable set

$$\mathcal{U} := \bigvee_{t \geqslant 0} \mathcal{U}^t$$

and a *defect subspace*

$$\mathcal{D} := \mathcal{H} \ominus \operatorname{clos} \mathcal{U}$$
.

The system α_{L_0} is said to be *controllable*, if the relation

$$\operatorname{clos} \mathcal{U} = \mathcal{H} \tag{1.15}$$

is valid or, equivalently, if $\mathcal{D} = \{0\}$.

Below we establish the necessary and sufficient conditions on the operator L_0 , which provide controllability of the system α_{L_0} . Taking into account the

well-known similar results and general principles of system theory [8], these conditions are quite expectable: L_0 has to be a *completely non-self-adjoint* (c.n.s.a.) operator.

Recall the definitions. We say that a symmetric operator A has a selfadjoint part in a (nonzero) subspace $\mathcal{N} \subset \mathcal{H}$ if

- the lineal set $\mathcal{N} \cap \text{Dom} A$ is dense in \mathcal{N}
- the embedding $A[\mathcal{N} \cap \text{Dom} A] \subset \mathcal{N}$ holds
- the operator $A|_{\mathcal{N}\cap \text{Dom }A}$ is self-adjoint in \mathcal{N} .

A symmetric operator A is said to be c.n.s.a. if it has a self-adjoint part in no nonzero subspace in \mathcal{H} .

Theorem 1 The system α_{L_0} is controllable if and only if L_0 is a c.n.s.a. operator.

Proof

Necessity Let L_0 have a self-adjoint part in $\mathcal{N} \subset \mathcal{H}$. Fix an $h \in \mathcal{K}$. Take a $g \in \mathcal{N}$ and represent $g = L_0 \tilde{g}$ with $\tilde{g} \in \mathcal{N} \cap \text{Dom } L_0$. The latter is possible because $L_0|_{\mathcal{N}}$ is a positive definite boundedly invertible operator in \mathcal{N} . In view of

$$(g,h) = (L_0 \widetilde{g}, h) = (\widetilde{g}, L_0^*) = 0$$

,
$$\mathcal{K} \subset \mathcal{N}^\perp = \mathcal{H} \ominus \mathcal{N}$$
(1.16)

holds.

we have $\mathcal{N} \perp \mathcal{K}$, i.e.

Recall that L is the Friedrichs extension: $L_0 \subset L \subset L_0^*$. For a $g \in \mathcal{N}$ we have

$$L^{-1}g = L^{-1}L_0\widetilde{g} = \widetilde{g} \in \mathcal{N}$$

that implies $L^{-1}\mathcal{N} \subset \mathcal{N}$. Since L^{-1} is self-adjoint, we have $L^{-1}\mathcal{N}^{\perp} \subset \mathcal{N}^{\perp}$, i.e., L^{-1} is reduced by \mathcal{N} . The latter leads to

$$L^{-\frac{1}{2}}\mathcal{N}\subset\mathcal{N},\qquad L^{-\frac{1}{2}}\mathcal{N}^{\perp}\subset\mathcal{N}^{\perp}.$$
 (1.17)

By (1.16), in the r.h.s. of (1.10), one has $h(t), h''(s) \in \mathcal{K} \subset \mathcal{N}^{\perp}$. By (1.17), the integral in (1.10) belongs to \mathcal{N}^{\perp} . As a result, $u^{h}(t) \in \mathcal{N}^{\perp}$ holds for all $t \ge 0$, i.e., trajectories u^{h} of the system $\alpha_{L_{0}}$ do not leave the subspace

 \mathcal{N}^{\perp} . Therefore, $\mathcal{U} \subset \mathcal{N}^{\perp}$ that leads to $\mathcal{D} \neq \{0\}$. So, the system α_{L_0} is not controllable.

Thus, if α_{L_0} is controllable then L_0 is c.n.s.a.

Sufficiency Assume that α_{L_0} is not controllable, i.e., $\mathcal{D} \neq \{0\}$. It will be shown that L_0 has to have a self-adjoint part in \mathcal{D} and, hence, is not a c.n.s.a. operator.

1. Take a nonzero $y \in \mathcal{D}$. For any (admissible) $h \in C^2_{\text{loc}}([0,\infty);\mathcal{K})$ and t > 0, we have

$$0 = (y, u^{h}(t)) = \langle \sec(1.10), (1.14) \rangle = \left(y, \int_{0}^{t} \left\{ -(t-s)h''(s) + L^{-\frac{1}{2}} \sin\left[(t-s)L^{\frac{1}{2}} \right] \right\} h''(s) \, ds \right) = \int_{0}^{t} (-(t-s)y + w^{y}(t-s), h''(s)) \, ds,$$
(1.18)

where

$$w^{y}(\eta) := L^{-\frac{1}{2}} \sin\left[\eta L^{\frac{1}{2}}\right] y, \qquad \eta \ge 0.$$

Fix a $k \in \mathcal{K}$. Choose a sequence of controls $h_j(s) = \varphi_j(s)k$, where $\varphi_j \in C_0^{\infty}(0,t)$ are such that $\varphi_j''(s) \to \delta(s)$ (the Dirac delta-function) as $j \to \infty$. For such $h_j(\cdot)$, the limit passage in (1.18) implies

$$0 = (-ty + w^y(t), k).$$

By arbitrariness of k, we conclude that

$$-ty + w^{y}(t) \in \mathcal{K}^{\perp}, \qquad t \ge 0.$$
(1.19)

Converting these considerations, we easily obtain the following result.

Proposition 1 The embedding $y \in \mathcal{D}$ holds if and only if (1.19) is valid. Denote $-ty + w^y(t) =: p(t) \in \mathcal{K}^{\perp}$ and represent

$$y = \frac{1}{t} L^{-\frac{1}{2}} \sin\left[tL^{\frac{1}{2}}\right] - \frac{p(t)}{t}$$

The operator $L^{-\frac{1}{2}} \sin\left[tL^{\frac{1}{2}}\right]$ is bounded. By the latter, tending $t \to \infty$ we get $y = -\lim t^{-1}p(t) \in \mathcal{K}^{\perp}$. Returning to (1.19), we conclude that

$$y, w^y(t) \in \mathcal{K}^\perp, \qquad t \ge 0.$$
 (1.20)

2. Consider an auxiliary dynamical system

$$w_{tt} + Lw = 0 \qquad \qquad \text{in } \mathcal{H}, \quad t > 0 \qquad (1.21)$$

$$w|_{t=0} = 0, \ w_t|_{t=0} = y$$
 in \mathcal{H} (1.22)

with $y \in \mathcal{D}$ chosen above. Its solution is of the well-known form

$$w^{y}(t) = L^{-\frac{1}{2}} \sin\left[tL^{\frac{1}{2}}\right] y$$

(see, e.g., [7]).

Denote $J := \int_0^t (\cdot) ds$. Applying J^2 in (1.21) with regard to (1.22), we have

$$w^{y}(t) - ty = -L\left(J^{2}w^{y}\right)(t).$$

This implies

$$L^{-1}(-ty + w^{y}(t)) = -(J^{2}w^{y})(t) \in \mathcal{K}^{\perp}$$
(1.23)

in view of (1.20). In the mean time, we have

$$L^{-1}w^{y}(t) = L^{-1}L^{-\frac{1}{2}}\sin\left[tL^{\frac{1}{2}}\right]y = L^{-\frac{1}{2}}\sin\left[tL^{\frac{1}{2}}\right]\left(L^{-1}y\right) = w^{L^{-1}y}(t).$$

Hence, (1.23) implies

$$-tL^{-1}y + w^{L^{-1}y}(t) \in \mathcal{K}^{\perp}, \qquad t \ge 0.$$

In accordance with Proposition 1, the latter is equivalent to $L^{-1}y \in \mathcal{D}$.

Thus, beginning with $y \in \mathcal{D}$, we arrive at $L^{-1}y \in \mathcal{D}$, i.e., the defect space reduces the operator L^{-1} :

$$L^{-1}\mathcal{D} \subset \mathcal{D}$$
 .

3. The part $L^{-1}|_{\mathcal{D}}$ is a self-adjoint invertible operator in \mathcal{D} . Therefore, $L^{-1}\mathcal{D}$ is dense in \mathcal{D} . Hence, the operator L has the part $L|_{\mathcal{D}}$, which is a (densely defined) self-adjoint operator in \mathcal{D} .

Show that $L|_{\mathcal{D}} = L_0$. Indeed, by (1.20) one has

$$\mathcal{D} \subset \mathcal{K}^{\perp} = \mathcal{H} \ominus \mathcal{K} = \operatorname{clos} \operatorname{Ran} \mathcal{L}_{\prime} = \operatorname{Ran} \mathcal{L}_{\prime}.$$

Therefore, by $L_0 \subset L$ we have

$$\operatorname{Dom} L|_{\mathcal{D}} = L^{-1}\mathcal{D} \subset L^{-1}\operatorname{Ran} L_0 = L_0^{-1}\operatorname{Ran} L_0 = \operatorname{Dom} L_0,$$

and $L|_{\mathcal{D}}x = L_0x$ for all $x \in L^{-1}\mathcal{D}$.

Thus, if the system α_{L_0} is not controllable then L_0 has a self-adjoint part in \mathcal{D} , i.e., is not a c.n.s.a. operator.

2 Wave spectrum

2.1 Inflation

We use the term 'lattice' in its general meaning [6]: a *lattice* is a partially ordered set provided with the operations $x \lor y = \sup \{x, y\}, x \land y = \inf \{x, y\}$. However, we deal with the concrete lattices endowed with additional structures (complement, topology, ets).

Definition Let $\mathfrak{L}(\mathcal{H})$ be the lattice of the (closed) subspaces of \mathcal{H} with the partial order \subseteq and operations $\mathcal{A} \wedge \mathcal{B} = \mathcal{A} \cap \mathcal{B}, \ \mathcal{A} \vee \mathcal{B} = \operatorname{clos} \{a + b \mid a \in \mathcal{A}, b \in \mathcal{B}\}, \ \mathcal{A} \mapsto \mathcal{A}^{\perp}$. Also, it possesses the least and greatest elements $\{0\}$ and \mathcal{H} .

By $P_{\mathcal{A}}$ we denote the (orthogonal) projection in \mathcal{H} onto \mathcal{A} . Topology on $\mathfrak{L}(\mathcal{H})$ is determined by convergence of the projections on the corresponding subspaces. Namely, we write $\mathcal{A}_{|} \to \mathcal{A}$ if $s - \lim P_{\mathcal{A}_{i}} = P_{\mathcal{A}}$.

An *inflation* is a family of maps $I = \{I^t\}_{t \ge 0}, I^t : \mathfrak{L}(\mathcal{H}) \to \mathfrak{L}(\mathcal{H})$ with the properties

- $I^0 = \mathrm{id}, \ I^t\{0\} = \{0\}$
- $\mathcal{A} \subseteq \mathcal{B}$ and $s \leq t$ imply $I^s \mathcal{A} \subseteq I^t \mathcal{B}$.

Inflation I_{L_0} As is shown in [5], with each operator L_0 of the class under consideration one associates an inflation I_{L_0} in the following way. Fix a subspace $\mathcal{A} \in \mathfrak{L}(\mathcal{H})$ and consider a dynamical system

$$\begin{aligned} v_{tt} + Lv &= a & \text{ in } \mathcal{H}, \ t > 0 \\ v|_{t=0} &= v_t|_{t=0} & \text{ in } \mathcal{H} \end{aligned}$$

where a = a(t) is an \mathcal{A} -valued function of time, $v = v^a(t)$ is a solution. By the well-known Duhamel formula, one has

$$v^{a}(t) = \int_{0}^{t} L^{-\frac{1}{2}} \sin\left[(t-s)L^{\frac{1}{2}}\right] a(s) \, ds, \qquad t \ge 0.$$

The reachable sets of the system are

$$\mathcal{V}_{\mathcal{A}}^{t} := \left\{ v^{a}(t) \mid a \in L_{2}^{\mathrm{loc}}\left([0,\infty);\mathcal{A}\right) \right\}.$$

As is easy to see, $\mathcal{V}_{\mathcal{A}}^t$ extends as \mathcal{A} extends and/or t grows. Define the family $I_{L_0} := \{I_{L_0}^t\}_{t \ge 0}$ by

$$I_{L_0}^t \mathcal{A} := \operatorname{clos} \mathcal{V}_{\mathcal{A}}^t \quad \text{for } t > 0, \qquad I_{L_0}^0 := \operatorname{id}.$$

Proposition 2 The family I_{L_0} is an inflation.

Proof see in [5].

Actually, I_{L_0} is determined not by L_0 but its extension L. As such, an inflation is well-defined for any bounded from bellow self-adjoint operator.

2.2 Set Ω_{L_0}

Lattice \mathfrak{L}_{L_0} Recall that a *sublattice* is a subset of $\mathfrak{L}(\mathcal{H})$, which is invariant with respect to all the lattice operations. Each sublattice necessarily contains $\{0\}$ and \mathcal{H} .

We say that a (sub)lattice $\mathfrak{L} \subset \mathfrak{L}(\mathcal{H})$ is invariant w.r.t. an inflation I if $I^t \mathfrak{L} \subseteq \mathfrak{L}$ holds for $t \ge 0$.

By \mathfrak{L}_{L_0} we denote the *minimal sublattice* in \mathfrak{L} , which

- contains all reachable subspaces $\operatorname{clos} \mathcal{U}^t, t \ge 0$
- is invariant with respect to the inflation I_{L_0}
- is closed in the above-mentioned topology on $\mathfrak{L}(\mathcal{H})$.

Lattice $\overline{I_{L_0}\mathfrak{L}_{L_0}}$ Let \mathcal{F} be a set of $\mathfrak{L}(\mathcal{H})$ -valued functions of $t \ge 0$. This set is also a lattice w.r.t. the point-wise order, operations, and topology:

$$\{f \leqslant g\} \Longleftrightarrow \{f(t) \subseteq g(t), \ t \ge 0\}, \quad (f \lor g)(t) := f(t) \lor g(t),$$
$$(f \land g)(t) := f(t) \land g(t), \ (f^{\perp})(t) := (f(t))^{\perp}, \ (\lim f_j)(t) := \lim(f_j(t)).$$

The least and greatest elements of \mathcal{F} are the functions equal $\{0\}$ and \mathcal{H} identically. We denote them by $0_{\mathcal{F}}$ and $1_{\mathcal{F}}$ respectively.

An inflation I can be regarded as a map from $\mathfrak{L}(\mathcal{H})$ to \mathcal{F} acting by the rule $(I\mathcal{A})(t) := I^t\mathcal{A}, t \ge 0$. If \mathfrak{L} is invariant w.r.t. I then the image $I\mathfrak{L}$, as well as its closure $\overline{I\mathfrak{L}}$ are sublattices in \mathcal{F} . Both of them contain $0_{\mathcal{F}}$.

The operator L_0 determines the lattice $I_{L_0} \mathfrak{L}_{L_0}$.

Atoms Let \mathcal{P} be a partially ordered set with the least element 0. An element $a \in \mathcal{P}$ is said to be an *atom* if $a \neq 0$ and $b \leq a$ implies b = a [6]. By At \mathcal{P} we denote the set of atoms.

The key object of the paper is the set

$$\Omega_{L_0} := \operatorname{At} \overline{I_{L_0} \mathfrak{L}_{L_0}}$$

that we call a *wave spectrum* of the operator L_0 .

Remark Certain additional assumptions on L_0 provide $\Omega_{L_0} \neq \emptyset$ [5]. There is L_0 such that its wave spectrum consists of a single point. A conjecture is that $\Omega_{L_0} \neq \emptyset$ does hold ever.

2.3 Space (Ω_{L_0}, β)

Here the wave spectrum is endowed with relevant structures.

Topology Recall that atoms are \mathfrak{L}_{L_0} -valued functions of time. Fix an atom $a \in \Omega_{L_0}$. A set

$$B_r[a] := \{ b \in \Omega_{L_0} \mid \exists t > 0 \text{ s.t. } \{ 0 \} \neq b(t) \leqslant a(r) \}, \qquad r > 0$$

is said to be a *ball*, a and r are its center and radius.

Lemma 3 The system of balls $\{B_r[a] \mid a \in \Omega_{L_0}, r > 0\}$ is a base of topology.

Proof One has to check the characteristic properties of a base:

- 1. for any $a \in \Omega_{L_0}$, there is a ball $B \ni a$
- 2. for an atom $a \in \Omega_{L_0}$ and the balls B_1 , B_2 such that $a \in B_1 \cap B_2$, there is a ball B such that $a \in B \subset B_1 \cap B_2$

(see, e.g., [9]).

1. Take an $a = a(\cdot) \in \Omega_{L_0}$. For any $r > t_0 := \inf \{t > 0 \mid a(t) \neq \{0\}\}$ and $t \in (t_0, r]$, one has $\{0\} \neq a(t) \leq a(r)$, i.e., $a \in B_r[a]$.

2. Let $a \in B^{r_1}[a_1] \cap B^{r_2}[a_2]$, so that both of $B^{r_i}[a_i]$ are nonempty. Choose t_i such that $\{0\} \neq a(t_i) \leq a_i(r_i)$ and denote $r := \min\{t_1, t_2\}$. By the choice, one has $\{0\} \neq a(r) \leq a_i(r_i)$ that implies $B_r[a] \neq \emptyset$.

For any $b \in B_r[a]$, there is a t > 0 such that $\{0\} \neq b(t) \leq a(r) \leq a_i(r_i)$. By the latter inequality, one has $b \in B_{r_i}[a_i]$. Hence $B_r[a] \subset B^{r_1}[a_1] \cap B^{r_2}[a_2]$.

The base $\{B_r[a] \mid a \in \Omega_{L_0}, r > 0\}$ determines the (unique) topology, in which an open set is a sum of balls [9]. We call it a *ball topology* and denote by β . So, we get a topological space (Ω_{L_0}, β) .

Boundary Return to the DSBC α_{L_0} . The family of reachable subspaces

$$\mathfrak{u}_{L_0} := \left\{ \operatorname{clos} \mathcal{U}^{\sqcup} \right\}_{t \ge 0}$$

can be regarded as an \mathfrak{L}_{L_0} -valued function of time. As such, \mathfrak{u}_{L_0} is an element of the lattice $\overline{I_{L_0}\mathfrak{L}_{L_0}} \subset \mathcal{F}$ and can be compared with its atoms. Thus, the set

$$\partial\Omega_{L_0} := \{ a \in \Omega_{L_0} \mid a \leq \mathfrak{u}_{L_0} \}$$

is well defined and said to be a *boundary* of the wave spectrum.

3 Illustration

3.1 Manifold

Let Ω be a C^{∞} -smooth compact Riemannian manifold of dimension $n \ge 2$ with the boundary $\partial \Omega$, g the metric tensor, $-\Delta$ the scalar Beltrami-Laplace operator. Recall that in local coordinates one has

$$-\Delta = -[\det g]^{-\frac{1}{2}} \frac{\partial}{\partial x_i} [\det g]^{\frac{1}{2}} g^{ij} \frac{\partial}{\partial x_j} \,.$$

By ν we denote an outward normal to $\partial\Omega$; ∂_{ν} is differentiation w.r.t. the normal.

The manifold is endowed with volume form dv, so that the (real) Hilbert space $\mathcal{H} := L_2(\Omega)$ with the inner product

$$(u,v) = \int_{\Omega} u \, v \, dv$$

is well defined.

The boundary $\partial\Omega$ is endowed with canonical (induced by the tensor $g|_{\partial\Omega}$) metric and volume (surface) element $d\sigma$. In the space $\mathcal{B} := L_2(\partial\Omega)$, the inner product is

$$(f,g)_{\mathcal{B}} = \int_{\partial\Omega} fg \, d\sigma$$

3.2 Operators

Our basic operator is the minimal Laplacian $L_0 : \mathcal{H} \to \mathcal{H}$, $\text{Dom } L_0 = \{y \in H^2(\Omega) \mid y = \partial_{\nu} y = 0 \text{ on } \partial\Omega\}$, $L_0 = -\Delta y$, where $H^2(\Omega)$ is the Sobolev class. Operator L_0 is positive definite and symmetric, its defect indexes are $n_{\pm} = \infty$.

The operator L_0^* is the maximal Laplacian, which is defined on Dom $L_0^* = \{y \in \mathcal{H} \mid \Delta y \in \mathcal{H}\}$ (here Δy is understood in the sense of distributions) and acts by $L_0^* y = -\Delta y$. Its null subspace consists of harmonic functions:

$$\mathcal{K} = \operatorname{Ker} L_0^* = \{ y \in \mathcal{H} \mid \Delta y = 0 \text{ in } \Omega \setminus \partial \Omega \}.$$

Also, we use the notation $L_0^* = -\Delta_{\max}$.

The Friedrichs extension $L \supset L_0$ is defined on Dom $L = \{y \in H^2(\Omega) \mid y = 0 \text{ on } \partial\Omega\}$ and acts by $Ly = -\Delta y$.

One more operator associated with the manifold is the harmonic continuation $\Pi : \mathcal{B} \to \mathcal{H}$ defined by the relations $\Delta \Pi f = 0$ in Ω , and $(\Pi f)|_{\partial\Omega} = f$. As is well-known in elliptic PDE theory, Π is a compact injective operator; its adjoint $\Pi^* : \mathcal{H} \to \mathcal{B}$ is

$$\Pi^* = \partial_{\nu} L^{-1}.$$

3.3 Green system

As one can check, for a $y \in \text{Dom } L_0^*$, the decomposition (1.3) is

$$y = y_0 + L^{-1}g_y + h_y$$

with

$$h_{y} = \Pi\left(y|_{\partial\Omega}\right), \quad g_{y} = \left(\Pi^{*}\right)^{-1} \left[\partial_{\nu}y - \Lambda\left(y|_{\partial\Omega}\right)\right], \quad (3.1)$$

where $\Lambda: \mathcal{B} \to \mathcal{B}$ is the *Dirichlet-to-Neumann map* defined by

$$\Lambda f := \partial_{\nu} \Pi f \qquad \text{on} \quad \partial \Omega.$$

Note that the right hand sides in (3.1) have to be understood properly; they are well defined for smooth enough y's and then extended on all $y \in \text{Dom } L_0^*$ by relevant continuity [13], [11].

By (3.1), we have

$$(L_0^* u, v)_{\mathcal{H}} - (u, L_0^* v)_{\mathcal{H}} = \langle \sec(1.4) \rangle = (g_u, h_v)_{\mathcal{H}} - (h_u, g_v)_{\mathcal{H}} = ((\Pi^*)^{-1} [\partial_\nu u - \Lambda (u|_{\partial\Omega})], \Pi (v|_{\partial\Omega}))_{\mathcal{H}} - (\Pi (u|_{\partial\Omega}), (\Pi^*)^{-1} [\partial_\nu v - \Lambda (v|_{\partial\Omega})])_{\mathcal{H}} = (\gamma_1 u, \gamma_2 v)_{\mathcal{B}} - (\gamma_2 u, \gamma_1 v)_{\mathcal{B}},$$
(3.2)

where $\gamma_i : \mathcal{H} \to \mathcal{B}$,

$$\gamma_1 := (\cdot)|_{\partial\Omega}, \quad \gamma_2 := [\partial_{\nu} - \Lambda] [(\cdot)|_{\partial\Omega}]$$

are the canonical (by Vishik) boundary operators: see [13], sec 6.

As a result, the canonical Green system associated with the manifold is

$$\mathfrak{Gr}_{\Omega} = \{ L_2(\Omega), L_2(\partial\Omega); -\Delta_{\max}, \gamma_1, \gamma_2 \}.$$
(3.3)

3.4 System α_{Ω}

In accordance with (3.2), (3.3), the system $\alpha_{L_0} =: \alpha_{\Omega}$ on the manifold takes the form

$$\begin{aligned} u_{tt} - \Delta u &= 0 & \text{in } (\Omega \setminus \partial \Omega) \times (0, \infty) \\ u|_{t=0} &= u_t|_{t=0} &= 0 & \text{in } \Omega \\ u &= f & \text{in } \partial \Omega \times (0, \infty), \end{aligned}$$

where $f \in L_2^{\text{loc}}(\partial \Omega \times (0, \infty))$ is a *boundary control*, $u = u^f(x, t)$ is a solution. The solution describes a *wave*, which is initiated by the boundary source and propagates into the manifold. The speed of propagation is finite (equal 1).

The system α_{Ω} is controllable [5]. Moreover, for a compact Ω one has

$$\operatorname{clos} \mathcal{U}^t = \mathcal{H}, \qquad t > \min_{x \in \Omega} \operatorname{dist} (x, \partial \Omega)$$

(see [3]).

3.5 Wave spectrum

In [5] a class of the so-called *simple manifolds* is introduced. Roughly speaking, simplicity means that the group of symmetries (isometries) of Ω is trivial.

This property is generic: any smooth compact manifold can be made simple by arbitrarily small variations of its boundary.

As is shown in [5], if Ω is simple then there is a canonical bijection

 $\Omega_{-\Delta_{\min}} \ni a_x \leftrightarrow x_a \in \Omega$

between atoms and points, which relates the balls and boundaries:

$$B_r[a_x] \leftrightarrow \{ x' \in \Omega \mid \operatorname{dist}(x', x_a) < r \}, \quad \partial \Omega_{-\Delta_{\min}} \leftrightarrow \partial \Omega.$$

Thus, a simple manifold is identical (isometric) to its wave spectrum. It is the fact, which is used in inverse problems for *reconstruction*.

Namely, each kind of traditional inverse data (response operator [3], Weyl function [11], characteristic function [12]) determines the operator $L_0 = -\Delta_{\min}$ up to unitary equivalence. By this, given the inverse data of a *simple* manifold, one can determine a unitary copy \tilde{L}_0 , find its wave spectrum $\Omega_{\tilde{L}_0}$ and thus recover the manifold up to isometry.

Note that a reconstruction up to isometry is the most that we can hope for. Assume that we are given with the boundary inverse data of a certain manifold Ω . Assume that another Ω' is isometric to Ω and has the same boundary: $\partial \Omega' = \partial \Omega$. As is easy to recognize, the boundary data of Ω' and Ω are identical. Therefore, in principle, these data do not determine the original Ω uniquely. In such a situation, the only relevant understanding of 'to recover' is to provide a representative of the class of manifolds with the given data. The wave spectrum $\Omega_{\tilde{L}_0}$ does provide such a representative.

References

- M.I.Belishev. Boundary control in reconstruction of manifolds and metrics (the BC method). *Inverse Problems*, 13(5): R1–R45, 1997.
- [2] M.I.Belishev. Dynamical systems with boundary control: models and characterization of inverse data. *Inverse Problems*, 17 (2001), 659–682.
- [3] M.I.Belishev. Recent progress in the boundary control method. *Inverse Problems*, 23 (2007), no 5, R1–R67.
- [4] M.I.Belishev. Geometrization of Rings as a Method for Solving Inverse Problems. Sobolev Spaces in Mathematics III. Applications in Mathematical Physics, Ed. V.Isakov., Springer, 2008, 5–24.

- [5] M.I.Belishev. A unitary invariant of semi-bounded operator in reconstruction of manifolds. http://www.arXiv:1208.3084v1 [math.FA] 15 Aug 2012
- [6] G.Birkhoff. Lattice Theory. Providence, Rhode Island, 1967.
- [7] M.S.Birman, M.Z.Solomyak. Spectral Theory of Self-Adjoint Operators in Hilbert Space. D.Reidel Publishing Comp., 1987.
- [8] R.Kalman, P.Falb, M.Arbib. Topics in Mathematical System Theory. New-York: McGraw-Hill, 1969.
- [9] J.L.Kelley. General Topology. D.Van Nostrand Company, Inc. Princeton, New Jersey, Toronto, London, New York, 1957.
- [10] A.N.Kochubei. On extension of symetric operators and symmetric binary relations. *Mat. Zametki*, 17 (1975), 41–48.
- [11] V.Ryzhov. A General Boundary Value Problem and its Weyl Function. Opuscula Math., 27 (2007), no. 2, 305–331.
- [12] A.V.Shtraus. Functional Models and Generalized Spectral Functions of Symmetric Operators. St.-Petersburg Math. Journal, 10(5): 1–76, 1998.
- [13] M.I.Vishik. On general boundary value problems for elliptic differential equations. *Proceedings of Moscow Math. Society*, 1 (1952), 187–246 (in Russian).