On almost surely periodic and almost periodic solutions of backward SPDEs

Nikolai Dokuchaev

Department of Mathematics & Statistics, Curtin University,

GPO Box U1987, Perth, 6845 Western Australia

Email N.Dokuchaev@curtin.edu.au

July 1, 2018

Abstract

We study linear backward stochastic partial differential equations of parabolic type with special boundary conditions in time. The standard Cauchy condition at the terminal time is replaced by a condition that holds almost surely and mixes the random values of the solution at different times, including the terminal time, initial time and continuously distributed times. Uniqueness, solvability and regularity results for the solutions are obtained. In particular, conditions of existence of periodic in time and "almost periodic" solutions are obtained for backward SPDEs.

AMS 1991 subject classification: Primary 60J55, 60J60, 60H10. Secondary 34F05, 34G10. Key words and phrases: backward SPDEs, periodic solutions, almost periodic solutions, Brownian bridge.

1 Introduction

Partial differential equations and stochastic partial differential equations (SPDEs) have fundamental significance for natural sciences, and various boundary value problems for them were widely studied. Usually, well-posedness of a boundary value depends on the choice of the boundary value conditions. For the deterministic parabolic equations, well-posedness requires the correct choice of the initial condition. For example, consider the heat equation $u'_t = u''_{xx}$, $t \in [0, T]$. For this equation, a boundary value problem with the Cauchy condition at initial time t = 0 is well-posed, and a boundary value problem with the Cauchy condition at terminal time t = T is ill-posed. It is known also that the problems for deterministic parabolic equation are well-posed for periodic type condition u(x,0) = u(x,T); see, e.g., Dokuchaev (1994, 1995), Fife (1964), Hess (1991), Lieberman (1999), Nakao (1984), Shelukhin (1993), Vejvoda (1982). Less is known for parabolic equation with more general non-local in time conditions and for SPDEs.

Boundary value problems for SPDEs are well studied in the existing literature for the case of forward parabolic Ito equations with the Cauchy condition at initial time (see, e.g., Alós et al (1999), Bally *et al* (1994), Da Prato and Tubaro (1996), Gyöngy (1998), Krylov (1999), Maslowski (1995), Pardoux (1993), Rozovskii (1990), Walsh (1986), Zhou (1992), and the bibliography there). Many results have been also obtained for the backward parabolic Ito equations with Cauchy condition at terminal time, as well as for pairs of forward and backward equations with separate Cauchy conditions at initial time and the terminal time respectively; see, e.g., Yong and Zhou (1999), and the author's papers (1992), (2005), (2011), (2012). Note that a backward SPDE cannot be transformed into a forward equation by a simple time change, unlike as for the case of deterministic equations. Usually, a backward SPDE is solvable in the sense that there exists a diffusion term being considered as a part of the solution that helps to ensure that the solution is adapted to the driving Brownian motions.

There are also results for SPDEs with boundary conditions that mix the solution at different times that may include initial time and terminal time. This category includes stationary type solutions for forward SPDEs (see, e.g., Dorogovtsev and Ortega (1988), Caraballo *et al* (2004), Chojnowska-Michalik (19987), Chojnowska-Michalik and Goldys (1995), Duan *et al* (2003), Mattingly (1999), Mohammed *et al* (2008), Sinai (1996), and the references here). There are also results for different types of the periodicity of the solutions of SPDEs; see, e.g., Chojnowska-Michalik (1990), Tudor (1992), Da Prato and Tudor (1995), Arnold and Tudor (1998), Klünger (2001), Bezandry and Diagana (2007), Mellah and de Fitte (2007), Feng and Zhao (2012), Bedouhene *et al* (2012), Crewe (2013). As was mentioned in Feng and Zhao (2012), it is difficult to expect that, in general, a SPDE has a periodic in time solution $u(\cdot, t)|_{t\in[0,T]}$ in a usual sense of exact equality $u(\cdot, t) = u(\cdot, t + T)$ that holds almost surely given that $u(\cdot, t)$ is adapted to some Brownian motion. However, there are important examples of stochastic processes with this property. In particular, this property holds at t = 0 for a Brownian bridge. Using this, the existence of almost surely periodic solutions was established in Rodkina (1992) for ordinary stochastic equations with the driving Brownian motion replaced by a Brownian bridge.

In a more typical setting with driving Brownian motion, the periodicity of the solutions of stochastic equations has to be interpreted differently. This periodicity was usually considered in the sense of the distributions. In Feng and Zhao (2012), the periodicity was established in a

stronger sense as a random periodic solution (see Definition 1.1 from Feng and Zhao (2012)); this definition does not assume the equality $u(\cdot, t) = u(\cdot, T)$. In Feng and Zhao (2012), semi-linear parabolic Ito equations with a self-adjoint main operator were considered. There are also results for almost periodic in mean-square sense solutions; see, e.g., Tudor (1992), Da Prato and Tudor (1995), Arnold and Tudor (1998), Bezandry and Diagana (2007), Bedouhene *et al* (2012), Crewe (2013).

The present paper addresses these and related problems again for a single period setting. We found examples of SPDEs where almost periodicity conditions and exact periodicity conditions hold almost surely a well as more general non-local boundary value conditions. It appears that this is possible with the replacement of forward SPDEs for backward SPDEs.

We consider linear Dirichlet condition at the boundary of the state domain; the equations are of a parabolic type and are not necessary self-adjoint. The standard boundary value Cauchy condition at the one fixed time is replaced by a condition that mixes in one equation the values of the solution at different times over given time interval, including the terminal time and continuously distributed times. This is a novel setting comparing with the periodic conditions for the distributions, or with conditions from Klünger (2001) and Feng and Zhao (2012), or with conditions for the expectations from Dokuchaev (2008), or mean-square almost periodicity from Tudor (1992). These conditions include, for instance, conditions $\kappa u(\cdot, 0) = u(\cdot, T)$ a.e. with $\kappa \in \mathbf{R}$ (Theorems 3.1-3.5). We present sufficient conditions for existence and regularity of solutions in L_2 -setting (Corollary 3.1 and Corollary 3.2). Corollary 3.2 establishes existence of modified almost periodic solution. It can be noted that we consider "almost periodicity" on a single time period; this setting is easier than a multiperiod setting on infinite interval since it does not require to formalize a periodic extension of the coefficients. However, with respect to a single period, our "almost periodicity" property from Corollary 3.2) is stronger than the mean square "almost periodicity". Finally, we present sufficient conditions for existence and regularity of almost surely exact periodicity (Theorem 3.5). The proofs is based on compactness and Fredholm theory in L_2 -spaces. The periodic solution obtained here can be considered as a generalization of a classical Brownian bridge for the case of an infinite dimensional state space.

It can be noted the almost surely periodicity was achieved for the purely backward SPDEs; the previous result of this kind was obtained for forward-backward SPDEs in Feng and Zhao (2012). Usually, backward SPDEs are usually associated with a Cauchy condition at the terminal time. With regards to the general theory of SPDEs, our results open a way to extend applications of backward SPDEs on the problems with periodic and mixed in time conditions.

Related problems were considered in Dokuchaev (2012b,c) for a less general backward SPDE

with $\bar{\beta}_i = 0$, in the notations of the present paper. In Dokuchaev (2012b), the approach was based on the contraction mapping theorem in a L_{∞} -space; this approach is not applicable for the more general SPDEs considered in the present paper. In Dokuchaev (2012c), related forward and backward SPDEs with $\bar{\beta}_i = 0$ were studied in an unified framework.

2 The problem setting and definitions

We are given a standard complete probability space $(\Omega, \mathcal{F}, \mathbf{P})$ and a right-continuous filtration \mathcal{F}_t of complete σ -algebras of events, $t \geq 0$. We are given also a N-dimensional Wiener process w(t) with independent components; it is a Wiener process with respect to \mathcal{F}_t .

Assume that we are given an open domain $D \subset \mathbf{R}^n$ such that either $D = \mathbf{R}^n$ or D is bounded with C^2 -smooth boundary ∂D . Let T > 0 be given, and let $Q \stackrel{\Delta}{=} D \times [0, T]$.

We will study the following boundary value problem in Q

$$d_t u + (\mathcal{A}u + \varphi) dt + \sum_{i=1}^N B_i \chi_i dt = \sum_{i=1}^N \chi_i(t) dw_i(t), \quad t \ge 0,$$
(2.1)

$$u(x,t,\omega)|_{x\in\partial D} = 0 \tag{2.2}$$

$$u(\cdot,T) - \Gamma u(\cdot) = \xi. \tag{2.3}$$

Here $u = u(x, t, \omega), \varphi = \varphi(x, t, \omega), \chi_i = \chi_i(x, t, \omega), (x, t) \in Q, \omega \in \Omega.$

In (2.3), Γ is a linear operator that maps functions defined on $Q \times \Omega$ to functions defines on $D \times \Omega$. For instance, the case where $\Gamma u = u(\cdot, 0)$ is not excluded; this case corresponds to the periodic type boundary condition

$$u(\cdot, T) - u(\cdot, 0) = \xi.$$
 (2.4)

In (2.1),

$$\mathcal{A}v \stackrel{\Delta}{=} \sum_{i=1}^{n} \frac{\partial}{\partial x_{i}} \sum_{j=1}^{n} \left(b_{ij}(x,t,\omega) \frac{\partial v}{\partial x_{j}}(x) \right) + \sum_{i=1}^{n} f_{i}(x,t,\omega) \frac{\partial v}{\partial x_{i}}(x) - \lambda(x,t,\omega)v(x), \tag{2.5}$$

where b_{ij}, f_i, x_i are the components of b, f, and x respectively, and

$$B_i v \stackrel{\Delta}{=} \frac{dv}{dx} (x) \beta_i(x, t, \omega) + \bar{\beta}_i(x, t, \omega) v(x), \quad i = 1, \dots, N.$$
(2.6)

We assume that the functions $b(x,t,\omega) : \mathbf{R}^n \times [0,T] \times \Omega \to \mathbf{R}^{n \times n}, \beta_j(x,t,\omega) : \mathbf{R}^n \times [0,T] \times \Omega \to \mathbf{R}^n, \overline{\beta}_i(x,t,\omega) : \mathbf{R}^n \times [0,T] \times \Omega \to \mathbf{R}, f(x,t,\omega) : \mathbf{R}^n \times [0,T] \times \Omega \to \mathbf{R}^n, \lambda(x,t,\omega) : \mathbf{R}^n \times [0,T] \times \Omega \to \mathbf{R}, \chi_i(x,t,\omega) : \mathbf{R}^n \times [0,T] \times \Omega \to \mathbf{R}, \text{ and } \varphi(x,t,\omega) : \mathbf{R}^n \times [0,T] \times \Omega \to \mathbf{R} \text{ are progressively}$ measurable with respect to \mathcal{F}_t for all $x \in \mathbf{R}^n$, and the function $\xi(x,\omega) : \mathbf{R}^n \times \Omega \to \mathbf{R}$ is \mathcal{F}_0 -measurable for all $x \in \mathbf{R}^n$. In fact, we will also consider φ and ξ from wider classes. In particular, we will consider generalized functions φ .

If the functions b, f, λ , φ , Γ , and ξ , are deterministic, then $\chi_i \equiv 0$ and equation (2.1) is deterministic.

Spaces and classes of functions

We denote by $\|\cdot\|_X$ the norm in a linear normed space X, and $(\cdot, \cdot)_X$ denote the scalar product in a Hilbert space X.

We introduce some spaces of real valued functions.

Let $G \subset \mathbf{R}^k$ be an open domain, then $W_q^m(G)$ denote the Sobolev space of functions that belong to $L_q(G)$ together with the distributional derivatives up to the *m*th order, $q \ge 1$.

We denote by $|\cdot|$ the Euclidean norm in \mathbf{R}^k , and \overline{G} denote the closure of a region $G \subset \mathbf{R}^k$. Let $H^0 \triangleq L_2(D)$, and let $H^1 \triangleq W_2^{(1)}(D)$ be the closure in the $W_2^{(1)}(D)$ -norm of the set of all smooth functions $u: D \to \mathbf{R}$ such that $u|_{\partial D} \equiv 0$. Let $H^2 = W_2^{(2)}(D) \cap H^1$ be the space equipped with the norm of $W_2^{(2)}(D)$. The spaces H^k and $W_2^k(D)$ are called Sobolev spaces, they are Hilbert spaces, and H^k is a closed subspace of $W_2^k(D)$, k = 1, 2.

Let H^{-1} be the dual space to H^1 , with the norm $\|\cdot\|_{H^{-1}}$ such that if $u \in H^0$ then $\|u\|_{H^{-1}}$ is the supremum of $(u, v)_{H^0}$ over all $v \in H^1$ such that $\|v\|_{H^1} \leq 1$. H^{-1} is a Hilbert space.

We shall write $(u, v)_{H^0}$ for $u \in H^{-1}$ and $v \in H^1$, meaning the obvious extension of the bilinear form from $u \in H^0$ and $v \in H^1$.

We denote by $\bar{\ell}_k$ the Lebesgue measure in \mathbf{R}^k , and we denote by $\bar{\mathcal{B}}_k$ the σ -algebra of Lebesgue sets in \mathbf{R}^k .

We denote by $\overline{\mathcal{P}}$ the completion (with respect to the measure $\overline{\ell}_1 \times \mathbf{P}$) of the σ -algebra of subsets of $[0,T] \times \Omega$, generated by functions that are progressively measurable with respect to \mathcal{F}_t .

We introduce the spaces

$$\begin{split} X^{k}(s,t) &\triangleq L^{2}([s,t] \times \Omega, \bar{\mathcal{P}}, \bar{\ell}_{1} \times \mathbf{P}; H^{k}), \\ Z^{k}_{t} &\triangleq L^{2}(\Omega, \mathcal{F}_{t}, \mathbf{P}; H^{k}), \\ \mathcal{C}^{k}(s,t) &\triangleq C\left([s,t]; Z^{k}_{T}\right), \qquad k = -1, 0, 1, 2, \\ \mathcal{X}^{k}_{c} &= L^{2}([0,T] \times \Omega, \bar{\mathcal{P}}, \bar{\ell}_{1} \times \mathbf{P}; \ C^{k}(\bar{D})), \quad k \geq 0 \end{split}$$

The spaces $X^k(s,t)$ and Z^k_t are Hilbert spaces.

We introduce the spaces

$$Y^{k}(s,t) \stackrel{\Delta}{=} X^{k}(s,t) \cap \mathcal{C}^{k-1}(s,t), \quad k = 1, 2,$$

with the norm $||u||_{Y^k(s,T)} \triangleq ||u||_{X^k(s,t)} + ||u||_{\mathcal{C}^{k-1}(s,t)}$. For brevity, we shall use the notations $X^k \triangleq X^k(0,T), \mathcal{C}^k \triangleq \mathcal{C}^k(0,T)$, and $Y^k \triangleq Y^k(0,T)$.

We also introduce spaces C_{PC}^k consisting of $u \in C^k$ such that either $u \in C^k$ or there exists $\theta = \theta(u) \in [0,T]$ such that $||u(\cdot,t)||_{Z_T^k}$ is bounded, $u(\cdot,t)$ is continuous in Z_T^k in $t \in [0,\theta]$, and $u(\cdot,t)$ is continuous in Z_T^k in $t \in [\theta + \varepsilon, T]$ for any $\varepsilon > 0$. We also introduce spaces $Y_{PC}^k = X^k \cap C_{PC}^{k-1}$, with the norms from Y^k .

Conditions for the coefficients

To proceed further, we assume that Conditions 2.1-2.3 remain in force throughout this paper.

Condition 2.1 The matrix $b = b^{\top}$ is symmetric and bounded. In addition, there exists a constant $\delta > 0$ such that

$$y^{\top}b(x,t,\omega)y - \frac{1}{2}\sum_{i=1}^{N}|y^{\top}\beta_i(x,t,\omega)|^2 \ge \delta|y|^2 \quad \forall y \in \mathbf{R}^n, \ (x,t) \in D \times [0,T], \ \omega \in \Omega.$$
(2.7)

Condition 2.2 The functions $f(x,t,\omega)$, $\lambda(x,t,\omega)$, $\beta_i(x,t,\omega)$, and $\bar{\beta}_i(x,t,\omega)$, are bounded.

Condition 2.3 The mapping $\Gamma: Y^1_{PC} \to Z^0_T$ is linear and continuous.

Condition 2.3 allows, for instance, to consider Γ such $\Gamma u = u(\cdot, 0)$, i.e., it covers periodic boundary value conditions (2.4). Another example includes the case where there exists an integer $m \geq 0$, a set $\{t_i\}_{i=1}^m \subset [0,T)$, and linear continuous operators $\tilde{\Gamma}_0 : L_2([0,T]; \mathcal{B}_1, \ell_1, H^0) \to H^0$, $\tilde{\Gamma}_i : H^0 \to H^0, i = 1, ..., N$, such that

$$\Gamma u = \tilde{\Gamma}_0 u + \sum_{i=1}^m \tilde{\Gamma}_i u(\cdot, t_i).$$

In particular, it includes

$$\tilde{\Gamma}_0 u = \int_0^T k_0(t) u(\cdot, t) dt, \quad \tilde{\Gamma}_i u(\cdot, t_i) = k_i u(\cdot, t_i),$$

where $k_0(\cdot) \in L_2(0,T)$ and $k_i \in \mathbf{R}$. It covers also Γ such that

$$\tilde{\Gamma}_0 u = \int_0^T dt \int_D k_0(x, y, t) u(y, t) dx, \quad \tilde{\Gamma}_i u(\cdot, t_i)(x) = \int_D k_i(x, y) u(y, t_i) dy,$$

where $k_i(\cdot)$ are some regular enough kernels.

We introduce the set of parameters

$$\mathcal{P} \stackrel{\Delta}{=} \left(n, D, T, \Gamma, \delta, \\ \operatorname{ess\,sup}_{x,t,\omega,i} \Big[|b(x,t,\omega)| + |f(x,t,\omega)| + |\lambda(x,t,\omega)| + |\beta_i(x,t,\omega)| + |\bar{\beta}_i(x,t,\omega)| \Big].$$

Sometimes we shall omit ω .

The definition of solution

Proposition 2.1 Let $\zeta \in X^0$, let a sequence $\{\zeta_k\}_{k=1}^{+\infty} \subset L^{\infty}([0,T] \times \Omega, \ell_1 \times \mathbf{P}; C(D))$ be such that all $\zeta_k(\cdot, t, \omega)$ are progressively measurable with respect to \mathcal{F}_t , and let $\|\zeta - \zeta_k\|_{X^0} \to 0$. Let $t \in [0,T]$ and $j \in \{1,\ldots,N\}$ be given. Then the sequence of the integrals $\int_0^t \zeta_k(x,s,\omega) dw_j(s)$ converges in Z_t^0 as $k \to \infty$, and its limit depends on ζ , but does not depend on $\{\zeta_k\}$.

Proof follows from completeness of X^0 and from the equality

$$\mathbf{E}\int_0^t \|\zeta_k(\cdot, s, \omega) - \zeta_m(\cdot, s, \omega)\|_{H^0}^2 \, ds = \int_D \, dx \, \mathbf{E} \left(\int_0^t \left(\zeta_k(x, s, \omega) - \zeta_m(x, s, \omega)\right) \, dw_j(s)\right)^2.$$

Definition 2.1 Let $\zeta \in X^0$, $t \in [0,T]$, $j \in \{1,\ldots,N\}$, then we define $\int_0^t \zeta(x,s,\omega) dw_j(s)$ as the limit in Z_t^0 as $k \to \infty$ of a sequence $\int_0^t \zeta_k(x,s,\omega) dw_j(s)$, where the sequence $\{\zeta_k\}$ is such as in Proposition 2.1.

Definition 2.2 Let $u \in Y^1$, $\chi_i \in X^0$, i = 1, ..., N, and $\varphi \in X^{-1}$. We say that equations (2.1)-(2.2) are satisfied if

$$u(\cdot, t, \omega) = u(\cdot, T, \omega) + \int_{t}^{T} \left(\mathcal{A}u(\cdot, s, \omega) + \varphi(\cdot, s, \omega) \right) ds$$
$$+ \sum_{i=1}^{N} \int_{t}^{T} B_{i}\chi_{i}(\cdot, s, \omega) ds - \sum_{i=1}^{N} \int_{t}^{T} \chi_{i}(\cdot, s) dw_{i}(s)$$

for all r, t such that $0 \le r < t \le T$, and this equality is satisfied as an equality in Z_T^{-1} .

Note that the condition on ∂D is satisfied in the sense that $u(\cdot, t, \omega) \in H^1$ for a.e. t, ω . Further, $u \in Y^1$, and the value of $u(\cdot, t, \omega)$ is uniquely defined in Z_T^0 given t, by the definitions of the corresponding spaces. The integrals with dw_i in (2.8) are defined as elements of Z_T^0 . The integral with ds in (2.8) is defined as an element of Z_T^{-1} . In fact, Definition 2.2 requires for (2.1) that this integral must be equal to an element of Z_T^0 in the sense of equality in Z_T^{-1} .

3 The main results

Theorem 3.1 There exist a number $\kappa = \kappa(\mathcal{P}) > 0$ such that problem (2.1)-(2.3) has an unique solution $(u, \chi_1, ..., \chi_N)$ in the class $Y^1 \times (X^0)^N$, for any $\varphi \in X^{-1}$, $\xi \in Z_T^0$, and any Γ such that $\|\Gamma\| \leq \kappa$, where $\|\Gamma\|$ is the norms of the operator $\Gamma : Y^1 \to Z_T^0$. In addition,

$$\|u\|_{Y^1} + \sum_{i=1}^N \|\chi_i\|_{X^0} \le C\left(\|\varphi\|_{X^{-1}} + \|\xi\|_{Z^0_T}\right),\tag{3.1}$$

where $C = C(\kappa, \mathcal{P}) > 0$ is a constant that depends only on κ and \mathcal{P} .

Let $\mathbb I$ denote the indicator function.

Theorem 3.2 Let Γ be such that there exists $\theta < T$ such that $\Gamma u = \Gamma(\mathbb{I}_{\{t \le \theta\}}u)$. Then

$$\|u\|_{Y^{1}} + \sum_{i=1}^{N} \|\chi_{i}\|_{X^{0}} \le C \left(\|\varphi\|_{X^{-1}} + \|u\|_{X^{-1}} + \|\xi\|_{Z^{0}_{T}} \right)$$
(3.2)

for all solutions $(u, \chi_1, ..., \chi_N)$ of problem (2.1)-(2.3) in the class $Y^1 \times (X^0)^N$, where $C = C(\mathcal{P}) > 0$ depends only on \mathcal{P} and does not depend on u, φ and ξ .

Starting from now and up to the end of this section, we assume that Condition 3.1 holds.

Condition 3.1 (i) The domain D is bounded.

- (ii) The functions $b(x,t,\omega)$, $f(x,t,\omega)$, $\lambda(x,t,\omega)$, $\beta_i(x,t,\omega)$ and $\bar{\beta}_i(x,t,\omega)$ are differentiable in x for a.e. t, ω , and the corresponding derivatives are bounded.
- (iii) $\beta_i(x,t,\omega) = 0$ for $x \in \partial D$, i = 1, ..., N.
- (iv) \mathcal{F}_0 is the **P**-augmentation of the set $\{\emptyset, \Omega\}$.
- (v) $\Gamma = \Gamma_0 + \Gamma_1$, where $\Gamma_0 : X^1 \to Z_0^1$ and $\Gamma_1 : Y_{PC}^1 \to Z_0^1$ are continuous linear operators such that there exists $\theta < T$ such that $\|\Gamma_1(\mathbb{I}_{\{t \le \theta\}}u)\|_{Z_0^1} \le \|u|_{t \le \theta}\|_{\mathcal{C}^1(0,\theta)}$ for all $u \in Y^1$ such that $u|_{t \le \theta} \in \mathcal{C}^1(0,\theta)$.

In particular, Condition 3.1(ii) implies that there exist modifications of β_i such that the functions $\beta_i(x, t, \omega)$ are continuous in x for a.e. t, ω . We assume that β_i are such functions.

Example 3.1 The assumptions on Γ in Condition 3.1 are satisfied, for instance, if there exists an integer $m \ge 0$, a set $\{t_i\}_{i=1}^m \subset [0,T)$, and linear continuous operators $\overline{\Gamma} : L_2(Q) \to H^0$, $\bar{\Gamma}_i: H^0 \to H^0, i = 0, 1, ..., N$, such that the operators $\bar{\Gamma}: L_2([0, T]; \mathcal{B}_1, \ell_1, H^1) \to W_2^1(D)$ and $\bar{\Gamma}_i: H^1 \to W_2^1(D)$ are continuous and

$$\Gamma u = \bar{\Gamma}_0 u(\cdot, 0) + \mathbf{E} \{ \bar{\Gamma} u + \sum_{i=1}^m \bar{\Gamma}_i u(\cdot, t_i) \}.$$

Theorem 3.3 Assume that problem (2.1)-(2.3) with $\varphi \equiv 0$, $\xi \equiv 0$, does not admit non-zero solutions $(u, \chi_1, ..., \chi_N)$ in the class $Y^1 \times (X^0)^N$. Then problem (2.1)-(2.3) has a unique solution $(u, \chi_1, ..., \chi_N)$ in the class $Y^1 \times (X^0)^N$, for any $\varphi \in X^{-1}$, and $\xi \in H^0$. In addition,

$$\|u\|_{Y^{1}} + \sum_{i=1}^{N} \|\chi_{i}\|_{X^{0}} \le C \left(\|\varphi\|_{X^{-1}} + \|\xi\|_{H^{0}}\right), \qquad (3.3)$$

where C > 0 does not depend on φ and ξ .

Theorem 3.4 There exists $\varepsilon_0 > 0$ such that, for any $\varepsilon \in (-\varepsilon_0, \varepsilon_0)$ such that $\varepsilon \neq 0$, problem (2.1)-(2.2) with the boundary value condition

$$u(\cdot, T) - (1 + \varepsilon)\Gamma u = \xi \tag{3.4}$$

has a unique solution $(u, \chi_1, ..., \chi_N)$ in the class $Y^1 \times (X^0)^N$ for any $\varphi \in X^{-1}$ and $\xi \in H^0$. In addition,

$$\|u\|_{Y^{1}} + \sum_{i=1}^{N} \|\chi_{i}\|_{X^{0}} \le C \left(\|\varphi\|_{X^{-1}} + \|\xi\|_{H^{0}}\right), \qquad (3.5)$$

where $C = C(\varepsilon) > 0$ does not depend on φ and ξ .

Corollary 3.1 For any $\kappa \in R$, there exists $\varepsilon_0 = \varepsilon_0(\mathcal{P}, \kappa) > 0$ such that, for any $\varepsilon \in (-\varepsilon_0, \varepsilon_0) \setminus \{0\}$, for any $\varphi \in X^{-1}$, there exists a unique solution $(u, \chi_1, ..., \chi_N)$ in the class $Y^1 \times (X^0)^N$, of problem (2.1), (2.2) with the boundary conditions

$$u(\cdot, T) = \kappa (1 + \varepsilon) u(\cdot, 0). \tag{3.6}$$

Corollary 3.2 Corollary 3.1 with $\kappa = 1$ implies that backward SPDE (2.1)-(2.2) can be regarded as an almost surely almost periodic solution on a single time period.

Note that Tudor (1992) and other authors considered almost periodic solution for SPDEs on a infinite time horizon, i.e., for many periods. We consider a single period only.

Remark 3.1 The "almost periodicity" in the mean-square sense of Tudor (1992) requires that, for any $\varepsilon > 0$, $\|u(\cdot, 0) - u(\cdot, T)\|_{Z^0_T} \le \varepsilon$; we establish that the equality $u(\cdot, 0) = (1 + \varepsilon)u(\cdot, T)$ can be achieved. It follows that $||u(\cdot,0) - u(\cdot,T)||_{Z_T^0} \leq \varepsilon ||u(\cdot,T)||_{Z_T^0} \leq \varepsilon C ||\varphi||_{X^{-1}}$. Therefore, for a single time period, the "almost periodicity" property in Corollary 3.2) implies the "almost periodicity" in the mean-square sense from Tudor (1992). Moreover, this condition is stronger than the mean-square almost periodicity since it requires that the shapes of $u(\cdot,T)$ and $u(\cdot,0)$ are the same up to proportionality.

The remaining part of the section devoted to an example where an exact periodic condition is satisfied almost surely; this corresponds to the case where $\varepsilon = 0$ in (3.6).

Let functions $\tilde{\beta}_i : Q \times \Omega \to \mathbf{R}^n, i = 1, \dots, M$, be such that

$$2b(x,t,\omega) = \sum_{i=1}^{N} \beta_i(x,t,\omega) \,\beta_i(x,t,\omega)^{\top} + \sum_{j=1}^{M} \tilde{\beta}_j(x,t,\omega) \,\tilde{\beta}_j(x,t,\omega)^{\top},$$

and $\tilde{\beta}_i$ has the similar properties as β_i . (Note that, by Condition 2.1, $2b > \sum_{i=1}^N \beta_i \beta_i^\top$).

Let $\tilde{w}(t) = (\tilde{w}_1(t), \dots, \tilde{w}_M(t))$ be a new Wiener process independent on w(t). Let $a \in L_2(\Omega, \mathcal{F}, \mathbf{P}; \mathbf{R}^n)$ be a vector such that $a \in D$. We assume also that a is independent from $(w(t) - w(t_1), \hat{w}(t) - \hat{w}(t_1))$ for all $t > t_1 > s$. Let $s \in [0, T)$ be given. Consider the following Ito equation

$$dy(t) = \tilde{f}(y(t), t) dt + \sum_{i=1}^{N} \beta_i(y(t), t) dw_i(t) + \sum_{j=1}^{M} \tilde{\beta}_j(y(t), t) d\tilde{w}_j(t),$$

$$y(s) = x.$$
 (3.7)

Here, $\tilde{f} = \hat{f} - \sum_i \bar{\beta}_i \beta_i$, $\tilde{f} : D \times [0, T] \times \Omega \to \mathbf{R}^n$ is a vector functions with the components \tilde{f}_i . Let $y(t) = y^{a,s}(t)$ be the solution of (3.7), and let $\tau^{a,s} \triangleq \inf\{t \ge s : y^{a,s}(t) \notin D\}$.

The following lemma is a modification for the case of random coefficients of Lemma 2.1 from Dokuchaev (2004).

Lemma 3.1 There exists $\nu \in (0,1)$ that depends only on \mathcal{P} such that $\mathbf{P}(\tau^{a,0} > T) \leq \nu$ for any random vector a such that $a \in D$ a.s. and a does not depend on w(t) - w(r) for all t > r > 0.

Theorem 3.5 Let the functions b, f and λ be such that the operator \mathcal{A} can be represented as

$$\mathcal{A}v = \sum_{i,j=1}^{n} b_{ij}(x,t,\omega) \frac{\partial^2 v}{\partial x_i \partial x_j}(x) + \sum_{i=1}^{n} \widehat{f}_i(x,t,\omega) \frac{\partial v}{\partial x_i}(x) - \widehat{\lambda}(x,t,\omega)v(x),$$

where $\widehat{\lambda}(x,t,\omega) \geq 0$ a.e., and where \widehat{f}_i are bounded functions.

Further, let at least one of the following conditions is satisfied:

(i) There exists $c_{\lambda} > 0$ such that $\widehat{\lambda}(x, t, \omega) \ge c_{\lambda}$ for all x, t, ω ; or

- (*ii*) $\kappa \in (-1, 1)$; or
- (iii) For $\nu \in (0, 1)$ from Lemma 3.1,

$$\frac{1}{2} \sum_{i=1}^{N} \int_{0}^{T} \sup_{x,\omega} \bar{\beta}_{i}(x,t,\omega)^{2} dt + \log \nu < 0.$$
(3.8)

Furthermore, let $b \in \mathcal{X}_c^3$, $\hat{f} \in \mathcal{X}_c^2$, $\hat{\lambda} \in \mathcal{X}_c^1$, $\beta_i \in \mathcal{X}_c^3$. Then there exists $\bar{\kappa} > 1$ such that, for any $\kappa \in [-\bar{\kappa}, \bar{\kappa}]$, problem (2.1)-(2.2) with the boundary condition

$$u(\cdot, T) - \kappa u(\cdot, 0) = \xi \tag{3.9}$$

has a unique solution $(u, \chi_1, ..., \chi_N)$ in the class $Y^1 \times (X^0)^N$ for any $\varphi \in X^{-1}$ and $\xi \in Z_T^0$. In addition, (3.3) holds with C > 0 that does not depend on φ and ξ .

4 Proofs

Let $s \in (0,T]$, $\varphi \in X^{-1}$ and $\Phi \in Z_s^0$. Consider the problem

$$d_{t}u + (\mathcal{A}u + \varphi) dt + \sum_{i=1}^{N} B_{i}\chi_{i}(t)dt = \sum_{i=1}^{N} \chi_{i}(t)dw_{i}(t), \quad t \leq s,$$

$$u(x, t, \omega)|_{x \in \partial D},$$

$$u(x, s, \omega) = \Phi(x, \omega).$$

(4.1)

The following lemma represents an analog of the so-called "the first energy inequality", or "the first fundamental inequality" known for deterministic parabolic equations (see, e.g., inequality (3.14) from Ladyzhenskaya (1985), Chapter III).

Lemma 4.1 Assume that Conditions 2.1–2.3 are satisfied. Then problem (4.1) has an unique solution a unique solution $(u, \chi_1, ..., \chi_N)$ in the class $Y^1 \times (X^0)^N$ for any $\varphi \in X^{-1}(0, s)$, $\Phi \in Z_s^0$, and

$$\|u\|_{Y^{1}(0,s)} + \sum_{i=1}^{N} \|\chi_{i}\|_{X^{0}} \le C\left(\|\varphi\|_{X^{-1}(0,s)} + \|\Phi\|_{Z^{0}_{s}}\right),$$
(4.2)

where $C = C(\mathcal{P})$ does not depend on φ and ξ .

(See, e.g., Dokuchaev (1991) or Theorem 4.2 from Dokuchaev (2010)).

Note that the solution $u = u(\cdot, t)$ is continuous in t in $L_2(\Omega, \mathcal{F}, \mathbf{P}, H^0)$, since $Y^1(0, s) = X^1(0, s) \cap \mathcal{C}^0(0, s)$.

Introduce operators $L_s : X^{-1}(0,s) \to Y^1(0,s)$ and $\mathcal{L}_s : Z_s^0 \to Y^1(0,s)$, such that $u = L_s \varphi + \mathcal{L}_s \Phi$, where $(u, \chi_1, ..., \chi_N)$ is the solution of problem (4.1) in the class $Y^2 \times (X^1)^N$. By Lemma 4.1, these linear operators are continuous.

Introduce operators $\mathcal{Q}: Z_T^0 \to Z_T^0$ and $\mathcal{T}: X^{-1} \to Z_T^0$ such that $\mathcal{Q}\Phi = \Gamma \mathcal{L}_T \Phi$ and $\mathcal{T}\varphi = \Gamma \mathcal{L}_T \varphi$, i.e., $\mathcal{Q}\Phi + \mathcal{T}\varphi = \Gamma u$, where u is the solution in Y^1 of problem (4.1) with s = T, $\varphi \in X^{-1}$, and $\Phi \in Z_T^0$. It is easy to see that if the operator $\Gamma: Y^1 \to Z_T^0$ is continuous, then the operators $\mathcal{Q}: Z_T^0 \to Z_T^0$ and $\mathcal{T}: X^{-1} \to Z_T^0$ are linear and continuous. In particular, $\|\mathcal{Q}\| \leq \|\Gamma\| \|\mathcal{L}_T\|$, where $\|\mathcal{Q}\|, \|\Gamma\|$, and $\|\mathcal{L}_T\|$, are the norms of the operators $\mathcal{Q}: Z_T^0 \to Z_T^0, \Gamma: Y^1 \to Z_T^0$, and $\mathcal{L}_T: Z_T^0 \to Y^1$, respectively.

Lemma 4.2 Assume that the operator $\Gamma : Y^1 \to Z_T^0$ is continuous. If the operator $(I - Q)^{-1} : Z_T^0 \to Z_T^0$ is also continuous then problem (4.1) has a unique solution $(u, \chi_1, ..., \chi_N)$ in the class $Y^1 \times (X^0)^N$ for any $\varphi \in X^{-1}$, $\Phi \in Z_T^0$. For this solution,

$$u = L_T \varphi + \mathcal{L}_T (I - \mathcal{Q})^{-1} (\xi + \mathcal{T} \varphi)$$
(4.3)

and

$$\|u\|_{Y^{1}(0,s)} + \sum_{i=1}^{N} \|\chi_{i}\|_{X^{0}} \leq C \left(\|\varphi\|_{X^{-1}(0,s)} + \|\Phi\|_{Z^{0}_{s}} \right),$$

where $C = C(\mathcal{P})$ does not depend on φ and ξ .

Proof of Lemma 4.2. For brevity, we denote $u(\cdot, t) = u(x, t, \omega)$. Clearly, $u \in Y^1$ is the solution of problem (2.1)-(2.3) with some $(\chi_1, ..., \chi_N) \in (X^0)^N$ if and only if

$$u = \mathcal{L}_T u(\cdot, T) + L_T \varphi, \tag{4.4}$$

$$u(\cdot, T) - \Gamma u = \xi. \tag{4.5}$$

Since $\Gamma u = \mathcal{Q}u(\cdot, T) + \mathcal{T}\varphi$, equation (4.5) can be rewritten as

$$u(\cdot, T) - \mathcal{Q}u(\cdot, T) - \mathcal{T}\varphi = \xi.$$
(4.6)

By the continuity of $(I - Q)^{-1}$, equation (4.6) can be rewritten as

$$u(\cdot, T) = (I - \mathcal{Q})^{-1}(\xi + \mathcal{T}\varphi).$$

Therefore, equations (4.4)-(4.5) imply that

$$u = L_T \varphi + \mathcal{L}_T u(\cdot, T) = L_T \varphi + \mathcal{L}_T (I - \mathcal{Q})^{-1} (\xi + \mathcal{T} \varphi).$$

Further, let us show that if (4.3) holds then equations (4.4)-(4.5) hold. Let u be defined by (4.3). Since $u = L_T \varphi + \mathcal{L}_T u(\cdot, T)$, it follows that $u(\cdot, T) = (I - \mathcal{Q})^{-1}(\xi + \mathcal{T}\varphi)$. Hence

$$u(\cdot, T) - \mathcal{Q}u(\cdot, T) = \xi + \mathcal{T}\varphi,$$

i.e., $u(\cdot, T) - \Gamma \mathcal{L}_T u(\cdot, T) = \xi + \mathcal{T} \varphi = \xi + \Gamma L_T \varphi$. Hence

$$u(\cdot, T) - \Gamma[\mathcal{L}_T u(\cdot, T) + L_T \varphi] = \xi.$$

This means that (4.4)-(4.5) hold. Then the proof of Lemma 4.2 follows. \Box

Proof of Theorem 3.1. Since the operator $\mathcal{Q}: Z_T^0 \to Z_T^0$ is continuous, the operator $(I-\mathcal{Q})^{-1}: Z_T^0 \to Z_T^0$ is continuous for small enough $\|\mathcal{Q}\|$, i.e. for a small enough $\kappa > 0$. Then the proof of Theorem 3.1 follows. \Box

Proof of Theorem 3.2. For a real q > 0, set $u_q(x,t,\omega) \stackrel{\Delta}{=} e^{q(T-t)}u(x,t,\omega)$. Then u_q is the solution of problem (2.1)-(2.3) with φ replaced by $e^{q(T-t)}\varphi(x,t) + qu_q(x,t)$, and with Γ replaced by the operator defined such that Γ_q , where

$$\Gamma_q u_q = \Gamma u_s$$

By the assumptions on Γ , we have that

$$\|\Gamma_{q}u_{q}\|_{Z_{T}^{0}} = \|\Gamma u\|_{Z_{T}^{0}} = \|\Gamma \mathbb{I}_{t \le \theta}u\|_{Z_{T}^{0}} = \left\|\Gamma \mathbb{I}_{\{t \le \theta\}}e^{-q(T-t)}u_{q}\right\|_{Z_{T}^{0}} \le e^{-q(T-\theta)}\|\Gamma\|\|u\|_{X^{0}(0,\theta)}.$$

It follows that $\|\Gamma_q\| \to 0$ as $q \to +\infty$, for the norm of the operator $\Gamma_q : Y^1 \to Z_T^T$. By Lemma 4.1 and Theorem 3.1, it follows that, for a large enough q > 0,

$$\begin{aligned} \|u_q\|_{Y^1} + \sum_{i=1}^N \|\chi_i\|_{X^0} &\leq C_1 \left(\|e^{q(T-t)}\varphi + qu_q\|_{X^{-1}} + \|\xi\|_{H_0} \right) \\ &\leq C_2 \left(\|\varphi\|_{X^{-1}} + \|u_q\|_{X^{-1}} + \|\xi\|_{H_0} \right), \end{aligned}$$

where $C_1 = C_1(\mathcal{P}) > 0$ and $C_2 = C_2(q, \mathcal{P}) > 0$ do not depend on u, φ, ξ . Then the proof of Theorem 3.2 follows. \Box

Starting from now, we assume that Condition 3.1 is satisfied, in addition to Conditions 2.1-2.3.

The following lemma represents an analog of the so-called "the second energy inequality", or "the second fundamental inequality" known for the deterministic parabolic equations (see, e.g., inequality (4.56) from Ladyzhenskaya (1985), Chapter III).

Lemma 4.3 Problem (4.1) has a unique solution $(u, \chi_1, ..., \chi_N)$ in the class $Y^2 \times (X^1)^N$ for any $\varphi \in X^0$, $\Phi \in Z_T^1$, and

$$\|u\|_{Y^2} + \sum_{i=1}^N \|\chi_i\|_{X^1} \le C\left(\|\varphi\|_{X^0} + \|\Phi\|_{Z^1_T}\right),\tag{4.7}$$

where C > 0 does not depend on φ and Φ ; it depends on \mathcal{P} and on the supremums of the derivatives listed in Condition 3.1(ii).

The lemma above represents a reformulation of Theorem 3.4 from Dokuchaev (2010) or Theorem 4.3 from Dokuchaev (2012a). In the cited paper, this result was obtained under some strengthened version of Condition 2.1; this was restrictive. In Du and Tang (2012), this result was obtained without this restriction, i.e., under Condition 2.1 only.

Lemma 4.4 The operator $Q: Z_T^0 \to Z_T^0$ is compact.

Proof of Lemma 4.4. Let $u = \mathcal{L}_0 \Phi$, where $\Phi \in Z_T^0$. By the semi-group property of backward SPDEs from Theorem 6.1 from Dokuchaev (2010), we obtain that $u|_{t \in [0,s]} = \mathcal{L}_s u(\cdot, s)$ for all $s \in (0,T]$. By Lemmas 4.1 and 4.3, we have that

$$\begin{split} \|\Gamma_1 u\|_{Z_0^1}^2 &\leq C_0 \sup_{\tau \in [0,\theta]} \|u(\cdot,\tau)\|_{Z_0^1}^2 \leq C_1 \sup_{\tau \in [0,T]} \inf_{t \in [\tau,T]} \|u(\cdot,t)\|_{Z_t^1}^2 \\ &\leq \sup_{\tau \in [0,T]} \frac{C_1}{T-\tau} \int_{\tau}^T \|u(\cdot,t)\|_{Z_t^1}^2 dt \leq \frac{C_2}{T-\theta} \|\Phi\|_{Z_T^0}^2 \end{split}$$

and

$$\|\Gamma_0 u\|_{Z_0^1}^2 \le C_3 \mathbf{E} \int_0^T \|u(\cdot, t)\|_{Z_T^1}^2 dt \le C_4 \|\Phi\|_{Z_T^0}.$$

for constants $C_i > 0$ which do not depend on Φ . Hence the operator $\mathcal{Q} : Z_T^0 \to Z_T^0$ represents a linear continuous operator $\mathcal{Q} : Z_T^0 \to W_2^1(D)$. Note that, by the definitions, $Z_T^0 = L_2(\Omega, \mathcal{F}_0, \mathbf{P}, L_2(D))$. Since \mathcal{F}_0 is a trivial σ -algebra, the convergence in Z_T^0 is equivalent to convergence in $H^0 = L_2(D)$. Since the embeddings of $W_2^1(D)$ into H^0 and into Z_T^0 are compact operators, the proof of Lemma 4.4 follows. \Box

Proof of Theorem 3.3. By the assumptions, the equation $\mathcal{Q}\Phi = \Phi$ has the only solution $\Phi = 0$ in H^0 . By Lemma 4.4 and by the Fredholm Theorem, the operator $(I - \mathcal{Q})^{-1} : H^0 \to H^0$ is continuous. Then the proof of Theorem 3.3 follows from representation (4.7). \Box

Proof of Theorem 3.4. By Lemma 4.4 and by the Fredholm Theorem again, for any $\varepsilon_0 \in (0, 1)$, there exists a finite set $\Lambda \subset \mathbf{C}$ such that the operator $(\lambda I - \mathcal{Q})^{-1} : H^0 \to H^0$ is continuous for all $\lambda \in (1 - \varepsilon_0, 1 + \varepsilon_0) \setminus \Lambda$. Then the proof of Theorem 3.4 follows from representation (4.7) again. \Box

Corollary 3.1 is a special case of Theorem 3.4 with $\Gamma u = u(\cdot, 0)$.

Proof of Lemma 3.1. This proof represents a modification of the proof for Lemma 2.1 from Dokuchaev (2004) for the case of random coefficients. For the case of $\bar{\beta} = 0$, the proof of Lemma 3.1 can be found in Dokuchaev (2012c).

Let $\mu = (\tilde{f}, \beta, x, s).$

Clearly, there exists a finite interval $D_1 \stackrel{\Delta}{=} (d_1, d_2) \subset \mathbf{R}$ and a bounded domain $D_{n-1} \subset \mathbf{R}^{n-1}$ such that $D \subset D_1 \times D_{n-1}$.

For $(x,s) \in D \times [0,T)$, let $\tau_1^{x,s} \stackrel{\Delta}{=} \inf\{t \geq s : y_1^{x,s}(t) \notin D_1\}$, where $y_1^{x,s}(t)$ is the first component of the vector $y^{x,s}(t) = (y_1^{x,s}(t), ..., y_n^{x,s}(t))$. We have that

$$\mathbf{P}(\tau^{x,0} > T) \le \mathbf{P}(\tau_1^{x,0} > T) = \mathbf{P}(y_1^{x,0}(t) \in D_1 \ \forall t \in [0,T]).$$
(4.8)

Let

$$M^{\mu}(t) \stackrel{\Delta}{=} \sum_{k=1}^{N} \int_{s}^{t} h_{k}(y^{x,0}(r), r) dw_{i}(r) + \sum_{k=N+1}^{N+M} \int_{s}^{t} h_{k}(y^{x,0}(r), r) d\tilde{w}_{i}(r), \quad t \ge s,$$

where $h = (h_1, ..., h_{N+M})$ is a vector that represents the first row of the matrix

$$(\beta_1, ..., \beta_N, \widehat{\beta}_1, ..., \widehat{\beta}_M)$$

with the values in $\mathbf{R}^{n \times (N+M)}$.

Let $\widehat{D}_1 \triangleq (d_1 + K_1, d_2 + K_2)$, where $K_1 \triangleq -d_2 - \vartheta \sup_{x,t,\omega} |\widehat{f}_1(x,t,\omega)|$, $K_2 \triangleq -d_1 + \vartheta \sup_{x,t} |\widehat{f}_1(x,t,\omega)|$. Clearly, \widehat{D}_1 depends only on n, D, and c_f . It is easy to see that

$$\mathbf{P}(y_1^{x,0}(t) \in D_1 \; \forall t \in [0,T]) \le \mathbf{P}(M^{\mu}(t) \in \widehat{D}_1 \; \forall t \in [0,T]).$$
(4.9)

Further,

$$h(y^{x,0}(t),t)^{\top}h(y^{x,0}(t),t) = |h(y^{x,0}(t),t)|^2 \in [\delta, c_{\beta}],$$
(4.10)

where

$$\delta = \inf_{x,0,\omega, \xi \in \mathbf{R}^n: |\xi|=1} 2\xi^{\top} b(x,t,\omega)\xi, \quad c_{\beta} = \sup_{x,0,\omega, \xi \in \mathbf{R}^n: |\xi|=1} 2\xi^{\top} b(x,t,\omega)\xi.$$

Clearly, $M^{\mu}(t)$ is a martingale vanishing at s with quadratic variation process

$$[M^{\mu}]_t \stackrel{\Delta}{=} \int_0^t |h(y^{x,0}(r), r)|^2 dr, \qquad t \ge 0.$$

Let $\theta^{\mu}(t) \triangleq \inf\{r \ge 0: [M^{\mu}]_r > t\}$. Note that $\theta^{\mu}(0) = 0$, and the function $\theta^{\mu}(t)$ is strictly increasing in t given x. By Dambis–Dubins–Schwarz Theorem (see, e.g., Revuz and Yor (1999)), the process $B^{\mu}(t) \triangleq M(\theta^{\mu}(t))$ is a Brownian motion vanishing at t = 0, i.e., $B^{\mu}(0) = 0$, and $M^{\mu}(t) = B^{\mu}([M^{\mu}]_t)$. Clearly,

$$\mathbf{P}(M^{\mu}(t) \in \widehat{D}_{1} \quad \forall t \in [0, s+T]) = \mathbf{P}(B^{\mu}([M^{\mu}]_{t}) \in \widehat{D}_{1} \quad \forall t \in [0, T])$$

$$\leq \mathbf{P}(B^{\mu}(r) \in \widehat{D}_{1} \quad \forall r \in [0, [M^{\mu}]_{T}]).$$
(4.11)

By (4.10), $[M^{\mu}]_T \ge \delta T$ a.s. for all x. Hence

$$\mathbf{P}(B^{\mu}(r) \in \widehat{D}_1 \quad \forall r \in [0, [M^{\mu}]_T]) \le \mathbf{P}(B^{\mu}(r) \in \widehat{D}_1 \quad \forall r \in [0, \delta T]).$$
(4.12)

By (4.8)-(4.9) and (4.11)-(4.12), it follows that

$$\sup_{\mu} \mathbf{P}(\tau^{x,0} > T) \le \nu \stackrel{\Delta}{=} \sup_{\mu} \mathbf{P}(B^{\mu}(r) \in \widehat{D}_{1} \quad \forall r \in [0, \delta T]),$$

and $\nu = \nu(\mathcal{P}) \in (0, 1)$. This completes the proof of Lemma 3.1. \Box

Proof of Theorem 3.5. Let us introduce operators

$$\mathcal{A}^* v \stackrel{\Delta}{=} \sum_{i,j=1}^n \frac{\partial^2}{\partial x_i \partial x_j} \left(b_{ij}(x,t) v(x) \right) - \sum_{i=1}^n \frac{\partial}{\partial x_i} \left(\tilde{f}_i(x,t) v(x) \right) - \widehat{\lambda}(x,t) v(x)$$

and

$$B_i^* v \stackrel{\Delta}{=} -\sum_{k=1}^n \frac{\partial}{\partial x_k} \left(\beta_{ik}(x,t,\omega) v(x)\right) + \bar{\beta}_i(x,t,\omega) v(x), \qquad i = 1, \dots, N.$$

Here b_{ij} , x_i , β_{ik} are the components of b, β_i , and x.

Let $\rho \in Z_s^0$, and let $p = p(x, t, \omega)$ be the solution of the problem

$$d_t p = \mathcal{A}^* p \, dt + \sum_{i=1}^N B_i^* p \, dw_i(t), \quad t \ge s,$$
$$p|_{t=s} = \rho, \qquad p(x, t, \omega)|_{x \in \partial D} = 0.$$

By Theorem 3.4.8 from Rozovskii (1990), this boundary value problem has an unique solution $p \in Y^1(s,T)$. Introduce an operator $\mathcal{M}_s : Z_s^0 \to Y^1(s,T)$ such that $p = \mathcal{M}_s \rho$, where $p \in Y^1(s,T)$ is the solution this boundary value problem.

The following lemma from Dokuchaev (2005) represents an analog of the so-called "the second energy inequality", or "the second fundamental inequality" known for the deterministic parabolic equations (see, e.g., inequality (4.56) from Ladyzhenskaya (1985), Chapter III).

Lemma 4.5 Problem (4.1) has an unique solution $p \in Y^2$ for any $\rho \in Z^1_s$, and

$$\|p\|_{Y^2(s,T)} \le C \|\rho\|_{Z^1_s},\tag{4.13}$$

where C > 0 does not depend on ρ . This C depends on \mathcal{P} and on the supremums of the derivatives in Condition 3.1.

By Theorem 4.2 from Dokuchaev (2010), we have that $\kappa p(\cdot, T) = \mathcal{Q}^* \rho$, i.e.,

$$(\rho, \mathcal{Q}\Phi)_{Z_0^0} = (\rho, \kappa v(\cdot, 0))_{Z_0^0} = (p(\cdot, T), \kappa v(\cdot, T))_{Z_T^0} = (\kappa p(\cdot, T), \Phi)_{Z_T^0}$$
(4.14)

for $v = \mathcal{L}_T \Phi$. (See also Lemma 6.1 from Dokuchaev (1991) and related results in Zhou (1992)).

Suppose that there exists $\Phi \in Z_T^0$ such that $\kappa v(\cdot, 0) = v(\cdot, T)$ for $v = \mathcal{L}_T \Phi$, i.e., $v(\cdot, 0) = \mathcal{Q}\Phi = \Phi$. Let us show that $\Phi = 0$ in this case.

Since $\mathcal{Q}\Phi \in Z_0^0$, it follows that $\Phi \in H^0 = Z_0^0$. Let $p = \mathcal{M}_0\rho$ and $\bar{p}(x,t,0) = \mathbf{E}p(x,t,\omega)$ (meaning the projection from Z_T^0 on $H^0 = Z_0^0$). Introduce an operator $\mathbf{Q}: H^0 \to H^0$ such that $\kappa \bar{p}(\cdot,T) = \mathbf{Q}\rho$. By (4.14), the properties of Φ lead to the equality

$$(\rho - \kappa p(\cdot, T), \Phi(\cdot, T))_{Z_T^0} = (\rho - \kappa \bar{p}(\cdot, T), \Phi(\cdot, T))_{H_0} = 0 \quad \forall \rho \in H^0.$$

$$(4.15)$$

It suffices to show that the set $\{\rho - \kappa \bar{p}(\cdot, T)\}_{\rho \in H^0}$ is dense in H^0 . For this, it suffices to show that the equation $\rho - \mathbf{Q}\rho = z$ is solvable in H^0 for any $z \in H^0$.

Let us show that the operator $\mathbf{Q}: H^0 \to H^0$ is compact. Let p be the solution of (4.13). This means that $\kappa \mathbf{E} p(\cdot, T) = \mathbf{Q} \rho$. By Lemma 4.3, it follows that

$$\|p(\cdot,\tau)\|_{Z^{1}_{\tau}} \le C \|p(\cdot,s)\|_{Z^{1}_{s}}, \quad \tau \in [s,T],$$
(4.16)

where $C_* > 0$ is a constant that does not depend on p, s, and τ .

We have that $p|_{t\in[s,T]} = \mathcal{M}_s p(\cdot, s)$ for all $s \in [0,T]$, and, for $\tau > 0$,

$$\begin{aligned} \|\bar{p}(\cdot,T)\|_{W_{2}^{1}(D)}^{2} &\leq C_{0} \|p(\cdot,T)\|_{Z_{T}^{1}}^{2} \leq C_{1} \inf_{t \in [0,T]} \|p(\cdot,t)\|_{Z_{t}^{1}}^{2} \\ &\leq \frac{C_{1}}{T} \int_{0}^{T} \|p(\cdot,t)\|_{Z_{t}^{1}}^{2} dt \leq \frac{C_{2}}{T} \|p\|_{X^{1}}^{2} \leq \frac{C_{3}}{T} \|\Phi\|_{H^{0}} \end{aligned}$$

for constants $C_i > 0$ that do not depend on Φ . Hence the operator $\mathbf{Q} : H^0 \to H^1$ is continuous. The embedding of H^1 into H^0 is a compact operator (see, e.g., Theorem 7.3 from Ladyzhenskaia (1985), Chapter I).

Let us show that if

$$\kappa \bar{p}(\cdot, T) = \kappa \mathbf{E} p(\cdot, T) = \mathbf{Q} \rho = p(\cdot, 0) \tag{4.17}$$

for some $\rho \in H^0$ then $\rho = 0$.

Let $\rho \in H^0$ be such that $\rho \geq 0$ a.e. and $\int_D \rho(x) dx = 1$. Let $a \in L_2(\Omega, \mathcal{F}, \mathbf{P}; \mathbf{R}^n)$ be independent from the process $(w(\cdot), \widehat{w}(\cdot))$ such that $a \in D$ a.s. and it has the probability density function ρ . Let $p = \mathcal{M}_0 \rho$, and let $y^{a,0}(t)$ be the solution of Ito equation (3.7) with the initial condition y(0) = a.

For
$$t \geq s$$
, set

$$\begin{split} \gamma_M^{x,s}(t) &\stackrel{\Delta}{=} \exp\left[\sum_{i=1}^N \int_s^t \bar{\beta}_i(y^{x,s}(s),s) \, dw_i(s) - \sum_{i=1}^N \frac{1}{2} \, \int_s^t \bar{\beta}_i(y^{x,s}(s),s)^2 \, ds\right],\\ \gamma^{x,s}(t) &\stackrel{\Delta}{=} \exp\left[-\int_s^t \widehat{\lambda}(y^{x,s}(t),t) \, dt\right] \gamma_M^{x,s}(t). \end{split}$$

By Theorem 6.1 from Dokuchaev (2011), for all bounded functions $\Phi \in Z_T^0$ and $u = \mathcal{L}_T \Phi$, we have that

$$\mathbf{E} \int_{D} p(x, T, \omega) \Phi(x, \omega) dx = \int_{D} p(x, 0) u(x, 0) dx = \mathbf{E} \mathbb{I}_{\{\tau^{a, 0} \ge T\}} \gamma^{a, 0}(T) \Phi(y^{a, 0}(T)) \quad \text{a.s.} \quad (4.18)$$

If $D = \mathbf{R}^n$ and $\mathbb{I}_{\{\tau^{a,0} \leq T\}} \equiv 1$, then this equality follows from Theorem 5.3.1 from Rozovskii (2001). Equality (4.18) means that $p(x, T, \omega)$ is the conditional (given \mathcal{F}_T) probability density function of the vector $y^{a,0}(T)$ if the process $y^{a,0}(t)$ is killed at ∂D and if it is killed inside Dwith the rate of killing $\hat{\lambda}$. In particular, it follows that $p(x, t, \omega) \geq 0$ a.e. and

$$\mathbf{E} \int_D p(x, T, \omega) dx = \mathbf{E} \mathbb{I}_{\{\tau^{a,0} \ge T\}} \gamma^{a,0}(T).$$

Assume first that $\widehat{\lambda}(x,t,\omega) \ge c_{\lambda} > 0$ for all x, t, ω , i.e., that condition (i) is satisfied. In this case, $0 \le \gamma^{a,0}(T) \le \nu_1 \gamma_M^{a,0}(T)$, where $\nu_1 \triangleq e^{-c_{\lambda}T}$, $\nu_1 \in (0,1)$. Hence

$$\mathbf{E} \int_{D} p(x, T, \omega) dx = \mathbf{E} \mathbb{I}_{\{\tau^{a,0} \ge T\}} \gamma^{a,0}(T) \le \nu_1 \mathbf{E} \gamma_M^{a,0}(T) = \nu_1.$$

$$(4.19)$$

Assume now that $|\kappa| < 1$, i.e., that condition (ii) is satisfied. This case can be reduced to the case of condition (i) as the following. The problem u(x,t) can be replaced by the problem for $u_q(x,t) = u(x,t)e^{q(T-t)}$ with $q = T^{-1}\log|\kappa| < 0$. The new boundary value condition for u_q is $\kappa e^{-qT}u_q(\cdot,0) + u_q(\cdot,T) = 0$, i.e.,

$$\frac{\kappa}{|\kappa|}u_q(\cdot,0) + u_q(\cdot,T) = 0.$$

In the new equation for $u_q(x,t)$, the coefficient λ has to be replaced by $\lambda - q$. It follows that condition (i) with $c_{\lambda} = -q$ is satisfied for the new problem.

Further, assume that (3.8) is satisfied, i.e. that condition (iii) is satisfied. Let p > 1 and q > 1 be such that 1/p + 1/q = 1. By Lemma 3.1, we have that it that

$$\begin{split} \mathbf{E} \int_{D} p(x,T,\omega) dx &= \mathbf{E} \mathbb{I}_{\{\tau^{a,0} \ge T\}} \gamma^{a,0}(T) \le \|\mathbb{I}_{\{\tau^{a,0} \ge T\}}\|_{L_{p}(\Omega)} \|\gamma_{M}^{a,0}(T)\|_{L_{q}(\Omega)} \\ &= \mathbf{P}(\tau^{a,0} \ge T)^{1/p} \|\gamma_{M}^{a,0}(T)\|_{L_{q}(\Omega)} \le \nu^{1/p} \|\gamma_{M}^{a,0}(T)\|_{L_{q}(\Omega)}. \end{split}$$

Clearly, we have that

$$\|\gamma_M^{a,0}(T)\|_{L_q(\Omega)}^q \le \exp\left(\frac{q^2-q}{2}\sum_{i=1}^N\int_0^T\sup_{x,\omega}\bar{\beta}_i(x,t,\omega)^2dt\right).$$

Hence

$$\|\gamma_M^{a,0}(T)\|_{L_q(\Omega)} \le \exp\left(\frac{q-1}{2}\sum_{i=1}^N \int_0^T \sup_{x,\omega} \bar{\beta}_i(x,t,\omega)^2 dt\right).$$

Further, (q-1)/q = 1/p and q-1 = q/p. Hence

$$\mathbf{E} \int_{D} p(x, T, \omega) dx \le \nu_2(q), \tag{4.20}$$

where

$$\nu_2(q) \stackrel{\Delta}{=} \nu^{1/p} \exp\left(\frac{1}{p} \left[\log \nu + \frac{q}{2} \sum_{i=1}^N \int_0^T \sup_{x,\omega} \bar{\beta}_i(x,t,\omega)^2 dt\right]\right).$$

By (3.8), there exists $q_0 > 1$ such that $\nu_2(q_0) < 1$.

Let $\nu_* \stackrel{\Delta}{=} \max(\nu_1, \nu_2(q_0))$. We have that $\nu_* \in (0, 1)$. By (4.19), (4.20), and by the linearity of problem (4.13), it follows that

$$\int_{D} \mathbf{E}p(x, T, \omega) dx \le \nu_* \int_{D} \rho(x) dx \tag{4.21}$$

for all non-negative $\rho(x)$.

Suppose that (4.17) holds for $\rho \in H^0$. Let

$$\rho_+(x) \stackrel{\Delta}{=} \max(0, \rho(x)), \quad \rho_-(x) \stackrel{\Delta}{=} \max(0, -\rho(x)).$$

Let p_+ and p_- be the solutions of (4.13) with s = 0 and with ρ replaced by ρ_{\pm} respectively. Let $\bar{p}_{\pm}(x,t) = \mathbf{E}p_{\pm}(x,t,\omega)$. By the definitions,

$$\bar{p}_+(\cdot,T) \stackrel{\Delta}{=} \mathbf{Q}\rho_+, \quad \bar{p}_-(\cdot,T) \stackrel{\Delta}{=} \mathbf{Q}\rho_-.$$

By (4.18), it follows that $\bar{p}_{\pm}(x,T) \geq 0$ for a.e. x. By (4.21), it follows that

$$\int_D \bar{p}_{\pm}(x,T)dx \le \nu_* \int_D \rho_{\pm}(x)dx.$$
(4.22)

Let us select any $\bar{\kappa} > 1$ such that $\bar{\kappa}\nu_* < 1$.

Let us assume first that $\rho_+ \neq 0$ and that $\kappa \in [0,1]$. It follows that there exist a measurable set $D_0 \subset D$ such that mes $(D_0) > 0$ and that $\rho(x) > 0$ and $\int_{D_0} \bar{p}_+(x,T) dx \leq \nu_* \int_{D_0} \rho(x) dx$ for all $x \in D_0$. It follows that $\int_{D_0} \bar{p}(x,T) dx = \int_{D_0} p_+(x,T) dx - \int_{D_0} p_-(x,t) dx \leq \nu_* \int_{D_0} \rho(x) dx$. Therefore, $\kappa \bar{p}(\cdot,T) \neq \rho(\cdot)$ in this case. Similarly, we can show that $\kappa \bar{p}(\cdot,T) \neq \rho$ if $\rho_- \neq 0$ and $\kappa \in [0,1]$.

Further, let us assume that $\kappa \in [-\bar{\kappa}, 0)$. Let $D_+ = \{x : \rho(x) \ge 0\}, D_- = \{x : \rho(x) < 0\}$. By the assumptions,

$$\int_{D_{+}} \rho(x) dx = \kappa \int_{D_{+}} \bar{p}(x,T) dx > 0, \qquad \int_{D_{-}} \rho(x) dx = \kappa \int_{D_{-}} \bar{p}(x,T) dx < 0.$$

We have that

$$0 \le -\int_{D_+} \bar{p}(x,T) dx \le -\nu_* \int_{D_-} \rho(x) dx, \qquad 0 \le \int_{D_-} \bar{p}(x,T) dx \le \nu_* \int_{D_+} \rho(x) dx.$$

Hence

$$-\int_{D_+} \bar{p}(x,T)dx \le -\nu_*\kappa \int_{D_-} \bar{p}(x,T)dx, \qquad \int_{D_-} \bar{p}(x,T)dx \le \nu_*\kappa \int_{D_+} \bar{p}(x,T)dx.$$

Hence

$$\int_{D_+} \bar{p}(x,T)dx \ge \nu_*\kappa \int_{D_-} \bar{p}(x,T)dx, \qquad \int_{D_-} \bar{p}(x,T)dx \le \nu_*\kappa \int_{D_+} \bar{p}(x,T)dx. \quad (4.23)$$

Hence

$$\left| \int_{D_+} \bar{p}(x,T) dx \right| \le |\nu_*\kappa| \left| \int_{D_-} \bar{p}(x,T) dx \right|, \qquad \left| \int_{D_-} \bar{p}(x,T) dx \right| \le |\nu_*\kappa| \left| \int_{D_+} \bar{p}(x,T) dx \right|.$$

The system of the last two inequalities can be satisfied only if the integrals there are zero. This means that $\rho = 0$.

We have proved that if (4.17) holds for $\rho \in H^0$ then $\rho = 0$. We had proved also that the operator **Q** is compact. By the Fredholm Theorem, it follows that the equation $\rho - \mathbf{Q}\rho = z$ is solvable in H^0 for any $z \in H^0$. By (4.15), it follows that $\Phi = 0$. Therefore, the condition $\kappa u(\cdot, 0) = u(\cdot, T)$ fails to be satisfied for $u \neq 0$, $\xi = 0$, and $\varphi = 0$. Thus, u = 0 is the unique solution of problem (2.1)-(2.3) for $\xi = 0$ and $\varphi = 0$. Then the proof of Theorem 3.5 follows from Theorem 3.3. \Box

Acknowledgment

This work was supported by ARC grant of Australia DP120100928 to the author.

References

Alós, E., León, J.A., Nualart, D. (1999). Stochastic heat equation with random coefficients, Probability Theory and Related Fields **115** (1), 41–94.

Arnold, L., and Tudor, C. (1998). Stationary and Almost Periodic Solutions of Almost Periodic Affine Stochastic Differential Equations, *Stochastics and Stochastic Reports* **64**, 177– 193.

Bally, V., Gyongy, I., Pardoux, E. (1994). White noise driven parabolic SPDEs with measurable drift. *Journal of Functional Analysis* **120**, 484–510. Bedouhene, F., Mellah, O., Raynaud de Fitte. P. (2012). Bochner-Almost Periodicity for Stochastic Processes. *Stochastic Analysis and Applications*, vol. 30, no. 2, pp. 322-342.

Bezandry, P. and Diagana, T. (2007). Square-mean almost periodic solutions nonautonomous stochastic differential equations. *Electron. J. Diff. Equ.* Vol. 2007, No. 117, pp. 1-10.

Caraballo, T., P.E. Kloeden, P.E., Schmalfuss, B. (2004). Exponentially stable stationary solutions for stochastic evolution equations and their perturbation, *Appl. Math. Optim.*, **50**, 183–207.

Chojnowska-Michalik, A. (1987). On processes of Ornstein-Uhlenbeck type in Hilbert space. Stochastics **21**, 251–286.

Chojnowska-Michalik, A. (1990). Periodic distributions for linear equations with general additive noise, *Bull. Pol. Acad. Sci. Math.* **38** (112) 23–33.

Chojnowska-Michalik, A., and Goldys, B. (1995). Existence, uniqueness and invariant measures for stochastic semilinear equations in Hilbert spaces, *Probability Theory and Related Fields*, **102**, No. 3, 331–356.

Crewe, P. (2013). Almost periodic solutions to stochastic evolution equations on Banach spaces. *Stoch. Dyn.* **13**, 1250027, 23 pages.

Da Prato, G., and Tudor, C. (1995). Periodic and Almost Periodic Solutions for Semilinear Stochastic Evolution Equations, *Stoch. Anal. Appl.* 13 (1), 1333.

Da Prato, G., and Tubaro, L. (1996). Fully nonlinear stochastic partial differential equations, *SIAM Journal on Mathematical Analysis* **27**, No. 1, 40–55.

Dokuchaev, N.G. (1992). Boundary value problems for functionals of Ito processes, *Theory* of Probability and its Applications **36** (3), 459-476.

Dokuchaev, N.G. (2004). Estimates for distances between first exit times via parabolic equations in unbounded cylinders. *Probability Theory and Related Fields*, **129**, 290 - 314.

Dokuchaev, N.G. (2005). Parabolic Ito equations and second fundamental inequality. *Stochastics* **77** (2005), iss. 4., pp. 349-370.

Dokuchaev N. (2008). Parabolic Ito equations with mixed in time conditions. *Stochastic Analysis and Applications* **26**, Iss. 3, 562–576.

Dokuchaev, N. (2010). Duality and semi-group property for backward parabolic Ito equations. *Random Operators and Stochastic Equations.* **18**, 51-72.

Dokuchaev, N. (2011). Representation of functionals of Ito processes in bounded domains. Stochastics 83, No. 1, 45–66.

Dokuchaev, N. (2012a). Backward parabolic Ito equations and second fundamental inequality. *Random Operators and Stochastic Equations* **20**, iss. 1, 69-102. Dokuchaev, N. (2012b). Backward SPDEs with non-local in time and space boundary conditions. Working paper, arXiv:1211.1460 (submitted).

Dokuchaev, N. (2012c). On forward and backward SPDEs with non-local boundary conditions. Working paper in arXiv (submitted).

Du K., and Tang, S. (2012). Strong solution of backward stochastic partial differential equations in C^2 domains. *Probability Theory and Related Fields*) **154**, 255–285.

Dorogovtsev, A.Ya., and Ortega, O.A. (1988). On the Existence of Periodic Solutions of a Stochastic Equation in a Hilbert Space. *Visnik Kiiv. Univ. Ser. Mat. Mekh.* **115** No. 30, 21–30.

Duan J., Lu K., Schmalfuss B. (2003). Invariant manifolds for stochastic partial differential equations. Ann. Probab. **31** 21092135.

Feng C., Zhao H. (2012). Random periodic solutions of SPDEs via integral equations and Wiener-Sobolev compact embedding. *Journal of Functional Analysis* **262**, 4377–4422.

Fife, P.C. (1964). Solutions of parabolic boundary problems existing for all time. Archive for Rational Mechanics and Analysis 16 155186.

Gyöngy, I. (1998). Existence and uniqueness results for semilinear stochastic partial differential equations. *Stochastic Processes and their Applications* **73** (2), 271–299.

Hess, P. (1991). Periodic Parabolic Boundary Value Problems and Positivity. Wiley, New York.

Klünger, M. (2001). Periodicity and Sharkovskys theorem for random dynamical systems, Stochastic and Dynamics 1, iss.3, 299–338.

Krylov, N. V. (1999). An analytic approach to SPDEs. Stochastic partial differential equations: six perspectives, 185–242, Mathematical Surveys and Monographs, 64, AMS., Providence, RI, pp.185-242.

Ladyzhenskaia, O.A. (1985). The Boundary Value Problems of Mathematical Physics. New York: Springer-Verlag.

Lieberman G.M. (1999). Time-periodic solutions of linear parabolic differential equations. Comm. Partial Differential Equations 24, 631663.

Liu, Y., Zhao, H.Z (2009). Representation of pathwise stationary solutions of stochastic Burgers equations, *Stochastics and Dynamics* **9** (4), 613–634.

Maslowski, B. (1995). Stability of semilinear equations with boundary and pointwise noise, Annali della Scuola Normale Superiore di Pisa - Classe di Scienze (4), **22**, No. 1, 55–93.

Mattingly. J. (1999). Ergodicity of 2D NavierStokes equations with random forcing and large viscosity. *Comm. Math. Phys.* 206 (2), 273288.

Mellah, O., and Raynaud de Fitte, P. (2007). Counterexamples to mean square almost periodicity of the solutions of some SDEs with almost periodic coefficients. *Electron. J. Differential Equations*, No. 117, 10 pp. (electronic).

Mohammed S.-E.A., Zhang T., Zhao H.Z. (2008). The stable manifold theorem for semilinear stochastic evolution equations and stochastic partial differential equations. *Mem. Amer. Math. Soc.* 196 (917) 1105.

Nakao, M. (1984). Periodic solution of some nonlinear degenerate parabolic equations. *Journal of Mathematical Analysis and Applications* **104** (2), 554–557.

Pardoux, E. (1993). Bulletin des Sciences Mathematiques, 2e Serie, 117, 29-47.

Revuz, D., and Yor, M. (1999). *Continuous Martingales and Brownian Motion*. Springer-Verlag: New York.

Rodkina, A.E. (1992). On solutions of stochastic equations with almost surely periodic trajectories. *Differ. Uravn.* 28, No.3, 534–536 (in Russian).

Rozovskii, B.L. (1990). Stochastic Evolution Systems; Linear Theory and Applications to Non-Linear Filtering. Kluwer Academic Publishers. Dordrecht-Boston-London.

Sinai, Ya. (1996). Burgers system driven by a periodic stochastic flows, in: Ito's Stochastic Calculus and Probability Theory, Springer, Tokyo, 1996, pp. 347353.

Shelukhin, V.V. (1993). Variation principle for non-local in time problems for linear evolutionary equations. *Siberian Math. J.* **34** (2), 191–207.

Tudor, C. (1992). Almost periodic solutions of affine stochastic evolutions equations. *Stochastics and Stochastics Reports* 38, 251–266.

Vejvoda, O. (1982). Partial Differential Equations: Time Periodic Solutions, Martinus Nijhoff, The Hague.

Walsh, J.B. (1986). An introduction to stochastic partial differential equations, Lecture Notes in Mathematics **1180**, Springer Verlag.

Yong, J., and Zhou, X.Y. (1999). Stochastic controls: Hamiltonian systems and HJB equations. New York: Springer-Verlag.

Zhou, X.Y. (1992). A duality analysis on stochastic partial differential equations, *Journal* of Functional Analysis **103**, No. 2, 275–293.