# On Singular Control Problems with State Constraints and Regime-Switching: A Viscosity Solution Approach

Qingshuo Song<sup>\*</sup> Chao Zhu<sup>†</sup>

September 17, 2018

#### Abstract

This paper investigates a singular stochastic control problem for a multi-dimensional regime-switching diffusion process confined in an unbounded domain. The objective is to maximize the total expected discounted rewards from exerting the singular control. Such a formulation stems from application areas such as optimal harvesting multiple species and optimal dividends payments schemes in random environments. With the aid of weak dynamic programming principle, we characterize the value function to be the unique constrained viscosity solution of a certain system of coupled nonlinear quasi-variational inequalities. Several examples are analyzed in details to demonstrate the main results.

Key words. constrained viscosity solution, regime-switching diffusion, singular stochastic control, weak dynamic programming principle, quasi-variational inequality.

AMS subject classification. 93E20, 60J60

# 1 Introduction

This paper is concerned with a class of singular stochastic control problems with state constraints. The controlled regime-switching diffusion process X and the singular control process Z take values in a convex cone  $S \subset \mathbb{R}^n$ . The control problem has the state process

$$X(t) = x + \int_0^t b(X(s), \alpha(s))ds + \int_0^t \sigma(X(s), \alpha(s))dW(s) - Z(t)dt$$

where W is a d-dimensional standard Brownian motion,  $\alpha$  is a continuous-time Markov chain with a finite state space  $\mathcal{M} = \{1, \ldots, m\}, Z = (Z_1, \ldots, Z_n)'$  is an n-dimensional adapted,

<sup>\*</sup>Department of Mathematics, City University of Hong Kong, 83 Tat Chee Avenue, Kowloon Tong, Hong Kong, song.qingshuo@cityu.edu.hk.

<sup>&</sup>lt;sup>†</sup>Department of Mathematical Sciences, University of Wisconsin-Milwaukee, Milwaukee, WI 53201, USA, zhu@uwm.edu.

nondecreasing, and càdlàg stochastic process, and  $b, \sigma$  are appropriate measurable functions. The income rates  $f_i$ , i = 1, ..., n, from exerting the singular control are allowed to be stateand regime-dependent. The objective is to maximize the total discounted reward

$$\mathbb{E}\left[\int_0^\infty e^{-rs} \sum_{i=1}^n f_i(X(s), \alpha(s)) dZ_i(s)\right],\tag{1.1}$$

where r > 0 is the discounting factor.

Such singular control problems (in various different settings) have been extensively studied in the literature. A partial list includes the monotone follower problems (Karatzas and Shreve (1984)), optimal harvesting problems (Alvarez and Shepp (1998), Song et al. (2011)), optimal dividend distribution schemes (Paulsen (2003)), portfolio selection management with transaction costs (Øksendal and Sulem (2002)), optimal partially reversible investment problem (Guo and Pham (2005)), and heavy traffic modeling and control problems (Lee and Weerasinghe (2011), Wein (1990)), etc. See also Haussmann and Suo (1995a,b) for a general singular stochastic control problem for a multidimensional Itô diffusion on a fixed time horizon, in which the existence of the optimal control and the characterization of the value function as the unique viscosity solution of a Hamilton-Jacobi-Bellman equation are established. Singular control problems with state constraints have drawn considerable interests in recent years; see, for example, Atar and Budhiraja (2006), Atar et al. (2007), Zariphopoulou (1992), among others.

Note that most, if not all, of the aforementioned literature on singular stochastic controls deal with Itô (jump) diffusions. One exception is our recent work Song et al. (2011), which studies an optimal harvesting problem of a single species living in random environments. Due to their capability of modeling complex systems with uncertainty, regime-switching models have drawn considerable attention from both researchers and practitioners in recent decades in a wide range of applications. Some of such examples can be found in mathematical finance (Zhang (2001)), ecosystem modeling (Slatkin (1978), Zhu and Yin (2009)), stochastic manufacturing systems (Sethi and Zhang (1994)), risk management (Elliott and Siu (2010), Zhu (2011)), to name just a few. In these systems, both continuous dynamics and discrete events coexist. In particular, the systems often display qualitative structural changes. Regime-switching models turn out to be quite versatile in capturing these inherent randomness. We refer to Mao and Yuan (2006) and Yin and Zhu (2010) for in-depth investigations of regime-switching diffusions.

This work aims to investigate the singular control problem (1.1) in the setting of multidimensional regime-switching diffusion with state constraints. First we recall the notion of constrained viscosity solution, illustrated by several simple yet nontrivial examples. Then we use the weak dynamic programming principle to show that the value function defined in (2.6) is a constrained viscosity solution to the coupled system of quasi-variational inequalities (2.11) in Theorem 4.5. Finally, we derive a strong comparison result in Theorem 5.3, from which we establish the uniqueness of the constrained viscosity solution to (2.11). Compared with the classical work on viscosity solution such as Crandall et al. (1992), Yong and Zhou (1999) and others, the novelty and contribution of this work can be summarized as follows. In lieu of a single differential equation studied in the literature, this work deals with a coupled system of nonlinear second-order differential equations with gradient constraints. The coupling effect is due to the presence of random environments or regime switching. This feature at one hand makes our model more appealing in real-world applications since it can naturally capture the qualitative structural changes of the systems; on the other hand, it adds much difficulty in the analysis. In particular, the function F defined in 2.10 is not proper with respect to the variable  $\xi$  in the sense of the User's Guide Crandall et al. (1992). Note that the properness was an essential assumption in the proof of strong comparison result in Crandall et al. (1992). Here we need to carefully handle the coupling effect; see the proof of Theorem (5.3) for more details. Another noteworthy feature of this work is that we introduce an exponential transformation which allows us to handle both the gradient constraints as well as the polynomial growth condition on an unbounded domain for the solution of the coupled system of quasi-variational inequalities (2.11).

The rest of the paper is arranged as follows. Section 2 presents the precise formulation of the problem, followed by some preliminary results in Section 3. We recall the notion of constrained viscosity solution in Section 4, followed by several examples for illustration. Further, in Section 4, we establish the existence by showing that the value function V defined in (2.6) is a constrained viscosity solution of (2.11). The strong comparison result is arranged in Section 5. A hierarchical PDE characterization of the boundary behavior is arranged in Section 5 as well. The paper is concluded with conclusions and remarks in Section 6.

To facilitate later presentation, we introduce some notations that will be used often in later sections. We say that a function from  $[0, \infty)$  to some Polish space E is càdlàg if it is right continuous and has left limits in E on  $[0, \infty)$ . When  $E = \mathbb{R}^n$  and  $\xi$  is càdlàg, then we write  $\Delta \xi(t) = \xi(t) - \xi(t-)$  for t > 0. As a convention, we set  $\Delta \xi(0) = \xi(0)$ . Throughout the paper, we use x'y or  $x \cdot y$  interchangably to denote the inner product of vectors x and y. For any vectors  $x, y \in \mathbb{R}^n$ ,  $x \leq y$  means  $x_i \leq y_i$  for every  $i = 1, 2, \ldots, n$ . The space of  $n \times n$ symmetric matrices is denoted by  $S_n$  and the family of positive definite symmetric matrices is denoted by  $S_n^+$ . If  $A, B \in S_n$  and  $A - B \in S_n^+$ , then we write A > B. If  $\phi : \mathbb{R}^n \to \mathbb{R}$ is sufficiently smooth, then  $D_{x_i}\phi = \frac{\partial \phi}{\partial x_i}$ ,  $D_{x_ix_j}\phi = \frac{\partial^2 \phi}{\partial x_i \partial x_j}$ , and  $D\phi = (D_{x_1}\phi, \ldots, D_{x_n}\phi)'$  is the gradient of  $\phi$  while  $D^2\phi = (D_{x_ix_j}\phi)$  denotes the Heissian of  $\phi$ . For any real-valued function f, we use  $f_*$  and  $f^*$  to denote the lower- and upper-semicontinuous envelopes of f, respectively. If B is a set, we use  $B^o$  and  $I_B$  to denote the interior and indicator function of B, respectively. Throughout the paper, we adopt the conventions that  $\sup \emptyset = -\infty$  and  $\inf \emptyset = +\infty$ .

### 2 Formulation

We consider singular control problems for a regime-switching diffusion

$$d\zeta(t) = b(\zeta(t), \alpha(t))dt + \sigma(\zeta(t), \alpha(t))dW(t), \quad \zeta(0) = x, \ \alpha(0) = \alpha, \tag{2.1}$$

where  $x \in \mathbb{R}^n, \alpha \in \mathcal{M} = \{1, \ldots, m\}$ , W is a d-dimensional standard Brownian motion,  $b : \mathbb{R}^n \times \mathcal{M} \mapsto \mathbb{R}^n, \sigma : \mathbb{R}^n \times \mathcal{M} \mapsto \mathbb{R}^{n \times d}$ , and  $\alpha(\cdot) \in \mathcal{M}$  is a continuous-time Markov chain that is independent of the Brownian motion W and is generated by  $Q = (q_{ij}) \in \mathbb{R}^{m \times m}$ :

$$\mathbb{P}\left\{\alpha(t+\Delta t)=j|\alpha(t)=i,\alpha(s),s\leq t\right\}=\begin{cases}q_{ij}\Delta t+o(\Delta t), & \text{if } j\neq i\\1+q_{ii}\Delta t+o(\Delta t), & \text{if } j=i,\end{cases}$$
(2.2)

where  $q_{ij} \ge 0$  for i, j = 1, ..., m with  $j \ne i$  and  $q_{ii} = -\sum_{j \ne i} q_{ij} < 0$  for each i = 1, ..., m.

Throughout the paper, we assume that the coefficients b and  $\sigma$  and the generator Q are such that for any initial condition  $(x, \alpha) \in \mathbb{R}^n \times \mathcal{M}$ , the solution  $\zeta^{x,\alpha}$  to (2.1) exists and is weakly unique. Sufficient condition for existence and uniqueness for stochastic differential equations with regime switching can be found in, for example, Mao and Yuan (2006), Yin and Zhu (2010).

We now introduce singular control into (2.1) with state constraint and suppose that the controlled dynamic is given by

$$dX(t) = b(X(t), \alpha(t))dt + \sigma(X(t), \alpha(t))dW(t) - dZ(t), \qquad (2.3)$$

with initial conditions

$$X(0-) = x \in S, \quad \alpha(0) = \alpha \in \mathcal{M}, \tag{2.4}$$

where  $Z \in \mathbb{R}^n$  is a singular control process to be specified below. Without loss of generality, we take  $S = \mathbb{R}^n_+ = \{x \in \mathbb{R}^n : x_i > 0, i = 1, ..., n\}$ . Note that X(0) may not be equal to X(0-) due to an instantaneous push Z(0) at time 0. Denote the solution to (2.3) with initial condition specified by (2.4) by  $X^{x,\alpha}(\cdot)$ .

Let  $\mathcal{A}_{x,\alpha}$  denote the collection of all *admissible controls* with initial conditions given by (2.4), where  $Z \in \mathcal{A}_{x,\alpha}$  satisfies

- (i) for each  $i = 1, ..., n, Z_i(t)$  is nonnegative, càdlàg and nondecreasing with respect to t,
- (ii)  $X(t) \in S$  for all  $t \ge 0$ , and

(iii) Z(t) is adapted to  $\mathfrak{F}_t := \sigma \{ W(s), \alpha(s), 0 \le s \le t \}$ , where  $\mathfrak{F}_0$  contains all  $\mathbb{P}$ -null sets. Moreover,

$$\mathbb{E}\int_{0}^{\infty}e^{-rs}d\left|Z\right|(s)<\infty.$$

Note that the state constraint is specified in condition (ii) above. Throughout the paper, we assume  $\mathcal{A}_{x,\alpha} \neq \emptyset$  for every  $(x,\alpha) \in S \times \mathcal{M}$ ; see Section 3 for a sufficient condition. For a fixed  $Z \in \mathcal{A}_{x,\alpha}$ , the discounted payoff is

$$J(x,\alpha,Z) := \mathbb{E} \int_0^\infty e^{-rs} f(X^{x,\alpha}(s-),\alpha(s-)) \cdot dZ(s), \qquad (2.5)$$

where  $f: S \times \mathcal{M} \mapsto \mathbb{R}^n$  with  $f_i$  representing the state- and regime-dependent instantaneous marginal yields accrued from exerting the singular control  $Z_i(t)$ . Assume  $f_i$  is continuous and non-increasing with respect to x in the sense that  $f_i(x, \alpha) \ge f_i(y, \alpha)$  for each  $\alpha \in \mathcal{M}$  if  $x \le y$ , where  $x = (x_1, \ldots, x_n)'$  and  $y = (y_1, \ldots, y_n)'$  satisfy  $x_j \le y_j$  for each  $j = 1, \ldots, n$ . Moreover, we assume  $0 < f_i(0, \alpha) < \infty$  for each  $i = 1, \ldots, n$  and  $\alpha \in \mathcal{M}$ . Such assumptions on f are motivated by considerations in optimal harvesting problems (Alvarez (2000) and Song et al. (2011)). The goal is to maximize the expected total discounted payoff and find an optimal control  $Z^*$ :

$$V(x,\alpha) = J(x,\alpha,Z^*) := \sup_{Z \in \mathcal{A}_{x,\alpha}} J(x,\alpha,Z).$$
(2.6)

In order to work with a well-formulated maximization problem, we assume throughout the paper that  $V(x, \alpha) < \infty$  for all  $(x, \alpha) \in \mathbb{R}^n_+ \times \mathcal{M}$ .

As usual, we shall rely on the dynamic programming principle (DPP) to deduce the behavior of the value function

$$V(x,\alpha) = \sup_{Z(\cdot)\in\mathcal{A}_{x,\alpha}} \mathbb{E}\bigg[\int_0^\eta e^{-rs} f(X^{x,\alpha}(s-),\alpha(s-)) \cdot dZ(s) + e^{-r\eta} V(X^{x,\alpha}(\eta),\alpha(\eta))\bigg]$$
(2.7)

for every  $(x, \alpha) \in S \times \mathcal{M}$  and stopping time  $\eta$ . A heuristic argument using the DPP (2.7) yields that V satisfies the following coupled system of quasi-variational inequalities

$$\min\left\{(r-\mathcal{L})V(x,\alpha),\min_{i=1,\dots,n}\left\{D_{x_i}V(x,\alpha)-f_i(x,\alpha)\right\}\right\}=0, \quad (x,\alpha)\in S\times\mathcal{M},$$
(2.8)

where for any  $h(\cdot, \alpha) \in C^2$ ,  $\alpha \in \mathcal{M}$ , we define

$$\mathcal{L}h(x,\alpha) = \frac{1}{2} \operatorname{tr}(\sigma\sigma'(x,\alpha)D^2h(x,\alpha)) + b(x,\alpha) \cdot Dh(x,\alpha) + \sum_{j=1}^m q_{\alpha j}h(x,j).$$
(2.9)

However, without a priori result on the continuity of the value function, a rigorous proof of (2.7) is nontrivial. Thanks to the state constraint as well as the generality of the set up

of the problem, it seems not easy to obtain the continuity of the value function V defined in (2.6). Also, in the current singular control setup with regime-switching diffusion, it appears that the DPP is not available from the literature. To overcome this difficulty, we will instead invoke the weak DPP (Bouchard and Touzi (2011)); see Section 3 for the precise statement. Also, the value function V is not necessarily sufficiently smooth to take first and second order partial derivatives. Therefore we aim to show in this work that V satisfies (2.8) in the weak sense using the notion of viscosity solution. We will show that the value function V is the unique viscosity solution to (2.8).

For convenience of later presentations, we define for any  $(x, \alpha, \xi, p, A) \in \mathbb{R}^n \times \mathcal{M} \times \mathbb{R}^m \times \mathbb{R}^n \times \mathcal{S}_n$ ,

$$F_{\alpha}(x,\xi,p,A) = F(x,\alpha,\xi,p,A) := r\xi_{\alpha} - \frac{1}{2} \operatorname{tr}(\sigma\sigma'(x,\alpha)A) - b(x,\alpha) \cdot p - \sum_{j=1}^{m} q_{\alpha j}\xi_{j}.$$
 (2.10)

Set  $\mathbf{V}(x) = (V(x, 1), \dots, V(x, m))' \in \mathbb{R}^m$ , then (2.8) can be rewritten as  $\min \left\{ F(x, \mathbf{V}(x), DV(x, \alpha), D^2 V(x, \alpha)) \right\} = 0$ 

$$\min\left\{F_{\alpha}(x, \mathbf{V}(x), DV(x, \alpha), D^{2}V(x, \alpha)), \min_{i=1,\dots,n}\left\{D_{x_{i}}V(x, \alpha) - f_{i}(x, \alpha)\right\}\right\} = 0, \quad (2.11)$$

for all  $x \in S$  and each  $\alpha = 1, \ldots, m$ .

As we indicated earlier, (2.11) is a coupled system of quasi-variational inequalities. Moreover, thanks to the term  $\sum_{j=1}^{m} q_{\alpha j} \xi_j$  with  $Q = (q_{ij})$  defined in (2.2), for each  $\alpha \in \mathcal{M}$ ,  $F_{\alpha}$  is not proper with respect to the variable  $\xi$  in the sense of equations (0.1) or (0.2) in the User's Guide Crandall et al. (1992). Note that properness assumption (and in particular equation (3.13) in Crandall et al. (1992)) enabled them to derive the strong comparison result and hence the uniqueness of the viscosity solution. Here for our analysis, special care has to be given to handle the fact that  $F_{\alpha}$  is not proper due to the coupling term. Also, instead of working on a bounded domain, we are dealing with unbounded domain S. These features make our analysis much more involved than the classical comparison result in Crandall et al. (1992).

# 3 Some Preliminary Results

We present some preliminary results in this section. The first one provides a sufficient condition for the assumption that  $\mathcal{A}_{x,\alpha} \neq \emptyset$  for all  $(x, \alpha) \in S \times \mathcal{M}$ .

**Proposition 3.1.** Assume there exists a function  $\Psi: S \times \mathcal{M} \mapsto \mathbb{R}_+$  satisfying

(i) for each  $i = 1, \ldots, n$  and  $\alpha \in \mathcal{M}$ ,

$$\lim_{|x|\to\infty}\Psi(x,\alpha)=\infty, \ and \ \lim_{x_i\downarrow 0}\Psi(x,\alpha)=\infty,$$

(ii)  $\Psi(\cdot, \alpha) \in C^2$  for each  $\alpha \in \mathcal{M}$  and

$$\mathcal{L}\Psi(x,\alpha) = b(x,\alpha) \cdot D\Psi(x,\alpha) + \frac{1}{2} \operatorname{tr}(\sigma\sigma'(x,\alpha)D^2\Psi(x,\alpha)) + \sum_{j=1}^m q_{\alpha j}\Psi(x,j) \le 0,$$

for all  $(x, \alpha) \in S \times M$ . Then, denoting by  $\zeta^{x, \alpha}$  the solution of (2.1) with initial condition  $(x, \alpha)$ , we have

$$\mathbb{P}\left\{\zeta^{x,\alpha}(t)\in S \text{ for all } t\geq 0\right\}=1, \quad \text{for any } (x,\alpha)\in S\times\mathcal{M}.$$
(3.1)

Consequently  $\mathcal{A}_{x,\alpha} \neq \emptyset$  for all  $(x, \alpha) \in S \times \mathcal{M}$ .

*Proof.* It suffices to prove (3.1), which leads to  $Z \equiv 0 \in \mathcal{A}_{x,\alpha}$  for any  $(x, \alpha) \in S \times \mathcal{M}$ . To this end, we consider  $(x, \alpha) \in S \times \mathcal{M}$  and define

$$\tau_k := \inf \left\{ t \ge 0 : |\zeta^{x,\alpha}(t)| \ge k \text{ or } \zeta_i^{x,\alpha}(t) \le \frac{1}{k} \text{ for some } i = 1, \dots, n \right\}.$$

Note that  $\{\tau_k\}$  is a nondecreasing sequence of  $\mathfrak{F}_t$ -stopping times. Now it is enough to show that  $\tau_k \to \infty$  with probability 1. Suppose on the contrary that

$$\liminf_{k \to \infty} \mathbb{P}\left\{\tau_k < \infty\right\} = \delta > 0. \tag{3.2}$$

Applying generalized Itô's formula to the function  $\Psi$  and using condition (ii), we obtain that for any  $t \ge 0$ ,

$$\mathbb{E}\left[\Psi(\zeta(\tau_k \wedge t), \alpha(\tau_k \wedge t)\right] = \Psi(x, \alpha) + \mathbb{E}\left[\int_0^{\tau_k \wedge t} \mathcal{L}\Psi(\zeta(s), \alpha(s))ds\right]$$
  
$$\leq \Psi(x, \alpha).$$

Then since  $\Psi \ge 0$ , it follows from condition (i) and (3.2) that

$$\Psi(x,\alpha) \ge \mathbb{E}\left[\Psi(\zeta(\tau_k \wedge t), \alpha(\tau_k \wedge t))\right] \ge \mathbb{E}\left[\Psi(\zeta(\tau_k), \alpha(\tau_k)I_{\{\tau_k \le t\}}\right]$$
$$\ge \Psi_k \mathbb{P}\left\{\tau_k \le t\right\} \to \infty, \text{ as } k \to \infty,$$

where

$$\Psi_k := \inf \left\{ \Psi(x, j), |x| = k \text{ or } x_i = \frac{1}{k} \text{ for some } i = 1, \dots, n, \text{ and } j \in \mathcal{M} \right\}.$$

This is a contradiction and hence  $\tau_k \to \infty$  with probability 1 as  $k \to \infty$ .

We will need the following proposition in the proof of Theorem 4.5.

**Proposition 3.2.** For each  $\alpha \in \mathcal{M}$  and any  $x, y \in S$  with  $y \leq x$ , we have

$$V(x,\alpha) \ge f(x,\alpha) \cdot (x-y) + V(y,\alpha), \tag{3.3}$$

$$V^*(x,\alpha) \ge f(x,\alpha) \cdot (x-y) + V^*(y,\alpha). \tag{3.4}$$

*Proof.* Equation (3.3) can be established using exactly the same arguments as those in Song et al. (2011), while (3.4) follows from (3.3) directly.  $\Box$ 

The next proposition can be established using similar arguments as those in Bouchard and Touzi (2011).

**Proposition 3.3.** Fix  $(x, \alpha) \in \mathbb{R}^n_+ \times \mathcal{M}$ . Then for any stopping time  $\tau$ , we have

$$V(x,\alpha) \le \sup_{Z \in \mathcal{A}_{x,\alpha}} \mathbb{E}\left[\int_0^\tau e^{-rs} f(X^{x,\alpha}(s-),\alpha(s-) \cdot dZ(s) + e^{-r\tau} V^*(X^{x,\alpha}(\tau),\alpha(\tau))\right], \quad (3.5)$$

and

$$V(x,\alpha) \ge \sup_{Z \in \mathcal{A}_{x,\alpha}} \mathbb{E}\left[\int_0^\tau e^{-rs} f(X^{x,\alpha}(s-),\alpha(s-) \cdot dZ(s) + e^{-r\tau}\varphi(X^{x,\alpha}(\tau),\alpha(\tau))\right], \quad (3.6)$$

for all upper-semicontinuous functions  $\varphi$  such that  $V \geq \varphi$  on  $\mathbb{R}^n_+ \times \mathcal{M}$ .

We finish this section with the verification theorem, whose proof is similar to those in Song et al. (2011).

**Theorem 3.4.** Suppose there exists a function  $\phi : S \times \mathcal{M} \mapsto \mathbb{R}_+$  solving (2.8).

- (a) Then  $\phi(x, \alpha) \ge V(x, \alpha)$  for every  $(x, \alpha) \in S \times \mathcal{M}$ .
- (b) *Define the* non-intervention region

$$\mathcal{C} = \left\{ (x, \alpha) \in S \times \mathcal{M} : \max_{i=1,\dots,n} \left\{ f_i(x, \alpha) - D_{x_i} \phi(x, \alpha) \right\} < 0 \right\}.$$

Assume there exists a strategy  $Z \in \mathcal{A}_{x,\alpha}$  such that

$$(X(t), \alpha(t)) \in \mathcal{C} \text{ for Lebesgue almost all } 0 \le t < \infty,$$
(3.7)

$$\int_0^t (D\phi(X(s), \alpha(s)) - f(X(s), \alpha(s))) \cdot d\widetilde{Z}^c(s) = 0, \text{ for any } 0 \le t < \infty,$$
(3.8)

$$\lim_{N \to \infty} \mathbb{E}\left[e^{-r(\tau \wedge N \wedge \beta_N)}\phi(X(\tau \wedge N \wedge \beta_N), \alpha(\tau \wedge N \wedge \beta_N))\right] = 0,$$
(3.9)

and that if  $X(s) \neq X(s-)$ , then

$$\phi(X(s),\alpha(s-)) - \phi(X(s-),\alpha(s-)) = -f(\widetilde{X}(s-),\alpha(s-)) \cdot \Delta \widetilde{Z}(s), \qquad (3.10)$$

where  $\beta_N := \inf\{t \ge 0 : |X(t)| \ge N\}$ , and  $X = X^{x,\alpha}$  denotes the solution of (2.3). Then  $\phi(x, \alpha) = V(x, \alpha)$  for every  $(x, \alpha) \in S \times \mathcal{M}$  and  $\widetilde{Z}$  is an optimal strategy.

### 4 Viscosity Solution: Existence

This section is devoted to the properties of the value function V. In particular, we aim to characterize V as a viscosity solution to the quasi variational inequality (2.11). Let's first recall the notion of viscosity solution.

**Definition 4.1.** A function  $\mathbf{u}(x) = (u(x, 1), \dots, u(x, m))'$  is said to be a viscosity subsolution of (2.11) on  $\bar{S} \times \mathcal{M}$ , if for any  $(x_0, \alpha_0) \in \bar{S} \times \mathcal{M}$  and functions  $\varphi(\cdot, \alpha) \in C^2(S), \alpha \in \mathcal{M}$ satisfying  $(u^* - \varphi)(x, \alpha) \leq (u^* - \varphi)(x_0, \alpha_0) = 0$  for all  $(x, \alpha) \in \bar{S} \times \mathcal{M}$ , we have

$$\min\left\{F_{\alpha_0}(x_0, \mathbf{u}^*(x_0), D\varphi(x_0, \alpha_0), D^2\varphi(x_0, \alpha_0)), \min_{i=1,\dots,n}\left\{D_{x_i}\varphi(x_0, \alpha_0) - f_i(x_0, \alpha_0)\right\}\right\} \le 0.$$

Similarly, a function  $\mathbf{u}(x) = (u(x,1), \dots, u(x,m))'$  is said to be a viscosity supersolution of (2.11) in  $S \times \mathcal{M}$ , if for any  $(x_0, \alpha_0) \in S \times \mathcal{M}$  and functions  $\varphi(\cdot, \alpha) \in C^2(S), \alpha \in \mathcal{M}$  satisfying  $(u_* - \varphi)(x, \alpha) \ge (u_* - \varphi)(x_0, \alpha_0) = 0$  for all  $(x, \alpha) \in S \times \mathcal{M}$ , we have

$$\min\left\{F_{\alpha_0}(x_0, \mathbf{u}_*(x_0), D\varphi(x_0, \alpha_0), D^2\varphi(x_0, \alpha_0)), \min_{i=1,\dots,n}\left\{D_{x_i}\varphi(x_0, \alpha_0) - f_i(x_0, \alpha_0)\right\}\right\} \ge 0.$$

The function u is said to be a *constrained viscosity solution*, if it is both a viscosity subsolution in  $\overline{S} \times \mathcal{M}$  and a viscosity supersolution in  $S \times \mathcal{M}$ .

Before presenting the main result of this section, we shall first study several examples to illustrate Definition 4.1. These examples will also help us to motivate later results.

**Example 4.2.** Consider the QVI

$$\min \{u(x) - u'(x), u'(x) - 1\} = 0, \quad x \in (0, \infty).$$
(4.1)

We claim that  $u(x) = Ke^x$ ,  $K \ge 1$  is a constrained viscosity solution of (4.1) on  $[0, \infty)$ . In fact, if x > 0, then we compute

$$\min \{u(x) - u'(x), u'(x) - 1\} = \min \{0, Ke^x - 1\} = 0.$$

Therefore it remains to verify that  $u(x) = Ke^x$  is a subsolution on  $[0, \infty)$  using Definition 4.1. Suppose  $\phi \in C^1$  and satisfies  $(u - \phi)(x) \leq (u - \phi)(0) = 0$  for  $x \in [0, \infty)$  in a neighborhood of 0. Then it follows that  $\phi'(0) \geq u'(0) = K$  and hence

$$\min\left\{\phi(0) - \phi'(0), \phi'(0) - 1\right\} = \min\left\{K - \phi'(0), \phi'(0) - 1\right\} = K - \phi'(0) \le 0.$$

Thus the claim follows.

Next we show that v(x) = x + 1 is also a constrained viscosity solution on  $[0, \infty)$ . In fact, it is easy to see that v(x) = x + 1 solves (4.1) for x > 0. Thus it remains to show that it is

also a subsolution on  $[0, \infty)$ . To this end, let  $\varphi \in C^1$  with  $(v - \varphi)(x) \leq (v - \varphi)(0) = 0$  for  $x \in [0, \infty)$  in a neighborhood of 0. Then we have  $\varphi'(0) \geq v'(0) = 1$  and therefore

$$\min \{\varphi(0) - \varphi'(0), \varphi'(0) - 1\} = \min \{1 - \varphi'(0), \varphi'(0) - 1\} = 1 - \varphi'(0) \le 0.$$

This shows that v is a subsolution and thus a constrained solution on  $[0, \infty)$ .

Note that the controlled process corresponding to (4.1) is  $dX(t) = 1 \cdot dt + 0 \cdot dW(t) - dZ(t)$ or X(t) = x + t - Z(t) for  $t \ge 0$  and the objective is to maximize  $J(x, Z) = \mathbb{E}_x \int_0^\infty e^{-t} dZ(t)$ . For this process, it is clear that  $\mathcal{A}_x \ne \emptyset$  for all  $x \in [0, \infty)$ . Moreover, from the state constraint,  $Z(t) \le x + t$  for all  $t \ge 0$ . Then it follows that

$$J(x,Z) = \mathbb{E}_x \int_0^\infty \int_t^\infty e^{-s} ds dZ(t) = \mathbb{E}_x \int_0^\infty \int_0^s dZ(t) e^{-s} ds \le \mathbb{E}_x \int_0^\infty e^{-s} (x+s) ds = x+1.$$

Thus the value function  $V(x) \leq x+1$ . In fact, V(x) = x+1 and  $Z^*(t) := xI_{\{t=0\}} + I_{\{t>0\}} \int_0^t s ds$  is an optimal control, since  $J(x, Z^*) = x + \int_0^\infty t e^{-t} dt = x + 1$ .

To conclude, the value function V is the unique constrained viscosity solution of (4.1) on  $[0, \infty)$  in the class of functions with polynomial growth rate.

**Example 4.3.** In this example, we demonstrate that the QVI

$$\min \{u(x) - u''(x), u'(x) - 1\} = 0, \quad x \in (0, \infty)$$
(4.2)

has no constrained viscosity solution on  $[0, \infty)$ .

First, one can show that u(x) = x + c is not a constrained viscosity solution of (4.2) on  $[0, \infty)$ , where c is a constant. Certainly it is the case if c < 0 since for  $x \in (0, -c)$ , we have  $\min \{u(x) - u''(x), u'(x) - 1\} = \min \{x + c, 1 - 1\} = x + c < 0$ . Now let's consider the case when  $c \ge 0$ . The function  $\phi(x) = c + 2x - (1 - \frac{c}{2})x^2$  satisfies

$$(u - \phi)(x) = -x + (1 - \frac{c}{2})x^2 \le (u - \phi)(0) = 0$$
, for  $x \ge 0$  sufficiently small,

and  $\phi'(x) = 2 - (2 - c)x$ ,  $\phi''(x) = -2 + c$ . Thus we have

$$\min \{\phi(0) - \phi''(0), \phi'(0) - 1\} = \min \{c - (-2 + c), 2 - 1\} > 0;$$

this shows that u(x) = x + c is not a subsolution on  $[0, \infty)$ .

Next we show that  $u(x) = c_1 e^x + c_2 e^{-x}$  is not a constrained viscosity solution of (4.2) on  $[0, \infty)$  either, where  $c_1, c_2$  are constants. Note that for  $x \ge 0$  small,

$$u(x) = c_1(1 + x + \frac{1}{2}x^2 + o(x^2)) + c_2(1 - x + \frac{1}{2}x^2 + o(x^2))$$
  
=  $(c_1 + c_2) + (c_1 - c_2)x + \frac{1}{2}(c_1 + c_2)x^2 + o(x^2).$ 

If  $c_1 + c_2 > 0$ , then we consider  $\phi(x) = (c_1 + c_2) + (|c_1 - c_2| + 2)x + \frac{1}{3}(c_1 + c_2)x^2$ . Clearly we have  $(u - \phi)(x) \le (u - \phi)(0) = 0$  for x small and

$$\min\left\{\phi(0) - \phi''(0), \phi'(0) - 1\right\} = \min\left\{(c_1 + c_2) - \frac{2}{3}(c_1 + c_2), |c_1 - c_2| + 2 - 1\right\} > 0$$

Thus u is not a constrained viscosity solution on  $[0, \infty)$ .

Now we consider the case when  $c_1 + c_2 \leq 0$ . Let  $\phi(x) = (c_1 + c_2) + (|c_1 - c_2| + 2)x + (c_1 + c_2 - 1)x^2$ . Then we can verify  $(u - \phi)(x) \leq (u - \phi)(0) = 0$  for x small and  $\phi'(x) = |c_1 - c_2| + 2 + 2(c_1 + c_2 - 1)x$  and  $\phi''(x) = 2(c_1 + c_2 - 1)$ . Then we compute

$$\min \{\phi(0) - \phi''(0), \phi'(0) - 1\} = \min \{c_1 + c_2 - 2(c_1 + c_2) + 2, |c_1 - c_2| + 2 - 1\}$$
$$= \min \{2 - (c_1 + c_2), |c_1 - c_2| + 1\} > 0;$$

which again demonstrates that u is not a constrained viscosity solution of (4.2) on  $[0, \infty)$ .

One observes that any linear combination of x+c and  $c_1e^x+c_2e^{-x}$  can not be a constrained viscosity solution of (4.2) on  $[0,\infty)$  either. In addition, functions of the form  $u(x) = (x + c)I_{\{x>a\}} + (c_1e^x+c_2e^{-x})I_{\{x\leq a\}}$  are not constrained viscosity solution of (4.2) on  $[0,\infty)$ , where  $a, c_1, c_2$  are appropriately selected constants so that  $u \in C^1([0,\infty)) \cap C^2([0,\infty) - \{a\})$  and solves (4.2) in  $(0,\infty)$ .

Finally we note that for the corresponding controlled process  $X(t) = x + \sqrt{2}W(t)$  and the reward functional  $\mathbb{E}_x \int_0^\infty e^{-t} dZ(t)$ ,  $\mathcal{A}_0 = \emptyset$ . The reason is that the Brownian motion W, starting from 0, changes sign infinitely many times and hence can not satisfy the state constraint in any time interval  $[0, \varepsilon]$ .

**Example 4.4.** In this example, we consider the system of coupled QVIs

$$\min\left\{ru(x,\alpha) - \mu_{\alpha}xu'(x,\alpha) - \frac{1}{2}\sigma_{\alpha}^{2}x^{2}u''(x,\alpha) - \lambda_{\alpha}u(x,\alpha) + \lambda_{\alpha}u(x,3-\alpha), \\ u'(x,\alpha) - 1\right\} = 0, \quad x \in (0,\infty), \ \alpha \in \{1,2\},$$

$$(4.3)$$

where for  $\alpha = 1, 2, \ \mu_{\alpha}, \sigma_{\alpha}$ , and  $\lambda_{\alpha} > 0$  are constants. Moreover, we assume  $\mu_1, \mu_2$  satisfy  $\mu_1 < r < \mu_2 \leq \frac{r\lambda_1 + (r-\mu_1)(r+\lambda_2)}{r+\lambda_1-\mu_1}$ . One can verify that the unique solution to (4.3) in  $(0, \infty) \times \mathcal{M}$  is

$$u(x,1) = x, \quad u(x,2) = \frac{\lambda_2}{\lambda_2 + r - \mu_2} x, \quad x > 0.$$
 (4.4)

Moreover, one can easily verify that  $u(\cdot, \alpha), \alpha = 1, 2$  satisfy the subsolution property at the point x = 0. Therefore u is the unique constrained solution on  $[0, \infty) \times \{1, 2\}$ .

The corresponding controlled dynamic is given by the regime-switching geometric Brownian motion:

$$dX(t) = \mu_{\alpha(t)}X(t)dt + \sigma_{\alpha(t)}X(t)dW(t) - dZ(t),$$

where  $\{\alpha(t), t \ge 0\}$  is a two-state continuous-time Markov chain with generator  $\begin{pmatrix} -\lambda_1 & \lambda_1 \\ \lambda_2 & -\lambda_2 \end{pmatrix}$ . The objective is maximize the reward  $J(x, \alpha, Z) = \mathbb{E}_{x,\alpha} \int_0^\infty e^{-rt} dZ(t)$ . Observe that  $\mathcal{A}_{x,\alpha} \neq \emptyset$  for all  $(x, \alpha) \in [0, \infty) \times \{1, 2\}$ . Moreover, as demonstrated in Song et al. (2011), the value function  $V(x, \alpha) = u(x, \alpha)$  for all  $(x, \alpha) \in [0, \infty) \times \{1, 2\}$ , where u is defined in (4.4).  $\Box$ 

Now let's present the main result of this section.

**Theorem 4.5.** Assume  $\mathcal{A}_{x,\alpha} \neq \emptyset$  and that the value function  $V(\cdot, \alpha)$  is finite for each  $(x, \alpha) \in \overline{S} \times \mathcal{M}$ . Then  $\mathbf{V}(x) = (V(x, 1), \ldots, V(x, m))'$  is a constrained viscosity solution of (2.11) on  $\overline{S} \times \mathcal{M}$ .

The proof of Theorem 4.5 is accomplished by the combination of Propositions 4.6 and 4.7: Proposition 4.6 shows that  $\mathbf{V}$  is a viscosity supersolution, while Proposition 4.7 establishes that  $\mathbf{V}$  is viscosity subsolution.

**Proposition 4.6.** The function **V** is a viscosity supersolution of (2.11) in  $S \times M$ . That is, for any  $(x_0, \alpha_0) \in S \times M$  and any  $C^2$  function  $\phi(\cdot, \cdot)$  satisfying  $\phi(x_0, \alpha_0) = V_*(x_0, \alpha_0)$  and that  $\phi(x, \alpha) \leq V_*(x, \alpha)$  for all x in a neighborhood of  $x_0$  and each  $\alpha \in M$ , we have

$$\min\left\{ (r - \mathcal{L})\phi(x_0, \alpha_0), \min_{i=1,\dots,n} \left\{ D_{x_i}\phi(x_0, \alpha_0) - f_i(x_0, \alpha_0) \right\} \right\} \le 0.$$
(4.5)

*Proof.* By the definition of  $V_*(x_0, \alpha_0)$ , there exists a sequence  $\{x_m\} \subset \mathbb{R}^n_+$  such that

$$x_m \to x_0$$
, and  $V(x_m, \alpha_0) \to V_*(x_0, \alpha_0)$ , as  $m \to \infty$ . (4.6)

This, together with the continuity of  $\phi$ , implies that

$$\gamma_m := V(x_m, \alpha_0) - \phi(x_m, \alpha_0) \to 0$$
, as  $m \to \infty$ .

Let  $B_{\varepsilon}(x_0) := \{x \in \mathbb{R}^n : |x - x_0| < \varepsilon\}$ , where  $\varepsilon > 0$  is sufficiently small so that (i)  $\overline{B}_{\varepsilon}(x_0) \subset S$ and (ii)  $\phi(x, \alpha) \leq V_*(x, \alpha)$  for all  $(x, \alpha) \in \overline{B}_{\varepsilon}(x_0) \times \mathcal{M}$ , where  $\overline{B}_{\varepsilon}(x_0) = \{x \in \mathbb{R}^n : |x| \leq \varepsilon\}$ denotes the closure of  $B_{\varepsilon}(x_0)$ . Choose Z such that Z(0-) = 0 and  $Z(t) = \eta$  for all  $t \geq 0$ , where  $0 \leq |\eta| < \varepsilon/2$ . Then thanks to (4.6),  $Z \in \mathcal{A}_{x_m,\alpha_0}$  for m sufficiently large. Let  $X(\cdot) = X^{x_m,\alpha_0}(\cdot; Z)$  be the corresponding controlled process with initial condition  $(x_m, \alpha_0)$ and control strategy  $Z(\cdot)$ . Put

$$\theta_m := \inf \left\{ t \ge 0 : X(t) \notin B_{\varepsilon}(x_0) \right\}.$$

Let  $\{h_m\}$  be a strictly positive sequence such that

$$h_m \to 0 \text{ and } \frac{\gamma_m}{h_m} \to 0 \text{ as } m \to \infty.$$

Note that the chosen control strategy Z guarantees that  $X(\cdot)$  has at most one jump at t = 0 and remains continuous on  $(0, \theta_m]$ . This, together with the choice of  $\varepsilon$ , implies that  $X(t) \in \overline{B}_{\varepsilon}(x_0)$  for all  $0 \le t \le \theta_m$ . Since  $\phi \le V_* \le V$ , we can apply the dynamic programming principle (3.6) to obtain

$$V(x_m, \alpha_0) \ge \mathbb{E} \left[ \int_0^{\theta_m \wedge h_m} e^{-rs} f(X(s-), \alpha(s-)) \cdot dZ(s) \right] + \mathbb{E} \left[ e^{-r(\theta_m \wedge h_m)} \phi(X(\theta_m \wedge h_m), \alpha(\theta_m \wedge h_m)) \right].$$
(4.7)

On the other hand, Itô's formula yields

$$\phi(x_m, \alpha_0) = \mathbb{E} \left[ e^{-r(\theta_m \wedge h_m)} \phi(X(\theta_m \wedge h_m), \alpha(\theta_m \wedge h_m)) \right] \\ + \mathbb{E} \left[ \int_0^{\theta_m \wedge h_m} e^{-rs} (r - \mathcal{L}) \phi(X(s), \alpha(s)) ds \right] \\ + \mathbb{E} \left[ \int_0^{\theta_m \wedge h_m} e^{-rs} D \phi(X(s), \alpha(s)) \cdot dZ^c(s) \right] \\ - \mathbb{E} \left[ \sum_{0 \le s \le \theta_m \wedge h_m} e^{-rs} [\phi(X(s), \alpha(s-)) - \phi(X(s-), \alpha(s-))] \right],$$

$$(4.8)$$

where in the above, we have used the fact that

$$\mathbb{E}\left[\int_{0}^{\theta_{m}\wedge h_{m}} e^{-rs} D\phi(X(s), \alpha(s)) \cdot \sigma(X(s), \alpha(s)) dW(s)\right] = 0.$$

A combination of (4.7) and (4.8) yields

$$\begin{aligned} \gamma_m &= V(x_m, \alpha_0) - \phi(x_m, \alpha_0) \\ &\geq \mathbb{E} \left[ \int_0^{\theta_m \wedge h_m} e^{-rs} \left( f(X(s-), \alpha(s-)) \cdot dZ(s) - (r-\mathcal{L})\phi(X(s), \alpha(s)) ds \right) \right] \\ &- \mathbb{E} \left[ \int_0^{\theta_m \wedge h_m} e^{-rs} D\phi(X(s), \alpha(s)) \cdot dZ^c(s) \right] \\ &+ \mathbb{E} \left[ \sum_{0 \leq s \leq \theta_m \wedge h_m} e^{-rs} [\phi(X(s), \alpha(s-)) - \phi(X(s-), \alpha(s-))] \right]. \end{aligned}$$

$$(4.9)$$

Now let  $\eta = 0$ , i.e.,  $Z(t) \equiv 0$  for any  $t \ge 0$ . Then (4.9) can be rewritten as

$$\frac{\gamma_m}{h_m} \ge -\frac{1}{h_m} \mathbb{E} \int_0^{\theta_m \wedge h_m} e^{-rs} (r - \mathcal{L}) \phi(X(s), \alpha(s)) ds$$
$$= -\frac{1}{h_m} \mathbb{E} \int_0^{h_m} e^{-rs} (r - \mathcal{L}) \phi(X(s), \alpha(s)) I_{\{s \le \theta_m\}} ds.$$

Note that as  $m \to \infty$  and hence  $h_m \to 0$ , by the right continuity of the trajectory  $(X(s), \alpha(s))$ ,  $e^{-rs}(r - \mathcal{L})\phi(X(s), \alpha(s))I_{\{s \le \theta_m\}} \to (r - \mathcal{L})\phi(x_0, \alpha_0)$  a.s. for  $s \in [0, h_m]$ . Thus by virtue of the bounded convergence theorem, we have

$$(r - \mathcal{L})\phi(x_0, \alpha_0) \ge 0. \tag{4.10}$$

On the other hand, if we choose  $\eta = \eta_i e_i$  with  $0 < \eta_i < \varepsilon/2$  and  $e_i = (0, \ldots, 1, \ldots, 0)'$ being the *i*th unit vector,  $i = 1, \ldots, n$ , then (4.9) reduces to

$$\gamma_m \ge -\mathbb{E} \int_0^{\theta_m \wedge h_m} e^{-rs} (r-\mathcal{L}) \phi(X(s), \alpha(s)) ds + f_i(x_m, \alpha_0) \eta_i + \phi(x_m - \eta, \alpha_0) - \phi(x_m, \alpha_0).$$

Now sending  $m \to \infty$ , we have

$$f_i(x_0, \alpha_0)\eta_i + \phi(x_0 - \eta, \alpha_0) - \phi(x_0, \alpha_0) \le 0.$$

Finally, dividing the above inequality by  $\eta_i$  and letting  $\eta_i \to 0$  lead to

$$D_{x_i}\phi(x_0,\alpha_0) - f_i(x_0,\alpha_0) \ge 0, \quad i = 1,\dots,n.$$
 (4.11)

Now (4.5) follows from a combination of (4.10) and (4.11).

**Proposition 4.7.** The function **V** is a viscosity subsolution of (2.11) in  $\overline{S} \times \mathcal{M}$ . That is, for any  $(x_0, \alpha_0) \in \overline{S} \times \mathcal{M}$  and any  $\varphi \in C^2$  such that  $\varphi(x_0, \alpha_0) = V^*(x_0, \alpha_0)$  and that  $\varphi(x, \alpha) \geq V^*(x, \alpha)$  for  $x \in \overline{S}$  in a neighborhood of  $x_0$  and each  $\alpha \in \mathcal{M}$ , we have

$$\min\left\{(r-\mathcal{L})\varphi(x_0,\alpha_0),\min_{i=1,\dots,n}\left\{D_{x_i}\varphi(x_0,\alpha_0)-f_i(x_0,\alpha_0)\right\}\right\} \le 0.$$
(4.12)

Proof. Suppose on the contrary that (4.12) was wrong, then there would exist some  $(x_0, \alpha_0) \in \overline{S} \times \mathcal{M}$ , a  $\varphi \in C^2$  with  $(V^* - \varphi)(x, \alpha) \leq (V^* - \varphi)(x_0, \alpha_0) = 0$ , and a constant A > 0 such that

$$\min\left\{ (r - \mathcal{L})\varphi(x_0, \alpha_0), \min_{i=1,\dots,n} \left\{ D_{x_i}\varphi(x_0, \alpha_0) - f_i(x_0, \alpha_0) \right\} \right\} \ge 2A > 0.$$
(4.13)

In what follows, we will derive a contradiction to (4.13). This is achieved in several steps. First we use the generalized Itô formula and (4.14) to obtain (4.15), from which we obtain (4.16) and (4.17). Next, detailed analysis using the monotonicity of the functions  $V^*$  and fleads to (4.22). Then we claim in (4.23) that the last term in (4.22) is bounded below by a positive constant, from which, with the aid of dynamic programming (3.5), we obtain a contradiction to (4.13). The final step of the proof is devoted to the proof of (4.23).

Step 1. As in the proof of Proposition 4.6, let  $\{x_m\} \subset \overline{S}$  be a sequence such that

$$x_m \to x_0$$
, and  $V(x_m, \alpha_0) \to V^*(x_0, \alpha_0)$  as  $m \to \infty$ ,

and

$$\gamma_m := V(x_m, \alpha_0) - \varphi(x_m, \alpha_0) \to 0, \text{ as } m \to \infty.$$

Choose *m* sufficiently large so that  $|x_m - x_0| < \varepsilon/2$ . Fix some  $Z \in \mathcal{A}_{x_m,\alpha_0}$  and let  $X(\cdot) = X^{x_m,\alpha_0}(\cdot, Z)$  be the corresponding controlled process. Define  $B_{\varepsilon}(x_0) := \{x \in \overline{S} : |x - x_0| < \varepsilon\}$ ,

where  $\varepsilon > 0$  is small enough so that (i)  $\varphi(x, \alpha) \ge V^*(x, \alpha) \ge V(x, \alpha)$  for all  $(x, \alpha) \in \overline{B}_{\varepsilon}(x_0) \times \mathcal{M}$ , and (ii)

$$\min\left\{(r-\mathcal{L})\varphi(x,\alpha),\min_{i=1,\dots,n}\left\{D_{x_{i}}\varphi(x,\alpha)-f_{i}(x,\alpha)\right\}\right\} \ge A>0, \quad \forall (x,\alpha)\in\overline{B}_{\varepsilon}(x_{0})\times\mathcal{M}.$$
(4.14)

Let  $\theta_m := \inf \{t \ge 0 : X(t) \notin B_{\varepsilon}(x_0)\}$ . Then for any t > 0, we have from the generalized Itô formula that

$$\varphi(x_m, \alpha_0) = \mathbb{E}\left[e^{-r(t \wedge \theta_m)}\varphi(X(t \wedge \theta_m - ), \alpha(t \wedge \theta_m - )) + \int_0^{t \wedge \theta_m -} e^{-rs}(r - \mathcal{L})\varphi(X(s), \alpha(s))ds\right] \\ + \mathbb{E}\left[\int_0^{t \wedge \theta_m -} e^{-rs} \sum_{i=1}^n D_{x_i}\varphi(X(s), \alpha(s))dZ_i^c(s)\right] \\ - \mathbb{E}\left[\sum_{0 \le s < t \wedge \theta_m} e^{-rs} \left[\varphi(X(s), \alpha(s - )) - \varphi(X(s - ), \alpha(s - ))\right]\right].$$

Note that

$$\varphi(X(s), \alpha(s-)) - \varphi(X(s-), \alpha(s-)) = \sum_{i=1}^{n} (X_i(s) - X_i(s-)) D_{x_i} \varphi(X(s-) + z(X(s) - X(s-)), \alpha(s-)) = -\sum_{i=1}^{n} \Delta Z_i(s) D_{x_i} \varphi(X(s-) + z(X(s) - X(s-)), \alpha(s-))$$

for some  $z \in [0, 1]$ . But by virtue of (4.14), for all  $0 \le s < t \land \theta_m$ , we have

$$D_{x_i}\varphi(X(s-) + z(X(s) - X(s-)), \alpha(s-)) \ge f_i(X(s-) + z(X(s) - X(s-)), \alpha(s-)) + A.$$

Further, since  $X(s) \leq X(s-) + z(X(s) - X(s-)) \leq X(s-)$  and that  $f_i(\cdot, \alpha)$  is non-increasing, we have

$$f_i(X(s-) + z(X(s) - X(s-)), \alpha(s-)) \ge f_i(X(s-), \alpha(s-)).$$

Then using (4.14) again, we obtain

$$\varphi(x_m, \alpha_0) \ge \mathbb{E}\left[e^{-r(t \wedge \theta_m)}\varphi(X(t \wedge \theta_m -), \alpha(t \wedge \theta_m -)) + \int_0^{t \wedge \theta_m -} e^{-rs}A(ds + \mathbf{1} \cdot dZ(s))\right] \\ + \mathbb{E}\left[\int_0^{t \wedge \theta_m -} e^{-rs}f(X(s-), \alpha(s-)) \cdot dZ(s)\right],$$

$$(4.15)$$

where 1 = (1, ..., 1)'. Now letting  $t \to \infty$  in (4.15), it follows that on the set  $\{\theta_m = \infty\}$ , we have  $V(x_m, \alpha_0) = \varphi(x_m, \alpha_0) + \gamma_m$ 

$$(x_m, \alpha_0) = \varphi(x_m, \alpha_0) + \gamma_m$$
  

$$\geq \mathbb{E}\left[\int_0^\infty e^{-rs} f(X(s-), \alpha(s-)) \cdot dZ(s)\right] + \frac{A}{r} + \gamma_m.$$
(4.16)

Step 2. On the set  $\{\theta_m < \infty\}$ , we have by letting  $t \to \infty$  in (4.15) that

$$\varphi(x_m, \alpha_0) \ge \mathbb{E}\left[e^{-r\theta_m}\varphi(X(\theta_m -), \alpha(\theta_m -)) + \int_0^{\theta_m -} e^{-rs} f(X(s), \alpha(s)) \cdot dZ(s)\right] + A\mathbb{E}\left[\int_0^{\theta_m -} e^{-rs} (ds + \mathbf{1} \cdot dZ(s))\right].$$
(4.17)

Note that  $X(\theta_m) \leq X(\theta_m-)$  and  $X(\theta_m-) \in B_{\varepsilon}(x_0)$ . Thus there exists some  $\lambda \in [0,1]$  such that

$$x_{\lambda} := X(\theta_m - ) + \lambda(X(\theta_m) - X(\theta_m - )) = X(\theta_m - ) - \lambda \Delta Z(\theta_m) \in \partial B_{\varepsilon}(x_0).$$

Moreover,  $X(\theta_m) \leq x_{\lambda} \leq X(\theta_m -)$ . Note that

$$\varphi(X(\theta_m-),\alpha(\theta_m-)) - \varphi(x_\lambda,\alpha(\theta_m-))$$
  
=  $(X(\theta_m-) - x_\lambda) \cdot D\varphi(X(\theta_m-) + z(X(\theta_m) - x_\lambda),\alpha(\theta_m-))$   
=  $\lambda \Delta Z(\theta_m) \cdot D\varphi(X(\theta_m-) + z(X(\theta_m) - x_\lambda),\alpha(\theta_m-)).$ 

But (4.14) and the monotonicity of  $f_i(\cdot, \alpha)$  imply that

$$D_{x_i}\varphi(X(\theta_m-)+z(X(\theta_m)-x_\lambda),\alpha(\theta_m-)) \ge f_i(X(\theta_m-)+z(X(\theta_m)-x_\lambda),\alpha(\theta_m-))+A$$
$$\ge f_i(X(\theta_m-),\alpha(\theta_m-))+A.$$

This, together with the fact that  $\Delta Z_i(\theta_m) \ge 0$ , leads to

$$\varphi(X(\theta_m -), \alpha(\theta_m -)) - \varphi(x_\lambda, \alpha(\theta_m -)) \ge \lambda \Delta Z(\theta_m) \cdot (f(X(\theta_m -), \alpha(\theta_m -)) + A1).$$
(4.18)

Combing (4.17) and (4.18), we obtain

$$V(x_{m}, \alpha_{0}) = \varphi(x_{m}, \alpha_{0}) + \gamma_{m}$$

$$\geq \mathbb{E} \left[ \int_{0}^{\theta_{m}-} e^{-rs} f(X(s-), \alpha(s-)) \cdot dZ(s) + A \int_{0}^{\theta_{m}-} e^{-rs} \left( ds + \mathbb{1} \cdot dZ(s) \right) \right]$$

$$+ \mathbb{E} \left[ \lambda e^{-r\theta_{m}} \Delta Z(\theta_{m}) \cdot f(X(\theta_{m}-), \alpha(\theta_{m}-)) \right]$$

$$+ \mathbb{E} \left[ e^{-r\theta_{m}} \left( \varphi(x_{\lambda}, \alpha(\theta_{m}-)) + \lambda A \Delta Z(\theta_{m}) \cdot \mathbb{1} \right) \right] + \gamma_{m}.$$

$$(4.19)$$

Note that  $x_{\lambda} \in \overline{B}_{\varepsilon}(x_0)$  and hence  $\varphi(x_{\lambda}, \alpha(\theta_m - )) \geq V^*(x_{\lambda}, \alpha(\theta_m - ))$ . On the other hand, since  $X(\theta_m) \leq x_{\lambda} \leq X(\theta_m - )$ , it follows from (3.4) and the monotonicity of f that

$$V^{*}(x_{\lambda}, \alpha(\theta_{m}-)) \geq V^{*}(X(\theta_{m}), \alpha(\theta_{m}-)) + (x_{\lambda} - X(\theta_{m})) \cdot f(x_{\lambda}, \alpha(\theta_{m}-))$$
  
$$\geq V^{*}(X(\theta_{m}), \alpha(\theta_{m}-)) + (1-\lambda)\Delta Z(\theta_{m}) \cdot f(X(\theta_{m}-), \alpha(\theta_{m}-)).$$
(4.20)

A similar argument as that in Song et al. (2011) yields that

$$\mathbb{E}\left[e^{-r\theta_m}V^*(X(\theta_m),\alpha(\theta_m-))\right] = \mathbb{E}\left[e^{-r\theta_m}V^*(X(\theta_m),\alpha(\theta_m))\right].$$
(4.21)

Now put (4.20) and (4.21) into (4.19) and we obtain

$$V(x_{m},\alpha_{0}) \geq \mathbb{E}\left[\int_{0}^{\theta_{m}-} e^{-rs} f(X(s-),\alpha(s-)) \cdot dZ(s) + e^{-r\theta_{m}}V^{*}(X(\theta_{m}),\alpha(\theta_{m}))\right] \\ + A\mathbb{E}\left[\int_{0}^{\theta_{m}-} e^{-rs} \left(ds + \mathbf{1} \cdot dZ(s)\right)\right] + \gamma_{m} \\ + (1-\lambda)\mathbb{E}\left[e^{-r\theta_{m}}\Delta Z(\theta_{m}) \cdot f(X(\theta_{m}-),\alpha(\theta_{m}-))\right] \\ + \lambda\mathbb{E}\left[e^{-r\theta_{m}}\Delta Z(\theta_{m}) \cdot \left(f(X(\theta_{m}-),\alpha(\theta_{m}-)) + A\mathbf{1}\right)\right] \\ = \mathbb{E}\left[\int_{0}^{\theta_{m}} e^{-rs} f(X(s-),\alpha(s-)) \cdot dZ(s) + e^{-r\theta_{m}}V^{*}(X(\theta_{m}),\alpha(\theta_{m}))\right] \\ + A\mathbb{E}\left[\int_{0}^{\theta_{m}-} e^{-rs} \left(ds + \mathbf{1} \cdot dZ(s)\right) + \lambda e^{-r\theta_{m}}\mathbf{1} \cdot \Delta Z(\theta_{m})\right] + \gamma_{m}.$$

$$(4.22)$$

We now claim that for some constant  $\kappa > 0$  that does not depend on m, we have

$$\mathbb{E}\left[\int_{0}^{\theta_{m}-} e^{-rs} \left(ds + \mathbb{1} \cdot dZ(s)\right) + \lambda e^{-r\theta_{m}} \mathbb{1} \cdot \Delta Z(\theta_{m})\right] \ge \kappa.$$
(4.23)

Step 3. Assume (4.23) for the moment. Then (4.22) can be rewritten as

$$V(x_m, \alpha_0) \ge \mathbb{E}\left[\int_0^{\theta_m} e^{-rs} f(X(s-), \alpha(s-)) \cdot dZ(s) + e^{-r\theta_m} V^*(X(\theta_m), \alpha(\theta_m))\right] + A\kappa + \gamma_m.$$
(4.24)

Combining (4.16) and (4.24), and then taking supremum over  $Z \in \mathcal{A}_{x_m,\alpha_0}$ , it follows from the weak dynamic programming principle (3.5) that

$$V(x_m, \alpha_0) \ge V(x_m, \alpha_0) + \frac{A}{r} \wedge A\kappa + \gamma_m > V(x_m, \alpha_0),$$

for m sufficiently large. This is a contradiction. So we must have (4.12) and hence V is a viscosity subsolution of (2.11).

Step 4. Now it remains to show (4.23). To this end, we consider the function  $\widetilde{W}(x,\alpha) := |x - x_0|^2 - \varepsilon^2$  for  $(x, \alpha) \in B_{\varepsilon}(x_0) \times \mathcal{M}$ . Then it follows that

$$(\mathcal{L}-r)\widetilde{W}(x,\alpha) = 2(x-x_0) \cdot b(x,\alpha) + \frac{1}{2}\operatorname{tr}(2I\sigma(x,\alpha)\sigma'(x,\alpha)) - r(|x-x_0|^2 - \varepsilon^2).$$

Since  $\widetilde{W}$ , b, and  $\sigma$  are continuous, and  $\mathcal{M}$  is a finite set, it is obvious that

$$|(\mathcal{L} - r)\widetilde{W}(x, \alpha)| \le K < \infty$$

for some positive constant K. Now let  $K_0 := \frac{1}{2\varepsilon + K}$  and define  $W(x, \alpha) = K_0 \widetilde{W}(x, \alpha)$  for  $(x, \alpha) \in B_{\varepsilon}(x_0) \times \mathcal{M}$ . Then it follows immediately that

$$|(\mathcal{L} - r)W(x, \alpha)| < 1, \quad (x, \alpha) \in B_{\varepsilon}(x_0) \times \mathcal{M}.$$
(4.25)

Moreover, we have

$$D_{x_i}W(x,\alpha) = 2K_0(x-x_0) \cdot e_i \ge -1,$$
(4.26)

where  $e_i = (0, \ldots, 1, \ldots, 0)'$  denotes the *i*th unit vector. Let  $x_m$ ,  $\theta_m$ ,  $Z \in \mathcal{A}_{x_m,\alpha_0}$  etc. as before. Using (4.25), (4.26), and generalized Itô's formula, detailed computations similar to those in Step 1 yield

$$\mathbb{E}\left[e^{-r\theta_m}W(X(\theta_m-),\alpha(\theta_m-))\right] - W(x_m,\alpha_0) \le \mathbb{E}\left[\int_0^{\theta_m-} e^{-rs}\left(ds + \mathbb{1} \cdot dZ(s)\right)\right].$$
(4.27)

Also, recall that  $X(\theta_m) \leq x_{\lambda} \leq X(\theta_m -)$ . It follows from (4.26) that

$$W(X(\theta_m - ), \alpha(\theta_m - )) - W(x_{\lambda}, \alpha(\theta_m - ))$$

$$= \sum_{i=1}^{n} D_{x_i}(x_{\lambda} + z(\hat{X}(\theta_m - ) - x_{\lambda}), \alpha(\theta_m - ))(X(\theta_m - ) - x_{\lambda}) \cdot e_i$$

$$= \lambda \sum_{i=1}^{n} D_{x_i}(x_{\lambda} + z(\hat{X}(\theta_m - ) - x_{\lambda}), \alpha(\theta_m - ))\Delta Z_i(\theta_m)$$

$$\geq -\lambda \sum_{i=1}^{n} \Delta Z_i(\theta_m) = -\lambda \mathbb{1} \cdot \Delta Z(\theta_m).$$
(4.28)

Combining (4.27) and (4.28), we have

$$\mathbb{E}\left[\int_{0}^{\theta_{m}-} e^{-rs} \left(ds + 1 \cdot dZ(s)\right) + \lambda e^{-r\theta_{m}} 1 \cdot \Delta Z(\theta_{m})\right] \ge \mathbb{E}\left[e^{-r\theta_{m}} W(x_{\lambda}, \alpha(\theta_{m}-))\right] - W(x_{m}, \alpha_{0}).$$

But  $x_{\lambda} \in \partial B_{\varepsilon}(x_0)$ , and consequently  $W(x_{\lambda}, \alpha(\theta_m - )) = 0$ . Also, it is immediate that  $W(x_m, \alpha_0) = K_0(|x_m - x_0|^2 - \varepsilon^2) \leq K_0((\frac{\varepsilon}{2})^2 - \varepsilon^2) = -\frac{3}{4}K_0\varepsilon^2$ . Hence it follows that

$$\mathbb{E}\left[\int_{0}^{\theta_{m}-} e^{-rs} \left(ds + \mathbb{1} \cdot dZ(s)\right) + \lambda e^{-r\theta_{m}} \mathbb{1} \cdot \Delta Z(\theta_{m})\right] \geq \frac{3}{4} K_{0} \varepsilon^{2} =: \kappa > 0.$$

This establishes (4.23) and hence finishes the proof of the theorem.

#### 

# 5 Viscosity Solution: Uniqueness

Our goal is to establish a strong comparison result for constrained viscosity solutions of (2.11). To this end, we need the following lemma.

**Lemma 5.1.** Let  $s(x) = x \cdot \mathbb{1} = \sum_{i=1}^{n} x_i$  and

$$\tilde{u}(x,\alpha) := e^{-\lambda s(x)} u(x,\alpha), \quad \tilde{v}(x,\alpha) := e^{-\lambda s(x)} v(x,\alpha), \quad \forall (x,\alpha) \in S \times \mathcal{M}, \tag{5.1}$$

where  $\lambda > 0$ . Then

(a)  $u(x, \alpha)$  is viscosity subsolution of (2.11) if and only if  $\tilde{u}(x, \alpha)$  is a viscosity subsolution of

$$\min\left\{r\tilde{u}(x,\alpha) - H_{\lambda}(x,\alpha,\tilde{u}(x,\alpha),D\tilde{u}(x,\alpha),D^{2}\tilde{u}(x,\alpha)) - Q\tilde{u}(x,\cdot)(\alpha), \\ \min_{i=1,\dots,n}\left\{e^{\lambda s(x)}[\lambda\tilde{u}(x,\alpha) + D_{x_{i}}\tilde{u}(x,\alpha)] - f_{i}(x,\alpha)\right\}\right\} = 0,$$
(5.2)

where for any  $(x, \alpha, q, p, A) \in \mathbb{R}^n \times \mathcal{M} \times \mathbb{R} \times \mathbb{R}^n \times \mathcal{S}_n$ ,

$$\begin{aligned} H_{\lambda}(x,\alpha,q,p,A) = &\frac{1}{2} \mathrm{tr}(\sigma\sigma'(x,\alpha)A) + \frac{\lambda}{2} \left( \mathbbm{1}'\sigma\sigma'(x,\alpha)p + p'\sigma\sigma'(x,\alpha)\mathbbm{1} \right) \\ &+ b(x,\alpha) \cdot p + \lambda q b(x,\alpha) \cdot \mathbbm{1} + \frac{\lambda^2}{2} q \left| \sigma'(x,\alpha)\mathbbm{1} \right|^2, \end{aligned}$$

and

$$Q\tilde{u}(x,\cdot)(\alpha) = \sum_{j=1}^{m} q_{\alpha j}\tilde{u}(x,j) = \sum_{j=1}^{m} q_{\alpha j}[\tilde{u}(x,j) - \tilde{u}(x,\alpha)]$$

(b) Similarly,  $v(x, \alpha)$  is viscosity supersolution of (2.11) if and only if  $\tilde{v}(x, \alpha)$  is a viscosity supersolution of (5.2).

Proof. We prove part (a) only; the proof of part (b) is similar. Suppose u is viscosity subsolution of (2.11). Let  $\tilde{\varphi}(\cdot, \alpha) \in C^2, \alpha \in \mathcal{M}$  and let  $(x_0, \alpha_0)$  be a maximum point of  $\tilde{u} - \tilde{\varphi}$  with  $(\tilde{u} - \tilde{\varphi})(x_0, \alpha_0) = 0$ . Put

$$\varphi(x,\alpha) := e^{\lambda s(x)} \tilde{\varphi}(x,\alpha).$$

Then it is easy to verify that  $\varphi(\cdot, \alpha) \in C^2$ ,  $\alpha \in \mathcal{M}$  and  $(x_0, \alpha_0)$  is a maximum point of  $u - \varphi$ with  $(u - \varphi)(x_0, \alpha_0) = 0$ . Since u is viscosity subsolution of (2.11), we obtain

$$r\varphi(x_0,\alpha_0) - \frac{1}{2} \operatorname{tr}(\sigma\sigma'(x_0,\alpha_0)D^2\varphi(x_0,\alpha_0)) - b(x_0,\alpha_0) \cdot D\varphi(x_0,\alpha_0) - \sum_{j=1}^m q_{\alpha_0j}\varphi(x_0,j) \le 0, \quad (5.3)$$

or

$$\min_{i=1,\dots,n} \left\{ D_{x_i} \varphi(x_0, \alpha_0) - f_i(x_0, \alpha_0) \right\} \le 0.$$
(5.4)

Since  $\varphi(x, \alpha) = e^{\lambda s(x)} \tilde{\varphi}(x, \alpha)$ , we compute

$$D_{x_i}\varphi(x,\alpha) = e^{\lambda s(x)} [\lambda \tilde{\varphi}(x,\alpha) + D_{x_i}\tilde{\varphi}(x,\alpha)],$$

and

$$D_{x_i x_j} \varphi(x, \alpha) = e^{\lambda s(x)} \left[ \lambda^2 \tilde{\varphi}(x, \alpha) + \lambda (D_{x_j} \tilde{\varphi}(x, \alpha) + D_{x_i} \tilde{\varphi}(x, \alpha)) + D_{x_i x_j} \tilde{\varphi}(x, \alpha) \right].$$

In other words,

$$D\varphi(x,\alpha) = e^{\lambda s(x)} \left[\lambda \tilde{\varphi}(x,\alpha) \mathbb{1} + D\tilde{\varphi}(x,\alpha)\right],$$

and

$$D^{2}\varphi(x,\alpha) = e^{\lambda s(x)} \left[ \lambda^{2} \tilde{\varphi}(x,\alpha) \mathbb{1} \mathbb{1}' + \lambda (\mathbb{1} D \tilde{\varphi}(x,\alpha)' + D \tilde{\varphi}(x,\alpha) \mathbb{1}') + D^{2} \tilde{\varphi}(x,\alpha) \right].$$

Then substituting  $D\varphi$  and  $D^2\varphi$  into (5.3) leads to

$$0 \ge r\tilde{\varphi}(x_0, \alpha_0) - \sum_{j=1}^m q_{\alpha_0 j} \tilde{\varphi}(x_0, j) - b(x_0, \alpha_0) \cdot (\lambda \tilde{\varphi}(x_0, \alpha_0) \mathbb{1} + D\tilde{\varphi}(x_0, \alpha_0)) - \frac{1}{2} \operatorname{tr} \left( \sigma \sigma'(x_0, \alpha_0) \left[ \lambda^2 \tilde{\varphi}(x_0, \alpha_0) \mathbb{1} \mathbb{1}' + \lambda (\mathbb{1} D \tilde{\varphi}(x_0, \alpha_0)' + D \tilde{\varphi}(x_0, \alpha_0) \mathbb{1}') + D^2 \tilde{\varphi}(x_0, \alpha_0) \right] \right),$$

which can be rewritten as

$$r\tilde{\varphi}(x_0,\alpha_0) - H_{\lambda}(x_0,\alpha_0,\tilde{\varphi}(x_0,\alpha_0), D\tilde{\varphi}(x_0,\alpha_0), D^2\tilde{\varphi}(x_0,\alpha_0)) - Q\tilde{\varphi}(x_0,\cdot)(\alpha_0) \le 0.$$
(5.5)

Similarly, (5.4) can be rewritten as

$$\min_{i=1,\dots,n} \left\{ e^{\lambda s(x_0)} [\lambda \tilde{\varphi}(x_0, \alpha_0) + D_{x_i} \tilde{\varphi}(x_0, \alpha_0)] - f_i(x_0, \alpha_0) \right\} \le 0.$$
(5.6)

Therefore in view of (5.5) and (5.6),  $\tilde{u}$  is a viscosity subsolution of (5.2).

Conversely, let  $\tilde{u}$  be a viscosity subsolution of (5.2). Recall  $u(x, \alpha) = e^{\lambda s(x)}\tilde{u}(x, \alpha)$ . Let  $\varphi(\cdot, \alpha) \in C^2, \alpha \in \mathcal{M}$  and  $(x_0, \alpha_0)$  be a maximum point of  $u - \varphi$  with  $(u - \varphi)(x_0, \alpha_0) = 0$ . Put  $\tilde{\varphi}(x, \alpha) := e^{-\lambda s(x)}\varphi(x, \alpha)$ . Detailed calculations as above show that u is a viscosity subsolution of (2.11).

**Lemma 5.2.** For every  $\xi \in \overline{S}$ , there exist  $\eta = \eta(\xi) \in \mathbb{R}^n$  and  $a = a(\xi) > 0$  such that

$$B_{ta}(x+t\eta) \subset S, \quad \forall x \in S \cap B_a(\xi), \ \forall t \in (0,1],$$

where  $B_a(x) = \{ y \in \mathbb{R}^n : |y - x| < a \}.$ 

*Proof.* See Atar and Budhiraja (2006).

With Lemmas 5.1 and 5.2 at our hands, we are now ready to establish the strong comparison result for the constrained viscosity solution of (2.11).

**Theorem 5.3.** Let  $\mathbf{u} \in USC(\bar{S} \times \mathcal{M}; \mathbb{R}^m)$  and  $\mathbf{v} \in LSC(\bar{S} \times \mathcal{M}; \mathbb{R}^m)$  be respectively viscosity subsolution on  $\bar{S} \times \mathcal{M}$  and supersolution in  $S \times \mathcal{M}$  of (2.11) and satisfy

$$|u(x,\alpha)| + |v(x,\alpha)| \le K(1+|x|^p), \quad \forall (x,\alpha) \in \bar{S} \times \mathcal{M},$$
(5.7)

where K and p are positive constants. Assume that for some positive constant  $\kappa_0$ , we have

$$|b(x,\alpha) - b(y,\alpha)| + |\sigma(x,\alpha) - \sigma(y,\alpha)| \le \kappa_0 |x-y|, \qquad (5.8)$$

$$b(x,\alpha)' \mathbb{1} \le \kappa_0 \text{ and } |\sigma(x,\alpha)' \mathbb{1}| \le \kappa_0,$$
(5.9)

for all  $(x, \alpha) \in \overline{S} \times \mathcal{M}$ . Then we have

$$u(x,\alpha) \le v(x,\alpha), \quad \forall (x,\alpha) \in \bar{S} \times \mathcal{M}.$$

*Proof.* Suppose on the contrary that

$$M := \max_{\alpha \in \mathcal{M}} \sup_{x \in \bar{S}} [u(x, \alpha) - v(x, \alpha)] > 0.$$
(5.10)

We will derive a contradiction in the following. Define  $\tilde{u}$  and  $\tilde{v}$  as in (5.1), where  $\lambda > 0$  is a constant to be determined later. Thanks to (5.7),  $\tilde{u}$  and  $\tilde{v}$  are uniformly bounded. Moreover, we have

$$\lim_{|x|\to\infty, x\in\bar{S}} (|\tilde{u}(x,\alpha)| + |\tilde{v}(x,\alpha)|) = 0, \ \forall \alpha \in \mathcal{M}.$$

Therefore in view of (5.10) and the facts that  $\mathcal{M}$  is finite and that  $\tilde{u} - \tilde{v}$  is upper semicontinuous, there exist some bounded set O of  $\bar{S}$  and  $(\hat{x}, \ell) \in O \times \mathcal{M}$ , such that

$$\tilde{M} := \max_{\alpha \in \mathcal{M}} \sup_{x \in \bar{S}} [\tilde{u}(x,\alpha) - \tilde{v}(x,\alpha)] = \max_{x \in O} [\tilde{u}(x,\ell) - \tilde{v}(x,\ell)] = \tilde{u}(\hat{x},\ell) - \tilde{v}(\hat{x},\ell) > 0.$$
(5.11)

Let  $\eta = \eta(\hat{x})$  be as in Lemma 5.2. For any  $\varepsilon, \delta \in (0, 1)$  and  $(x, y) \in O \times O$ , define

$$\Phi(x,y) = \Phi_{\varepsilon,\delta,\lambda}(x,y) := \tilde{u}(x,\ell) - \tilde{v}(y,\ell) - \phi(x,y),$$
  

$$\phi(x,y) := \left|\frac{1}{\varepsilon}(y-x) - \delta\eta\right|^2 + \delta |x-\hat{x}|^2.$$
(5.12)

Note that  $\Phi$  is USC and hence achieves its maximum  $M = M_{\varepsilon,\delta,\lambda}$  on the compact set  $\bar{O}^2$  at  $(\tilde{x}, \tilde{y}) := (x_{\varepsilon,\delta,\lambda}, y_{\varepsilon,\delta,\lambda})$ . By virtue of Lemma 5.2,  $\hat{x} + \varepsilon \delta \eta \in S^o$ . Also, since

$$\Phi(\tilde{x}, \tilde{y}) = \tilde{u}(\tilde{x}, \ell) - \tilde{v}(\tilde{y}, \ell) - \left|\frac{1}{\varepsilon}(\tilde{y} - \tilde{x}) - \delta\eta\right|^2 - \delta |\tilde{x} - \hat{x}|^2$$
  
 
$$\geq \Phi(\hat{x}, \hat{x} + \varepsilon\delta\eta) = \tilde{u}(\hat{x}, \ell) - \tilde{v}(\hat{x} + \varepsilon\delta\eta, \ell),$$

we have

$$\tilde{u}(\tilde{x},\ell) - \tilde{v}(\tilde{y},\ell) - \tilde{u}(\hat{x},\ell) + \tilde{v}(\hat{x} + \varepsilon\delta\eta,\ell) \ge \left|\frac{1}{\varepsilon}(\tilde{y} - \tilde{x}) - \delta\eta\right|^2 + \delta |\tilde{x} - \hat{x}|^2.$$

Multiplying  $\varepsilon^2$  on both sides of the above equation, we see that for each  $\delta$  and  $\lambda$ ,  $\tilde{x} - \tilde{y} \to 0$  as  $\varepsilon \to 0$ . Further, by virtue of (5.11), we have

$$\limsup_{\varepsilon \to 0} \left| \frac{1}{\varepsilon} (\tilde{y} - \tilde{x}) - \delta \eta \right|^2 + \delta \left| \tilde{x} - \hat{x} \right|^2 \le 0;$$

and therefore

$$\tilde{x} \to \hat{x}, \text{ and } \frac{1}{\varepsilon} (\tilde{y} - \tilde{x}) \to \delta\eta, \text{ as } \varepsilon \to 0.$$
 (5.13)

In particular, it follows that

$$\tilde{y} = \tilde{x} + \varepsilon \delta \eta + o(\varepsilon) = \hat{x} + \varepsilon \delta \eta + o(\varepsilon), \text{ as } \varepsilon \to 0,$$
(5.14)

and hence  $\tilde{y} \in S^o$  for  $\varepsilon$  sufficiently small.

The function  $x \mapsto \tilde{u}(x, \ell) - \phi_1(x)$  achieves its maximum at  $\tilde{x}$ , where

$$\phi_1(x) = \tilde{v}(\tilde{y}, \ell) + \left|\frac{1}{\varepsilon}(\tilde{y} - x) - \delta\eta\right|^2 + \delta |x - \hat{x}|^2.$$

Moreover, we compute

$$D\phi_1(x) = -\frac{2}{\varepsilon} \left( \frac{1}{\varepsilon} (\tilde{y} - x) - \delta \eta \right) + 2\delta(x - \hat{x}), \text{ and } D^2\phi_1(x) = \frac{2}{\varepsilon^2} I + 2\delta I.$$

Hence it follows from Lemma 5.1, the definition of viscosity subsolution, and Ishii's lemma that for some  $M \in S_n$ ,  $(-D\phi_1(\tilde{x}), M) \in \bar{\mathcal{P}}^{2,+}\tilde{u}(\tilde{x}, \ell)$ , such that

$$\min\left\{r\tilde{u}(\tilde{x},\ell) - H_{\lambda}(\tilde{x},\ell,\tilde{u}(\tilde{x},\ell), D\phi_{1}(\tilde{x}), M) - Q\tilde{u}(\tilde{x},\cdot)(\ell), \\ \min_{i=1,\dots,n} \left\{e^{\lambda s(\tilde{x})} [\lambda \tilde{u}(\tilde{x},\ell) + D\phi_{1}(\tilde{x}) \cdot e_{i}] - f_{i}(\tilde{x},\ell)\right\}\right\} \leq 0.$$

Thus either

$$\min_{i=1,\dots,n} \left\{ e^{\lambda s(\tilde{x})} [\lambda \tilde{u}(\tilde{x},\ell) + D\phi_1(\tilde{x}) \cdot e_i] - f_i(\tilde{x},\ell) \right\} \le 0,$$
(5.15)

or

$$r\tilde{u}(\tilde{x},\ell) - H_{\lambda}(\tilde{x},\ell,\tilde{u}(\tilde{x},\ell), D\phi_1(\tilde{x}), M) - Q\tilde{u}(\tilde{x},\cdot)(\ell) \le 0.$$
(5.16)

On the other hand, the function  $y \mapsto \tilde{v}(y, \ell) - \phi_2(y)$  achieves its minimum at  $\tilde{y}$ , where

$$\phi_2(y) = \tilde{u}(\tilde{x}, \ell) - \left( \left| \frac{1}{\varepsilon} (y - \tilde{x}) - \delta \eta \right|^2 + \delta \left| \tilde{x} - \hat{x} \right|^2 \right).$$

Direct calculations reveal that

$$D\phi_2(y) = -\frac{2}{\varepsilon} \left( \frac{1}{\varepsilon} (y - \tilde{x}) - \delta \eta \right), \text{ and } D^2 \phi_2(y_\varepsilon) = -\frac{2}{\varepsilon^2} I.$$

Hence the definition of supersolution and Ishii's lemma imply that for some  $N \in S_n$ , we have  $(D\phi_2(\tilde{y}), N) \in \bar{\mathcal{P}}^{2,-}\tilde{v}(\tilde{y}, \ell)$  and

$$\min\left\{r\tilde{v}(\tilde{y},\ell) - H_{\lambda}(\tilde{y},\ell,\tilde{v}(\tilde{y},\ell), D\phi_{2}(\tilde{y}), N) - Q\tilde{v}(\tilde{y},\cdot)(\ell), \\ \min_{i=1,\dots,n}\left\{e^{\lambda s(\tilde{y})}[\lambda\tilde{v}(\tilde{y},\ell) + D\phi_{2}(\tilde{y})\cdot e_{i}] - f_{i}(\tilde{y},\ell)\right\}\right\} \geq 0.$$
(5.17)

Case 1. Now suppose (5.15) is true. Recall  $\tilde{u}(x,\alpha) = e^{-\lambda s(x)}u(x,\alpha)$  and  $\tilde{v}(x,\alpha) = e^{-\lambda s(x)}v(x,\alpha)$ . Then we have from (5.15) and (5.17) that

$$0 \ge \min_{i=1,\dots,n} \left\{ e^{\lambda s(\tilde{x})} [\lambda \tilde{u}(\tilde{x},\ell) + D\phi_1(\tilde{x}) \cdot e_i] - f_i(\tilde{x},\ell) - e^{\lambda s(\tilde{y})} [\lambda \tilde{v}(\tilde{y},\ell) + D\phi_2(\tilde{y}) \cdot e_i] + f_i(\tilde{y},\ell) \right\} = \min_{i=1,\dots,n} \left\{ \lambda (u(\tilde{x},\ell) - v(\tilde{x},\ell)) + (e^{\lambda s(\tilde{x})} D\phi_1(\tilde{x}) - e^{\lambda s(\tilde{y})} D\phi_2(\tilde{y})) \cdot e_i - (f_i(\tilde{x},\ell) - f_i(\tilde{y},\ell)) \right\}.$$

Hence it follows that

$$\begin{aligned} \lambda(u(\tilde{x},\ell) - v(\tilde{x},\ell)) \\ &\leq \max_{i=1,\dots,n} \left\{ (f_i(\tilde{x},\ell) - f_i(\tilde{y},\ell)) - (e^{\lambda s(\tilde{x})} D\phi_1(\tilde{x}) - e^{\lambda s(\tilde{y})} D\phi_2(\tilde{y})) \cdot e_i \right\} \\ &= \max_{i=1,\dots,n} \left\{ (f_i(\tilde{x},\ell) - f_i(\tilde{y},\ell)) - e_i \cdot \left[ e^{\lambda s(\tilde{x})} \left( -\frac{2}{\varepsilon} \left( \frac{1}{\varepsilon} (\tilde{y} - \tilde{x}) - \delta\eta \right) + 2\delta(\tilde{x} - \hat{x}) \right) + e^{\lambda s(\tilde{y})} \frac{2}{\varepsilon} \left( \frac{1}{\varepsilon} (\tilde{y} - \tilde{x}) - \delta\eta \right) \right] \right\}. \end{aligned}$$
(5.18)

Thanks to (5.13), (5.14), and the continuity of  $f_i$ , the right-hand-side of (5.18) converges to 0 as  $\varepsilon \downarrow 0$ . Note that for any  $x \in O$ , we have

$$\begin{split} \tilde{u}(x,\ell) - \tilde{v}(x,\ell) &= \Phi(x,x) \le \Phi(\tilde{x},\tilde{y}) \\ &= \tilde{u}(\tilde{x},\ell) - \tilde{v}(\tilde{y},\ell) - \phi(\tilde{x},\tilde{y}) \le \tilde{u}(\tilde{x},\ell) - \tilde{v}(\tilde{y},\ell). \end{split}$$

In particular, taking  $x = \hat{x}$  leads to  $\tilde{u}(\hat{x}, \ell) - \tilde{v}(\hat{x}, \ell) \leq 0$ , which gives a contradiction to (5.11). Hence, Case 1 is impossible.

Case 2. Now suppose (5.16) is true. Then it follows from (5.16) and (5.17) that

$$r(\tilde{u}(\tilde{x},\ell) - \tilde{v}(\tilde{y},\ell)) - [Q\tilde{u}(\tilde{x},\cdot)(\ell) - Q\tilde{v}(\tilde{y},\cdot)(\ell)] - [H_{\lambda}(\tilde{x},\ell,\tilde{u}(\tilde{x},\ell), D\phi_1(\tilde{x}), M) - H_{\lambda}(\tilde{y},\ell,\tilde{v}(y_{\varepsilon},\ell), D\phi_2(\tilde{y}), N)] \le 0.$$
(5.19)

Using the definition of  $H_{\lambda}$ ,

$$\begin{aligned} H_{\lambda}(\tilde{x},\ell,\tilde{u}(\tilde{x},\ell),D\phi_{s}(\tilde{x}),M) &- H_{\lambda}(\tilde{y},\ell,\tilde{v}(\tilde{y},\ell),D\phi_{2}(\tilde{y}),N) \\ &= \frac{1}{2}\left(\operatorname{tr}(\sigma\sigma'(\tilde{x},\ell)M) - \operatorname{tr}(\sigma\sigma'(\tilde{y},\ell)N)\right) + \lambda\left(\tilde{u}(\tilde{x},\ell)b(\tilde{x},\ell) - \tilde{v}(\tilde{y},\ell)b(\tilde{y},\ell)\right) \cdot \mathbb{1} \\ &+ \frac{\lambda}{2}\left(\left[\mathbb{1}'\sigma\sigma'(\tilde{x},\ell)D\phi_{1}(\tilde{x}) + D\phi_{1}(\tilde{x})'\sigma\sigma'(\tilde{x},\ell)\mathbb{1}\right] - \left[\mathbb{1}'\sigma\sigma'(\tilde{y},\ell)D\phi_{2}(\tilde{y}) + D\phi_{2}(\tilde{y})'\sigma\sigma'(\tilde{y},\ell)\mathbb{1}\right]\right) \\ &+ b(\tilde{x},\ell) \cdot D\phi_{1}(\tilde{x}) - b(\tilde{y},\ell) \cdot D\phi_{2}(\tilde{y}) + \frac{\lambda^{2}}{2}\left(\tilde{u}(\tilde{x},\ell)\left|\sigma'(\tilde{x},\ell)\mathbb{1}\right|^{2} - \tilde{v}(\tilde{y},\ell)\left|\sigma'(\tilde{y},\ell)\mathbb{1}\right|^{2}\right).\end{aligned}$$

Hence it follows that

$$\begin{bmatrix}
r - \lambda b(\tilde{x}, \ell)' \mathbb{1} - \frac{\lambda^2}{2} |\sigma'(\tilde{x}, \ell) \mathbb{1}|^2 \\
\leq [Q\tilde{u}(\tilde{x}, \cdot)(\ell) - Q\tilde{v}(\tilde{y}, \cdot)(\ell)] + \lambda \tilde{v}(\tilde{y}, \ell)(b(\tilde{x}, \ell) - b(\tilde{y}, \ell)) \cdot \mathbb{1} \\
+ \frac{1}{2} [\operatorname{tr}(\sigma\sigma'(\tilde{x}, \ell)M) - \operatorname{tr}(\sigma\sigma'(\tilde{y}, \ell)N)] \\
+ \frac{\lambda}{2} ([\mathbb{1}'\sigma\sigma'(\tilde{x}, \ell)D\phi_1(\tilde{x}) + D\phi_1(\tilde{x})'\sigma\sigma'(\tilde{x}, \ell)\mathbb{1}] - [\mathbb{1}'\sigma\sigma'(\tilde{y}, \ell)D\phi_2(\tilde{y}) + D\phi_2(\tilde{y})'\sigma\sigma'(\tilde{y}, \ell)\mathbb{1}]) \\
+ b(\tilde{x}, \ell) \cdot D\phi_1(\tilde{x}) - b(\tilde{y}, \ell) \cdot D\phi_2(\tilde{y}) + \frac{\lambda^2}{2} \tilde{v}(\tilde{y}, \ell) \left( |\sigma'(\tilde{x}, \ell)\mathbb{1}|^2 - |\sigma'(\tilde{y}, \ell)\mathbb{1}|^2 \right).$$
(5.20)

Using (5.9) and the fact that r > 0, we choose  $\lambda > 0$  sufficiently small so that

$$r - \lambda b(\tilde{x}, \ell)' 1 - \frac{\lambda^2}{2} |\sigma'(\tilde{x}, \ell) 1|^2 > 0.$$
(5.21)

Next we analyze the terms on the right-hand side of (5.20). Recall that

$$\Phi(\tilde{x},\tilde{y}) = \tilde{u}(\tilde{x},\ell) - \tilde{v}(\tilde{y},\ell) - \phi(\tilde{x},\tilde{y}) \ge \Phi(\hat{x},\hat{x}) = \tilde{u}(\hat{x},\ell) - \tilde{v}(\hat{x},\ell) - \delta^2 |\eta|^2.$$

Note also  $q_{\ell\ell} < 0$ . Thus it follows that

$$q_{\ell\ell}\left[\tilde{u}(\tilde{x},\ell) - \tilde{v}(\tilde{y},\ell)\right] \le q_{\ell\ell}\left[\tilde{u}(\hat{x},\ell) - \tilde{v}(\hat{x},\ell) + \phi(\tilde{x},\tilde{y}) - \delta^2 |\eta|^2\right].$$

This, together with (5.11), (5.13), (5.14), and the fact that  $\tilde{u} - \tilde{v}$  is USC, lead to

$$\begin{split} &\lim_{\epsilon \downarrow 0} \sup \left[ Q \tilde{u}(\tilde{x}, \cdot)(\ell) - Q \tilde{v}(\tilde{y}, \cdot)(\ell) \right] \\ &= \limsup_{\epsilon \downarrow 0} \sum_{j \neq \ell} q_{\ell j} (\tilde{u}(\tilde{x}, j) - \tilde{v}(\tilde{y}, j)) + q_{\ell \ell} (\tilde{u}(\tilde{x}, \ell) - \tilde{v}(\tilde{y}, \ell)) \\ &\leq \sum_{j \neq \ell} q_{\ell j} (\tilde{u}(\hat{x}, j) - \tilde{v}(\hat{x}, j)) + q_{\ell \ell} (\tilde{u}(\hat{x}, \ell) - \tilde{v}(\hat{x}, \ell)) - \delta^2 q_{\ell \ell} |\eta|^2 \\ &\leq \sum_{j \neq \ell} q_{\ell j} (\tilde{u}(\hat{x}, \ell) - \tilde{v}(\hat{x}, \ell)) + q_{\ell \ell} (\tilde{u}(\hat{x}, \ell) - \tilde{v}(\hat{x}, \ell)) - \delta^2 q_{\ell \ell} |\eta|^2 = -\delta^2 q_{\ell \ell} |\eta|^2 \,. \end{split}$$
(5.22)

By virtue of Ishii's lemma,

$$\lim_{\varepsilon \downarrow 0} \frac{1}{2} \left( \operatorname{tr}(\sigma \sigma'(x_{\varepsilon}, \ell)M) - \operatorname{tr}(\sigma \sigma'(y_{\varepsilon}, \ell)N) \right) = 0.$$
(5.23)

Next, using (5.8) and (5.13), and noting that  $\tilde{v}$  is bounded, we have

$$\lim_{\varepsilon \downarrow 0} \left[ \lambda \tilde{v}(\tilde{y},\ell) (b(\tilde{x},\ell) - b(\tilde{y},\ell)) \cdot 1 + \frac{\lambda^2}{2} \tilde{v}(\tilde{y},\ell) \left( \left| \sigma'(\tilde{x},\ell) 1 \right|^2 - \left| \sigma'(\tilde{y},\ell) 1 \right|^2 \right) \right] = 0.$$
(5.24)

Similarly (5.8) and (5.13) imply that

$$\begin{aligned} \left[ \mathbb{1}' \sigma \sigma'(\tilde{x}, \ell) D \phi_{1}(\tilde{x}) + D \phi_{1}(\tilde{x})' \sigma \sigma'(\tilde{x}, \ell) \mathbb{1} \right] &- \left[ \mathbb{1}' \sigma \sigma'(\tilde{y}, \ell) D \phi_{2}(\tilde{y}) + D \phi_{2}(\tilde{y})' \sigma \sigma'(\tilde{y}, \ell) \mathbb{1} \right] \\ &= -\frac{2}{\varepsilon} \mathbb{1}' \left( \sigma \sigma'(\tilde{x}, \ell) - \sigma \sigma'(\tilde{y}, \ell) \right) \cdot \left( \frac{1}{\varepsilon} (\tilde{y} - \tilde{x}) - \delta \eta \right) \\ &- \frac{2}{\varepsilon} \left( \frac{1}{\varepsilon} (\tilde{y} - \tilde{x}) - \delta \eta \right)' \left( \sigma \sigma'(\tilde{x}, \ell) - \sigma \sigma'(\tilde{y}, \ell) \right) \mathbb{1} \\ &+ 2\delta \left( \mathbb{1}' \sigma \sigma'(\tilde{x}, \ell) (\tilde{x} - \hat{x}) + (\tilde{x} - \hat{x})' \sigma \sigma'(\tilde{x}, \ell) \mathbb{1} \right) \\ &\to 0, \quad \text{as } \varepsilon \to 0; \end{aligned}$$
(5.25)

and

$$b(\tilde{x},\ell) \cdot D\phi_1(\tilde{x}) - b(\tilde{y},\ell)D\phi_2(\tilde{y})$$
  
=  $-\frac{2}{\varepsilon}(b(\tilde{x},\ell) - b(\tilde{y},\ell)) \cdot \left(\frac{1}{\varepsilon}(\tilde{y} - \tilde{x}) - \delta\eta\right) + 2\delta b(\tilde{x},\ell) \cdot (\tilde{x} - \hat{x})$  (5.26)  
 $\rightarrow 0, \text{ as } \varepsilon \rightarrow 0.$ 

Now letting  $\varepsilon \downarrow 0$  and using (5.21)–(5.26) in (5.20), we conclude that for sufficiently small  $\lambda$ ,

$$\limsup_{\varepsilon \to 0} \tilde{u}(\tilde{x}, \ell) - \tilde{v}(\tilde{y}, \ell) \le -\delta^2 q_{\ell\ell} |\eta|^2.$$

But  $\delta > 0$  can be arbitrarily small, as argued in Case 1, it follows that

$$\tilde{u}(\hat{x},\ell) - \tilde{v}(\hat{x},\ell) \le \tilde{u}(x_{\varepsilon},\ell) - \tilde{v}(y_{\varepsilon},\ell) \to 0$$
, as  $\varepsilon \to 0$  and  $\delta \to 0$ ,

which again contradicts (5.11). Therefore for any  $x \in S$  and  $\alpha \in \mathcal{M}$ , we have  $u(x, \alpha) \leq v(x, \alpha)$ , as desired.

**Remark 5.4.** Note that under condition (5.9), the value function is bounded above by an affine function. In fact, for any  $Z \in \mathcal{A}_{x,\alpha}$  with  $(x, \alpha) \in S \times \mathcal{M}$ , we have

$$d(e^{-rt}X(t)) = e^{-rt} \left[ (b(X(t), \alpha(t)) - rX(t))dt + \sigma(X(t), \alpha(t))dW(t) - dZ(t) \right].$$

Thus

$$e^{-rt}dZ(t) = e^{-rt}(b(X(t), \alpha(t)) - rX(t))dt + e^{-rt}\sigma(X(t), \alpha(t))dW(t) - d(e^{-rt}X(t)),$$

from which it follows that

$$\int_0^\infty e^{-rt} \mathbb{1} \cdot dZ(t) \le \int_0^\infty e^{-rt} \mathbb{1} \cdot \left[ (b(X(t), \alpha(t)) - rX(t))dt + \sigma(X(t), \alpha(t))dW(t) \right] + \mathbb{1} \cdot x.$$

Taking expectations on both sides and using (5.9), we have

$$\mathbb{E}\int_0^\infty e^{-rt}\mathbb{1} \cdot dZ(t) \le \mathbb{E}\int_0^\infty e^{-rt}(\kappa_0 dt + \mathbb{1} \cdot \sigma(X(t), \alpha(t))dW(t)) + \mathbb{1} \cdot x = \frac{\kappa_0}{r} + \mathbb{1} \cdot x.$$

In the above,  $\mathbb{E} \int_0^\infty e^{-rt} \mathbb{1} \cdot \sigma(X(t), \alpha(t)) dW(t) = 0$  since  $\mathbb{1} \cdot \sigma$  is uniformly bounded. Hence, it follows that

$$V(x,\alpha) = \sup_{Z \in \mathcal{A}_{x,\alpha}} \mathbb{E} \int_0^\infty e^{-rt} f(X(t-),\alpha(t-) \cdot dZ(t) \le \|f\|_\infty \left(\frac{\kappa_0}{r} + \mathbf{1} \cdot x\right)$$

Finally we summarize the main result of this paper from Theorems 4.5 and 5.3.

**Theorem 5.5.** Assume (5.8), (5.9), and that  $\mathcal{A}_{x,\alpha} \neq \emptyset$  for every  $(x, \alpha) \in S \times \mathcal{M}$ . Then the value function V defined in (2.6) is the unique constrained viscosity solution of the system of coupled quasi-variational inequalities (2.11) on  $\bar{S} \times \mathcal{M}$ .

**Remark 5.6.** At first look, condition (5.9) seems rather restrictive. Simple models such as regime-switching geometric Brownian motion considered in Example 4.4 are excluded. However, the following example indicates that in general, one can not remove (5.9); otherwise, uniqueness may not hold.

Example 5.7. Let's consider a 1-dimensional squared Bessel process subject to control

$$dX(t) = dt + 2\sqrt{|X(t)|}dW(t) - dZ(t),$$
(5.27)

with reward functional  $J(x, Z) = \mathbb{E}_x \int_0^\infty e^{-t} dZ(t)$ , where x > 0. It is well known (see, e.g., Revuz and Yor (1999)) that the stochastic differential equation

$$\xi(t) = x + t + 2 \int_0^t \sqrt{|\xi(s)|} dW(s)$$

has a unique strong solution  $\xi^x$ , and for all  $t \ge 0$ ,  $\xi^x(t) = x + |W(t)|^2 \ge 0$  if  $x \ge 0$ . Moreover, using Song et al. (2012), it follows that  $\mathbb{P}_0 \{\tau = 0\} = 1$ , where  $\tau := \inf \{t > 0 : \xi^0(t) > 0\}$ . Hence it follows that  $\mathcal{A}_x \neq \emptyset$  for all  $x \in [0, \infty)$ .

The corresponding QVI is

$$\min \{ u(x) - u'(x) - 2xu''(x), u'(x) - 1 \} = 0, \quad x \in (0, \infty).$$
(5.28)

One can easily check that v(x) = x + 1 is a constrained viscosity solution to (5.28) on  $[0, \infty)$ . In fact, for x > 0,

$$\min \{v(x) - v'(x) - 2xv''(x), v'(x) - 1\} = \min \{x + 1 - 1 - 2x \cdot 0, 1 - 1\} = \min \{x, 0\} = 0.$$

Moreover, the subsolution property holds at the point x = 0 since for any  $\phi \in C^2$  with  $(v - \phi)(x) \leq (v - \phi)(0) = 0$ , we have  $\phi'(0) \geq 1$  and hence

$$\min\left\{\phi(0) - \phi'(0) - 2 \cdot 0 \cdot \phi''(0), \phi'(0) - 1\right\} = \min\left\{1 - \phi'(0), \phi'(0) - 1\right\} \le 0.$$

Thus v(x) = x + 1 is indeed a constrained viscosity solution to (5.28) on  $[0, \infty)$ .

Next, we demonstrate that (5.28) has at least another constrained viscosity solution on  $[0,\infty)$ . First we note that the function  $\psi(x) := \sinh(\sqrt{2x})$  is increasing and solves the equation u(x) - u'(x) - 2xu''(x) = 0 in  $(0,\infty)$ . Further, straightforward calculations reveal that

$$\psi'(x) = \frac{\cosh(\sqrt{2x})}{\sqrt{2x}}, \quad \psi''(x) = \frac{1}{2\sqrt{2}} \left[ -\frac{\cosh(\sqrt{2x})}{x^{3/2}} + \frac{\sqrt{2}\sinh(\sqrt{2x})}{x} \right]$$

The equation  $\psi''(x) = 0$  or equivalently  $\frac{\cosh(\sqrt{2x})}{\sinh(\sqrt{2x})} = \sqrt{2x}$  has a unique positive root, denoted by z. Now we claim that the function defined by

$$u(x) = \sinh(\sqrt{2x}) \frac{\sqrt{2z}}{\cosh(\sqrt{2z})} I_{(0,z]}(x) + \left(x - z + \sinh(\sqrt{2z}) \frac{\sqrt{2z}}{\cosh(\sqrt{2z})}\right) I_{(z,\infty)}(x)$$

is the only constrained viscosity solution to (5.28) on  $[0, \infty)$ . In fact, one can directly verify that u(x) is a solution to (5.28) for x > 0. As in Example 4.2, it remains to verify the

subsolution property at the point x = 0. To this end, let  $\phi \in C^2$  with  $(u - \phi)(x) \leq (u - \phi)(0) = 0$  for  $x \in [0, \infty)$  in a neighborhood of 0. Then  $\phi(0) = 0$  and for x > 0,  $\phi(x) \geq u(x) > 0$ . Thus we must have  $\phi'(0) \geq 0$  and hence

$$\min\left\{\phi(0) - \phi'(0) - 0 \cdot \phi''(0), \phi'(0) - 1\right\} \le 0.$$

The desired conclusion follows. Note that u also satisfies the polynomial growth condition (5.7).

In terms of the singular control problem (5.27), it turns out that the value function V(x) = v(x) = x + 1. In fact, from the state constraint, we have  $Z(t) \le x + W(t)^2$  for any  $t \ge 0$ . Therefore

$$J(x,Z) = \mathbb{E}_x \int_0^\infty e^{-t} dZ(t) = \mathbb{E}_x \int_0^\infty \int_t^\infty e^{-s} ds dZ(t) = \mathbb{E}_x \int_0^\infty \int_0^s dZ(t) e^{-s} ds$$
$$\leq \mathbb{E}_x \int_0^\infty e^{-s} (x+W^2(s)) ds = \int_0^\infty e^{-s} (x+s) ds = x+1.$$

Furthermore, it is easy to check that the control  $Z^*(t) = x + W^2(t)$  is optimal and  $J(x, Z^*) = x + 1$ . Hence V(x) = x + 1 as claimed.

We finish the section with a hierarchical PDE characterization for the boundary behavior of the solution to (2.11). Let  $\ell \subset \{1, 2, ..., n\}$  be an index subset. For a vector v = $(v_1, ..., v_n) \in \mathbb{R}^n$ , we induce a smaller vector  $v^{\ell} := (v_i)_{i \in \ell} \in \mathbb{R}^{|\ell|}$ , i.e.  $v_i^{\ell} = v_{\ell_i}$  for all  $i \in \{1, ..., n\}$ . Typically, this notation will be used for  $v = b, \sigma, f, \xi, X, Z$ .

In a reverse direction, for a vector  $v \in \mathbb{R}^{|\ell|}$ , we define a larger vector  $v^{-\ell} \in \mathbb{R}^n$  by

$$(v^{-\ell})_j = \begin{cases} v_i, & \text{if } j = \ell_i, \\ 0, & \text{otherwise.} \end{cases}$$

For a function  $g : \mathbb{R}^n \times \mathcal{M} \mapsto \mathbb{R}^n$ , we induce another function  $g^{\ell} : \mathbb{R}^{\ell} \times \mathcal{M} \mapsto \mathbb{R}^n$  such that  $g_i^{\ell}(x, \alpha) = g_{\ell_i}(x^{-\ell}, \alpha).$ 

The following assumption is imposed.

(H1) 
$$b_i(x,\alpha) = \sigma_i(x,\alpha) = 0$$
 on  $\{x \in \overline{\mathbb{R}^n_+} \mid x_i = 0\}$ 

This basically means that, in the content of ecosystem modeling, once the *i*th species becomes extinct, it will never revive, i.e. if  $(\zeta_i)_t = 0$  for some t, then  $(\zeta_i)_s = 0$  for all  $s \ge t$ .

Thanks to (H1), (2.1) implies following sub-dynamics:

$$d\zeta^{\ell}(t) = b^{\ell}(\zeta^{\ell}(t), \alpha(t))dt + \sigma^{\ell}(\zeta^{\ell}(t), \alpha(t))dW^{\ell}(t), \ \zeta^{\ell}(0) = x^{\ell}, \alpha(0) = \alpha.$$
(5.29)

Therefore, we can look at following subsystem. Suppose the survived species are indexed by  $\ell$  with its remaining amount  $x \in \mathbb{R}^{|\ell|}$ , then the associated value function can be defined as

$$J^{\ell}(x,\alpha,Z) := \mathbb{E} \int_0^\infty e^{-rs} f^{\ell}(X^{x,\alpha,\ell}(s-),\alpha(s-)) \cdot dZ^{\ell}(s)$$

and

$$V^{\ell}(x, \alpha) = \sup_{Z \in \mathcal{A}_{x, \alpha}} J^{\ell}(x, \alpha, Z).$$

For all  $x \in \mathbb{R}^{|\ell|}, \xi \in \mathbb{R}^m, p \in \mathbb{R}^{|\ell|}, A \in \mathcal{S}^{|\ell|}$ , we define a function

$$G^{\ell}(x,\alpha,\xi,p,A) = \min\{r - \xi_{\alpha} - \frac{1}{2}tr(\sigma^{\ell}(\sigma^{\ell})'(x^{\ell},\alpha)A) - b^{\ell}(x^{\ell},\alpha) \cdot p - \sum_{j=1}^{m} q_{ij}\xi_{j}, \\ \min_{i=1,\dots,|\ell|}\{p_{i} - f_{i}^{\ell}(x,\alpha)\}\}$$

Then, one can apply induction to the previous results to show that,  $\mathbf{V}(x) = (V(x, \alpha))_{\alpha}$ is the unique solution of

$$\begin{cases} G(x,\alpha,\mathbf{V}(x),DV(x,\alpha),D^2V(x,\alpha)) = 0, & (x,\alpha) \in \mathbb{R}^n_+ \times \mathcal{M} \\ V(x^{-\ell},\alpha) = V^{\ell}(x,\alpha), & (x,\alpha) \in \mathbb{R}^{n-1}_+ \times \mathcal{M}, |\ell| = n-1. \end{cases}$$
(5.30)

# 6 Conclusions and Remarks

In this work, we considered a class of singular control problems with state constraints and regime-switching. The controlled dynamic is given by a regime-switching diffusion confined in the unbounded domain  $S = \mathbb{R}^n_+$  and the objective is to maximize the total expected discounted rewards from exerting the singular control. Using the weak dynamic programming principle, we showed that the value function is the unique constrained viscosity solution of the system of coupled nonlinear quasi-variational inequalities (2.11).

Throughout our analysis, the discount rate r was fixed. It is interesting to ask how the solution, with appropriate scaling of the cost, will behave as  $r \to 0$ ; and how the limit, if it exists, relates to that of the average cost control problem. A number of other questions deserve further investigations. In particular, it is worth studying the case when the random environment or the Markov chain  $\alpha$  is unobservable.

# References

- Alvarez, L. H. R. (2000). Singular stochastic control in the presence of a state-dependent yield structure. *Stochastic Process. Appl.*, 86(2):323–343.
- Alvarez, L. H. R. and Shepp, L. A. (1998). Optimal harvesting of stochastically fluctuating populations. J. Math. Biol., 37(2):155–177.
- Atar, R. and Budhiraja, A. (2006). Singular control with state constraints on unbounded domain. Ann. Probab., 34(5):1864–1909.

- Atar, R., Budhiraja, A., and Williams, R. J. (2007). HJB equations for certain singularly controlled diffusions. Ann. Appl. Probab., 17(5-6):1745–1776.
- Bouchard, B. and Touzi, N. (2011). Weak dynamic programming principle for viscosity solutions. *SIAM Journal on Control and Optimization*, 49(3):948–962.
- Crandall, M. G., Ishii, H., and Lions, P.-L. (1992). User's guide to viscosity solutions of second order partial differential equations. *Bull. Amer. Math. Soc.* (N.S.), 27(1):1–67.
- Elliott, R. J. and Siu, T. K. (2010). On risk minimizing portfolios under a Markovian regime-switching Black-Scholes economy. *Ann. Oper. Res.*, 176:271–291.
- Guo, X. and Pham, H. (2005). Optimal partially reversible investment with entry decision and general production function. *Stochastic Process. Appl.*, 115(5):705–736.
- Haussmann, U. G. and Suo, W. (1995a). Singular optimal stochastic controls. I. Existence. SIAM J. Control Optim., 33(3):916–936.
- Haussmann, U. G. and Suo, W. (1995b). Singular optimal stochastic controls. II. Dynamic programming. SIAM J. Control Optim., 33(3):937–959.
- Karatzas, I. and Shreve, S. E. (1984). Connections between optimal stopping and singular stochastic control. I. Monotone follower problems. SIAM J. Control Optim., 22(6):856–877.
- Lee, C. and Weerasinghe, A. (2011). Convergence of a queueing system in heavy traffic with general patience-time distributions. *Stochastic Process. Appl.*, 121(11):2507–2552.
- Mao, X. and Yuan, C. (2006). *Stochastic differential equations with Markovian switching*. Imperial College Press, London.
- Øksendal, B. and Sulem, A. (2002). Optimal consumption and portfolio with both fixed and proportional transaction costs. *SIAM J. Control Optim.*, 40(6):1765–1790 (electronic).
- Paulsen, J. (2003). Optimal dividend payouts for diffusions with solvency constraints. Finance Stoch., 7(4):457–473.
- Pham, H. (2009). Continuous-time stochastic control and optimization with financial applications, volume 61 of Stochastic Modelling and Applied Probability. Springer-Verlag, Berlin.
- Revuz, D. and Yor, M. (1999). Continuous martingales and Brownian motion, volume 293 of Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]. Springer-Verlag, Berlin, third edition.

- Sethi, S. P. and Zhang, Q. (1994). Hierarchical decision making in stochastic manufacturing systems. Systems & Control: Foundations & Applications. Birkhäuser Boston Inc., Boston, MA.
- Slatkin, M. (1978). The dynamics of a population in a Markovian environment. *Ecology*, 59:249–256.
- Song, Q. S., Stockbridge, R. H., and Zhu, C. (2011). On optimal harvesting problems in random environments. *SIAM J. Control Optim.*, 49(2):859–889.
- Song, Q., Yin, G., and Zhu, C. (2012). Optimal switching with constraints and utility maximization of an indivisible market. SIAM Journal on Control and Optimization, 50(2):629– 651.
- Wein, L. M. (1990). Optimal control of a two-station Brownian network. *Math. Oper. Res.*, 15(2):215–242.
- Yin, G. G. and Zhu, C. (2010). Hybrid Switching Diffusions: Properties and Applications, volume 63 of Stochastic Modelling and Applied Probability. Springer, New York.
- Yong, J. and Zhou, X. Y. (1999). Stochastic Controls: Hamiltonian Systems and HJB Equations, volume 43 of Applications of Mathematics (New York). Springer-Verlag, New York.
- Zariphopoulou, T. (1992). Investment-consumption models with transaction fees and Markov-chain parameters. SIAM J. Control Optim., 30(3):613–636.
- Zhang, Q. (2001). Stock trading: an optimal selling rule. *SIAM J. Control Optim.*, 40(1):64–87 (electronic).
- Zhu, C. (2011). Optimal control of risk process in a regime switching environment. Automatica, 47(8):1570–1579.
- Zhu, C. and Yin, G. (2009). On competitive Lotka-Volterra model in random environments. J. Math. Anal. Appl., 357(1):154–170.