

EXTREMAL CROSS-POLYTOPES AND GAUSSIAN VECTORS

GERGELY AMBRUS

ABSTRACT. For $n \geq 1$, let ξ_1, \dots, ξ_n be independent, identically distributed standard normal variables. Among nonnegative real vectors $u = (u_1, \dots, u_n)$ of norm 1, the quantity $\mathbb{E}\|(u_1\xi_1, \dots, u_n\xi_n)\|_\infty$ is maximised when u has at most two non-zero entries, and it is minimised when u is proportional to $(1, \dots, 1)$. Further generalisations of this result are also discussed. As a corollary, a lower bound on the mean width of a general convex body K is derived in terms of the successive inner radii of K .

1. CONTEXT AND MOTIVATION

Let K be a convex body in \mathbb{R}^n . A natural way to measure how close K is to a ball is to relate its volume to that of the largest ball inscribed in K , or to the smallest ball circumscribed about K . Equivalently, one may as well relate $\text{Vol}(K)$ to the inradius of K and to the circumradius of K . However, the best possible estimates in this case are the trivial ones.

More interesting inequalities can be obtained by taking into account the successive inner and outer radii of K . These are defined as follows. Let $r(K)$ and $R(K)$ denote the inradius and the circumradius of K . Furthermore, let \mathcal{A}_i^n denote the set of i -dimensional affine subspaces of \mathbb{R}^n , and for a subspace $L \in \mathcal{A}_i^n$, denote by $K|L$ the orthogonal projection of K onto L . The successive inner and outer radii of K for $1 \leq i \leq n$ are given by

$$r_i(K) = \max_{L \in \mathcal{A}_i^n} r(K \cap L) \quad \text{and} \quad R_i(K) = \min_{L \in \mathcal{A}_i^n} R(K|L).$$

Note that $r_n(K) = r(K)$, $R_n(K) = R(K)$, $2r_1(K)$ is the diameter of K , and $2R_1(K)$ is the minimum width of K .

We also introduce the intrinsic volumes of K for $0 \leq i \leq n$ by

$$(1) \quad V_i(K) = \frac{\binom{n}{i} \kappa_n}{\kappa_i \kappa_{n-i}} \int_{\mathcal{L}_i^n} \text{Vol}_i(K|L) d\mu_i(L),$$

where \mathcal{L}_i^n is the Grassmannian of all i -dimensional linear subspaces of \mathbb{R}^n , equipped with the unique Haar probability measure μ_i , and κ_n is the volume of B^n , the unit ball of dimension n :

$$\kappa_n = \frac{\pi^{n/2}}{\Gamma(1 + \frac{n}{2})}.$$

2010 *Mathematics Subject Classification.* 52A40(primary), and 60D05(secondary).

Key words and phrases. Orthogonal crosspolytopes, Gaussian vectors.

Research was supported by OTKA grants 75016 and 76099.

Alternatively, $V_i(K)$ may be expressed as the coefficients in Steiner's formula

$$\text{Vol}(K + \lambda B^n) = \sum_{i=0}^d \lambda^{n-i} \kappa_{n-i} V_i(K).$$

The most well-known special cases are: $V_n(K) = \text{Vol}(K)$; $2V_{n-1}(K)$ is the surface area of K ; $2\kappa_{n-1}/(n\kappa_n)V_1(K)$ is the mean width of K ; and $V_0(K) = 1$ is the Euler characteristic. For further references, see Gruber [4] or Schneider [7].

Since the intrinsic volumes are the average volumes of projections of K onto lower dimensional subspaces, it is natural to expect a relationship between these and the successive radii of K . This link was established by M. Henk and M. Hernández Cifre [5] who proved that the following inequalities hold:

$$(2) \quad \begin{aligned} V_i(K) &\leq 2^i s_i(R_1(K), \dots, R_n(K)) \quad \text{for every } 0 \leq i \leq n; \\ V_{n-1}(K) &\geq \frac{2^{n-1}}{(n-1)!} \sqrt{s_{n-1}(r_1(K)^2, \dots, r_n(K)^2)}; \\ V_{n-2}(K) &\geq \frac{2\sqrt{2}}{\pi} \frac{2^{n-2}}{(n-2)!} \sqrt{s_{n-2}(r_1(K)^2, \dots, r_n(K)^2)}, \end{aligned}$$

where s_i stands for the i th elementary symmetric polynomial:

$$s_i(\lambda_1, \dots, \lambda_n) = \sum_{1 \leq k_1 < \dots < k_i \leq n} \lambda_{k_1} \dots \lambda_{k_i}.$$

The upper bound (2) is sharp. In order to derive the lower bounds on $V_i(K)$, the authors of [5] apply a sequence of Steiner symmetrisations to K , leading to a convex body \tilde{K} which contains the orthogonal cross-polytope spanned by $\pm r_1(K) e_1, \dots, \pm r_n(K) e_n$, where $(e_i)_1^n$ is the standard basis of \mathbb{R}^n . Using that Steiner symmetrisations do not increase the intrinsic volumes (see e.g. [4]), one arrives at the following bound:

$$(3) \quad V_i(K) \geq V_i(C_n(r_1(K), \dots, r_n(K))),$$

where

$$C_n(\lambda_1, \dots, \lambda_n) = \text{conv}(\pm \lambda_i e_i : i = 1, \dots, n).$$

Thus, in order to derive the best bounds provided by this method, one faces the following question:

Problem 1. For $1 \leq i \leq n$, determine the vectors $u = (u_1, \dots, u_n) \in \mathbb{R}_+^n$ minimising

$$(4) \quad \frac{V_i(C_n(u_1, \dots, u_n))}{\sqrt{s_i(u_1^2, \dots, u_n^2)}}.$$

We note that the quantity (4) is invariant under scaling (see e.g. Corollary 2.1. of [5]), thus we may assume that $u \in S^{n-1}$. It is proved in [5] that for $i = n-1$ and $i = n-2$, the minimum of (4) is attained when u is a multiple of $(1, \dots, 1)$, that is, when the cross-polytope is regular. The authors also conjecture that the same statement should hold for every i .

In the present note, we settle the $i = 1$ case of Problem 1, showing that the minimum is attained in the regular case, whereas the maximum

is attained when the cross-polytope is at most 2 dimensional. Perhaps it is more convenient to formulate an equivalent question about Gaussian vectors.

Problem 2. *Let $n \geq 1$, and ξ_1, \dots, ξ_n be independent, identically distributed standard normal variables. Determine the unit vector $u = (u_1, \dots, u_n) \in \mathbb{R}^n$ with non-negative coordinates which minimises*

$$(5) \quad \mathbb{E} \max_{1 \leq i \leq n} \{|u_i \xi_i|\}.$$

Denoting the random vector (ξ_1, \dots, ξ_n) by ξ and introducing the Hadamard product $v \odot w$ of $v, w \in \mathbb{R}^n$ by

$$(v \odot w)_i = v_i w_i,$$

the quantity in (5) becomes $\mathbb{E}\|u \odot \xi\|_\infty$. Note that $u \odot \xi$ is an n -dimensional Gaussian random vector with independent coordinates, whose covariance matrix has trace 1.

The equivalence of Problem 2 and the $i = 1$ case of Problem 1 follows by the following standard transformation. For a convex body K in \mathbb{R}^n and $x \in S^{n-1}$, let $h_K(x)$ denote the support function and $\rho_{K^*}(x)$ the radial function of K . If K is symmetric, $\|\cdot\|_K$ denotes the assigned norm with unit ball K . K^* stands for the polar body of K . Furthermore, $\sigma(x) = \sigma_{k-1}(x)$ denotes the $(k-1)$ -dimensional surface (Lebesgue) measure on S^{k-1} ; note that σ is a scaled copy of the rotationally invariant probability measure on S^{k-1} , with total mass $k\kappa_k$. With these conventions, using (1),

$$\kappa_{n-1} V_1(K) = \int_{S^{n-1}} h_K(x) d\sigma(x) = \int_{S^{n-1}} \frac{1}{\rho_{K^*}(x)} d\sigma(x) = \int_{S^{n-1}} \|x\|_{K^*} d\sigma(x).$$

Since $C_n^*(u_1, \dots, u_n)$ is a rectangular box of half-axes $e_1/u_1, \dots, e_n/u_n$,

$$\begin{aligned} V_1(C_n(u_1, \dots, u_n)) &= \frac{1}{\kappa_{n-1}} \int_{S^{n-1}} \|u \odot x\|_\infty d\sigma(x) \\ &= \frac{1}{(2\pi)^{(n-1)/2}} \int_{S^{n-1}} \|u \odot x\|_\infty \int_0^\infty e^{-r^2/2} r^n dr d\sigma(x) \\ (6) \quad &= \frac{\sqrt{2\pi}}{(2\pi)^{n/2}} \int_{\mathbb{R}^n} e^{-|x|^2/2} \|u \odot x\|_\infty dx \\ &= \sqrt{2\pi} \mathbb{E}\|u \odot \xi\|_\infty. \end{aligned}$$

Relaxing the independence condition of Problem 2, we can ask the following, more general question. We call a multivariate random variable *centred*, if the mean values of its coordinate variables are 0.

Problem 3. *Among the n -dimensional centred Gaussian random vectors X satisfying $\text{tr Cov } X = 1$, which ones minimise and maximise $\mathbb{E}\|X\|_\infty$?*

Problem 2 is a special case of Problem 3 when $\text{Cov } X$ is assumed to be diagonal.

We answer Problems 2 and 3 formulated above. By *standard normal vector* we understand a vector with i.i.d. standard normal coordinate variables.

Theorem 1. *Let $\xi = (\xi_i)_1^n$ be an n -dimensional standard normal vector, and let $u \in \mathbb{R}^n$ be a unit vector. For $n = 2$, the expectation $\mathbb{E}\|u \odot \xi\|_\infty$ is independent of the choice of u . For $n \geq 3$, the expectation is maximised*

when at most two coordinates of u are non-zero, and it is minimised when $u = (\pm 1/\sqrt{n}, \dots, \pm 1/\sqrt{n})$.

Thus, the regular cross-polytope is the minimiser for Problem 1.

Theorem 2. *Among the n -dimensional centred Gaussian random vectors X satisfying $\text{tr Cov } X = 1$, $\mathbb{E}\|X\|_\infty$ is maximal when $\text{Cov } X$ is diagonal with at most two non-zero entries, and minimal when the absolute values of the coordinates of X are identical almost everywhere.*

We note that among symmetric convex bodies K in \mathbb{R}^n in John position (that is, B_2^n is the maximal volume ellipsoid inscribed in K), $\int_{S^{n-1}} \|x\|_K d\sigma(x)$ is minimal for the cube [6], see also [1], pp. 52–53. It also follows that the cube has minimal mean width among its affine images of the same volume [3]. This, however, does not imply the above results, as the cube has the smallest volume among the rectangular boxes to be considered in the present problem.

Next, we derive a lower estimate for $V_1(K)$ from Theorem 1. Let μ be the median of $\|x\|_\infty$ on \mathbb{R}^n with respect to the standard Gaussian measure. For the reader's convenience, we illustrate how to estimate μ , following Ball [1], pp. 52. On the one hand, (6) implies that

$$V_1\left(C_n\left(\frac{1}{\sqrt{n}}, \dots, \frac{1}{\sqrt{n}}\right)\right) = \sqrt{\frac{2\pi}{n}} \frac{1}{(2\pi)^{n/2}} \int_{\mathbb{R}^n} e^{-|x|^2/2} \|x\|_\infty dx > \sqrt{\frac{\pi}{2n}} \mu.$$

On the other hand, μ satisfies that

$$\frac{1}{2} = \frac{1}{(2\pi)^{n/2}} \int_{[-\mu, \mu]^n} e^{-|x|^2/2} dx = \left(\sqrt{\frac{2}{\pi}} \int_0^\mu e^{-s^2/2} ds\right)^n \approx (1 - e^{-\mu^2/2})^n,$$

thus, from $2^{-1/n} \approx 1 - (\log 2)/n$ we deduce that $\mu \approx \sqrt{2 \log n}$, and

$$(7) \quad V_1\left(C_n\left(\frac{1}{\sqrt{n}}, \dots, \frac{1}{\sqrt{n}}\right)\right) \approx \sqrt{\pi} \sqrt{\frac{\log n}{n}}.$$

Taking (3) into account, we arrive at the following estimate.

Corollary 1. *There exists an absolute constant c , so that for any convex body $K \subset \mathbb{R}^n$,*

$$V_1(K) \geq c \sqrt{\frac{\log n}{n}} \sqrt{r_1(K)^2 + \dots + r_n(K)^2}.$$

Numerical calculations show that the value of c can be chosen to be 1.74. This estimate, however, is not optimal, because the approximating orthogonal cross-polytope does not cover K (e.g. the successive inner radii of the regular cross-polytope form a strictly decreasing sequence). In fact, the authors of [5] conjecture that the sharp lower bound in terms of the inner radii should be $2\sqrt{r_1(K)^2 + \dots + r_n(K)^2}$; setting $K = C_n(\mu, \mu^2, \dots, \mu^n)$ for μ large shows that this bound would be the best possible.

2. PROOFS

We start with a technical lemma.

Lemma 1. *For any $0 < q \leq 2$, the function*

$$F(x) = \frac{e^{x^2/2}}{x^{q-1}} \int_0^x e^{-t^2/2} dt$$

is strictly increasing for $x > 0$.

Proof. It is easy to obtain that

$$F'(x) = \frac{1}{x^q} \left(x + (x^2 - q + 1)e^{x^2/2} \int_0^x e^{-t^2/2} dt \right) =: \frac{1}{x^q} f(x).$$

Here $f(0) = 0$ and

$$f'(x) = x^2 + 2 - q + x(x^2 + 3 - q) e^{x^2/2} \int_0^x e^{-t^2/2} dt$$

which is positive for $x > 0$. Thus, $F'(x) > 0$ for every $x > 0$. \square

Proof of Theorem 1. Since $\mathbb{E}\|u \odot \xi\|_\infty$ is a continuous function of u on the unit sphere, it suffices to find the extremum values among the critical points on S^{n-1} (the compactness of the unit sphere implies that the extrema exist). First, we show that the absolute values of the non-zero coordinates of the critical points are all equal. Because of symmetry, we may and do assume that $u_i \geq 0$ for every $i = 1, \dots, n$.

Using that for a non-negative random variable X

$$\mathbb{E}X = \int_0^\infty \mathbb{P}(X > t) dt = \int_0^\infty (1 - \mathbb{P}(X \leq t)) dt,$$

we can express the expectation in question as

$$\begin{aligned} \mathbb{E}\|u \odot \xi\|_\infty &= \int_0^\infty \left(1 - \prod_{i=1}^n \mathbb{P}(|u_i \xi_i| \leq t) \right) dt \\ (8) \quad &= \int_0^\infty \left(1 - \left(\frac{2}{\pi} \right)^{n/2} \int_0^{t/u_1} e^{-s^2/2} ds \dots \int_0^{t/u_n} e^{-s^2/2} ds \right) dt \\ &= \int_0^\infty (1 - \phi(t/u_1) \dots \phi(t/u_n)) dt, \end{aligned}$$

where

$$\phi(a) = \operatorname{Erf} \left(\frac{a}{\sqrt{2}} \right) = \sqrt{\frac{2}{\pi}} \int_0^a e^{-s^2/2} ds,$$

also using the convention that $c/0 = \infty$ for $c \geq 0$.

When $u_1 \neq 0$, integrating by parts leads to

$$\begin{aligned}
\frac{\partial \mathbb{E}\|u \odot \xi\|_\infty}{\partial u_1} &= \sqrt{\frac{2}{\pi}} \int_0^\infty \frac{t e^{-t^2/2u_1^2}}{u_1^2} \phi(t/u_2) \dots \phi(t/u_n) dt \\
(9) \qquad &= \left[-\sqrt{\frac{2}{\pi}} e^{-t^2/2u_1^2} \phi(t/u_2) \dots \phi(t/u_n) \right]_0^\infty + \\
&\quad + \frac{2}{\pi} \int_0^\infty e^{-t^2/2u_1^2} \sum_{i=2}^n \left(\frac{e^{-t^2/2u_i^2}}{u_i} \prod_{\substack{j=2 \\ j \neq i}}^n \phi(t/u_j) \right) dt.
\end{aligned}$$

Note that on the right hand side, the first summand vanishes.

Next, assume that u is a critical point of $\mathbb{E}\|u \odot \xi\|_\infty$ on S^{n-1} with at least two non-zero coordinates, say, $u_1 \geq u_2 > 0$. Keeping all the other coordinates fixed and applying the Lagrange multiplier method to the restricted function, we obtain that

$$\frac{1}{u_1} \frac{\partial \mathbb{E}\|u \odot \xi\|_\infty}{\partial u_1} = \frac{1}{u_2} \frac{\partial \mathbb{E}\|u \odot \xi\|_\infty}{\partial u_2}.$$

Along with (9), this implies that

$$\begin{aligned}
(10) \qquad &\int_0^\infty \frac{e^{-t^2/2u_1^2}}{u_1} \sum_{i=3}^n \left(\frac{e^{-t^2/2u_i^2}}{u_i} \prod_{\substack{j=2 \\ j \neq i}}^n \phi(t/u_j) \right) dt \\
&= \int_0^\infty \frac{e^{-t^2/2u_2^2}}{u_2} \sum_{i=3}^n \left(\frac{e^{-t^2/2u_i^2}}{u_i} \phi(t/u_1) \prod_{\substack{j=3 \\ j \neq i}}^n \phi(t/u_j) \right) dt.
\end{aligned}$$

The quotient of the above integrands is

$$\frac{u_2 e^{-t^2/2u_1^2} \phi(t/u_2)}{u_1 e^{-t^2/2u_2^2} \phi(t/u_1)} = \frac{s}{\phi(s)} e^{-s^2/2} \frac{e^{(\mu s)^2/2}}{\mu s} \phi(\mu s),$$

where $s = t/u_1$ and $\mu = u_1/u_2$. Setting $q = 2$ in Lemma 1 implies that for any fixed $s > 0$, this is a strictly increasing function of μ . In particular, $\mu > 1$ would imply that the quotient is strictly greater than 1 for every $s > 0$. Thus, equality in (10) can hold only if $\mu = 1$, that is, $u_1 = u_2$.

Therefore, all the non-zero coordinates of the extremal vectors u must be equal, and in order to find the minimum and the maximum values of $\mathbb{E}\|u \odot \xi\|_\infty$, it suffices to compute the expectations for the set of points

$$u^k = (\underbrace{1/\sqrt{k}, \dots, 1/\sqrt{k}}_k, \underbrace{0, \dots, 0}_{n-k}), \quad k = 1, \dots, n.$$

Introducing the notation

$$E_k = \frac{1}{\sqrt{k}} \mathbb{E}\|(\xi_1, \dots, \xi_k)\|_\infty,$$

our goal is to show that

$$E_1 = E_2 > E_3 > \dots > E_n.$$

Note that (6) and (7) imply that the above inequality is asymptotically true, as $E_n \approx \sqrt{\log n/2n}$.

Fix $n \geq 1$, and for $0 \leq \rho \leq 1/\sqrt{n+1}$, introduce

$$u(\rho) = \left(\sqrt{\frac{1-\rho^2}{n}}, \dots, \sqrt{\frac{1-\rho^2}{n}}, \rho \right) \in \mathbb{R}^{n+1}.$$

Then, by (8),

$$\mathbb{E}\|u(\rho) \odot \xi\|_\infty = \int_0^\infty \left(1 - \phi\left(\frac{t}{\rho}\right) \phi\left(t\sqrt{\frac{n}{1-\rho^2}}\right)^n \right) dt =: R(\rho).$$

This leads to

$$\begin{aligned} R'(\rho) &= \sqrt{\frac{2}{\pi}} \int_0^\infty \frac{t}{\rho^2} e^{-t^2/2\rho^2} \phi\left(t\sqrt{\frac{n}{1-\rho^2}}\right)^n dt \\ &\quad - \sqrt{\frac{2}{\pi}} \int_0^\infty \frac{t\rho n^{3/2}}{(1-\rho^2)^{3/2}} e^{-nt^2/2(1-\rho^2)} \phi\left(\frac{t}{\rho}\right) \phi\left(t\sqrt{\frac{n}{1-\rho^2}}\right)^{n-1} dt. \end{aligned}$$

By partial integration,

$$\begin{aligned} \int_0^\infty \frac{t}{\rho^2} e^{-t^2/2\rho^2} \phi\left(t\sqrt{\frac{n}{1-\rho^2}}\right)^n dt &= \left[-e^{-t^2/2\rho^2} \phi\left(t\sqrt{\frac{n}{1-\rho^2}}\right)^n \right]_0^\infty \\ &\quad + \sqrt{\frac{2}{\pi}} \int_0^\infty e^{-t^2/2\rho^2} n\sqrt{\frac{n}{1-\rho^2}} e^{-nt^2/2(1-\rho^2)} \phi\left(t\sqrt{\frac{n}{1-\rho^2}}\right)^{n-1} dt, \end{aligned}$$

where the first term on the right hand side vanishes. Similarly,

$$\begin{aligned} \int_0^\infty \frac{t\rho n^{3/2}}{(1-\rho^2)^{3/2}} e^{-nt^2/2(1-\rho^2)} \phi\left(\frac{t}{\rho}\right) \phi\left(t\sqrt{\frac{n}{1-\rho^2}}\right)^{n-1} dt \\ &= \left[-e^{-nt^2/2(1-\rho^2)} \rho\sqrt{\frac{n}{1-\rho^2}} \phi\left(\frac{t}{\rho}\right) \phi\left(t\sqrt{\frac{n}{1-\rho^2}}\right)^{n-1} \right]_0^\infty \\ &\quad + \sqrt{\frac{2}{\pi}} \int_0^\infty e^{-nt^2/2(1-\rho^2)} \sqrt{\frac{n}{1-\rho^2}} e^{-t^2/2\rho^2} \phi\left(t\sqrt{\frac{n}{1-\rho^2}}\right)^{n-1} dt \\ &\quad + \sqrt{\frac{2}{\pi}} \int_0^\infty e^{-nt^2/2(1-\rho^2)} \rho \frac{n(n-1)}{1-\rho^2} \phi\left(\frac{t}{\rho}\right) \phi\left(t\sqrt{\frac{n}{1-\rho^2}}\right)^{n-2} dt. \end{aligned}$$

The above equations lead to

$$(11) \quad R'(\rho) = \frac{2}{\pi} \int_0^\infty e^{-nt^2/2(1-\rho^2)} (n-1) \sqrt{\frac{n}{1-\rho^2}} \phi\left(t\sqrt{\frac{n}{1-\rho^2}}\right)^{n-2} \cdot \left[e^{-t^2/2\rho^2} \phi\left(t\sqrt{\frac{n}{1-\rho^2}}\right) - e^{-nt^2/2(1-\rho^2)} \rho\sqrt{\frac{n}{1-\rho^2}} \phi\left(\frac{t}{\rho}\right) \right] dt.$$

Since $1/\rho \geq \sqrt{n/(1-\rho^2)}$, Lemma 1 ($q=2$) implies that for every $t > 0$

$$\frac{1}{t\sqrt{n/(1-\rho^2)}} e^{nt^2/2(1-\rho^2)} \phi\left(t\sqrt{\frac{n}{1-\rho^2}}\right) \leq \frac{\rho}{t} e^{t^2/2\rho^2} \phi\left(\frac{t}{\rho}\right),$$

which, by (11), shows that for any $t > 0$, the function $R(\rho)$ is decreasing on the interval $[0, 1/\sqrt{n+1}]$. Hence, for every $n \geq 1$,

$$E_n = R(0) \geq R\left(\frac{1}{\sqrt{n+1}}\right) = E_{n+1}.$$

Furthermore, (11) shows that equality holds above if and only if $n = 1$. \square

Proof of Theorem 2. Let $u_i = \sqrt{(\text{Cov}X)_{ii}}$. For the minimum inequality, we may and do assume that u_1^2 is the largest diagonal entry of $\text{Cov}X$, hence $u_1 \geq 1/\sqrt{n}$. Since $\|X\|_\infty \geq |X_1|$, and $\mathbb{E}|X_1| = \sqrt{2/\pi} u_1$, we obtain the lower bound $\mathbb{E}\|X\|_\infty \geq \sqrt{2/(n\pi)}$. This bound is sharp only if for all i , $u_i = 1/\sqrt{n}$ and $\|X\|_\infty = |X_i|$ almost everywhere, which yields that for every i and j , $|X_i| = |X_j|$ almost everywhere.

For the upper bound, let ξ_1, \dots, ξ_n be i.i.d standard normal variables. By a theorem of Šidák ([8], [9]), which is a relative of Slepian's lemma,

$$\begin{aligned} \mathbb{P}(\|X\|_\infty \leq t) &= \mathbb{P}(|X_1| \leq t, \dots, |X_n| \leq t) \\ &\geq \prod_{i=1}^n \mathbb{P}(|X_i| \leq t) = \prod_{i=1}^n \mathbb{P}(|u_i \xi_i| \leq t). \end{aligned}$$

Thus, the question reduces to Problem 2, and the upper bound provided by Theorem 1 is sharp if the coordinate variables of X are independent. \square

3. FURTHER REMARKS

Theorem 1 may also be proved by induction on n ; the inductive statement asserts that for any constant $C \geq 0$, the quantity

$$\int_{\mathbb{R}^n} \max\{C, \|u \odot x\|_\infty\} d\gamma(x),$$

where γ is the standard n -variate Gaussian distribution, is maximal for $u = (1, 0, \dots, 0)$, and minimal for $u = (1/\sqrt{n}, \dots, 1/\sqrt{n})$. The initial step is the $n = 2$ case. After determining the possible extremum points using Lagrange multipliers, the remaining statement amounts to the following.

Lemma 2. *For any $c \geq 0$, the following inequality holds:*

$$c + \int_c^\infty (1 - (\phi(\sqrt{2}t))^2) dt \leq c + \int_c^\infty (1 - \phi(t)) dt = \sqrt{\frac{2}{\pi}} e^{-c^2/2} + c\phi(c).$$

We choose not to include the somewhat technical proof here; the interested reader can verify the statement by taking second derivatives and analysing the functions.

There are two natural directions to generalise the above results. First, Problem 1 is open for $2 \leq i \leq n-3$. A method similar to the one presented here may be applied to these cases as well; however, when computing the mixed volumes of cross-polytopes, one faces a formula (see e.g. Corollary 2.1 of [5]) which is too complicated to carry out the necessary analysis. It may be possible to express the mixed volumes in a more suitable way; in that respect, it is illustrative that (6) differs from the $i = 1$ case of the above cited formula.

The other direction is to generalise Problem 2 the following way.

Problem 4. Let $n \geq 1$, $p, q \in (1, \infty]$, and let ξ be an n -variate Gaussian vector with i.i.d. standard normal coordinate variables. Determine the maximum and minimum of $\mathbb{E}\|u \odot \xi\|_p$ subject to the condition $\|u\|_q = 1$.

Clearly, in the $p = q = 1$ case the expectation is independent of the choice of u , whereas Theorem 1 provides the answer to the $q = 2, p = \infty$ case. We now extend this for $0 < q < 2$ as well.

Theorem 3. For $p = \infty$ and $0 < q \leq 2$, the answer to Problem 4 is given as follows: $\mathbb{E}\|u \odot \xi\|_\infty$ is maximised by $u = (1, 0, \dots, 0)$, and it is minimised by $u = (n^{-1/q}, \dots, n^{-1/q})$.

Proof. The argument applied in the course of the proof of Theorem 1, together with the general case of Lemma 1 imply that the extremal vectors are among the u_q^k given by

$$u_q^k = (\underbrace{k^{-1/q}, \dots, k^{-1/q}}_k, \underbrace{0, \dots, 0}_{n-k}), \quad k = 1, \dots, n.$$

Introducing

$$E_{k,q} = k^{-1/q} \mathbb{E}\|(\xi_1, \dots, \xi_k)\|_\infty,$$

the chain of inequalities

$$E_{1,q} > E_{2,q} > \dots > E_{n,q}$$

easily follows by

$$\left(\frac{k+1}{k}\right)^{1/q} > \left(\frac{k+1}{k}\right)^{1/2} \geq \frac{\mathbb{E}\|(\xi_1, \dots, \xi_{k+1})\|_\infty}{\mathbb{E}\|(\xi_1, \dots, \xi_k)\|_\infty}. \quad \square$$

It would be natural to expect that for every p and q , the behaviour of the extremal vectors is similar to the above case. However, this is very far from being true. We illustrate this phenomenon by the $n = 2$ and $p = 2$ case, where by elementary but tedious calculations one can show the following. Let $q_L = 3/2$, $q_M = \log 2 / (\log \pi - \log 2) \approx 1.53$ and $q_U = 2$. For $1 \leq q \leq q_L$, the expectation is maximal when $u = (1, 0)$, and there is exactly one local (and global) minimum at $(2^{-1/q}, 2^{-1/q})$. For $q_L < q < q_U$, there is a local maximum at both of these directions, and there are two further local (and global) minimum points. When $q = q_M$, these two maxima are equal; for $q < q_M$, the global maximum is at $(1, 0)$, whereas for $q > q_M$, the global maximum is at $(2^{-1/q}, 2^{-1/q})$. For $q \geq q_U$, the vector $(1, 0)$ becomes a global minimum point, and the only local extremum points are $(1, 0)$, $(0, 1)$ and $(2^{-1/q}, 2^{-1/q})$. For general p , a similar pattern holds with $q_U = 2$, and $q_L \rightarrow 2$ as $p \rightarrow \infty$.

The same situation occurs in higher dimensions, with q_U depending on n . Thus, in particular, it is not true in general that the extremal vectors u are necessarily of the form $(\alpha, \dots, \alpha, 0, \dots, 0)$, up to permutation and sign changes of the coordinates.

For $p = \infty$ and $q > 2$, the distribution of the minimum and maximum points is unclear. Theorem 1 and the discussion above show that for $q \leq 2$, the maximum is achieved at $(1, 0, \dots, 0)$, whereas the minimum is taken when u is parallel to $(1, \dots, 1)$. However, for $q = \infty$, the role of these two directions is clearly swapped. Thus, there must be a transition phase as

$q \rightarrow \infty$, and it is plausible to expect that the behaviour of the extremal points depends heavily on the dimension as well. That for any $q > 2$, whether all the extremal points have equal absolute values of the non-zero coordinates remains an open question.

4. ACKNOWLEDGEMENTS

I would like to thank F. Fodor and V. Vígih for communicating the problem to me and for the fruitful discussions and ideas; K. Ball, M. Henk, A. Litvak, and G. Pisier for the useful advice; and the anonymous referee for the valuable suggestions. I am grateful for the hospitality of the Mathematical Sciences Research Institute, Berkeley, CA.

REFERENCES

- [1] K. Ball, *An elementary introduction to convex geometry*. In: Flavors of geometry, ed. S. Levy, Cambridge University Press, Cambridge, 1997.
- [2] U. Betke, M. Henk, *Intrinsic volumes and lattice points of crosspolytopes*. Monatsch. Math. **115** (1993), no. 1–2, 27–33.
- [3] A. A. Giannopoulos, V. D. Milman and M. Rudelson, *Convex bodies with minimal mean width*. GAFA, Lecture Notes in Mathematics, **1745** (2000), 81–93.
- [4] P. Gruber, *Convex and Discrete Geometry*. Springer-Verlag, 2007.
- [5] M. Henk, M. Hernández Cifre, *Intrinsic volumes and successive radii*. J. Math. Analysis Appl. **343** (2008), no. 2, 733–742.
- [6] G. Schechtman, M. Schmuckenschlger, *A concentration inequality for harmonic measures on the sphere*. Geometric aspects of functional analysis (Israel, 1992–1994), 255–273. Oper. Theory Adv. Appl. **77**, Birkhuser, Basel, 1995.
- [7] R. Schneider, *Convex bodies: the Brunn-Minkowski theory*. Cambridge University Press, 1993.
- [8] Z. Šidák, *Rectangular Confidence Regions for the Means of Multivariate Normal Distributions*. J. Amer. Stat. Association, **62** (1967), no. 318, 626–633.
- [9] Z. Šidák, *On Multivariate Normal Probabilities of Rectangles: Their Dependence on Correlations*. Ann. Math. Stat., **39** (1968), no. 5, 1425–1434.

E-mail address, G. Ambrus: ambrus@renyi.hu

ALFRÉD RÉNYI INSTITUTE OF MATHEMATICS, HUNGARIAN ACADEMY OF SCIENCES,
REÁLTANODA U. 13-15, 1053 BUDAPEST, HUNGARY