## Another proof of $\zeta(2) = \frac{\pi^2}{6}$ using double integrals

Daniele Ritelli \*

Abstract

Starting from the double integral

$$\int_0^\infty \int_0^\infty \frac{\mathrm{d}x\mathrm{d}y}{(1+y)(1+x^2y)}$$

we give another solution to the Basel Problem

$$\zeta(2) = \sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{\pi^2}{6}.$$

2000 AMS Classification 40A25

KEYWORD: Basel Problem, Double integrals, Geometric Series Monotone Convergence Theorem

NOTE: To appear on American Mathematical Monthly

The celebrated Euler identity, known as the Basel Problem,

$$\sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{\pi^2}{6} \tag{1}$$

has been proved in many different ways. In this note, we focus on the derivation of (1), taking advantage of the nice interplay between a double integral and a geometric series as has appeared in several articles on this subject, [1, 5, 3]. As Sir Michael Atiyah declared in an interview [8]:

Any good theorem should have several proofs, the more the better. For two reasons: usually, different proofs have different strengths and weaknesses, and they generalize in different directions: they are not just repetitions of each other

we find that there is always something worthy of attention in a new proof of a known result. Here, we provide a proof that uses a rational function with the lowest degree among the functions used in different proofs of the same kind.

The author who inaugurated this approach was Apostol, [1], inspired by Beukers' paper [2], where the double integral

$$\int_{0}^{1} \int_{0}^{1} \frac{dxdy}{1 - xy}$$
(2)

was utilized to prove the irrationality of  $\zeta(2)$ . Apostol, instead, evaluated (2) in two ways. First, by expanding 1/(1 - xy) into a geometric series and then, with a change of variables corresponding to the rotation of the coordinate axes through the angle  $\pi/4$  radians. By equating the expression so obtained, the value of  $\zeta(2)$  is found. It is worth noting that for the second evaluation, one has to compute the elaborate integrals

$$I_{1} = \int_{0}^{1/\sqrt{2}} \frac{1}{\sqrt{2 - u^{2}}} \arctan\left(\frac{u}{\sqrt{2 - u^{2}}}\right) du, \text{ and}$$
$$I_{2} = \int_{1/\sqrt{2}}^{\sqrt{2}} \frac{1}{\sqrt{2 - u^{2}}} \arctan\left(\frac{\sqrt{2} - u}{\sqrt{2 - u^{2}}}\right) du$$

<sup>\*</sup>Dipartimento di Scienze Statistiche, Università di Bologna daniele.ritelli@unibo.it

Another proof of  $\zeta(2) = \frac{\pi^2}{6}$  using double integrals

using suitable trigonometric changes of variables. The evaluation of Beukers, Calabi and Kolk, [3], is similar. They expand  $1/(1 - x^2y^2)$  into a geometric series to obtain the  $\zeta(2)$  series, after which they introduce the two dimensional trigonometric changes of variables

$$x = \frac{\sin u}{\cos v}, \quad y = \frac{\sin v}{\cos u}$$

to evaluate the double integral. The proof of Hirschhon [6] stems from the double inequality

$$2\left(\arctan\frac{a}{1+\sqrt{1-a^2}}\right)^2 < \sum_{n=0}^{\infty} \frac{a^{4n+2}}{(2n+1)^2} < 2\left(\arctan a\right)^2, \ 0 < a < 1,$$
(3)

and a passage to the limit as  $a \to 1$ . The derivation of (3) is based on an integral inequality with regard again to the function  $f(x, y) = 1/(1 - x^2y^2)$ . To obtain (3), two integrals of f(x, y) over two different regions of the plane are computed. All these approaches have in common the need to remove a singularity at the point (1,1) of the integrand. Our proof is inspired by [5], where another definite integral

$$\int_{0}^{+\infty} \int_{0}^{1} \frac{x}{(1+x^2)(1+x^2y^2)} \mathrm{d}x \mathrm{d}y \tag{4}$$

is computed, first by integrating with respect to x and then with respect to y and vice versa. The same integral is considered in the probabilistic proof given in [7], where integral (4) comes from the product of two positive Cauchy random variables. The proof of [5], as well our proof, uses functions with no singularity in the domain of integration, so we can consider that in same sense these proofs are simpler. Moreover, our proof uses a lower degree rational function than the one used in [5].

Our starting point, as with most of the papers on this subject, is that (1) is equivalent to

$$\sum_{n=0}^{\infty} \frac{1}{(2n+1)^2} = \frac{\pi^2}{8}.$$
(5)

In our proof we will show (5) starting from the double integral

$$\int_0^\infty \int_0^\infty \frac{dxdy}{(1+y)(1+x^2y)}.$$
 (6)

If we integrate (6) first with respect to *x* and then to *y*, we find that:

$$\int_{0}^{\infty} \left(\frac{1}{1+y} \int_{0}^{\infty} \frac{dx}{1+x^{2}y}\right) dy = \int_{0}^{\infty} \left(\frac{1}{1+y} \left[\frac{\arctan(\sqrt{y}x)}{\sqrt{y}}\right]_{x=0}^{x=\infty}\right) dy$$

$$= \frac{\pi}{2} \int_{0}^{\infty} \frac{dy}{\sqrt{y}(1+y)} = \frac{\pi}{2} \int_{0}^{\infty} \frac{2u}{u(1+u^{2})} du = \frac{\pi^{2}}{2}$$
(7)

where we used the change of variable  $y = u^2$  in the last step. Reversing the order of integration yields

$$\int_{0}^{\infty} \left( \int_{0}^{\infty} \frac{\mathrm{d}y}{(1+y)(1+x^{2}y)} \right) \mathrm{d}x = \int_{0}^{\infty} \frac{1}{1-x^{2}} \left( \int_{0}^{\infty} \left( \frac{1}{1+y} - \frac{x^{2}}{1+x^{2}y} \right) \mathrm{d}y \right) \mathrm{d}x$$
$$= \int_{0}^{\infty} \frac{1}{1-x^{2}} \ln \frac{1}{x^{2}} \mathrm{d}x = 2 \int_{0}^{\infty} \frac{\ln x}{x^{2}-1} \mathrm{d}x.$$
(8)

Hence, equating (7) and (8) we get

$$\int_0^\infty \frac{\ln x}{x^2 - 1} \mathrm{d}x = \frac{\pi^2}{4}.$$
(9)

Now split the integration domain in (9) between [0, 1] and  $[1, \infty)$  and change the variable x = 1/u in the second integral, so that

$$\int_{0}^{\infty} \frac{\ln x}{x^{2} - 1} dx = \int_{0}^{1} \frac{\ln x}{x^{2} - 1} dx + \int_{1}^{\infty} \frac{\ln x}{x^{2} - 1} dx$$

$$= \int_{0}^{1} \frac{\ln x}{x^{2} - 1} dx + \int_{0}^{1} \frac{\ln u}{u^{2} - 1} du.$$
(10)

From (9) and (10) we get

$$\int_0^1 \frac{\ln x}{x^2 - 1} \mathrm{d}x = \frac{\pi^2}{8}.$$
(11)

Equation (5) now follows, expanding, as in [5], the denominator of the integrand on the left hand side of (11) into a geometric series and using the Monotone Convergence Theorem (see [4] pp. 95-96). Thus, we have:

$$\int_0^1 \frac{\ln x}{x^2 - 1} dx = \int_0^1 \frac{-\ln x}{1 - x^2} dx = \sum_{n=0}^{+\infty} \int_0^1 (-x^{2n} \ln x) dx.$$
 (12)

Integrating by parts yields

$$\int_0^1 (-x^{2n} \ln x) \, \mathrm{d}x = \left[ -\frac{x^{2n+1}}{2n+1} \ln x \right]_0^1 + \int_0^1 \frac{x^{2n}}{2n+1} \, \mathrm{d}x = \frac{1}{(2n+1)^2} \tag{13}$$

so that considering (13), we can write (12) as

$$\int_0^1 \frac{\ln x}{x^2 - 1} \mathrm{d}x = \sum_{n=0}^{+\infty} \frac{1}{(2n+1)^2} \tag{14}$$

and we are done equating (11) and (14).

## References

- [1] T. Apostol, A proof that Euler missed: Evaluating  $\zeta(2)$  the easy way, *Math. Intelligencer* **5** (1983) 59–60.
- [2] F. Beukers, A note on the irrationality of  $\zeta(2)$  and  $\zeta(3)$ , Bull. London Math. Soc. 11 (1979) 268–272
- [3] F. Beukers; E. Calabi; J.A.C. Kolk, Sums of generalized harmonic series and volumes, *Nieuw Arch. Wisk.* 11 (1993) 217–224.
- [4] M. Capiński; E. Kopp, Measure, Integral and Probability second edition, Springer, New York, 2004.
- [5] J.D. Harper, Another Simple Proof of  $1 + \frac{1}{2^2} + \frac{1}{3^2} + \cdots = \frac{\pi^2}{6}$ , Amer. Math. Monthly **110** (2003) 540–541.
- [6] M.D. Hirschhon, A simple Proof that  $\zeta(2) = \frac{\pi^2}{6}$ , Math. Intelligencer 22 (2011) 81–82.
- [7] L. Pace, Probabilistically Proving that  $\zeta(2) = \frac{\pi^2}{6}$ , Amer. Math. Monthly **118** (2011) 641–643.
- [8] M. Raussen; C. Skau, Interview with Michael Atiyah and Isadore Singer, *Notices Amer. Math. Soc.* **52** (2005) 225–233.

Dipartimento di Statistica, Università di Bologna, Viale Filopanti, 5 Bologna daniele.ritelli@unibo.it