# The distribution function of the distance between two random points in a right-angled triangle

Uwe Bäsel

#### Abstract

In this paper we obtain the density function and the distribution function of the distance between two uniformly and independently distributed random points in any right-angled triangle. The density function is derived from the chord length distribution function using Piefke's formula. We conclude results for random distances between two congruent right-angled triangles together forming a rectangle.

2010 Mathematics Subject Classification: 60D05, 52A22

**Keywords:** Geometric probability, random sets, integral geometry, chord length distribution function, random distances, distance distribution function, right-angled triangles, right triangle, Piefke formula

#### **1** Introduction

A random line g intersecting a convex set  $\mathcal{K}$  in the plane produces a chord of  $\mathcal{K}$ . The length s of this chord is a random variable. The chord length distribution function of a regular triangle was calculated by Sulanke [14, p. 57]. Harutyunyan and Ohanyan [8] calculated the chord length distribution function for regular polygons (see also Bäsel [2, pp. 2-9]).

The distance t between two points chosen independently and uniformly at random from  $\mathcal{K}$  is also a random variable. Borel [3] considered random distances in elementary geometric figures such as triangles, squares and so on (see [11, p. 163]). The expectations of the distance between two random points for an equilateral triangle and a rectangle are to be found in [13, p. 49]. Ghosh [6] derived the distance density function for a rectangle. Bäsel [2, pp. 10-16] derived the density function and the distribution function of the distance for regular polygons from its chord length distribution function using Piefke's formula [12, p. 130]. There are a lot of results concerning the distance t within a convex set or in two convex sets, see Chapter 2 in [10].

For practical applications in physics, material sciences, operations research, communication networks and other fields see [7], [9] and [15].

The first aim of the present paper is to derive the chord length distribution function for any right-angled triangle. The second aim is to conclude the distribution function of the distance between two points chosen independently and uniformly from a right-angled triangle.

 $A = (0, b), B = (a, 0) \text{ and } C = (0, 0) \text{ are the vertices of the right-angled triangle (see Fig. 1b). We assume <math>|\overline{AC}| = b \ge a = |\overline{BC}|$  and denote this triangle by  $\mathcal{T}_{a,b}$ . Clearly, it is possible to bring every right-angled triangle in such a position by use of Euclidean motions (translations and rotations) and, if necessary, a reflection. These operations do not influence the chord length distribution and the point distance distribution.  $c = |\overline{AB}|$  is the length of the hypotenuse and  $\alpha = \arctan(a/b)$  the inner angle between  $\overline{AB}$  and  $\overline{AC}$ . u = a + b + c is the length of the perimeter of  $\mathcal{T}_{a,b}, \tilde{A} = ab/2$  its area and  $h = a \cos \alpha$  the height over  $\overline{AB}$ . The hypotenuse  $\overline{AB}$  of the triangle  $\mathcal{T}_{a,b}$  is the diameter of its circumcircle (converse of Thales' theorem).

## 2 Chord length distribution function

A straight line g in the plane is determined by the angle  $\psi$ ,  $0 \le \psi < 2\pi$ , that the direction perpendicular to g makes with a fixed direction (e.g. the x-axis) and by its distance  $p, 0 \le p < \infty$ , from the origin O:

$$g = g(p, \psi) = \{(x, y) \in \mathbb{R}^2 : x \cos \psi + y \sin \psi = p\}.$$

The measure  $\mu$  of a set of lines  $g(p, \psi)$  is defined by the integral, over the set, of the differential form  $dg = dp d\psi$ . Up to a constant factor, this measure is the only one that is invariant under motions in the Euclidean plane [13, p. 28].

We use the chord length distribution function of  $\mathcal{T}_{a,b}$  in the form

$$F(s) = \frac{1}{u} \mu(\{g : g \cap \mathcal{T}_{a,b} \neq \emptyset, |\chi(g)| \le s\}),$$

where  $\chi(g) = g \cap \mathcal{T}_{a,b}$  is the chord of  $\mathcal{T}_{a,b}$ , produced by the line g, and  $|\chi(g)|$  the length of  $\chi(g)$  (see [5, p. 863] and [1, p. 161]). (The measure of all lines g that intersect a convex set is equal to its perimeter [13, p. 30].) So it remains to calculate the measure of all lines that produce a chord of length  $|\chi(g)| \leq s$ . Using the abbreviation

$$\mathcal{S}(g,s) := \{g: g \cap \mathcal{T}_{a,b} \neq \emptyset, |\chi(g)| \le s\},\$$

we have

$$\mu(\mathcal{S}(g,s)) = \int_{\mathcal{S}(g,s)} \mathrm{d}g = \int_{\mathcal{S}(g,s)} \mathrm{d}p \,\mathrm{d}\psi.$$

Instead of the angle  $\psi$  it is possible to use the angle  $\phi$  that a line makes with the positive x-axis, hence

$$\mu(\mathcal{S}(g,s)) = \int_{\mathcal{S}(g,s)} \mathrm{d}p \,\mathrm{d}\phi \,.$$

**Theorem 1.** The chord length distribution function F of  $\mathcal{T}_{a,b}$  is given by

$$F(s) = \begin{cases} 0 & \text{if } s < 0, \\ H_1(s)/u & \text{if } 0 \le s < h, \\ H_2(s)/u & \text{if } h \le s < a, \\ H_3(s)/u & \text{if } a \le s < b, \\ H_4(s)/u & \text{if } b \le s < c, \\ 1 & \text{if } s \ge c, \end{cases}$$

where

$$\begin{split} H_1(s) &= \frac{\Theta_1 s}{4} \,, \\ H_2(s) &= \frac{\Theta_2 s}{4} + \Theta_3 L_1(s,h) + c L_2(s,h) \,, \\ H_3(s) &= a + \Theta_4 s + \frac{\Theta_3}{2} L_1(s,h) + \frac{c}{2} L_2(s,h) + \frac{b}{2a} L_1(s,a) + \frac{b}{2} L_2(s,a) \,, \\ H_4(s) &= \Theta_5 + \frac{\Theta_6 s}{4} + \frac{b}{2a} L_1(s,a) + \frac{b}{2} L_2(s,a) + \frac{a}{2b} L_1(s,b) + \frac{a}{2} L_2(s,b) \end{split}$$

with

$$\Theta_{1} = \frac{a}{b} (2\alpha + \pi) + \frac{2b}{a} (\pi - \alpha) + 6 , \quad \Theta_{2} = \frac{a}{b} (2\alpha - \pi) - \frac{2b}{a} \alpha + 6 ,$$
  
$$\Theta_{3} = \frac{a}{b} + \frac{b}{a} , \quad \Theta_{4} = 1 - \frac{b}{a} \alpha , \quad \Theta_{5} = a + b , \quad \Theta_{6} = \frac{a}{b} (2\alpha - \pi) - \frac{2b}{a} \alpha + 2$$

and

$$L_1(s,a) = s \arcsin \frac{a}{s}$$
,  $L_2(s,a) = \sqrt{1 - \left(\frac{a}{s}\right)^2}$ .

*Proof.* In the following we need the width w of  $\mathcal{T}_{a,b}$  in the direction perpendicular to the angle  $\phi$  that is given by:

$$w(\phi) = \begin{cases} w_1(\phi) & \text{if } 0 \le \phi < \pi/2, \\ w_2(\phi) & \text{if } \pi/2 \le \phi < \pi/2 + \alpha, \\ w_3(\phi) & \text{if } \pi/2 + \alpha \le \phi \le \pi, \end{cases}$$

where

$$w_1(\phi) = a \frac{\cos(\phi - \alpha)}{\sin \alpha}$$
,  $w_2(\phi) = a \sin \phi$ ,  $w_3(\phi) = -a \frac{\cos \phi}{\tan \alpha}$ .

We have to distinguish the cases  $0 \le s < h$ ,  $h \le s < a$ ,  $a \le s < b$  and  $b \le s < c$ , and put

$$\begin{split} H_1(s) &= \mu(\mathcal{S}(g,s)) \quad \text{if} \quad 0 \leq s < h \ , \quad H_2(s) = \mu(\mathcal{S}(g,s)) \quad \text{if} \quad h \leq s < a \ , \\ H_3(s) &= \mu(\mathcal{S}(g,s)) \quad \text{if} \quad a \leq s < b \ , \quad H_4(s) = \mu(\mathcal{S}(g,s)) \quad \text{if} \quad b \leq s < c \ . \end{split}$$

We demonstrate the calculation of  $H_2$ . Here it is necessary to distinguish the following intervals of  $\phi$  (see Fig. 1):

a)  $0 \leq \phi < \phi_1 = \alpha - \arccos((a/s) \cos \alpha)$ : All lines g with fixed angle  $\phi$  produce a chord  $\chi$  of length  $|\chi(g)| \leq s$  if they are lying in one of the two strips of breadth

$$b_1(s,\phi) = \frac{s}{2} \frac{\sin(2\phi - \alpha) + \sin\alpha}{\cos\alpha}$$
 and  $b_2(s,\phi) = \frac{s}{2} \frac{\cos(2\phi - \alpha) + \cos\alpha}{\sin\alpha}$ 

respectively.

- b)  $\phi_1 \leq \phi < \phi_2 = \alpha + \arccos((a/s) \cos \alpha)$ : All lines g with fixed angle  $\phi$ and  $g \cap \mathcal{T}_{a,b} \neq \emptyset$  produce a chord of length  $|\chi(g)| \leq s$ . These lines are lying in a strip of breadth  $w_1(\phi)$ .
- c)  $\phi_2 \leq \phi < \pi/2$ : See  $0 \leq \phi < \phi_1$ .
- d)  $\pi/2 \leq \phi < \pi/2 + \alpha$ : All lines g with fixed angle  $\phi$  produce a chord of length  $\leq s$  if they are lying in one of the two strips of breadth  $b_1(s, \phi)$  and

$$b_3(s,\phi) = -\frac{s}{2}\sin 2\phi$$

respectively.

e)  $\pi/2 + \alpha \leq \phi < \pi$ : All lines g with fixed angle  $\phi$ , lying in one of the two strips of breadth  $b_2(s, \phi)$  and  $b_3(s, \phi)$  repectively, produce a chord of length  $\leq s$ .

Therefore, using the abbreviations  $b_i$  and  $w_i$  for  $b_i(s, \phi)$  and  $w_i(\phi)$  respectively,

$$H_2(s) = \int_{\mathcal{S}(g,s)} dp \, d\phi = \int_0^{\phi_1} (b_1 + b_2) \, d\phi + \int_{\phi_1}^{\phi_2} w_1 \, d\phi + \int_{\phi_2}^{\pi/2} (b_1 + b_2) \, d\phi + \int_{\pi/2}^{\pi/2+\alpha} (b_1 + b_3) \, d\phi + \int_{\pi/2+\alpha}^{\pi} (b_2 + b_3) \, d\phi.$$

With the angles

$$\phi_3 = \pi - \arccos \frac{a}{s}$$
 and  $\phi_4 = \pi - \arcsin \left(\frac{a}{s} \cot \alpha\right)$ 

we analogously get

$$H_{1}(s) = \int_{0}^{\pi/2} (b_{1} + b_{2}) d\phi + \int_{\pi/2}^{\pi/2+\alpha} (b_{1} + b_{3}) d\phi + \int_{\pi/2+\alpha}^{\pi} (b_{2} + b_{3}) d\phi$$
$$H_{3}(s) = \int_{0}^{\phi_{2}} w_{1} d\phi + \int_{\phi_{2}}^{\pi/2} (b_{1} + b_{2}) d\phi + \int_{\pi/2}^{\pi/2+\alpha} (b_{1} + b_{3}) d\phi$$
$$+ \int_{\pi/2+\alpha}^{\phi_{3}} (b_{2} + b_{3}) d\phi + \int_{\phi_{3}}^{\pi} w_{3} d\phi,$$




Figure 1: Calculation of  $H_2$ 

$$H_4(s) = \int_0^{\pi/2} w_1 \,\mathrm{d}\phi + \int_{\pi/2}^{\phi_4} w_2 \,\mathrm{d}\phi + \int_{\phi_4}^{\pi/2+\alpha} (b_1 + b_3) \,\mathrm{d}\phi + \int_{\pi/2+\alpha}^{\phi_3} (b_2 + b_3) \,\mathrm{d}\phi + \int_{\phi_3}^{\pi} w_3 \,\mathrm{d}\phi \,.$$

Using the indefinite integrals

$$\int b_1(s,\phi) d\phi = \frac{s}{4} \frac{2\phi \sin \alpha - \cos(2\phi - \alpha)}{\cos \alpha},$$
  
$$\int b_2(s,\phi) d\phi = \frac{s}{4} \frac{2\phi \cos \alpha + \sin(2\phi - \alpha)}{\sin \alpha},$$
  
$$\int b_3(s,\phi) d\phi = \frac{s}{2} \cos^2 \phi,$$
  
$$\int w_1(\phi) d\phi = \frac{a \sin(\phi - \alpha)}{\sin \alpha} = a(\cot \alpha \sin \phi - \cos \phi),$$
  
$$\int w_2(\phi) d\phi = -a \cos \phi, \quad \int w_3(\phi) d\phi = -\frac{a \sin \phi}{\tan \alpha},$$

one finds the result of Theorem 1.

**Remark.** It is easy to derive the chord length density function from Theorem 1.

# 3 Point distances

**Theorem 2.** The density function g of the distance t between two random points in  $\mathcal{T}_{a,b}$  is given by

$$g(t) = \begin{cases} \frac{2t}{\tilde{A}} \left[ \pi + \frac{1}{\tilde{A}} \left( J^*(t) - ut \right) \right] & \text{if } 0 \le t \le c \,, \\ 0 & \text{else} \,, \end{cases}$$

where

$$J^{*}(t) = \begin{cases} J_{1}^{*}(0,t) & \text{if } 0 \leq t < h, \\ J_{1}^{*}(0,h) + J_{2}^{*}(h,t) & \text{if } h \leq t < a, \\ J_{1}^{*}(0,h) + J_{2}^{*}(h,a) + J_{3}^{*}(a,t) & \text{if } a \leq t < b, \\ J_{1}^{*}(0,h) + J_{2}^{*}(h,a) + J_{3}^{*}(a,b) + J_{4}^{*}(b,t) & \text{if } b \leq t \leq c \end{cases}$$

with

$$J_k^*(s,t) = H_k^*(t) - H_k^*(s) \,,$$

where

$$\begin{split} H_1^*(t) &= \frac{\Theta_1 t^2}{8}, \\ H_2^*(t) &= \frac{\Theta_2 t^2}{8} + \Theta_3 L_1^*(t,h) + c L_2^*(t,h), \\ H_3^*(t) &= at + \frac{\Theta_4 t^2}{2} + \frac{\Theta_3}{2} L_1^*(t,h) + \frac{c}{2} L_2^*(t,h) + \frac{b}{2a} L_1^*(t,a) + \frac{b}{2} L_2^*(t,a), \\ H_4^*(t) &= \Theta_5 t + \frac{\Theta_6 t^2}{8} + \frac{b}{2a} L_1^*(t,a) + \frac{b}{2} L_2^*(t,a) + \frac{a}{2b} L_1^*(t,b) + \frac{a}{2} L_2^*(t,b) \end{split}$$

with

$$L_1^*(t,a) = \frac{1}{2} \left( a\sqrt{t^2 - a^2} + t^2 \arcsin\frac{a}{t} \right), \quad L_2^*(t,a) = \sqrt{t^2 - a^2} + a \arcsin\frac{a}{t}.$$

*Proof.* According to a theorem of Piefke [12, p. 130], the density function of the distance between two random points is given by

$$g(t) = \frac{2ut}{\tilde{A}^2} \int_t^c (s-t)f(s) \,\mathrm{d}s \,,$$

where f is the density function of the chord length. Following [2, pp. 11/12], we obtain

$$g(t) = \frac{2t}{\tilde{A}} \left[ \pi - \frac{u}{\tilde{A}} \left( t - \int_0^t F(s) \, \mathrm{d}s \right) \right] = \frac{2t}{\tilde{A}} \left[ \pi + \frac{1}{\tilde{A}} \left( J^*(t) - ut \right) \right]$$

with

$$J^*(t) := u \int_0^t F(s) \,\mathrm{d}s \,.$$

This yields for  $0 \le t < h$ 

$$J^{*}(t) = \int_{0}^{t} H_{1}(s) \, \mathrm{d}s = H_{1}^{*}(t) - H_{1}^{*}(0) = J_{1}^{*}(0, t) \, .$$

and for  $h \leq t < a$ 

$$J^*(t) = u \int_0^t F(s) \, \mathrm{d}s = \int_0^h H_1(s) \, \mathrm{d}s + \int_h^t H_2(s) \, \mathrm{d}s$$
$$= H_1^*(h) - H_1^*(0) + H_2^*(t) - H_2^*(h) = J_1^*(0,h) + J_2^*(h,t)$$

The calculations of  $J^*(t)$  for the intervals  $a \le t < b$  and  $b \le t < c$  can be carried out in the same manner.

Now we determine the indefinite integrals of  $L_1(t, a)$  and  $L_2(t, a)$ . Using integration by parts, we find

$$L_1^*(t,a) = \int L_1(t,a) dt = \int t \arcsin\frac{a}{t} dt$$
$$= \frac{1}{2} \left( t^2 \arcsin\frac{a}{t} + a \int \frac{t}{\sqrt{t^2 - a^2}} dt \right).$$

Clearly,

$$\int \frac{t}{\sqrt{t^2 - a^2}} \, \mathrm{d}t = \sqrt{t^2 - a^2}$$

and therefore

$$L_1^*(t,a) = \frac{1}{2} \left( a \sqrt{t^2 - a^2} + t^2 \arcsin \frac{a}{t} \right).$$

Since  $t \ge a > 0$  in the present cases,

$$L_2^*(t,a) = \int L_2(t,a) \, \mathrm{d}t = \int \sqrt{1 - \left(\frac{a}{t}\right)^2} \, \mathrm{d}t = \int \frac{\sqrt{t^2 - a^2}}{t} \, \mathrm{d}t \,,$$

and (cf. [4, p. 48, Eq. 217])

$$L_2^*(t,a) = \sqrt{t^2 - a^2} + a \arcsin \frac{a}{t}$$
.

**Corollary 1.** The distribution function G of the distance t between two random points in  $\mathcal{T}_{a,b}$  is given by

$$G(t) = \begin{cases} 0 & \text{if } t < 0, \\ \frac{1}{\tilde{A}} \left[ t^2 \left( \pi - \frac{2u}{3\tilde{A}} t \right) + \frac{2}{\tilde{A}} J^{\natural}(t) \right] & \text{if } 0 \le t < c, \\ 1 & \text{if } t \ge c \end{cases}$$

with

$$J^{\natural}(t) = \begin{cases} K_1(t) & \text{if } 0 \le t < h \,, \\ K_1(h) + K_2(t) & \text{if } h \le t < a \,, \\ K_1(h) + K_2(a) + K_3(t) & \text{if } a \le t < b \,, \\ K_1(h) + K_2(a) + K_3(b) + K_4(t) & \text{if } b \le t < c \,, \end{cases}$$

where

$$\begin{split} K_1(t) &= J_1^{\natural}(0,t) \,, \\ K_2(t) &= \frac{1}{2} \left( t^2 - h^2 \right) \left[ J_1^*(0,h) - H_2^*(h) \right] + J_2^{\natural}(h,t) \,, \\ K_3(t) &= \frac{1}{2} \left( t^2 - a^2 \right) \left[ J_1^*(0,h) + J_2^*(h,a) - H_3^*(a) \right] + J_3^{\natural}(a,t) \,, \\ K_4(t) &= \frac{1}{2} \left( t^2 - b^2 \right) \left[ J_1^*(0,h) + J_2^*(h,a) + J_3^*(a,b) - H_4^*(b) \right] + J_4^{\natural}(b,t) \end{split}$$

with  $J_k^*$  and  $H_k^*$  according to Theorem 2 and

$$J_k^{\natural}(s,t) = H_k^{\natural}(t) - H_k^{\natural}(s) \,,$$

where

$$\begin{split} H_1^{\natural}(t) &= \frac{\Theta_1 t^4}{32} \,, \\ H_2^{\natural}(t) &= \frac{\Theta_2 t^4}{32} + \Theta_3 \, L_1^{\natural}(t,h) + c \, L_2^{\natural}(t,h) \,, \\ H_3^{\natural}(t) &= \frac{a t^3}{3} + \frac{\Theta_4 t^4}{8} + \frac{\Theta_3}{2} \, L_1^{\natural}(t,h) + \frac{c}{2} \, L_2^{\natural}(t,h) + \frac{b}{2a} \, L_1^{\natural}(t,a) + \frac{b}{2} \, L_2^{\natural}(t,a) \,, \\ H_4^{\natural}(t) &= \frac{\Theta_5 t^3}{3} + \frac{\Theta_6 t^4}{32} + \frac{b}{2a} \, L_1^{\natural}(t,a) + \frac{b}{2} \, L_2^{\natural}(t,a) + \frac{a}{2b} \, L_1^{\natural}(t,b) + \frac{a}{2} \, L_2^{\natural}(t,b) \end{split}$$

$$\begin{split} L_1^{\natural}(t,a) &= \frac{1}{8} \left[ \frac{5a}{3} \left( t^2 - a^2 \right)^{3/2} + a^3 \sqrt{t^2 - a^2} + t^4 \arcsin \frac{a}{t} \right] \,, \\ L_2^{\natural}(t,a) &= \frac{1}{3} \left( t^2 - a^2 \right)^{3/2} + \frac{a}{2} \left( a \sqrt{t^2 - a^2} + t^2 \arcsin \frac{a}{t} \right) \,. \end{split}$$

Proof. For  $0 \le t < c$  one gets

$$\begin{aligned} G(t) &= \int_0^t g(\tau) \,\mathrm{d}\tau = \int_0^t \left( \frac{2\pi\tau}{\tilde{A}} - \frac{2u\tau^2}{\tilde{A}^2} + \frac{2u\tau}{\tilde{A}^2} \int_0^\tau F(s) \,\mathrm{d}s \right) \mathrm{d}\tau \\ &= \frac{\pi t^2}{\tilde{A}} - \frac{2ut^3}{3\tilde{A}^2} + \frac{2u}{\tilde{A}^2} \int_0^t \tau \left( \int_0^\tau F(s) \,\mathrm{d}s \right) \mathrm{d}\tau \\ &= \frac{\pi t^2}{\tilde{A}} - \frac{2ut^3}{3\tilde{A}^2} + \frac{2}{\tilde{A}^2} \int_0^t \tau J^*(\tau) \,\mathrm{d}\tau = \frac{1}{\tilde{A}} \left[ t^2 \left( \pi - \frac{2u}{3\tilde{A}} t \right) + \frac{2}{\tilde{A}} J^{\natural}(t) \right] \end{aligned}$$

with

$$J^{\natural}(t) := \int_0^t s J^*(s) \,\mathrm{d}s \,.$$

It remains to calculate  $J^{\natural}(t)$ . For  $0 \leq t < h$  we have

$$J^{\natural}(t) = \int_{0}^{t} s J_{1}^{*}(0,s) \, \mathrm{d}s = \int_{0}^{t} s \left[ H_{1}^{*}(s) - H_{1}^{*}(0) \right] \mathrm{d}s$$
  
=  $H_{1}^{\natural}(t) - H_{1}^{\natural}(0) - \frac{1}{2} t^{2} H_{1}^{*}(0) = H_{1}^{\natural}(t) - H_{1}^{\natural}(0) - 0 = J_{1}^{\natural}(0,t) = K_{1}(t) \, .$ 

For  $h \leq t < a$  we find

$$J^{\natural}(t) = K_1(h) + \int_h^t s \left[ J_1^*(0,h) + J_2^*(h,s) \right] ds$$
  
=  $K_1(h) + \frac{1}{2} \left( t^2 - h^2 \right) J_1^*(0,h) + \int_h^t s J_2^*(h,s) ds$ 

With

$$\int_{h}^{t} s J_{2}^{*}(h,s) \, \mathrm{d}s = \int_{h}^{t} s \left[ H_{2}^{*}(s) - H_{2}^{*}(h) \right] \mathrm{d}s = \int_{h}^{t} s H_{2}^{*}(s) \, \mathrm{d}s - H_{2}^{*}(h) \int_{h}^{t} s \, \mathrm{d}s$$

$$\begin{split} &= H_2^{\natural}(t) - H_2^{\natural}(h) - \frac{1}{2} \left( t^2 - h^2 \right) H_2^{*}(h) \\ &= J_2^{\natural}(h,t) - \frac{1}{2} \left( t^2 - h^2 \right) H_2^{*}(h) \end{split}$$

it follows that

$$J^{\natural}(t) = K_1(h) + \frac{1}{2} \left( t^2 - h^2 \right) \left[ J_1^*(0,h) - H_2^*(h) \right] + J_2^{\natural}(h,t) = K_1(h) + K_2(t) \,.$$

The calculation applies analogously to the intervals  $a \le t < b$  and  $b \le t < c$ .

Now we calculate the indefinite integrals of  $L_1^*(t, a)$  and  $L_2^*(t, a)$ . We have

$$L_{1}^{\natural}(t,a) = \int t L_{1}^{*}(t,a) \, \mathrm{d}t = \frac{1}{2} \int t \left( a \sqrt{t^{2} - a^{2}} + t^{2} \arcsin \frac{a}{t} \right) \mathrm{d}t$$
$$= \frac{1}{2} \left( a \int t \sqrt{t^{2} - a^{2}} \, \mathrm{d}t + \int t^{3} \arcsin \frac{a}{t} \, \mathrm{d}t \right)$$

with

$$\int t \sqrt{t^2 - a^2} \, \mathrm{d}t = \frac{1}{3} \left( t^2 - a^2 \right)^{3/2} \quad [4, \, \mathrm{p.} \, 47, \, \mathrm{Eq.} \, 214] \,.$$

Using integration by parts, we find

$$\int t^3 \arcsin\frac{a}{t} \, \mathrm{d}t = \frac{1}{4} \left( t^4 \arcsin\frac{a}{t} + a \int \frac{t^2}{\sqrt{1 - (a/t)^2}} \, \mathrm{d}t \right).$$

Since  $t \ge a > 0$  in the present cases,

$$\int \frac{t^2}{\sqrt{1 - (a/t)^2}} \, \mathrm{d}t = \int \frac{t^3}{\sqrt{t^2 - a^2}} \, \mathrm{d}t \,,$$

and

$$\int \frac{t^3}{\sqrt{t^2 - a^2}} \, \mathrm{d}t = \frac{1}{3} \left( t^2 - a^2 \right)^{3/2} + a^2 \sqrt{t^2 - a^2} \quad [4, \, \mathrm{p.} \, 48, \, \mathrm{Eq.} \, 223] \,.$$

This yields

$$L_1^{\natural}(t,a) = \frac{1}{8} \left[ \frac{5a}{3} \left( t^2 - a^2 \right)^{3/2} + a^3 \sqrt{t^2 - a^2} + t^4 \arcsin \frac{a}{t} \right].$$

Further,

$$L_{2}^{\natural}(t,a) = \int t L_{2}^{*}(t,a) dt = \int t \sqrt{t^{2} - a^{2}} dt + a \int t \arcsin \frac{a}{t} dt$$
$$= \frac{1}{3} \left( t^{2} - a^{2} \right)^{3/2} + \frac{a}{2} \left( a \sqrt{t^{2} - a^{2}} + t^{2} \arcsin \frac{a}{t} \right)$$

(see the calculations of  $L_1^{\natural}(t,a)$  and  $L_1^{*}(t,a)$ ).

### 4 Two triangles

Now considering a rectangle  $\mathcal{R}_{a,b}$  of side lengths a and b ( $a \leq b$ ) as made up of two congruent copies of the triangle  $\mathcal{T}_{a,b}$  with different orientation and sharing their hypotenuses. Let  $g_2$  and  $G_2$  denote respectively the density function and the distribution function of the distance between two uniformly and independently distributed points  $P_1$  and  $P_2$ , one in each of the triangles.

Using the *indirect method* (see Ghosh [6, pp. 22/23]), it is possible to calculate  $g_2$  from the density function  $g_{\mathcal{R}}$  of the distance between two random points inside  $\mathcal{R}_{a,b}$  and the density function g of  $\mathcal{T}_{a,b}$  (according to Theorem 2). The probability of  $P_1$  and  $P_2$  belonging to the same triangle is equal to the probability of  $P_1$  and  $P_2$  belonging to different triangles, each probability being 1/2. Therefore,

$$g_{\mathcal{R}} = \frac{1}{2} (g + g_2)$$
, whence  $g_2 = 2 g_{\mathcal{R}} - g$  and  $G_2 = 2 G_{\mathcal{R}} - G$ ,

where  $G_{\mathcal{R}}$  denotes the corresponding distribution function of  $g_{\mathcal{R}}$ . As already mentioned,  $g_{\mathcal{R}}$  was found by Ghosh [6] (Theorem 2, pp. 18/19). Denoting by  $A_{\mathcal{R}} = ab$  the area of  $\mathcal{R}_{a,b}$  and by  $u_{\mathcal{R}} = 2(a+b)$  its perimeter, we can write Ghosh's result as

$$g_{\mathcal{R}}(t) = \begin{cases} \frac{2t}{A_{\mathcal{R}}^2} J_{\mathcal{R}}^*(t) & \text{if } 0 \le t \le c = \sqrt{a^2 + b^2}, \\ 0 & \text{else}, \end{cases}$$

where

$$J_{\mathcal{R}}^{*}(t) = \begin{cases} H_{\mathcal{R},1}^{*}(t) & \text{if } 0 \le t < a, \\ H_{\mathcal{R},2}^{*}(t) & \text{if } a \le t < b, \\ H_{\mathcal{R},3}^{*}(t) & \text{if } b \le t \le c \end{cases}$$

with

$$\begin{aligned} H^*_{\mathcal{R},1}(t) &= \pi A_{\mathcal{R}} - u_{\mathcal{R}}t + t^2 \,, \\ H^*_{\mathcal{R},2}(t) &= -a^2 - 2bt + 2bL^*_2(t,a) \,, \\ H^*_{\mathcal{R},3}(t) &= -(\pi A_{\mathcal{R}} + c^2) - t^2 + 2bL^*_2(t,a) + 2aL^*_2(t,b) \end{aligned}$$

 $(L_2^\ast$  as in our Theorem 2). From this, one easily concludes the distribution function

$$G_{\mathcal{R}}(t) = \int_{-\infty}^{t} g_{\mathcal{R}}(\tau) \,\mathrm{d}\tau = \begin{cases} 0 & \text{if } t < 0, \\ \frac{2}{A_{\mathcal{R}}^2} J_{\mathcal{R}}^{\natural}(t) & \text{if } 0 \le t < c, \\ 1 & \text{if } t \ge c, \end{cases}$$

where

$$J_{\mathcal{R}}^{\natural}(t) = \begin{cases} J_{\mathcal{R},1}^{\natural}(0,t) & \text{if } 0 \le t < a , \\ J_{\mathcal{R},1}^{\natural}(0,a) + J_{\mathcal{R},2}^{\natural}(a,t) & \text{if } a \le t < b , \\ J_{\mathcal{R},1}^{\natural}(0,a) + J_{\mathcal{R},2}^{\natural}(a,b) + J_{\mathcal{R},3}^{\natural}(b,t) & \text{if } b \le t < c \end{cases}$$

with

$$J_{\mathcal{R},k}^{\natural}(s,t) = H_{\mathcal{R},k}^{\natural}(t) - H_{\mathcal{R},k}^{\natural}(s) \,,$$

where

$$\begin{aligned} H_{\mathcal{R},1}^{\natural}(t) &= \int t H_{\mathcal{R},1}^{*}(t) \, \mathrm{d}t = \frac{\pi A_{\mathcal{R}} t^{2}}{2} - \frac{u_{\mathcal{R}} t^{3}}{3} + \frac{t^{4}}{4} \,, \\ H_{\mathcal{R},2}^{\natural}(t) &= \int t H_{\mathcal{R},2}^{*}(t) \, \mathrm{d}t = -\frac{a^{2} t^{2}}{2} - \frac{2bt^{3}}{3} + 2bL_{2}^{\natural}(t,a) \,, \\ H_{\mathcal{R},3}^{\natural}(t) &= \int t H_{\mathcal{R},3}^{*}(t) \, \mathrm{d}t = -\frac{(\pi A_{\mathcal{R}} + c^{2})t^{2}}{2} - \frac{t^{4}}{4} + 2bL_{2}^{\natural}(t,a) + 2aL_{2}^{\natural}(t,b) \end{aligned}$$

with  $L_2^{\natural}$  as in Corollary 1. Now it is easy to calculate the density function  $g_2$  and the distribution function  $G_2$ . Examples are to be found in subsection 5.2.

# 5 Examples and simulation

#### 5.1 One triangle

Fig. 2 shows some density functions cg. The representation in this diagram has the advantage that cg is independent from the maximum distance c, it depends only on the ratio a/b. Fig. 3 shows the comparison between distribution functions G and corresponding simulation results (plot markers) with c = 1 and  $10^7$  pairs of random points.





Figure 2:  $\mathcal{T}_{a,a}$  (thick),  $\mathcal{T}_{a,2a}$  (thick and dashed),  $\mathcal{T}_{a,5a}$  (thin and dashed) and  $\mathcal{T}_{a,20a}$  (thin)



\_\_\_\_\_

#### 5.2 Two triangles

Fig. 4 shows the density function  $cg_2$  for b = 5a, and the corresponding functions  $cg_{\mathcal{R}}$  and cg. Fig. 5 shows the distribution function  $G_2$  for b = 5a, and the corresponding functions  $G_{\mathcal{R}}$  and G. The plot markers result from a simulation with c = 1 and  $10^7$  pairs of random points.




Figure 4:  $cg_{\mathcal{R}}$  (short-dashed), cg (long-dashed) and  $cg_2$  for b = 5a




Figure 5:  $G_{\mathcal{R}}$  (short-dashed), G (long-dashed) and  $G_2$  for b = 5a

## References

- Ambartzumjan, R. V.; Mecke, J.; Stoyan, D.: Geometrische Wahrscheinlichkeiten und Stochastische Geometrie, Akademie Verlag, Berlin, 1993.
- [2] Bäsel, U.: Random chords and point distances in regular polygons, arXiv: 1204.2707v2 [math.PR] 5 Sep 2012.
- Borel, E.: Principes et Formules Classiques du Calcul des Probabilités, Gauthier-Villars, Paris, 1925.
- [4] Bronstein, I. N.; Semendjajew, K. A.: Taschenbuch der Mathematik, Verlag Nauka, Moskau und BSB B. G. Teubner Verlagsgesellschaft, Leipzig, 1989.
- [5] Gates, J.: Some properties of chord length distributions, J. Appl. Prob., 24 (1987), 863-874.
- [6] Ghosh, B.: Random distances within a rectangle and between two rectangles, Bull. Calcutta Math. Soc., 43 (1951), 17-24.
- [7] Gille, W.: The chord length distribution of parallelepipeds with their limiting cases, *Exp. Techn. Phys.*, **36** (1988), 197-208.

- [8] Harutyunyan, H. S.; Ohanyan, V. K.: The chord length distribution function for regular polygons, Adv. Appl. Prob. (SGSA), 41 (2009), 358-366.
- [9] Marsaglia, G.; Narasimhan, B. G.; Zaman, A.: The distance between random points in rectangles, *Commun. Statist. - Theory Meth.*, **19(11)** (1990), 4199-4212.
- [10] Mathai, A. M.: An Introduction to Geometrical Probability, Gordon and Breach, Australia, 1999.
- [11] Mathai, A. M.; Moschopoulos, P.; Pederzoli, G.: Random points associated with rectangles, *Rend. Circ. Mat. Palermo*, Serie II, Tomo XLVIII (1999), 163-190.
- [12] Piefke, F.: Beziehungen zwischen der Sehnenlängenverteilung und der Verteilung des Abstandes zweier zufälliger Punkte im Eikörper, Z. Wahrscheinlichkeitstheorie verw. Gebiete, 43 (1978), 129-134.
- [13] Santaló, L. A.: Integral Geometry and Geometric Probability, Addison-Wesley, London, 1976.
- [14] Sulanke, R.: Die Verteilung der Sehnenlängen an ebenen und räumlichen Figuren, Math. Nachr., 23 (1961), 51-74.
- [15] Zhuang, Y.; Pan, J.: A geometrical probability approach to locationcritical network performance metrics, *INFOCOM*, *Proceedings IEEE*, (2012), 1817-1825.

Uwe Bäsel

HTWK Leipzig, University of Applied Sciences, Faculty of Mechanical and Energy Engineering, PF 30 11 66, 04251 Leipzig, Germany,