

Asymptotical Bounds for Complete Elliptic Integrals of the Second Kind

Miao-Kun Wang, Yu-Ming Chu

Abstract: In this paper, we establish several asymptotical bounds for the complete elliptic integrals of the second kind $\mathcal{E}(r)$, and improve the well-known conjecture $\mathcal{E}(r) > \pi[(1 + (1 - r^2)^{3/4})/2]^{2/3}/2$ for all $r \in (0, 1)$ proposed by M. Vuorinen.

Keywords: Elliptic integrals, Toader mean, Monotonicity

2010 Mathematics Subject Classification: 33E05

1. Introduction

For $0 < r < 1$, the well-known complete elliptic integrals of the first and second kinds [15, 16] are defined by

$$\mathcal{K} = \mathcal{K}(r) = \int_0^{\pi/2} (1 - r^2 \sin^2 \theta)^{-1/2} d\theta, \quad \mathcal{K}(0^+) = \pi/2, \quad \mathcal{K}(1^-) = +\infty$$

and

$$\mathcal{E} = \mathcal{E}(r) = \int_0^{\pi/2} (1 - r^2 \sin^2 \theta)^{1/2} d\theta, \quad \mathcal{E}(0^+) = \pi/2, \quad \mathcal{E}(1^-) = 1,$$

respectively.

M.-K. Wang

Department of Mathematics, Huzhou Teachers College, Huzhou 313000, China
e-mail: wmk000@126.com

Y.-M. Chu (Corresponding author)

Department of Mathematics, Huzhou Teachers College, Huzhou 313000, China
e-mail: chuyuming@hutczj.cn; tel: +86 572 2321510; fax: +86 572 2321165

It is well known that the complete elliptic integrals have many important applications in physics, engineering, geometric function theory, quasiconformal analysis, theory of mean values, number theory and other related fields [4-6, 9, 16, 24, 26, 27].

Recently, the complete elliptic integrals have attracted the attention of numerous mathematicians. In particular, many remarkable properties and inequalities for the complete elliptic integrals can be found in the literature [1-4, 7, 8, 10, 12-14, 19, 21, 22, 26, 29-33].

Let $L(1, r)$ be the arc length of an ellipse with semiaxes 1 and $r \in (0, 1)$, then

$$L(1, r) = 4\mathcal{E}(r'),$$

where $r' = \sqrt{1 - r^2}$, and in what follows we use the symbol r' to stand for $\sqrt{1 - r^2}$ for $r \in [0, 1]$. Inspired by a result of T. Muri [23], who pointed out in 1883 that $L(1, r)$ can be approximated by $2\pi[(1 + r^{3/2})/2]^{2/3}$, M. Vuorinen [28] conjectured that inequality

$$\mathcal{E}(r) > \frac{\pi}{2} \left(\frac{1 + r^{3/2}}{2} \right)^{2/3} \quad (1.1)$$

holds for all $r \in (0, 1)$. The conjecture was proved by R. W. Barnard et al. in [13]. Later, they also provided an upper bound for $\mathcal{E}(r)$ [14]

$$\mathcal{E}(r) < \frac{\pi}{2} \left(\frac{1 + r'^2}{2} \right)^{1/2}, \quad 0 < r < 1. \quad (1.2)$$

In [2], H. Alzer and S.-L. Qiu announced that (without proof) the inequality

$$\mathcal{E}(r) < \frac{\pi}{4} \left(\sqrt{1 - \bar{\alpha}r^2} + \sqrt{1 - \bar{\beta}r^2} \right) \quad (1.3)$$

holds for all $r \in (0, 1)$ with $\bar{\alpha} = 1/2 - \sqrt{2}/4$ and $\bar{\beta} = 1/2 + \sqrt{2}/4$, and stated that inequality (1.3) was offered by one of the referees. Later, H. Kazi and E. Neuman [22] gave a full proof of inequality (1.3) based on the two-point Gauss-Chebyshev quadrature formula with the remainder [11].

The purpose of this paper is to establish several asymptotical upper and lower bounds for complete elliptic integrals of the second kind, which improve inequalities (1.1) and (1.2), and give an lower bound of inequality (1.3). Our main results are the following Theorems 1.1 and 1.2.

Theorem 1.1. If $\alpha, \beta \in (0, 1/2]$, then the double inequality

$$\begin{aligned} \frac{\pi}{4} \left(\sqrt{\beta + (1 - \beta)r'^2} + \sqrt{(1 - \beta) + \beta r'^2} \right) &< \mathcal{E}(r) \\ &< \frac{\pi}{4} \left(\sqrt{\alpha + (1 - \alpha)r'^2} + \sqrt{(1 - \alpha) + \alpha r'^2} \right) \end{aligned} \quad (1.4)$$

holds for all $r \in (0, 1)$ if and only if $\beta \leq 1/2 - 2\sqrt{2(\pi^2 - 8)}/\pi^2$ and $\alpha \geq 1/2 - \sqrt{2}/4$.

Theorem 1.2. If $t_1, t_2 \in [1/2, 1]$ and $p \in [1/2, 2]$, then the double inequality

$$\begin{aligned} 2^{p-2}\pi(1+r')^{1-2p} \left\{ [t_1 + (1-t_1)r']^2 + [(1-t_1) + t_1r']^2 \right\}^p &< \mathcal{E}(r) \\ &< 2^{p-2}\pi(1+r')^{1-2p} \left\{ [t_2 + (1-t_2)r']^2 + [(1-t_2) + t_2r']^2 \right\}^p \end{aligned} \quad (1.5)$$

holds for all $r \in (0, 1)$ if and only if $t_1 \leq 1/2 + \sqrt{1/(4p)}/2$ and $t_2 \geq 1/2 + \sqrt{(4/\pi)^{1/p} - 1}/2$.

2. Lemmas

In order to prove our main results we need several formulas and lemmas, which we present in this section.

For $r \in (0, 1)$, the following formulas were presented in [9, Appendix E, pp. 474-475, and Theorem 3.21(7)].

$$\begin{aligned} \frac{d\mathcal{K}}{dr} &= \frac{\mathcal{E} - r'^2\mathcal{K}}{rr'^2}, \quad \frac{d\mathcal{E}}{dr} = \frac{\mathcal{E} - \mathcal{K}}{r}, \quad \frac{d(\mathcal{E} - r'^2\mathcal{K})}{dr} = r\mathcal{K}, \\ \frac{d(\mathcal{K} - \mathcal{E})}{dr} &= \frac{r\mathcal{E}}{r'^2}, \quad \mathcal{E} \left(\frac{2\sqrt{r}}{1+r} \right) = \frac{2\mathcal{E} - r'^2\mathcal{K}}{1+r}, \quad \lim_{r \rightarrow 1} r'^\alpha \mathcal{K}(r) = 0 \quad (\alpha > 0). \end{aligned}$$

Lemma 2.1 (See [9, Theorem 1.25]). For $-\infty < a < b < \infty$, let $f, g : [a, b] \rightarrow \mathbb{R}$ be continuous on $[a, b]$, and be differentiable on (a, b) , let $g'(x) \neq 0$ on (a, b) . If $f'(x)/g'(x)$ is increasing (decreasing) on (a, b) , then so are

$$\frac{f(x) - f(a)}{g(x) - g(a)} \quad \text{and} \quad \frac{f(x) - f(b)}{g(x) - g(b)}.$$

If $f'(x)/g'(x)$ is strictly monotone, then the monotonicity in the conclusion is also strict.

Lemma 2.2. Let $r \in (0, 1)$, then

- (1) the function $r \rightarrow (\mathcal{E} - r'^2\mathcal{K})/r^2$ is strictly increasing from $(0, 1)$ onto $(\pi/4, 1)$;
- (2) the function $r \rightarrow \mathcal{E}/r'^{1/2}$ is strictly increasing from $(0, 1)$ onto $(\pi/2, \infty)$;
- (3) the function $r \rightarrow (\mathcal{K} - \mathcal{E})/(r^2\mathcal{K})$ is strictly increasing from $(0, 1)$ onto $(1/2, 1)$;
- (4) the function $r \rightarrow (\mathcal{E} - r'^2\mathcal{K})/(r^2\mathcal{K})$ is strictly decreasing from $(0, 1)$ onto $(0, 1/2)$;
- (5) the function $r \rightarrow r'^{3/4}(\mathcal{K} - \mathcal{E})/r^2$ is strictly decreasing from $(0, 1)$ onto $(0, \pi/4)$;
- (6) the function $r \rightarrow (\mathcal{E} - r'^2\mathcal{K})^2/(\mathcal{E}^2 - r'^2\mathcal{K}^2)$ is strictly decreasing from $(0, 1)$ onto $(1, 2)$;
- (7) the function $r \rightarrow [4(2\mathcal{E} - r'^2\mathcal{K})^2 - \pi^2]/r^2$ is strictly increasing from $(0, 1)$ onto $(\pi^2/2, 16 - \pi^2)$.

Proof. Parts (1)-(5) can be found in [9, Theorem 3.21(1) and (8), Exercises 3.43(32) and (46)] and [2, Theorem 15].

For part (6), let $I_1(r) = (\mathcal{E} - r'^2\mathcal{K})^2$, $I_2(r) = \mathcal{E}^2 - r'^2\mathcal{K}^2$ and $I(r) = (\mathcal{E} - r'^2\mathcal{K})^2/(\mathcal{E}^2 - r'^2\mathcal{K}^2)$. Then $I(r) = I_1(r)/I_2(r)$, $I_1(0) = I_2(0) = 0$ and

$$\frac{I_1'(r)}{I_2'(r)} = \left(\frac{r^2\mathcal{K}}{\mathcal{K} - \mathcal{E}} \right)^2 \cdot \frac{\mathcal{E} - r'^2\mathcal{K}}{r^2\mathcal{K}}. \quad (2.1)$$

It follows from parts (3) and (4) together with Lemma 2.1 and (2.1) that $I(r)$ is strictly decreasing in $(0, 1)$. By l'Hôpital's rule, we have $I(0^+) = 2$, while $I(1^-) = 1$ is clear.

For part (7), let $J_1(r) = 4(2\mathcal{E} - r'^2\mathcal{K})^2 - \pi^2$, $J_2(r) = r^2$ and $J(r) = [4(2\mathcal{E} - r'^2\mathcal{K})^2 - \pi^2]/r^2$. Then $J(r) = J_1(r)/J_2(r)$, $J_1(0) = J_2(0) = 0$ and

$$\frac{J_1'(r)}{J_2'(r)} = \frac{4(2\mathcal{E} - r'^2\mathcal{K})(\mathcal{E} - r'^2\mathcal{K})}{r^2}. \quad (2.2)$$

It follows from part (1) and (2.2) together with Lemma 2.1 that $J(r)$ is strictly increasing in $(0, 1)$. By l'Hôpital's rule, we get $J(0^+) = \pi^2/2$, while $J(1^-) = 16 - \pi^2$ is clear. \square

Lemma 2.3. Let $r \in (0, 1)$ and

$$g(r) = \frac{(\mathcal{K} - \mathcal{E})(\mathcal{E} - r'^2\mathcal{K}) + \mathcal{E}[(\mathcal{K} - \mathcal{E}) - (\mathcal{E} - r'^2\mathcal{K})]}{(\mathcal{E} - r'^2\mathcal{K})^2}.$$

Then $g(r)$ is strictly increasing from $(0, 1)$ onto $(3/2, \infty)$.

Proof. Differentiating $g(r)$ and elaborated computations lead to

$$\begin{aligned} g'(r) &= \frac{\frac{r\mathcal{E}}{r'^2}(\mathcal{E} - r'^2\mathcal{K}) + (\mathcal{K} - \mathcal{E})r\mathcal{K} + \frac{\mathcal{E}-\mathcal{K}}{r}[\mathcal{K} - \mathcal{E} - (\mathcal{E} - r'^2\mathcal{K})] + \mathcal{E}(\frac{r\mathcal{E}}{r'^2} - r\mathcal{K})}{(\mathcal{E} - r'^2\mathcal{K})^2} \\ &\quad - \frac{2r\mathcal{K} \{(\mathcal{K} - \mathcal{E})(\mathcal{E} - r'^2\mathcal{K}) + \mathcal{E}[(\mathcal{K} - \mathcal{E}) - (\mathcal{E} - r'^2\mathcal{K})]\}}{(\mathcal{E} - r'^2\mathcal{K})^3} \\ &= \frac{2r^2\mathcal{E}(\mathcal{E}^2 - r'^2\mathcal{K}^2) - 2r'^2(\mathcal{K} - \mathcal{E})^2(\mathcal{E} - r'^2\mathcal{K})}{rr'^2(\mathcal{E} - r'^2\mathcal{K})^3} \\ &= \frac{2r\mathcal{E}(\mathcal{E}^2 - r'^2\mathcal{K}^2)}{r'^2(\mathcal{E} - r'^2\mathcal{K})^3} [1 - g_1(r)] \end{aligned} \quad (2.3)$$

where

$$g_1(r) = \left[\frac{r'^{3/4}(\mathcal{K} - \mathcal{E})}{r^2} \right]^2 \cdot \frac{r'^{1/2}}{\mathcal{E}} \cdot \frac{r^2}{\mathcal{E} - r'^2\mathcal{K}} \cdot \frac{(\mathcal{E} - r'^2\mathcal{K})^2}{\mathcal{E}^2 - r'^2\mathcal{K}^2}. \quad (2.4)$$

It follows from Lemma 2.2(1), (2), (5) and (6) together with (2.4) that $g_1(r)$ is strictly decreasing from $(0, 1)$ onto $(0, 1)$. Then equation (2.3) leads to the conclusion that $g(r)$ is strictly increasing in $(0, 1)$. Making use of l'Hôpital's rule one has

$$\begin{aligned} \lim_{r \rightarrow 0^+} g(r) &= \lim_{r \rightarrow 0^+} \frac{(\mathcal{K} - \mathcal{E})(\mathcal{E} - r'^2\mathcal{K})/r^4 + \mathcal{E}[(\mathcal{K} - \mathcal{E}) - (\mathcal{E} - r'^2\mathcal{K})]/r^4}{(\mathcal{E} - r'^2\mathcal{K})^2/r^4} \\ &= \frac{\pi^2/16 + \pi/2 \times \pi/16}{\pi^2/16} = \frac{3}{2}, \end{aligned}$$

while $g(1^-) = +\infty$ is clear. \square

Lemma 2.4. Let $r \in (0, 1)$, $p \geq 1/2$ and

$$h(r) = (2p - 1)r^2 + 2p \frac{r^2\mathcal{E}}{\mathcal{E} - r'^2\mathcal{K}}.$$

Then $h(r)$ is strictly decreasing from $(0, 1)$ onto $(4p - 1, 4p)$ if and only if $p \leq 2$.

Proof. Differentiating $h(r)$ leads to

$$\begin{aligned} h'(r) &= 2(2p - 1)r + 2p \frac{r(3\mathcal{E} - \mathcal{K})(\mathcal{E} - r'^2\mathcal{K}) - r^3\mathcal{K}\mathcal{E}}{(\mathcal{E} - r'^2\mathcal{K})^2} \\ &= 2pr \left[2 - \frac{1}{p} - g(r) \right], \end{aligned} \quad (2.5)$$

where $g(r)$ is defined as in Lemma 2.3.

Therefore, Lemma 2.4 follows easily from Lemma 2.3 and (2.5) together with the limiting values $h(0^+) = 4p$ and $h(1^-) = 4p - 1$. \square

Lemma 2.5. The inequality

$$\frac{1}{4p} < \left(\frac{4}{\pi} \right)^{1/p} - 1 < \frac{1}{4p - 1}$$

holds for $p \in [1/2, 2]$.

Proof. Let $f_1(p) = [(4p + 1)/(4p)]^p$ and $f_2(p) = [4p/(4p - 1)]^p$. Then it is not difficult to verify that $f_1(p)$ is strictly increasing in $[1/2, 2]$ and $f_2(p)$ is strictly decreasing in $[1/2, 2]$. Hence, we get

$$f_1(p) < f_1(2) = \frac{81}{64} = 1.265625 \cdots < \frac{4}{\pi} = 1.273238 \cdots \quad (2.6)$$

and

$$f_2(p) > f_2(2) = \frac{64}{49} = 1.306122 \cdots > \frac{4}{\pi} = 1.273238 \cdots \quad (2.7)$$

for $p \in [1/2, 2]$.

Therefore, Lemma 2.5 follows from (2.6) and (2.7). \square

Lemma 2.6. Let $u \in [0, 1]$, $p \in [1/2, 2]$, $r \in (0, 1)$ and

$$f_{u,p}(r) = p \log(1 + ur^2) - \log \left[\frac{2}{\pi} (2\mathcal{E} - r'^2\mathcal{K}) \right]. \quad (2.8)$$

Then $f_{u,p}(r) < 0$ for $r \in (0, 1)$ if and only if $u \leq 1/(4p)$, and $f_{u,p}(r) > 0$ for $r \in (0, 1)$ if and only if $u \geq (4/\pi)^{1/p} - 1$.

Proof. From (2.8) we get

$$f_{u,p}(0^+) = 0, \quad (2.9)$$

$$f_{u,p}(1^-) = p \log(1+u) + \log\left(\frac{\pi}{4}\right) \quad (2.10)$$

and

$$\begin{aligned} f_{u,p}'(r) &= \frac{2pur}{1+ur^2} - \frac{\mathcal{E} - r'^2\mathcal{K}}{r(2\mathcal{E} - r'^2\mathcal{K})} \\ &= \frac{(2p-1)r^2(\mathcal{E} - r'^2\mathcal{K}) + 2pr^2\mathcal{E}}{r(1+ur^2)(2\mathcal{E} - r'^2\mathcal{K})} \left[u - \frac{1}{h(r)} \right], \end{aligned} \quad (2.11)$$

where $h(r)$ is defined as in Lemma 2.4.

It follows from Lemma 2.4 that $1/h(r)$ is strictly increasing from $(0, 1)$ onto $(1/(4p), 1/(4p-1))$ if $1/2 \leq p \leq 2$.

Making use of Lemma 2.5, we divide the proof into four cases.

Case 1 $u \leq 1/(4p)$. Then (2.11) and the monotonicity of $1/h(r)$ lead to the conclusion that $f_{u,p}(r)$ is strictly decreasing in $(0, 1)$. Therefore, $f_{u,p}(r) < 0$ for $r \in (0, 1)$ follows from (2.9) and the monotonicity of $f_{u,p}(r)$.

Case 2 $u \geq 1/(4p-1)$. Then (2.11) and the monotonicity of $1/h(r)$ lead to the conclusion that $f_{u,p}(r)$ is strictly increasing in $(0, 1)$. Therefore, $f_{u,p}(r) > 0$ for $r \in (0, 1)$ follows from (2.9) and the monotonicity of $f_{u,p}(r)$.

Case 3 $(4/\pi)^{1/p} - 1 < u < 1/(4p-1)$. Then from (2.10) and (2.11) together with the monotonicity of $1/h(r)$ we clearly see that there exists $\lambda \in (0, 1)$ such that $f_{u,p}(r)$ is strictly increasing in $(0, \lambda]$ and strictly decreasing in $[\lambda, 1)$, and

$$f_{u,p}(1^-) \geq 0. \quad (2.12)$$

Therefore, $f_{u,p}(r) > 0$ for $r \in (0, 1)$ follows from (2.9) and (2.12) together with the piecewise monotonicity of $f_{u,p}(r)$.

Case 4 $1/(4p) < u < (4/\pi)^{1/p} - 1$. Then from (2.10) and (2.11) together with the monotonicity of $1/h(r)$ we clearly see that there exists $0 < \mu < 1$ such that $f_{u,p}(r)$ is strictly increasing in $(0, \mu]$ and strictly decreasing in $[\mu, 1)$, and

$$f_{u,p}(1^-) < 0. \quad (2.13)$$

Therefore, equation (2.9) and inequality (2.13) together with the piecewise monotonicity of $f_{u,p}(r)$ lead to the conclusion that there exists $0 < \mu < \eta < 1$ such that $f_{u,p}(r) > 0$ for $r \in (0, \eta)$ and $f_{u,p}(r) < 0$ for $r \in (\eta, 1)$. \square

Lemma 2.7. If $r \in (0, 1)$, then the function

$$F(r) = (2\mathcal{E} - r'^2\mathcal{K})^2 \left[1 + \frac{\pi^2 - 4(2\mathcal{E} - r'^2\mathcal{K})^2}{\pi^2 r^2} \right]$$

is strictly increasing from $(0, 1)$ onto $(\pi^2/8, 8(\pi^2 - 8)/\pi^2)$.

Proof. Differentiating $F(r)$ leads to

$$\begin{aligned} f'(r) &= \frac{2(2\mathcal{E} - r'^2\mathcal{K})(\mathcal{E} - r'^2\mathcal{K})}{r} \left[1 + \frac{\pi^2 - 4(2\mathcal{E} - r'^2\mathcal{K})^2}{\pi^2 r^2} \right] \\ &\quad + (2\mathcal{E} - r'^2\mathcal{K})^2 \frac{8\mathcal{E}(2\mathcal{E} - r'^2\mathcal{K}) - 2\pi^2}{\pi^2 r^3} \\ &= \frac{2r'^2(2\mathcal{E} - r'^2\mathcal{K})(\mathcal{E} - r'^2\mathcal{K})}{r^3} \left[\frac{4(2\mathcal{E} - r'^2\mathcal{K})^2 - \pi^2}{\pi^2 r^2} \frac{r^2\mathcal{K}}{(\mathcal{E} - r'^2\mathcal{K})} - 1 \right]. \end{aligned} \tag{2.14}$$

It follows from Lemma 2.2(4) and (7) together with (2.14) that $F(r)$ is strictly increasing in $(0, 1)$. Moreover, the limiting values $F(0^+) = \pi^2/8$ and $F(1^-) = 8(\pi^2 - 8)/\pi^2$ follows easily from Lemma 2.2(7). \square

3. Proof of Theorems 1.1 and 1.2

Proof of Theorem 1.1. Let $q \in (0, 1/2]$ and $t = (1 - r')/(1 + r') \in (0, 1)$. Then

$$\mathcal{E}(r) = \mathcal{E} \left(\frac{2\sqrt{t}}{1+t} \right) = \frac{2\mathcal{E}(t) - t'^2\mathcal{K}(t)}{1+t} \tag{3.1}$$

and

$$\begin{aligned} &\sqrt{q + (1 - q)r'^2} + \sqrt{(1 - q) + qr'^2} \\ &= \frac{\sqrt{1 + t^2 + 2(1 - 2q)t} + \sqrt{1 + t^2 - 2(1 - 2q)t}}{1 + t}. \end{aligned} \tag{3.2}$$

From Lemma 2.2(7) we get

$$\frac{8}{\pi^2} \left[2\mathcal{E}(t) - t'^2\mathcal{K}(t) \right]^2 > 1 + t^2. \tag{3.3}$$

Equations (3.1) and (3.2) together with inequality (3.3) lead to the following equivalence relations:

$$\begin{aligned}
& \frac{4}{\pi} \mathcal{E}(r) > \sqrt{q + (1-q)r'^2} + \sqrt{(1-q) + qr'^2} \\
\iff & \frac{4}{\pi} \left[2\mathcal{E}(t) - t'^2 \mathcal{K}(t) \right] > \sqrt{1+t^2 + 2(1-2q)t} + \sqrt{1+t^2 - 2(1-2q)t} \\
\iff & \frac{8}{\pi^2} \left[2\mathcal{E}(t) - t'^2 \mathcal{K}(t) \right]^2 - (1+t^2) > \sqrt{(1+t^2)^2 - 4(1-2q)^2 t^2} \\
\iff & \frac{64}{\pi^4} \left[2\mathcal{E}(t) - t'^2 \mathcal{K}(t) \right]^4 - \frac{16}{\pi^2} (1+t^2) \left[2\mathcal{E}(t) - t'^2 \mathcal{K}(t) \right]^2 > -4(1-2q)^2 t^2 \\
\iff & \frac{4}{\pi^2} \left[2\mathcal{E}(t) - t'^2 \mathcal{K}(t) \right]^2 \left[1 + \frac{\pi^2 - 4(2\mathcal{E}(t) - t'^2 \mathcal{K}(t))^2}{\pi^2 t^2} \right] < (1-2q)^2.
\end{aligned} \tag{3.4}$$

Similarly, we have

$$\begin{aligned}
& \frac{4}{\pi} \mathcal{E}(r) < \sqrt{q + (1-q)r'^2} + \sqrt{(1-q) + qr'^2} \iff \\
& \frac{4}{\pi^2} \left[2\mathcal{E}(t) - t'^2 \mathcal{K}(t) \right]^2 \left[1 + \frac{\pi^2 - 4(2\mathcal{E}(t) - t'^2 \mathcal{K}(t))^2}{\pi^2 t^2} \right] > (1-2q)^2.
\end{aligned} \tag{3.5}$$

Therefore, Theorem 1.1 follows easily from (3.4) and (3.5) together with Lemma 2.7. \square

In order to prove Theorem 1.2, we need to introduce the Toader mean which is closely related to the complete elliptic integral of the second kind. For $a, b > 0$ with $a \neq b$, the Toader mean $T(a, b)$ [17, 18, 20, 25] is defined by

$$T(a, b) = \frac{2}{\pi} \int_0^{\pi/2} \sqrt{a^2 \cos^2 \theta + b^2 \sin^2 \theta} d\theta = \begin{cases} 2a\mathcal{E}(\sqrt{1 - (b/a)^2})/\pi, & a > b, \\ 2b\mathcal{E}(\sqrt{1 - (a/b)^2})/\pi, & a < b. \end{cases} \tag{3.6}$$

In particular,

$$\mathcal{E}(r) = \pi T(1, r')/2 \tag{3.7}$$

for $r \in (0, 1)$.

Proof of Theorem 1.2. Let $a, b > 0$ with $a \neq b$, $t, t_1, t_2 \in [1/2, 1]$, $p \in [1/2, 2]$ and

$$Q_{t,p}(a, b) = C^p(ta + (1-t)b, tb + (1-t)a)A^{1-p}(a, b), \tag{3.8}$$

where $A(a, b) = (a + b)/2$ and $C(a, b) = (a^2 + b^2)/(a + b)$ are the arithmetic and contraharmonic means of a and b , respectively. Then $Q_{t,p}(a, b)$ is strictly increasing with respect to $t \in [1/2, 1]$ for fixed $a, b > 0$ with $a \neq b$.

Since both $Q_{t,p}(a, b)$ and $T(a, b)$ are symmetric and homogeneous of degree 1. Without loss of generality, we assume that $a > b$. Let $x = (a - b)/(a + b) \in (0, 1)$. Then (3.6)-(3.8) lead to

$$\begin{aligned} \log \left(\frac{Q_{t,p}(a, b)}{T(a, b)} \right) &= \log \left(\frac{Q_{t,p}(a, b)}{A(a, b)} \right) - \log \left(\frac{T(a, b)}{A(a, b)} \right) \\ &= p \log [1 + (1 - 2t)^2 x^2] - \log \left[\frac{T(1, b/a)}{A(1, b/a)} \right] \\ &= p \log [1 + (1 - 2t)^2 x^2] - \log \left[\frac{2}{\pi} (1 + x) \mathcal{E} \left(\frac{2\sqrt{x}}{1 + x} \right) \right] \\ &= p \log [1 + (1 - 2t)^2 x^2] - \log \frac{2}{\pi} [2\mathcal{E}(x) - x'^2 \mathcal{K}(x)]. \quad (3.9) \end{aligned}$$

From Lemma 2.6 and equation (3.9) we clearly see that the double inequality

$$Q_{t_1,p}(a, b) < T(a, b) < Q_{t_2,p}(a, b) \quad (3.10)$$

holds if and only if $t_1 \leq 1/2 + \sqrt{1/(4p)}/2$ and $t_2 \geq 1/2 + \sqrt{(4/\pi)^{1/p} - 1}/2$.

In particular, if $a = 1$ and $b = r'$, then inequality (3.10) becomes

$$\begin{aligned} 2^{p-2} \pi (1 + r')^{1-2p} \left\{ [t_1 + (1 - t_1)r']^2 + [(1 - t_1) + t_1 r']^2 \right\}^p &< \mathcal{E}(r) \\ &< 2^{p-2} \pi (1 + r')^{1-2p} \left\{ [t_2 + (1 - t_2)r']^2 + [(1 - t_2) + t_2 r']^2 \right\}^p. \end{aligned}$$

The proof of Theorem 1.2 is therefore complete. \square

Let $p = 2$ and $p = 1/2$ in the first and second inequality of (1.5), respectively. Then we get

Corollary 3.1. The double inequality

$$\begin{aligned} \frac{\pi}{(1 + r')^3} \left\{ [\lambda + (1 - \lambda)r']^2 + [(1 - \lambda) + \lambda r']^2 \right\}^2 &< \mathcal{E}(r) \\ &< \frac{\pi}{2} \left\{ \frac{[\mu + (1 - \mu)r']^2 + [(1 - \mu) + \mu r']^2}{2} \right\}^{1/2} \end{aligned}$$

holds for all $r \in (0, 1)$ with $\lambda = 1/2 + \sqrt{2}/8$ and $\mu = 1/2 + \sqrt{(4/\pi)^2 - 1}/2$.

4. Comparison of the bounds for $\mathcal{E}(r)$

In this section, we compare our bounds for $\mathcal{E}(r)$ with the bounds in (1.1)-(1.3).

Remark 4.1. Simple computations lead to that the upper bound in (1.4) is exactly the bound in (1.3).

Remark 4.2. Let $x \in (0, 1)$ and $\mu = 1/2 + \sqrt{(4/\pi)^2 - 1}/2$. Then simple computation leads to

$$(1 + x^2) - \{[\mu + (1 - \mu)x]^2 + [(1 - \mu) + \mu x]^2\} = (1 - \frac{8}{\pi^2})(1 - x)^2 > 0.$$

Therefore, the upper bound in Corollary 3.1 is better than the bound in (1.2).

Remark 4.3. Let $\bar{\alpha} = 1/2 - \sqrt{2}/4$, $\bar{\beta} = 1/2 + \sqrt{2}/4$ and $\mu = 1/2 + \sqrt{(4/\pi)^2 - 1}/2$. Then we have

$$\begin{aligned} & \lim_{r \rightarrow 1} \left[\frac{\pi}{4} \left(\sqrt{1 - \bar{\alpha}r^2} + \sqrt{1 - \bar{\beta}r^2} \right) \right] \\ &= \frac{\pi}{8} \left(\sqrt{2 + \sqrt{2}} + \sqrt{2 - \sqrt{2}} \right) = 1.026172 \dots \end{aligned} \quad (4.1)$$

and

$$\lim_{r \rightarrow 1} \frac{\pi}{2} \left\{ \frac{[\mu + (1 - \mu)r']^2 + [(1 - \mu) + \mu r']^2}{2} \right\}^{1/2} = 1. \quad (4.2)$$

Equations (4.1) and (4.2) show that the upper bound in Corollary 3.1 is asymptotically precise when $r \rightarrow 1$, and there exists $0 < \delta_1 < 1$ such that the upper bound in Corollary 3.1 is better than the bound in (1.3) when $r \in (1 - \delta_1, 1)$.

Remark 4.4. Let $\beta = 1/2 - 2\sqrt{2(\pi^2 - 8)}/\pi^2$, then

$$\lim_{r \rightarrow 1} \frac{\pi}{4} \left(\sqrt{\beta + (1 - \beta)r'^2} + \sqrt{(1 - \beta) + \beta r'^2} \right) = 1 \quad (4.3)$$

and

$$\lim_{r \rightarrow 1} \frac{\pi}{2} \left(\frac{1 + r'^{3/2}}{2} \right)^{2/3} = 2^{-5/3}\pi = 0.989539 \dots \quad (4.4)$$

Equations (4.3) and (4.4) imply that the lower bound in Theorem 1.1 with $\beta = 1/2 - 2\sqrt{2(\pi^2 - 8)}/\pi^2$ is asymptotically precise when $r \rightarrow 1$, and there exists $0 < \delta_2 < 1$ such that the lower bound in Theorem 1.1 is better than the bound in (1.1) when $r \in (1 - \delta_2, 1)$.

Remark 4.5. Let $\lambda = 1/2 + \sqrt{2}/8$, $r \in (0, 1)$ and $x = (1 - r^2)^{1/4} \in (0, 1)$. Then the following equivalence relations lead to the conclusion that the lower bound in Corollary 3.1 is better than the bound in inequality (1.1).

$$\begin{aligned}
& \frac{\pi}{(1+r')^3} \left\{ [\lambda + (1-\lambda)r']^2 + [(1-\lambda) + \lambda r']^2 \right\}^2 > \frac{\pi}{2} \left(\frac{1+r'^{3/2}}{2} \right)^{2/3} \\
\iff & \frac{[9 + 9(1-r^2) + 14\sqrt{1-r^2}]^2}{128(1+\sqrt{1-r^2})^3} > \left[\frac{1+(1-r^2)^{3/4}}{2} \right]^{2/3} \\
\iff & \frac{(9+9x^4+14x^2)^2}{128(1+x^2)^3} > \left(\frac{1+x^3}{2} \right)^{2/3} \\
\iff & (9+9x^4+14x^2)^6 - 524288(1+x^2)^9(1+x^3)^2 > 0 \\
\iff & (x-1)^4 [7153x^{20} + 28612x^{19} + 313054x^{18} + 60580x^{17} + 2074909x^{16} \\
& - 277424x^{15} + 6613736x^{14} - 1390192x^{13} + 12597746x^{12} - 2615880x^{11} \\
& + 15507060x^{10} - 2615880x^9 + 12597746x^8 - 1390192x^7 + 6613736x^6 \\
& - 277424x^5 + 2074909x^4 + 60580x^3 + 313054x^2 + 28612x + 7153] > 0.
\end{aligned}$$

Acknowledgements

This research was supported by the Natural Science Foundation of China under Grant 11071069, and the Innovation Team Foundation of the Department of Education of Zhejiang Province under Grant T200924.

References

- [1] G. Almkvist, B. Berndt, Gauss, Landen, Ramanujan, the arithmetic-geometric mean, ellipses, π , and the Ladies diary, *Amer. Math. Monthly* **95** (1988), 585-608
- [2] H. Alzer, S.-L. Qiu, Monotonicity theorems and inequalities for the complete elliptic integrals, *J. Comput. Appl. Math.* **172** (2004) 289-312.

- [3] G. D. Anderson, R. W. Barnard, K. C. Richards, M. K. Vamanamurthy, M. Vuorinen, Inequalities for zero-balanced hypergeometric functions, *Trans. Amer. Math. Soc.* **347** (1995) 1713-1723.
- [4] G. D. Anderson, S.-L. Qiu, M. K. Vamanamurthy, Elliptic integral inequalities, with applications, *Constr. Approx.* **14** (1998) 195-207.
- [5] G. D. Anderson, S.-L. Qiu, M. K. Vamanamurthy, M. Vuorinen, Generalized elliptic integrals and modular equations, *Pacific J. Math.* **192** (2000) 1-37.
- [6] G. D. Anderson, M. K. Vamanamurthy, M. Vuorinen, Distortion functions for plane quasiconformal mappings, *Israel J. Math.* **62** (1988) 1-16.
- [7] G. D. Anderson, M. K. Vamanamurthy, M. Vuorinen, Functional inequalities for complete elliptic integrals and their ratios, *SIAM J Math. Anal.* **21** (1990) 536-549.
- [8] G. D. Anderson, M. K. Vamanamurthy, M. Vuorinen, Functional inequalities for hypergeometric functions and complete elliptic integrals, *SIAM J. Anal.* **23** (1992) 512-524.
- [9] G. D. Anderson, M. K. Vamanamurthy, M. Vuorinen, *Conformal Invariants, Inequalities, and Quasiconformal Maps*, John Wiley & Sons, New York, 1997.
- [10] S. András, Á. Baricz, Bounds for complete elliptic integrals of the first kind, *Expo. Math.* **28** (2010) 357-364.
- [11] K. E. Atkinson, *An Introduction to Numerical Analysis*, John Wiley & Sons, New York, 1989.
- [12] Á. Baricz, Turán type inequalities for generalized complete elliptic integrals, *Math. Z.* **256** (2007) 895-911.
- [13] R. W. Barnard, K. Pearce, K. C. Richards, An inequality involving the generalized hypergeometric function and the arc length of an ellipse, *SIAM J. Math. Anal.* **31** (2000) 693-699.
- [14] R. W. Barnard, K. Pearce, K. C. Richards, A monotonicity property involving ${}_3F_2$ and comparisons of the classical approximations of elliptical arc length, *SIAM J. Math. Anal.* **32** (2000) 403-419.

- [15] F. Bowman, Introduction to Elliptic Functions with Applications, Dover Publications, New York, 1961.
- [16] P. F. Byrd, M. D. Friedman, Handbook of Elliptic Integrals for Engineers and Scientists, Springer-Verlag, New York, 1971.
- [17] Y.-M. Chu, M.-K. Wang, Optimal Lehmer mean bounds for the Toader mean, Results Math. **61** (2012) 223-229.
- [18] Y.-M. Chu, M.-K. Wang, Inequalities between arithmetic-geometric, Gini, and Toader means, Abstr. Appl. Anal. **2012**, Art. ID 830585, 11 pages.
- [19] Y.-M. Chu, M.-K. Wang, Y.-P. Jiang, S.-L. Qiu, Concavity of the complete elliptic integrals of the second kind with respect to Hölder means, J. Math. Anal. Appl. **395** (2012) 637-642.
- [20] Y.-M. Chu, M.-K. Wang, S.-L. Qiu, Optimal combinations bounds of root-square and arithmetic means for Toader mean, Proc. Indian Acad. Sci. Math. Sci. **121** (2011) 495-501.
- [21] Y.-M. Chu, M.-K. Wang, Y.-F. Qiu, On Alzer and Qiu's conjecture for complete elliptic integrals and inverse hyperbolic tangent function, Abstr. Appl. Anal. **2011**, Art. ID 697547, 7 pages.
- [22] H. Kazi, E. Neuman, Inequalities and bounds for elliptic integrals, J. Approx. Theory **146** (2007) 212-226.
- [23] T. Muir, On the perimeter of an ellipse, Mess. Math. **12** (1883) 149-151.
- [24] S.-L. Qiu, M.K. Vamanamurthy, M. Vuorinen, Some inequalities for the Hersch-Pfluger distortion function, J. Inequal. Appl. **4** (1999) 115-139.
- [25] Gh. Toader, Some mean values related to the arithmetic-geometric mean, J. Math. Anal. Appl. **218** (1998) 358-368.
- [26] M. K. Vamanamurthy, M. Vuorinen, Inequalities for means, J. Math. Anal. Appl. **183** (1994) 155-166.
- [27] M. Vuorinen, Singular values, Ramanujan modular equations, and Landen transformations, Studia Math. **121** (1996) 221-230.

- [28] M. Vuorinen, Hypergeometric functions in geometric function theory, in: Special Functions and Differential Equations, Proceedings of a Workshop held at The Institute of Mathematical Sciences, Madras, India, January 13-24, 1997, Allied Publ., New Delhi, 1998, pp. 119-126.
- [29] G.-D. Wang, X.-H. Zhang, Y.-M. Chu, Inequalities for the generalized elliptic integrals and modular functions, *J. Math. Anal. Appl.* **331** (2007) 1275-1283.
- [30] M.-K. Wang, Y.-M. Chu, S.-L. Qiu, Convexity of the complete elliptic integrals of the first and kind with respect to Hölder means, *J. Math. Anal. Appl.* **388** (2012) 1141-1146.
- [31] M.-K. Wang, Y.-M. Chu, Y.-F. Qiu, S.-L. Qiu, An optimal power mean inequality for the complete elliptic integrals, *Appl. Math. Lett.* **24** (2011) 887-890.
- [32] M.-K. Wang, S.-L. Qiu, Y.-M. Chu, Y.-P. Jiang, Generalized Hersch-Pfluger distortion function and complete elliptic integrals, *J. Math. Anal. Appl.* **385** (2012) 221-229.
- [33] X.-H. Zhang, G.-D. Wang, Y.-M. Chu, Remark on generalized elliptic integrals, *Proc. Roy. Soc. Edinburgh Sect. A* **139** (2009) 417-426.