An exponential integrator for finite volume discretization of nonlinear parabolic partial differential equation

Antoine Tambue^{a,b}

^a The African Institute for Mathematical Sciences(AIMS) and Stellenbosh University, 6-8 Melrose Road, Muizenberg 7945, South Africa

^bCenter for Research in Computational and Applied Mechanics (CERECAM), and Department of Mathematics and Applied Mathematics, University of Cape Town, 7701 Rondebosch, South Africa.

Abstract

We consider the numerical approximation of a general second order semi-linear parabolic partial differential equation. Equations of this type arise in many contexts, such as transport in porous media which is fundamental in many geo-engineering applications, including oil and gas recovery from subsurface. Using the finite volume with two-point flux approximation on regular mesh combined with exponential time differencing of order one (ETD1) for temporal discretization, we derive the L^2 estimate under the assumption that the non linear term is locally Lipschitz. Numerical simulations to sustain the theoretical results are provided.

Keywords: Parabolic partial differential equation, Finite volume method, Exponential integrators, Errors estimate

1. Introduction

Flow and transport are fundamental in many geo-engineering applications, including oil and gas recovery from hydrocarbon reservoirs, groundwater contamination and sustainable use of groundwater resources, storing greenhouse gases (e.g. CO_2) or radioactive waste in the subsurface, or mining heat from geothermal reservoirs. In porous media, a non-degenerated advection-diffusion-reaction is given by

$$\frac{\partial X}{\partial t} = \nabla \cdot (\mathbf{D}(\mathbf{x})\nabla X) - \nabla \cdot (\mathbf{q}(\mathbf{x})X) + f(\mathbf{x},X) \quad (\mathbf{x},t) \in \Omega \times [0,T],$$
(1)

where Ω is an open domain of \mathbb{R}^d , $d \in \{2,3\}$, **D** is the symmetric dispersion tensor, X is the unknown concentration of the contaminant, **q** the Darcy's velocity and f the reaction and source term. For the sake of simplicity, without loss of generality, we assume that f is explicitly independent of time. The model equation (1) finds interest in many engineering problems with specific coefficients. Finite element, finite volume or the combination finite

Email address: antonio@aims.ac.za (Antoine Tambue)

element-finite volume methods are mostly used for space discretization of the problem (1) while explicit, semi implicit and fully implicit methods are usually used for time discretization (see [27, 6, 8, 1, 9]). Due to time step size constraints, fully implicit schemes are more popular for time discretization for quite a long time compared to explicit Euler schemes. However, implicit schemes need at each time step a solution of large systems of nonlinear equations. This can be the bottleneck in computations. In recent years, exponential integrators have become an attractive alternative in many evolutions equations (see [4, 10, 11, 5, 7, 12, 17, 16, 12, 13, 15]). In contrast to classical methods, they are robust with respect to the Péclet number, they do not require the solution of large linear systems. Instead they make explicit use of the matrix exponential and related matrix functions. The price to pay is the computing of the matrix exponential functions of the non diagonal matrices, which has revived interest and significance progresses nowadays (see [7, 5, 12, 16, 12, 13]).

In this work, we combine a finite volume method with the first order exponential time differencing scheme of order 1 (ETD1). Although both discretization techniques have been together used for solving evolutionary problems like (1) (see [4, 10, 11]), a proper combination of rigorous convergence proof of them has been lacking so far. Furthermore the nonlinear term is assumed to be locally Lipschitz, which covers many reaction functions in geo-engineering applications.

The paper is organised as follows. In Section 2, we present the semi group formulation of (1), the existence and uniqueness of the solution along with some proprieties of the mild solution. In Section 4, we present the finite volume space discretization of (1), the existence and the uniqueness of the corresponding semi-discrete problem, and the L^2 error estimate between the exact solution and the semi-discrete solution. We end by presenting in Section 5 The time discretization of the semi-discrete problem based on ETD1 scheme is presented in Section 5, along with the convergence proof of the fully discrete scheme based on finite volume method and ETD1 scheme. We end by providing numerical simulations to sustain the theoretical results in Section 6. These results also show the efficiency of the ETD1 scheme compared to the standard time integrators, from which ETD1 scheme is ten times faster that the standard implicit scheme.

2. Semi group formulation and well posedness

Let us start by presenting briefly the notation of the main function spaces and norms that we will use in this paper. We denote by $\|\cdot\|$ the norm associated to the inner product (\cdot, \cdot) of the Hilbert space $H = L^2(\Omega)$. The norms in the Sobolev spaces $H^m(\Omega), m \ge 0$ will be denoted by $\|.\|_m$. The space $H^{-1}(\Omega)$ is the dual of $H_0^1(\Omega)$ equipped with the norm $\|u\|_{-1} = \sup_{v \in H_0^1(\Omega)} \frac{|(u, v)|}{\|u\|_1}$. For a Banach space \mathcal{V} we denote by $L(\mathcal{V})$ the set of bounded linear mapping from \mathcal{V} to \mathcal{V} . We assume that the domain Ω is bounded, has a smooth boundary or is a convex polygon. For the sake of simplicity, without loss of generality, we use the homogeneous Dirichlet boundary condition. We also assume that the Darcy velocity \mathbf{q} is known, and satisfies the mass conservation for incompressible fluids without internal source, For a given initial solution $X_0 \in H$, the model problem (1) is reformulated as: find the function $X(t) \in H^1(\Omega)$ such that

$$\begin{cases} \partial X/\partial t + \mathcal{A}X = f(\mathbf{x}, X) & (\mathbf{x}, t) \in \Omega \times [0, T] \\ X(\mathbf{x}, 0) = X_0 & \mathbf{x} \in \Omega \\ X(\mathbf{x}, t) = 0 & (\mathbf{x}, t) \in \partial\Omega \times [0, T] , \end{cases}$$
(2)

where

$$\mathcal{A}X = \mathcal{A}(\mathbf{x})X = -\nabla \cdot (\mathbf{D}\nabla X) + \nabla \cdot (\mathbf{q}(\mathbf{x})X)$$

$$= -\sum_{i,j=1}^{d} \frac{\partial}{\partial x_{i}} \left(D_{i,j}(\mathbf{x}) \frac{\partial X}{\partial x_{j}} \right) + \sum_{i=1}^{d} q_{i}(\mathbf{x}) \frac{\partial X}{\partial x_{i}} + (\nabla \cdot \mathbf{q}) X$$

$$= -\sum_{i,j=1}^{d} \frac{\partial}{\partial x_{i}} \left(D_{i,j}(\mathbf{x}) \frac{\partial X}{\partial x_{j}} \right) + \sum_{i=1}^{d} q_{i}(\mathbf{x}) \frac{\partial X}{\partial x_{i}}.$$

For well posedness of (2), we assume that **D** is symmetric, $D_{i,j} \in L^{\infty}(\Omega), q_i \in L^{\infty}(\Omega)$ and there exists a positive constant $c_1 > 0$ such that

$$\sum_{i,j=1}^{d} D_{i,j}(\mathbf{x})\xi_i\xi_j \ge c_1|\xi|^2 \qquad \forall \xi \in \mathbb{R}^d \quad \mathbf{x} \in \overline{\Omega} \quad c_1 > 0,$$
(3)

and

$$|f(\mathbf{x},u) - f(\mathbf{x},v)| \le L \left(1 + |u|^{\gamma} + |v|^{\gamma}\right) |u - v| \quad \forall u, v \in \mathbb{R} \ x \in \overline{\Omega}, \ t \in [0,T],$$
(4)

or

$$|f(\mathbf{x},u) - f(\mathbf{x},v)| \le L\left(1 + u + v + |u|^{\gamma} + |v|^{\gamma}\right)|u - v| \quad \forall u, v \in \mathbb{R} \ x \in \overline{\Omega}, \ t \in [0,T],$$
(5)

with $\gamma = 2$ for d = 3 and $\gamma \in [0, \infty)$ for d = 2.

Set $V = H_0^1(\Omega)$, the bilinear form associated to the operator \mathcal{A} is given by

$$a(u,v) = \int_{\Omega} \left(\sum_{i,j=1}^{d} D_{i,j} \frac{\partial u}{\partial x_j} \frac{\partial v}{\partial x_i} + \sum_{i=1}^{d} q_i \frac{\partial u}{\partial x_j} v \right) dx \qquad u,v \in V.$$
(6)

According to Gårding's inequality (see [5, 3]), there exists two positive constants c_0 and λ_0 such that

$$a(v,v) + c_0 \|v\|^2 \ge \lambda_0 \|v\|_1^2 \qquad \forall v \in V.$$
 (7)

By adding $c_0 X$ in both side of the first equation of (1) we have a new operator that we still call \mathcal{A} corresponding to the new bilinear form that we still call a such that the following coercivity property holds

$$a(v,v) \ge \lambda_0 \|v\|_1^2 \quad \forall v \in V.$$
(8)

For sake of simplicity, we will still call the right hand side of the first equation of (2) f. We define the following Nemytskii operator $F: H \to H$ by

$$(F(X))(x) = f(x, X).$$
 (9)

Using the Green's formula, the weak form of (2) consists to find the function $X(t) \in V$ such that

$$\begin{cases} (X_t, \chi) + a(X, \chi) = (F(X), \chi) & \forall \chi \in V, \quad t \in [0, T] \\ X(0) = X_0. \end{cases}$$
(10)

Note that a(,) is bounded in $V \times V$, so the following operator $A: V \to V^*$ is well defined by Riez's representation Theorem

$$a(u,v) = \langle Au, v \rangle, \quad \forall u, v \in V, \tag{11}$$

where V^* is the adjoint space (or dual space) of V and \langle,\rangle the duality pairing between V^* and V. By identifying H to its adjoint space H^* , we get the following continuous and dense inclusions

$$V \subset H \subset V^*. \tag{12}$$

So, we have

$$(u,v) = \langle u,v \rangle \qquad \qquad \forall u \in H, \, \forall v \in V.$$
(13)

The domain of A denoted by $\mathcal{D}(A)$ is defined by

$$\mathcal{D}(A) = \{ u \in V, \, Au \in H \}.$$
(14)

We write the restriction of $A: V \to V^*$ to $\mathcal{D}(A)$ again by A, which is therefore regarded as an operator of H (more precisely the H realization of \mathcal{A} [3, p. 812]). In the abstract setting, equation (10) is equivalent to find the function $X(t) \in V$ such that

$$\begin{cases} X_t + AX = F(X), & t \in [0, T] \\ X(0) = X_0, & \end{cases}$$
(15)

where equation (15) is understood in the space V^* using (12) and (13). As the domain Ω has a smooth boundary or is a convex polygon, we therefore have (see [3])

$$\mathcal{D}(A) = H_0^1(\Omega) \cap H^2(\Omega).$$
(16)

The V-ellipticity (8) implies that -A is a sectorial on $H = L^2(\Omega)$ (see [2, 3]) i.e. there exists $C_1, \theta \in (\frac{1}{2}\pi, \pi)$ such that

$$\|(\lambda I + A)^{-1}\|_{L(H)} \le \frac{C_1}{|\lambda|} \qquad \lambda \in S_{\theta},$$
(17)

where $S_{\theta} = \{\lambda \in \mathbb{C} : \lambda = \rho e^{i\phi}, \ \rho > 0, \ 0 \le |\phi| \le \theta \}.$

Then -A is the infinitesimal generator of bounded analytic semigroups $S(t) := e^{-tA}$ on $L^2(\Omega)$ such that

$$S(t) := e^{-tA} = \frac{1}{2\pi i} \int_{\mathcal{C}} e^{t\lambda} (\lambda I + A)^{-1} d\lambda, \qquad t > 0$$
(18)

where \mathcal{C} denotes a path that surrounds the spectrum of -A.

The coercivity property in (8) implies also that the set of the real part of the spectrum of A is non negative, which allows the definition of the fractional power of A as: for any $\alpha > 0$

$$\begin{cases} A^{-\alpha} = \frac{1}{\Gamma(\alpha)} \int_0^\infty t^{\alpha - 1} e^{-At} dt \\ A^{\alpha} = (A^{-\alpha})^{-1} \end{cases}$$
(19)

where $\Gamma(\alpha)$ is the Gamma function of α (see [2]). It is well known that $\|.\|_{\alpha} \equiv \|A^{\alpha/2}.\|$ in the space $\mathcal{D}(A^{\alpha/2})$, and that $V = \mathcal{D}(A^{1/2}) = \mathcal{D}(A^{*1/2})$ (see [27, 28]). Note that $A^{*1/2}$ is the adjoint of $A^{1/2}$.

For the nonlinear reaction term we make the following assumption

Proposition 2.1. [Lipschitz condition for F]

Under the assumption (4) or (5) on the nonlinear function f, let F the Nemytskii operator corresponding to f defined by (9). For each bounded set $\mathcal{B} \subset V$ there is a constant $C(\mathcal{B})$ such that

$$\|F(u) - F(v)\|_{-1} \le C(\mathcal{B}) \|u - v\|, \quad \forall u, v \in \mathcal{B}$$

$$\tag{20}$$

$$\|F(u) - F(v)\| \le C(\mathcal{B}) \|u - v\|_1, \quad \forall u, v \in \mathcal{B}.$$
(21)

Proof. The proofs of (20) and (21) can be found in [27] for function f of type (4). The proofs for function f of type (5) can easily be deducted. Indeed using Holder inequality yields

$$\begin{aligned} \|F(u) - F(v)\| & (22) \\ &\leq C\left(\left(\|u\|_{L^{q_1}(\Omega)} + \|u\|_{L^{q_1}(\Omega)}\right) \|u - v\|_{L^{p_1}(\Omega)} + \left(1 + \|u\|_{L^{q}(\Omega)}^{\gamma} + \|v\|_{L^{q}(\Omega)}^{\gamma}\right) \|u - v\|_{L^{p}(\Omega)}\right) \\ &\forall u, v \in \mathcal{B}, \end{aligned}$$

where $\frac{1}{p} + \frac{\gamma}{q} = \frac{1}{2}$ with p = q = 6 if d = 3, and arbitrary $p \in (1, \infty)$ if d = 2, $\frac{1}{p_1} + \frac{1}{q_1} = \frac{1}{2}$ with $p_1 \in [3, 6]^1$ if d = 3 and arbitrary $p_1 \in (2, \infty)$ if d = 2. Since Ω is bounded, $L^r(\Omega) \hookrightarrow L^s(\Omega)$ for $r \ge s$, combining with Sobolev the embedding theorem, we therefore have

$$\|F(u) - F(v)\| \leq C \left(1 + \|u\|_{1} + \|v\|_{1} + \|u\|_{1}^{\gamma} + \|v\|_{1}^{\gamma}\right) \|u - v\|_{1}.$$
(23)

Lipschitz condition (20) is proved in the same manner using the one in [27] for function f of type (4).

¹The corresponding q_1 is in the same interval

By Duhamel's principle we may represent the solution of (15) by the following integral equation

$$X(t) = S(t)X_0 + \int_0^t S(t-s)F(X(s))ds, \quad t \in [0,T].$$
(24)

Theorem 2.1. Assume that -A is the infinitesimal generator of bounded analytic semigroup (**D** is symmetric, $D_{i,j} \in L^{\infty}(\Omega), q_i \in L^{\infty}(\Omega)$ and the inequality (3) is fulfilled) and (4) (or (5)) is satisfied. For any bounded set $\mathcal{B}_0 \subset V$ there is $t^* = t^*(\mathcal{B}_0)$ such that equation (24) has an unique solution $X \in C([0, t^*], H^1(\Omega) \text{ for any } X_0 \in \mathcal{B}_0.$

Proof. Applying the contraction mapping principle in the topology of the Banach space $C([0,T], H^1(\Omega))$ to the integral equation (24) [2, Theorem 3.3.3] or [21, Theorem 6.3.1] ensure the existence and uniqueness of X.

Remark 2.1. The regularity of the solution X depends of the regularity of the coefficients $\mathbf{D} = (D_{i,j})_{1 \le i,j \le d}, \mathbf{q} = (q_i)_{1 \le i \le d}$ as we can observe in [26].

The following proposition will be largely used in this work

Proposition 2.2. [Smoothing properties of the semi group [2]] Let $\beta \ge 0$ and $0 \le \gamma \le 1$, then there exists C > 0 such that

$$\begin{aligned} \|A^{\beta}S(t)\|_{L(H)} &\leq Ct^{-\beta} \quad for \quad t > 0, \\ \|A^{-\gamma}(I - S(t))\|_{L(H)} &\leq Ct^{\gamma} \quad for \quad t \ge 0. \end{aligned}$$

In addition, the following results hold

$$A^{\beta}S(t) = S(t)A^{\beta} \quad on \quad \mathcal{D}(A^{\beta}).$$

$$If \quad \beta \ge \alpha \quad then \quad \mathcal{D}(A^{\beta}) \subset \mathcal{D}(A^{\alpha}).$$

$$\|D_{t}^{l}S(t)v\|_{\beta} \le Ct^{-l-(\beta-\alpha)/2} \|v\|_{\alpha}, \quad t > 0, \ v \in \mathcal{D}(A^{\alpha/2}) \quad l = 0, 1,$$

where $D_t^l := \frac{d^l}{dt^l}, \, \|.\|_{\alpha} := \|A^{\alpha/2}.\|$

The following lemma will be also used in our errors estimates.

Lemma 2.1. Let X be the mild solution of (2) given in (24). Let $\mathcal{B} \subset V$ be a bounded set such that $\forall t \in [0, t^*(\mathcal{B})], X(t) \in \mathcal{B}$. Let $t_1, t_2 \in [0, T] \subset [0, t^*(\mathcal{B})], t_1 < t_2$, the following estimates hold :

• (i) If $X_0 \in \mathcal{D}(A)$ then

$$||X(t_2) - X(t_1)|| \leq C(\mathcal{B})(t_2 - t_1)^{1-\epsilon} (||X_0||_2 + 1),$$

for $\epsilon \in (0, 1/4)$ small enough.

• (ii) If $X_0 \in \mathcal{D}(A)$ and F satisfies the Lipschitz condition in (20) then

$$||X(t_2) - X(t_1)|| \leq C(\mathcal{B})(t_2 - t_1) (||X_0||_2 + 1)).$$

Proof. Part (i).

Consider the difference

$$X(t_2) - X(t_1) = (S(t_2) - S(t_1)) X_0 + \left(\int_0^{t_2} S(t_2 - s) F(X(s)) ds - \int_0^{t_1} S(t_1 - s) F(X(s)) ds\right)$$

= $I + II$, (25)

so that $||X(t_2) - X(t_1)|| \le ||I|| + ||II||$. We estimate each of the terms ||I|| and ||II||. For ||I||, using Proposition 2.2 yields

$$||I|| = ||S(t_1)A^{-1}(I - S(t_2 - t_1))A^{1}X_0|| \leq C(t_2 - t_1)||X_0||_2.$$

For the term II, we have

$$II = \int_{0}^{t_{1}} (S(t_{2} - s) - S(t_{1} - s))F(X(s))ds + \int_{t_{1}}^{t_{2}} S(t_{2} - s)F(X(s))ds$$

= $II_{1} + II_{2}$,

with

$$\|II\| \le \|II_1\| + \|II_2\|.$$

We now estimate each term $||II_1||$ and $||II_2||$. For $||II_1||$

$$\begin{aligned} \|II_1\| &= \|\int_0^{t_1} (S(t_2 - s) - S(t_1 - s))F(X(s))ds\| \\ &\leq \int_0^{t_1} \|(S(t_2 - s) - S(t_1 - s))F(X(s))\|ds \\ &\leq \left(\int_0^{t_1} \|(S(t_2 - s) - S(t_1 - s))\|_{L(H)}ds\right) \left(\sup_{0 \le s \le T} \|F(X(s))\|\right). \end{aligned}$$

For $\epsilon \in (0, 1/4)$ small enough, using Proposition 2.2 yields

$$\begin{split} \|II_1\| &\leq \left(\int_0^{t_1} \|S(t_1 - s)A^{1 - \epsilon}A^{-1 + \epsilon}(\mathbf{I} - S(t_2 - t_1))\|_{L(H)} ds \right) \left(\sup_{0 \le s \le T} \|F(X(s))\| \right) \\ &\leq \left(\int_0^{t_1} \|A^{1 - \epsilon}S(t_1 - s)A^{-1 + \epsilon}(\mathbf{I} - S(t_2 - t_1))\|_{L(H)} ds \right) \left(\sup_{0 \le s \le T} \|F(X(s))\| \right) \\ &\leq C(t_2 - t_1)^{1 - \epsilon} \left(\int_0^{t_1} (t_1 - s)^{-1 + \epsilon} ds \right) \left(\sup_{0 \le s \le T} \|F(X(s))\| \right) \\ &\leq C(t_2 - t_1)^{1 - \epsilon} \left(\sup_{0 \le s \le T} \|F(X(s))\| \right). \end{split}$$

For $||II_2||$, using the fact that the semigroup is bounded, we have

$$\begin{aligned} \|II_2\| &= \|\int_{t_1}^{t_2} S(t_2 - s) F(X(s)) ds\| \\ &\leq \left(\int_{t_1}^{t_2} \|S(t_2 - s) F(X(s))\| ds\right) \\ &\leq \left(\int_{t_1}^{t_2} \|F(X(s))\| ds\right) \\ &\leq C(t_2 - t_1) \left(\sup_{0 \le s \le T} \|F(X(s))\|\right) \end{aligned}$$

Hence

$$||II|| \le ||II_1|| + ||II_2||) \le C(t_2 - t_1)^{1-\epsilon} \left(\sup_{0 \le s \le T} (||F(X(s))||) \right).$$

Using the fact that F satisfies (21), we therefore have

$$|F(X(t))|| \leq ||F(X_0)|| + ||F(X(t)) - F(X_0)|| \leq ||F(X_0)|| + L||X(t) - X_0||_1 \leq C(\mathcal{B}, X_0, F).$$
(26)

Combining (26) and previous estimations of ||I|| and ||II|| ends the proof of part (i).

Proof of part (ii). We consider again the difference in (25). The difference with the proof of part (i) comes from the estimation of II_1 . This time we write

$$II_{1} = \int_{0}^{t_{1}} (S(t_{2} - s) - S(t_{1} - s))F(X(s))ds$$

= $\int_{0}^{t_{1}} (S(t_{2} - s) - S(t_{1} - s)) (F(X(s)) - F(X(t_{1})))ds$
+ $\int_{0}^{t_{1}} (S(t_{2} - s) - S(t_{1} - s))F(X(t_{1}))ds$
= $II_{11} + II_{12}.$

Remember that $\|.\|_{-1} = \|A^{-1/2}.\|$ in V^* , since $H^{-1}(\Omega) = V^* = \mathcal{D}(A^{-1/2})$ as $V = \mathcal{D}(A^{1/2})$. If F satisfies the Lipschitz condition given in (20), then using the result in part (i) together with Proposition 2.2 yields

$$\begin{aligned} \|II_{11}\| &\leq \left(\int_{0}^{t_{1}} \|\left(S(t_{2}-s)-S(t_{1}-s)\right)A^{1/2}\|_{L(H)}\|A^{-1/2}\left(F(X(s))-F(X(t_{1})\right)\|ds\right) \\ &\leq C(\mathcal{B})\left(\int_{0}^{t_{1}} \|A^{1/2}\left(S(t_{2}-s)-S(t_{1}-s)\right)\|_{L(H)}\|X(s)-X(t_{1})\|ds\right) \\ &\leq C(\mathcal{B})\left(\int_{0}^{t_{1}} \|A^{3/2}S(t_{1}-s)A^{-1}\left(I-S(t_{2}-t_{1})\right)\|_{L(H)}\|X(s)-X(t_{1})\|ds\right) \\ &\leq C(\mathcal{B})\left((t_{2}-t_{1})\int_{0}^{t_{1}}(t_{1}-s)^{-\epsilon-1/2}ds\right) \\ &\leq C(\mathcal{B})\left(t_{2}-t_{1}\right). \end{aligned}$$

We also have

$$||II_{12}|| \leq ||F(X(t_1)||| \int_0^{t_1} (S(t_2 - s) - S(t_1 - s))ds||_{L(H)}$$

$$\leq C(\mathcal{B})||\int_0^{t_1} S(t_2 - s) - S(t_1 - s)ds||_{L(H)}.$$

Using the two transformations $y = t_2 - s$, $y = t_1 - s$, we find

$$\begin{aligned} \|II_{12}\| &= C(\mathcal{B}) \| \int_{t_2-t_1}^{t_2} S(y) dy - \int_0^{t_1} S(y) dy \|_{L(L^2(\Omega))} \\ &= C(\mathcal{B}) \| \int_{t_2-t_1}^{t_1} S(y) dy + \int_{t_1}^{t_2} S(y) dy - \int_0^{t_1} S(y) dy \|_{L(H)} \\ &= C(\mathcal{B}) \| \int_{t_1}^{t_2} S(y) dy - \int_0^{t_2-t_1} S(y) dy \|_{L(H)} \\ &\leq C(\mathcal{B})(t_2-t_1). \end{aligned}$$

The estimate of II_1 combined with (26) in the estimate of II_2 ends the proof.

3. Finite Element method for semi-linear parabolic problem

Finite element method for space discretization has been used in [27] for semilinear problem (1), time discretization has been performed using first order implicit and semi-implicit methods under the locally Lipschitz condition (4) or (5). The convergence proofs of parabolic stochastic partial differential equations with multiplicative or additive noise are provided in [20, 18, 19] where the space and time discretizations are performed using respectively finite element method and exponential integrators schemes. The convergence proof for deterministic problem (1) using locally Lipschitz condition (4) can easily be deducted by canceling the noise term in [20, 18, 19] and combined with different results in [27].

The keys features while using finite element method for space discretization comes from the fact that the corresponding semi-discrete problem shares the same bilinear form (6) with the continuous problem (1). The convergence proof for deterministic problem (1) using locally Lipschitz condition (4) or (5) will be more difficult with finite volume method (or finite difference method) for space discretization since the corresponding bilinear form of the semi-discrete problem (37) is different with the one of the continuous problem.

4. Finite volume for space discretization

4.1. Admissible mesh

A cell–centred finite volume methods for heterogeneous and anisotropic diffusion problems remains a challenging problem. An active area of research consists to make the approximation of the diffusion flux more efficient and simple as possible (see [24] for the references). The finite volume method is widely applied when the differential equations are in divergence form. To obtain a finite volume discretization, the domain Ω is subdivided into subdomains $(A_i)_{i \in \mathcal{I}}$, \mathcal{I} being the corresponding set of indices, called control volumes or control domains such that the collection of all those subdomains forms a partition of Ω . The common feature of all finite volume methods is to integrate the equation over each control volume A_i , $i \in \mathcal{I}$ and apply Gauss's divergence theorem to convert the volume integral to a surface integral. For our parabolic problem (2), finite volume methods differ in the way they approximate the diffusion flux $\mathcal{F} = -\mathbf{D}\nabla X$. Two techniques are mostly used: the finite volume with two-point flux approximation (TPFA) (see [6, 24]) and the finite volume with multi-point flux approximations (MPFA)([22, 23]).

An advantage of the two-point approximation is that it provides monotonicity properties, under the form of a local maximum principle. It is efficient and mostly used in industrial simulations. In this paper we use the TPFA as developed in [6]. The main drawback of TPFA is that it is applicable in the so called " admissible mesh " or "**D**-orthogonal mesh" and not in general mesh.

Definition 4.1. [Admissible mesh]

An admissible mesh \mathcal{T} for problem (2) with the full diffusion tensor **D** is defined by:

• A set $\{A_i\}_{i \in \mathcal{I}}$ of control volumes such that $\overline{\Omega} = \bigcup_{i \in \mathcal{I}} \overline{A_i}$ with the corresponding local inner product induced by $\mathbf{D}_{A_i}^{-1}$ where

$$\mathbf{D}_{A_i} = \frac{1}{\max(A_i)} \int_{A_i} \mathbf{D}(\mathbf{x}) d\mathbf{x}.$$

- The corresponding set of center points $\{\mathbf{x}_i\}_{i \in \mathcal{I}}$ such that
 - (a) $\mathbf{x}_i \in \overline{A}_i, i \in \mathcal{I}.$
 - (b) \mathbf{x}_i is the intersection of the straight lines perpendicular to the boundary of A_i with respect to the inner product induced by $\mathbf{D}_{A_i}^{-1}$.

Let $h = \operatorname{size}(\mathcal{T})$ be the maximum mesh size of \mathcal{T} . We denote by \mathcal{T}_h a dual Delaunay triangulation of \mathcal{T} i.e. a Delaunay triangulation where $\{\mathbf{x}_i\}_{i\in\mathcal{I}}$ is the set of vertices (2-D delaunay Triangulation with triangular mesh or 3-D delaunay Triangulation with tetrahedal mesh). For a given set $\{\mathbf{x}_i\}_{i\in\mathcal{I}}$, a dual mesh \mathcal{T}_h can easily be constructed with the Matlab function **delaunayTriangulation.m**.

Let us illustre Definition 4.1 to make it more understandable.

Example 1. In the case where the diffusion tensor **D** is diagonal and Ω is a rectangular or parallelepiped domain, any rectangular grid (d = 2) or parallelepiped grid (d = 3) is an admissible mesh. The set $\{\mathbf{x}_i\}$ is the set of centers of gravity of the rectangular grid or parallelepiped grid (see Figure 1). The inner product induced locally by $\mathbf{D}_{A_i}^{-1}$ is equivalent to the standard inner product corresponding to the Euclidean norm |.|. This mesh will yield a 5-point scheme (d = 2) and 7-point scheme (d = 3) for our model problem (2).



Figure 1: Example of admissible mesh in d = 2 for diagonal diffusion tensor **D**. The mesh \mathcal{T} is the set of rectangular grid (in black line) and a corresponding dual Delaunay triangulation \mathcal{T}_h (triangle mesh) is in red line. For d = 3, this figure can represent the upper view of the set of parallelepiped grid with the corresponding dual Delaunay triangulation (tetrahedal mesh). Note that the dual triangulation \mathcal{T}_h is not unique here.

If d = 2, for isotropic and heterogeneous media (**D**(**x**) = b(**x**)I₂ **x** ∈ Ω, I₂ being the identity matrix of dimension 2) we can define a triangular admissible mesh *T* to be a family of open triangular disjoint subsets of Ω such that two triangles having a common edge have also two common vertices. The angles of the triangles are assumed to be less than π/2 to allow the orthogonal bisectors to intersect inside each triangle, thus naturally defining the center point **x**_i of the control volume A_i. The finite volume scheme defined on such mesh will yield a 4-point scheme for our model problem (2). The inner product induced locally by **D**⁻¹_{A_i} is equivalence to the standard inner product corresponding to the Euclidean norm |.|.

4.2. Finite volume space discretization and semi-discrete solution

Consider the modified model problem of (2) where $c_0 X$ is added on both sides of the first equation of problem (2), c_0 is defined in (7). Consider an admissible mesh \mathcal{T} in the sense of Definition 4.1. Denote by \mathcal{E} the set of edges of control volume of \mathcal{T} , \mathcal{E}_{int} the set of interior edges of control volume of \mathcal{T} , $X_i(t)$ the approximation of X at time t at the center (or at any point) of the control volume $A_i \in \mathcal{T}$ and $X_{\sigma}(t)$ the approximation of X at time t at the center (or at any point) of the edge $\sigma \in \mathcal{E}$. For a control volume $A_i \in \mathcal{T}$, denote by \mathcal{E}_i the set of edges of A_i , mes (A_i) the Lebesgue measure of the control volume $A_i \in \mathcal{T}$.

As in [6, 25], integration over any control volume $A_i \in \mathcal{T}$, using the divergence theorem to convert the integral over A_i to a surface integral, finite differences for the diffusion flux approximation [6] and the upwind technique for the advection flux approximation yields

$$\begin{aligned}
\operatorname{mes}(A_{i})\frac{dX_{i}(t)}{dt} + \sum_{\sigma \in \mathcal{E}_{i}} \left(F_{i,\sigma}(t) + q_{i,\sigma}X_{\sigma,+}(t)\right) + c_{0}\operatorname{mes}(A_{i})X_{i}(t) \\
&= \operatorname{mes}(A_{i})f(\mathbf{x}_{i}, X_{i}(t)), \\
D_{i,\sigma} = |\mathbf{D}_{A_{i}} \mathbf{n}_{i,\sigma}|, \quad \mathbf{D}_{A_{i}} = \frac{1}{\operatorname{mes}(A_{i})}\int_{A_{i}}\mathbf{D}(\mathbf{x})d\mathbf{x}, \\
F_{i,\sigma}(t) = \operatorname{mes}(\sigma) D_{i,\sigma}\frac{X_{\sigma}(t) - X_{i}(t)}{d_{i,\sigma}}, \\
q_{i,\sigma} = \int_{\sigma} \mathbf{q} \cdot \mathbf{n}_{i,\sigma}d\sigma \qquad \qquad \forall A_{i} \in \mathcal{T}, \ \forall \sigma \in \mathcal{E}_{i}.
\end{aligned}$$
(27)

Here $\mathbf{n}_{i,\sigma}$ is the normal unit vector to σ outward to A_i , $\operatorname{mes}(\sigma)$ is the Lebesgue measure of the edge $\sigma \in \mathcal{E}_i$ and $d_{i,\sigma}$ the distance between the center of A_i and the edge σ .

Since the flux is continuous at the interface of two control volumes A_i and A_j (denoted by $i \mid j$) we therefore have $F_{i,\sigma}(t) = -F_{j,\sigma}(t)$ for $\sigma = i \mid j$, which yields

$$\begin{cases} F_{i,\sigma}(t) = -\tau_{\sigma} \left(X_{j}(t) - X_{i}(t) \right) = -\frac{\mu_{\sigma} \operatorname{mes}(\sigma)}{d_{i,j}} \left(X_{j}(t) - X_{i}(t) \right), \ \sigma = i \mid j \\ \tau_{\sigma} = \operatorname{mes}(\sigma) \frac{D_{i,\sigma} D_{j,\sigma}}{D_{i,\sigma} \ d_{i,\sigma} + D_{j,\sigma} d_{j,\sigma}} \quad (\text{transmissibility through } \sigma) \end{cases}$$
(28)

with

$$\mu_{\sigma} = d_{i,j} \frac{D_{i,\sigma} D_{j,\sigma}}{D_{i,\sigma} d_{i,\sigma} + D_{j,\sigma} d_{j,\sigma}},\tag{29}$$

where $d_{i,j}$ is the distance between the center of A_i and center of A_j . We will set $d_{i,j} = d_{i,\sigma}$ for $\sigma = \mathcal{E}_i \cap \partial \Omega$. For $\sigma \subset \partial \Omega$, we also write

$$F_{i,\sigma}(t) = -\tau_{\sigma} \left(X_{j}(t) - X_{i}(t) \right)$$
$$= -\frac{\operatorname{mes}(\sigma)\mu_{\sigma}}{d_{i,\sigma}} \left(X_{j}(t) - X_{i}(t) \right)$$

with

$$\begin{cases} X_j(t) = X_{\sigma}(t) = 0\\ \tau_{\sigma} = \frac{\operatorname{mes}(\sigma)D_{i,\sigma}}{d_{i,\sigma}}\\ \mu_{\sigma} = D_{i,\sigma}. \end{cases}$$
(30)

The upwind term for advection flux $X_{\sigma,+}$ is defined as

$$X_{\sigma,+}(t) = \begin{cases} X_i(t) & \text{if } q_{i,\sigma} \ge 0\\ & & \text{for } \sigma = i \mid j \end{cases}$$
(31)

$$X_{\sigma,+}(t) = \begin{cases} X_i(t) & \text{if } q_{i,\sigma} \ge 0\\ & & \text{for } \sigma \in \mathcal{E}_i \cap \partial \Omega. \end{cases}$$
(32)
$$X_{\sigma}(t) & \text{if } q_{i,\sigma} < 0 \end{cases}$$

We can write $X_{\sigma,+}$ as

$$X_{\sigma,+} = r_{\sigma} X_i(t) + (1 - r_{\sigma}) X_j(t), \quad \sigma = i \mid j$$
(33)

where $r_{\sigma} = \frac{1}{2}(\operatorname{sign}(q_{i,\sigma}) + 1)$. Note that according to (9), we have

$$(F(X_i))(\mathbf{x}_i) = f(\mathbf{x}_i, X_i(t)). \tag{34}$$

Using previous approximations, the finite volume space discretization for the model problem (2) is given by

$$\begin{cases} \operatorname{mes}(A_i) \frac{dX_i(t)}{dt} + \sum_{\sigma \in \mathcal{E}_i} \left(-\frac{\operatorname{mes}(\sigma) \mu_{\sigma}}{d_{i,j}} \left(X_j(t) - X_i(t) \right) \right. \\ \left. + q_{i,\sigma} \left(r_{\sigma} X_i(t) + (1 - r_{\sigma}) X_j(t) \right) \right) + c_0 \operatorname{mes}(A_i) X_i(t) = \operatorname{mes}(A_i) F(X_i(t)) \\ \left. X_i(t) = 0, \ d_{i,j} = d_{i,\sigma} \right. \\ \left. \operatorname{if} \sigma \subset \partial \Omega, \quad \forall A_i \in \mathcal{T}. \end{cases}$$

$$(35)$$

The scheme (35) clearly indicates the affinity of the finite volume method to the finite difference method. However, for the subsequent analysis it is more convenient to rewrite scheme (35) in a discrete variational form.

Multiplying the first equation of (35) by arbitrary numbers $v_i \in \mathbb{R}$ and summing the results over all control volume in \mathcal{T} yields

$$\begin{cases} \sum_{A_i \in \mathcal{T}} \left[\operatorname{mes}(A_i) \frac{dX_i(t)}{dt} + \sum_{\sigma \in \mathcal{E}_i} \left(\frac{\operatorname{mes}(\sigma) \mu_{\sigma}}{d_{i,j}} \left(X_i(t) - X_j(t) \right) \right. \\ \left. + q_{i,\sigma} \left(r_{\sigma} X_i(t) + (1 - r_{\sigma}) X_j(t) \right) \right] v_i = \sum_{A_i \in \mathcal{T}} \operatorname{mes}(A_i) F(X_i(t)) v_i. \end{cases}$$
(36)

Let V_h denote the space of continuous functions that are piecewise linear over the Delaunay triangulation \mathcal{T}_h (dual of \mathcal{T}), and $X(\mathcal{T})$ be the space of the functions constant in each control volume of \mathcal{T} , the following lemma creates a one-to-one correspondence between V_h and $X(\mathcal{T})$.

Lemma 4.1. There exists a one-to-one correspondence between the space V_h and $X(\mathcal{T})$, more precisely we have :

- For any $U_h \in V_h$ corresponds the unique function $U = (U_h(\mathbf{x}_i))_{i \in \mathcal{I}} \in X(\mathcal{T})$.
- For any function $U = (U_i)_{i \in \mathcal{I}} \in X(\mathcal{T})$, there exists an unique function $U_h \in V_h$ such that $U_h(\mathbf{x}_i) = U_i$.

Proof. The first correspondence is obvious. The proof of the second correspondence can be found in [25, Lemma 2.10, p.58].

Let us consider equation (36), according to Lemma 4.1, there are unique functions $X_h(t), v_h \in V_h$ such that $X_h(t)(\mathbf{x}_i) = X_i(t)$ and $v_h(\mathbf{x}_i) = v_i$ for all $A_i \in \mathcal{T}$, where \mathbf{x}_i is a center of the control volume $A_i \in \mathcal{T}$ (\mathbf{x}_i is also a vertex in \mathcal{T}_h).

Denote by a_h the bilinear form defined by

$$\begin{cases}
 a_h(u_h, v_h) = \sum_{A_i \in \mathcal{T}_{\sigma} \in \mathcal{E}_i} \left(-\frac{\operatorname{mes}(\sigma) \mu_{\sigma}}{d_{i,j}} \left(u_j - u_i \right) + q_{i,\sigma} \left(r_{\sigma} u_i + (1 - r_{\sigma}) u_j \right) \right) v_i \\
 + c_0 \operatorname{mes}(A_i) u_i v_i \qquad \qquad \forall u_h, v_h \in V_h,
\end{cases}$$
(37)

and by $\langle ., . \rangle_{0,h}$ the scalar product on $C(\overline{\Omega}) \supset V_h$ defined by

$$\langle u, v \rangle_{0,h} = \sum_{i \in \mathcal{T}} \operatorname{mes}(A_i) u_i v_i, \quad u_i = u(\mathbf{x}_i), \quad v_i = v(\mathbf{x}_i), \quad u, v \in C(\overline{\Omega}).$$
 (38)

Note that this scalar product can be extended in $L^2(\Omega)$ (see [30]) by

$$\langle u, v \rangle_{0,h} = \sum_{i \in \mathcal{T}} \operatorname{mes}(A_i) u_i v_i, \quad u_i = \frac{1}{\operatorname{mes}(A_i)} \int_{A_i} u dx, \quad v_i = \frac{1}{\operatorname{mes}(A_i)} \int_{A_i} v dx, \quad u, v \in L^2(\Omega).(39)$$

Note also that when $u \in C(\overline{\Omega})$ we will use $u_i = u(\mathbf{x}_i)$ in (39). We can easily observe for $u, v \in X(\mathcal{T})^2 \langle u, v \rangle_{0,h} = (u, v)$. The corresponding norm of $\langle ., . \rangle_{0,h}$ is the discrete $L^2(\Omega)$ norm denoted by $\|.\|_{0,h}$. We therefore have the following variational form of our finite volume scheme (36).

$$\begin{cases} \langle \frac{d}{dt} X_h, \varphi \rangle_{0,h} + a_h(X_h(t), \varphi) = \langle F(X_h(t)), \varphi \rangle_{0,h}, \quad \forall \varphi \in V_h, \quad t \in (0, T], \\ X_h(0) = X_{h\,0}. \end{cases}$$
(40)

Consider the operator $A_h: V_h \to V_h$ such that

$$\langle A_h \psi, \chi \rangle_{0,h} = a_h(\psi, \chi) \quad \forall \psi, \chi \in V_h.$$
 (41)

The semidiscrete solution in V_h is then given by: find $X_h(t) \in V_h$ such that

$$\begin{cases} \frac{dX_h}{dt} + A_h X_h = P_h F(X_h) & t \in (0,T] \\ X_h(0) = X_{0h} \end{cases}$$

$$\tag{42}$$

 $^{^2}$ Remember that this is the space of the functions constant in each control volume of \mathcal{T}

where P_h is the orthogonal projection defined from $L^2(\Omega)$ to V_h by

$$\langle P_h u, \chi \rangle_{0,h} = \langle u, \chi \rangle_{0,h} \quad \forall \chi \in V_h, \ u \in L^2(\Omega).$$
 (43)

In order to provide the corresponding mild form of (42), let us define the discrete $H_0^1(\Omega)$ norm.

Definition 4.2. [Discrete $H_0^1(\Omega)$ norm [6]]

Let \mathcal{T} be an admissible finite volume mesh in the sense of Definition 4.1, and $X(\mathcal{T})$ the space of the functions constant in each control volume of \mathcal{T} . For $u \in X(\mathcal{T})$ corresponding to $u_h \in V_h$ (according to Lemma 4.1), the discrete $H_0^1(\Omega)$ norm of u and u_h is defined by

$$\|u_h\|_{1,\mathcal{T}} := \|u\|_{1,\mathcal{T}} := \left(\sum_{\sigma \in \mathcal{E}} \tau'_\sigma \left(D_\sigma u\right)^2\right)^{1/2} \tag{44}$$

where

$$\tau'_{\sigma} = \frac{mes(\sigma)}{d_{\sigma}}$$

$$D_{\sigma}u = |u_i - u_j| \quad if \quad \sigma = i|j \in \mathcal{E}_{int}$$

$$D_{\sigma}u = |u_i| \quad if \quad \sigma \in \partial\Omega.$$

Note that this norm is equivalent in $V = H_0^1(\Omega) = \mathcal{D}(A^{1/2}) = \mathcal{D}(A^{*1/2})$ to the natural norm of $H^1(\Omega)$ which is $\|.\|_1$. Following closely [27, 28] we also have $\mathcal{D}(A_h^{1/2}) = \mathcal{D}(A_h^{*1/2})$ with the following norm equivalence

$$||A_h^{1/2}u|| \equiv ||A_h^{*1/2}u|| \equiv ||u||_1, \qquad u \in V_h.$$

We make the following assumption as in [6, Theorem 3.8], very useful for our convergence proof.

Assumption 4.1. [Regularity of D, q and \mathcal{T}]

We assume that **D** is bounded³, the restriction of **D** to any $A_i \in \mathcal{T}$ belongs to $C^1(A_i, \mathbb{R}^{d \times d}), q_j \in C^1(\overline{\Omega})$, the discontinuities of **D** coincide with the interfaces of the mesh, and that there exists $\zeta_1 > 0$ such that

$$\zeta_1 h \le d_{i,\sigma}, \qquad \forall \ A_i \in \mathcal{T}, \ \forall \sigma \in \mathcal{E}_i, \tag{45}$$

where $h = size(\mathcal{T})$.

The inequality (45) is called regularity property of the mesh \mathcal{T} .

Remark 4.1. The regularity property of the dual mesh \mathcal{T}_h given in [25, Definition 3.28, p 138] states that there exists some constant c > 0 such that

$$\frac{h_K}{\rho_K} \leqslant c, \qquad \forall K \in \mathcal{T}_h$$

$$\tag{46}$$

³From (3), \mathbf{D} is bounded below, to be bounded \mathbf{D} also need to be bounded above.

where $h_K = diam(K) = \sup_{(x,y)\in K^2} d(x,y)$ and $\rho_K = \sup\{diam(S) \mid S \text{ is a ball in } \mathbb{R}^d \text{ and } S \subset \mathbb{R}^d$ $K\}.$

As we are dealing in \mathcal{T}_h with triangle or tetrahedron, h_K denotes the longest edge and ρ_K the diameter of the inscribed circle (d = 2) or sphere (d = 3). Using Heron's formula and its consequences, one can prove that the regularity of mesh \mathcal{T} given by (45) implies the regularity of the dual mesh \mathcal{T}_h given by (46).

Assumption 4.1 allows the following V_h – ellipticity of a_h .

Theorem 4.1. Under the regularity of the admissible mesh \mathcal{T} in Assumption 4.1, there exists a constant $\alpha > 0$ independent of h such that

$$a_h(v_h, v_h) \ge \alpha \|v_h\|_{1,\mathcal{T}}^2 \quad \forall v_h \in V_h.$$

$$\tag{47}$$

Proof. Let b_h^1 , b_h^2 and b_h^3 the bilinear forms defined in $V_h \times V_h$ by

$$b_h^1(u_h, v_h) = \sum_{A_i \in \mathcal{T}\sigma \in \mathcal{E}_i} - \frac{\operatorname{mes}(\sigma) \,\mu_\sigma}{d_{i,j}} \left(u_j - u_i \right) v_i, \tag{48}$$

$$b_h^2(u_h, v_h) = \sum_{A_i \in \mathcal{T}\sigma \in \mathcal{E}_i} q_{i,\sigma} \left(r_\sigma u_i + (1 - r_\sigma) u_j \right) v_i = \sum_{A_i \in \mathcal{T}\sigma \in \mathcal{E}_i} q_{i,\sigma} u_{\sigma,+} v_i, \tag{49}$$

$$b_h^3(u_h, v_h) = c_0 \sum_{A_i \in \mathcal{T}} \operatorname{mes}(A_i) \, u_i v_i.$$
(50)

Note that according to Lemma 4.1, we have identified $u_h \in V_h$ and $v_h \in V_h$ to their correspondent $(u_i)_{i \in \mathcal{T}} = (u_h(\mathbf{x}_i))_{i \in \mathcal{T}} \in X(\mathcal{T})$ and $(v_i)_{i \in \mathcal{T}} = (v_h(\mathbf{x}_i)) \in X(\mathcal{T})$ in the definition of b_h^1, b_h^2 and b_h^3 .

Using Assumption 4.1, mainly the regularity of \mathcal{T} ($\zeta_1 h \leq d_{i,\sigma} \leq h$) and the fact that the coefficients of the diffusion tensor **D** are bounded, there exists two constants $C_5(\Omega, \zeta_1, \mathbf{D})$ and $C'_5(\Omega,\zeta_1,\mathbf{D})$ such that

$$C_5 \le \mu_{\sigma} = d_{i,j} \frac{D_{i,\sigma} D_{j,\sigma}}{D_{i,\sigma} d_{i,\sigma} + D_{j,\sigma} d_{j,\sigma}} \le C'_5, \qquad \sigma = i|j,$$
(51)

and

$$C_5 \le \mu_{\sigma} = D_{i,\sigma} \le C'_5, \qquad \sigma \in \mathcal{E}_i \cap \partial\Omega,$$
(52)

so that

$$C_5 \le \mu_\sigma \le C_5', \quad \forall \sigma \in \mathcal{E},$$
(53)

where μ_{σ} is defined in (30) and (29).

Using the fact that the transmissibility given in (28) is symmetric, i.e. $\tau_{i|j} = \tau_{j|i}$ and reorganizing the summation, we therefore have

$$C_5 \|v_h\|_{1,\mathcal{T}}^2 \le b_h^1(v_h, v_h) \le C_5' \|v_h\|_{1,\mathcal{T}}^2.$$
(54)

Let use some important results from [6]. Indeed as in [6] reordering the summation over the set of edges yields

$$b_h^2(v_h, v_h) = \sum_{\sigma \in \mathcal{E}} q_\sigma \left(v_{\sigma, +} - v_{\sigma, -} \right) v_{\sigma, +}$$
(55)

where

$$v_{\sigma,-} = \begin{cases} v_i & \text{if } q_{i,\sigma} \leq 0 \\ v_j & \text{(or } v_\sigma) & \text{if } q_{i,\sigma} > 0 \end{cases} \quad \sigma \in \mathcal{E}_{int} (\text{or } \sigma \in \mathcal{E}_i \cap \partial \Omega), \tag{56}$$

$$q_{\sigma} = |\int_{\sigma} \mathbf{q} \cdot \mathbf{n}_{i,\sigma} d\sigma|.$$
(57)

Note that

$$\sum_{\sigma \in \mathcal{E}} q_{\sigma} \left(v_{\sigma,+} - v_{\sigma,-} \right) v_{\sigma,+} = \frac{1}{2} \sum_{\sigma \in \mathcal{E}} q_{\sigma} \left(\left(v_{\sigma,+} - v_{\sigma,-} \right)^2 + \left(v_{\sigma,+}^2 - v_{\sigma,-}^2 \right) \right).$$
(58)

As we have assumed divergence-free flow, we therefore have

$$\sum_{\sigma \in \mathcal{E}} q_{\sigma} \left(v_{\sigma,+}^2 - v_{\sigma,-}^2 \right) = \sum_{A_i \in \mathcal{T}} \left(\int_{A_i} \mathbf{q} \cdot \mathbf{n}_{i,\sigma} d\sigma \right) v_i^2 = \int_{\Omega} \nabla \cdot \mathbf{q}(\mathbf{x}) v_h^2(\mathbf{x}) dx = 0.$$
(59)

Then from (58), we have

$$b_h^2(v_h, v_h) \ge 0.$$
 (60)

We also have

$$b_h^3(v_h, v_h) = c_0 \|v_h\|_{0,h} \ge 0,$$
(61)

Combining (54),(60) and (61) yields

$$a_h(v_h, v_h) \ge C_5 \|v_h\|_{1,\mathcal{T}}^2 \quad \forall v_h \in V_h.$$

$$\tag{62}$$

So we should take $\alpha = C_5$.

The following V_h – ellipticity of a_h implies that $-A_h$ is a sectorial on $H = L^2(\Omega)$ (uniformly in h) i.e. there exists $C_1, \theta \in (\frac{1}{2}\pi, \pi)$, such that

$$\|(\lambda I + A_h)^{-1}\|_{L(H)} \le \frac{C_1}{|\lambda|}, \qquad \lambda \in S_{\theta},$$
(63)

where $S_{\theta} = \{\lambda \in \mathbb{C} : \lambda = \rho e^{i\phi}, \ \rho > 0, \ 0 \le |\phi| \le \theta \}.$

The discrete operator $-A_h$ therefore is the infinitesimal generator of bounded analytic semigroup (or exponential operator) $S_h(t) := e^{-tA_h}$ on V_h such that

$$S_h(t) := e^{-tA_h} = \frac{1}{2\pi i} \int_{\mathcal{C}'} e^{t\lambda} (\lambda I + A_h)^{-1} d\lambda , \qquad t > 0 \qquad (64)$$

where C' denotes a path that surrounds the spectrum of $-A_h$. As for the continuous case, Theorem 2.1 and Duhamel's principle ensure the existence and uniqueness of the solution of (42) represented by the following integral equations (mild form)

$$X_h(t) = S_h(t)X_{0h} + \int_0^t S_h(t-s)P_hF(X_h(s))ds, \qquad t \in [0,T].$$
(65)

The solution X converges to the semi-discrete solution X_h according to the following theorem.

Theorem 4.2. Let $\mathcal{B} \subset V$ be bounded, consider the solution X of (2) and the semi-discrete solution X_h of (42) represented by (35) or (65) in the interval $[0, t^*]$, $t^* = t^*(\mathcal{B})$ defined in Theorem 2.1, such that $X(t) \in \mathcal{B}$ and $X_h(t) \in \mathcal{B} \bigcap V_h$ for all $t \leq T \leq t^*$. We assume that the unique mild solution X of (2) is the classical solution (i.e. X is twice continuously differentiable with respect to \mathbf{x} and differentiable with respect to t), Assumption 4.1 is satisfied and the reaction function F satisfies (20). Furthermore assume that $X_0 \in C(\overline{\Omega}) \bigcap \mathcal{B}$, $X_{0h} \in$ $V_h \bigcap \mathcal{B}$ and $f(\mathbf{x}, u)$ is differentiable respect to \mathbf{x} and u with

$$|f_{\mathbf{x}}(\mathbf{x}, u)| + |f_{u}(\mathbf{x}, u)| \le C(1 + |u|^{\gamma}), \quad \forall \mathbf{x} \in \Omega, \ u \in \mathbb{R},$$
(66)

for function of type f of type (4) and

$$|f_{\mathbf{x}}(\mathbf{x}, u)| + |f_{u}(\mathbf{x}, u)| \le C(1 + |u| + |u|^{\gamma}), \quad \forall \mathbf{x} \in \Omega, \ u \in \mathbb{R},$$
(67)

for function of type f of type (5), then the following estimate holds

$$||X(t) - X_h(t)||_{0,h} \le C(\mathcal{B})(||X_0 - X_{0h}||_{0,h} + h), \qquad \forall t \in [0,T],$$

where $C = C(\mathcal{B}, \Omega, X, F, \mathbf{D}, \mathbf{q}, T, \zeta_1).$

Before given the proof, let us give this lemma which will be very useful in the proof.

As for the elliptic case [6, Proof of Theorem 2.3 or Theorem 3.8] or linear parabolic case [6, Proof of Theorem 4.1], we have the following fluxes consistency.

Lemma 4.2. / Fluxes consistency

Let $R_{i,\sigma}(t)$ and $r_{i,\sigma}(t)$ be respectively the errors of diffusion flux and advective flux through the the edge σ (interface of control volume A_i and control volume A_j , or edge of control volume A_i if $\sigma \subset \partial \Omega$) at time t given by

$$R_{i,\sigma}(t) = \frac{1}{\operatorname{mes}(\sigma)} \left[\frac{\operatorname{mes}(\sigma)\mu_{\sigma}}{d(i,j)} \left(X(\mathbf{x}_{i},t) - X(\mathbf{x}_{j},t) \right) - \int_{\sigma} -\mathbf{D}\nabla X \cdot \mathbf{n}_{i,\sigma} d\sigma \right],$$
(68)
(69)

$$r_{i,\sigma}(t) = \frac{1}{\operatorname{mes}(\sigma)} \left[q_{i,\sigma} X(\mathbf{x}_{\sigma,+},t) - \int_{\sigma} \mathbf{q} X(t) \cdot \mathbf{n}_{i,\sigma} \right],$$

 4 where

$$\begin{cases} \mathbf{x}_{\sigma,+} = \begin{cases} \mathbf{x}_{i} & if \, \mathbf{q} \cdot \mathbf{n}_{\sigma} \ge 0, \\ \mathbf{x}_{j} & if \, \mathbf{q} \cdot \mathbf{n}_{\sigma} < 0, \end{cases} & \sigma = i | j, \\ \mathbf{x}_{j} & if \, \mathbf{q} \cdot \mathbf{n}_{\sigma} < 0, \end{cases} \\ \mathbf{x}_{\sigma,+} = \begin{cases} \mathbf{x}_{i} & if \, \mathbf{q} \cdot \mathbf{n}_{\sigma} \ge 0, \\ \mathbf{x}_{\sigma}, & \mathbf{x}_{\sigma} \in \partial\Omega & if \, \mathbf{q} \cdot \mathbf{n}_{\sigma} < 0 \end{cases} & \sigma \in \mathcal{E}_{i} \cap \partial\Omega. \end{cases}$$
(70)

Under the Assumption 4.1, if X is twice continuously differentiable with respect to \mathbf{x} , there exists three positive constants C_2 , C'_2 and C_3 such that

$$\begin{cases}
|R_{i,\sigma}(t)| \leq C_2 \left(\mathbf{D}, X, T, \Omega \right) h, \\
|r_{i,\sigma}(t)| \leq C'_2 \left(\mathbf{q}, X, T, \Omega \right) h, \\
|R_{i,\sigma}(t)| + |r_{i,\sigma}(t)| \leq C_3 \left(\mathbf{q}, \mathbf{D}, X, T\Omega \right) h.
\end{cases}$$
(71)

Proof. The proof of this lemma can be done in the same manner as the one in [6] in 1 D. Let us provide some details. By setting

$$\overline{F}_{i,\sigma} = \int_{\sigma} -\mathbf{D}\nabla X \cdot \mathbf{n}_{i,\sigma} d\sigma, \quad F_{i,\sigma}^* = \frac{\operatorname{mes}(\sigma)\mu_{\sigma}}{d(i,j)} \left(X(\mathbf{x}_i, t) - X(\mathbf{x}_j, t) \right) = -\tau_{\sigma} \left(X(\mathbf{x}_j, t) - X(\mathbf{x}_i, t) \right), (72)$$

let us prove that there exists $C_2 = C_2(\mathbf{D}, X, T, \Omega)$ such that

$$F_{i,\sigma}^* = \overline{F}_{i,\sigma} + R_{i,\sigma}(t), \text{ where } |R_{i,\sigma}| \le C_2 \operatorname{mes}(\sigma)h.$$
 (73)

Case 1: $\sigma = A_i \mid A_j \in \mathcal{E}_i$. Let $\mathbf{y}_{\sigma} = \sigma \bigcap \mathcal{D}_{i,\sigma} = \sigma \bigcap \mathcal{D}_{j,\sigma}$, where $\mathcal{D}_{i,\sigma}$ and $\mathcal{D}_{j,\sigma}$ are respectively the straight lines perpendicular to $\sigma = A_i \mid A_j$ with respect to the inner product induced by $\mathbf{D}_{A_i}^{-1}$ and $\mathbf{D}_{A_j}^{-1}$. Let us set

$$F_{i,\sigma}^{*,i} = -D_{i,\sigma}\operatorname{mes}(\sigma)\frac{X(\mathbf{y}_{\sigma},t) - X(\mathbf{x}_{i},t)}{d_{i,\sigma}}, \quad F_{j,\sigma}^{*,j} = -D_{j,\sigma}\operatorname{mes}(\sigma)\frac{X(\mathbf{y}_{\sigma},t) - X(\mathbf{x}_{j},t)}{d_{i,\sigma}}.$$
 (74)

Since X is twice continuously differentiable with respect to \mathbf{x} , using Assumption 4.1 and Definition 4.1 and combined with Taylor expansion yields

$$F_{i,\sigma}^{*,i} = \overline{F}_{i,\sigma} + t_{i,\sigma}^{i} \quad \text{with} \quad |t_{i,\sigma}^{i}| \le \alpha_1 \text{mes}(\sigma)h \tag{75}$$

$$F_{j,\sigma}^{*,j} = \overline{F}_{j,\sigma} + t_{j,\sigma}^j \quad \text{with} \quad |t_{j,\sigma}^j| \le \alpha_1 \text{mes}(\sigma)h \tag{76}$$

First of all let us proof the (75). As the restriction of \mathbf{D} to A_i is differentiable, there exists a positive constant $C = C(\mathbf{D})$ such that

$$\mathbf{D}(\mathbf{x}) = \mathbf{D}_{A_i} + \mathbf{m}_i \qquad \forall \, \mathbf{x} \in \overline{A_i}, \quad \text{with} \quad \|\mathbf{m}_i\|_{\mathbb{R}^{d \times d}} \le Ch, \tag{77}$$

⁴Remember that $X(\mathbf{x}_j, t) = X(\mathbf{x}_\sigma, t) = 0$ if $\sigma \subset \partial \Omega$.

So, using the fact that X is differentiable yields

$$\int_{\sigma} \mathbf{D}\nabla X \cdot \mathbf{n}_{i,\sigma} d\sigma = \int_{\sigma} \mathbf{D}_{A_i} \nabla X \cdot \mathbf{n}_{i,\sigma} d\sigma + M_i, \text{ with } |M_i| \le C(\Omega, T, \mathbf{D}) \operatorname{mes}(\sigma) h.$$
(78)

As X is twice continuously differentiable with respect to \mathbf{x} , Taylor expansion yields

$$X(\mathbf{y}_{\sigma},t) - X(\mathbf{x}_{i},t) = \nabla X \cdot (\mathbf{y}_{\sigma} - \mathbf{x}_{i}) + \int_{0}^{1} H(X)(t\mathbf{x}_{i} + (1-t)\mathbf{y}_{\sigma})(\mathbf{y}_{\sigma} - \mathbf{x}_{i}) \cdot (\mathbf{y}_{\sigma} - \mathbf{x}_{i})tdt.$$

where H(X)(z) denotes the Hessian matrix of X at point z. Note that

$$\mathbf{y}_{\sigma} - \mathbf{x}_i = d_{i,\sigma} \mathbf{n}_{i,\sigma}^*,\tag{79}$$

where $\mathbf{n}_{i,\sigma}^*$ is the normal unit vector to σ outward to A_i with respect to the inner product induced by $\mathbf{D}_{A_i}^{-1}$, which is different to $\mathbf{n}_{i,\sigma}$ (the normal unit vector to σ outward to A_i with the with respect to the inner product of \mathbb{R}^d). We therefore have

$$\frac{X(\mathbf{y}_{\sigma},t) - X(\mathbf{x}_{i},t)}{d_{i,\sigma}} = \nabla X \cdot \mathbf{n}_{i,\sigma}^{*} + \frac{1}{d_{i,\sigma}} \int_{0}^{1} H(X)(t\mathbf{x}_{i} + (1-t)\mathbf{y}_{\sigma})(\mathbf{y}_{\sigma} - \mathbf{x}_{i}) \cdot (\mathbf{y}_{\sigma} - \mathbf{x}_{i})tdt.$$
(80)

Note that by definition of the scalar product induced by $\mathbf{D}_{A_i}^{-1}$ (Definition 4.1), we have

$$(\mathbf{y}_{\sigma} - \mathbf{a})^T \mathbf{D}_{A_i}^{-1} \mathbf{n}_{i,\sigma}^* = 0, \quad \forall \mathbf{a} \in \sigma.$$
(81)

Since

$$(\mathbf{y}_{\sigma} - \mathbf{a})^T \mathbf{D}_{A_i}^{-1} (\mathbf{D}_{A_i} \mathbf{n}_{i,\sigma}) = (\mathbf{y}_{\sigma} - \mathbf{a})^T \mathbf{n}_{i,\sigma} = 0, \quad \forall \mathbf{a} \in \sigma,$$
(82)

we can therefore take

$$\mathbf{n}_{i,\sigma}^{*} = \frac{\mathbf{D}_{A_{i}}\mathbf{n}_{i,\sigma}}{|\mathbf{D}_{A_{i}}\mathbf{n}_{i,\sigma}|}.$$
(83)

So, (80) becomes

$$|\mathbf{D}_{A_i}\mathbf{n}_{i,\sigma}| \frac{X(\mathbf{y}_{\sigma}, t) - X(\mathbf{x}_i, t)}{d_{i,\sigma}}$$
(84)

$$= \nabla X \cdot \mathbf{D}_{A_i} \mathbf{n}_{i,\sigma} + \frac{|\mathbf{D}_{A_i} \mathbf{n}_{i,\sigma}|}{d_{i,\sigma}} \int_0^1 H(X) (t\mathbf{x}_i + (1-t)\mathbf{y}_{\sigma}) (\mathbf{y}_{\sigma} - \mathbf{x}_i) \cdot (\mathbf{y}_{\sigma} - \mathbf{x}_i) t dt.$$
(85)

Using the fact that \mathbf{D}_{A_i} is symmetric, we have

$$\mathbf{D}_{A_i} \nabla X \cdot \mathbf{n}_{i,\sigma} = (\mathbf{D}_{A_i} \nabla X)^T \mathbf{n}_{i,\sigma} = (\nabla X)^T (\mathbf{D}_{A_i})^T \mathbf{n}_{i,\sigma} = (\nabla X)^T \mathbf{D}_{A_i} \mathbf{n}_{i,\sigma} = \nabla X \cdot \mathbf{D}_{A_i} \mathbf{n}_{i,\sigma}.$$
 (86)

By integrating (84) over σ , using the fact that X is twice differentiable respect to **x**, combining (78) and (86) yields

$$F_{i,\sigma}^{*,i} = \overline{F}_{i,\sigma} + t_{i,\sigma}^{i} \quad \text{with} \quad |t_{i,\sigma}^{i}| \le \alpha_{1}(\mathbf{D}, X, T, \Omega) \operatorname{mes}(\sigma)h,$$
(87)

and this conclude the proof of (75).

Let us continue with the proof of the lemma. The continuity of the fluxes, i.e. $\overline{F}_{i,\sigma} = -\overline{F}_{j,\sigma}$ implies

$$X(\mathbf{y}_{\sigma},t) = -\frac{t_{i,\sigma}^{i} + t_{j,\sigma}^{j}}{\left(\frac{D_{i,\sigma}}{d_{i,\sigma}} + \frac{D_{j,\sigma}}{d_{j,\sigma}}\right)\operatorname{mes}(\sigma)} + \frac{1}{\left(\frac{D_{i,\sigma}}{d_{i,\sigma}} + \frac{D_{j,\sigma}}{d_{j,\sigma}}\right)} \left(\frac{D_{i,\sigma}}{d_{i,\sigma}}X(\mathbf{x}_{i},t) + \frac{D_{j,\sigma}}{d_{j,\sigma}}X(\mathbf{x}_{j},t)\right).$$
(88)

We therefore have

$$F_{i,\sigma}^{*,i} = \frac{D_{i,\sigma}}{d_{i,\sigma}} \frac{t_{i,\sigma}^i + t_{j,\sigma}^j}{\left(\frac{D_{i,\sigma}}{d_{i,\sigma}} + \frac{D_{j,\sigma}}{d_{j,\sigma}}\right)} - \tau_{\sigma}(X(\mathbf{x}_j, t) - X(\mathbf{x}_i, t)).$$
(89)

Combining (75) and (89) yields

$$\overline{F}_{i,\sigma} = F_{i,\sigma}^{*,i} - t_{i,\sigma}^{i} = \frac{D_{i,\sigma}}{d_{i,\sigma}} \frac{t_{i,\sigma}^{i} + t_{j,\sigma}^{j}}{\left(\frac{D_{i,\sigma}}{d_{i,\sigma}} + \frac{D_{j,\sigma}}{d_{i,\sigma}}\right)} + F_{i,\sigma}^{*} - t_{i,\sigma}^{i}$$
(90)

$$= F_{i,\sigma}^* - R_{i,\sigma} \tag{91}$$

where

$$-R_{i,\sigma} = \frac{D_{i,\sigma}}{d_{i,\sigma}} \frac{t_{i,\sigma}^{i} + t_{j,\sigma}^{j}}{\left(\frac{D_{i,\sigma}}{d_{i,\sigma}} + \frac{D_{j,\sigma}}{d_{j,\sigma}}\right)} - t_{i,\sigma}^{i}$$

$$(92)$$

$$|R_{i,\sigma}| \leq |t_{i,\sigma}^{i}| + \left|\frac{D_{i,\sigma}}{d_{i,\sigma}}\frac{t_{i,\sigma}^{i} + t_{j,\sigma}^{j}}{\left(\frac{D_{i,\sigma}}{d_{i,\sigma}} + \frac{D_{j,\sigma}}{d_{j,\sigma}}\right)}\right|$$
(93)

$$\leq |t_{i,\sigma}^{i}| + \frac{D_{i,\sigma}d_{j,\sigma}}{d_{j,\sigma}D_{i,\sigma} + d_{i,\sigma}D_{j,\sigma}} \left(|t_{i,\sigma}^{i}| + |t_{j,\sigma}^{j}|\right)$$

$$\tag{94}$$

$$\leq 3\alpha_1 \operatorname{mes}(\sigma)h,$$
 (95)

since

$$\frac{D_{i,\sigma}d_{j,\sigma}}{d_{j,\sigma}D_{i,\sigma} + d_{i,\sigma}D_{j,\sigma}} \le 1.$$
(96)

Case 2: $\sigma \in \partial \Omega \bigcap \mathcal{E}_i$. As we have in **Case 1**, since X is twice continuously differentiable with respect to **x**, using Assumption 4.1 and Definition 4.1 combined with Taylor expansion yields

$$F_{i,\sigma}^{*} = \frac{\operatorname{mes}(\sigma)D_{i,\sigma}}{d_{i,\sigma}}\left(X(\mathbf{x}_{i},t)\right) = \tau_{\sigma}X(\mathbf{x}_{i},t) = \overline{F}_{i,\sigma} + R_{i,\sigma}, \text{ with } |R_{i,\sigma}| \le \alpha \operatorname{mes}(\sigma)h.$$
(97)

To conclude the proof of the diffusion error $R_{i,\sigma}(t)$, we can take $C_2 = 3\alpha_1$.

The proof of the convection error $r_{i,\sigma}(t)$ is done by using Tayor expansion of X and **q** (since **q** is assumed to be differentiable according to Assumption 4.1). Some details can be found in [6].

Let us now prove our first main result in Theorem 4.2.

Proof. Integrating the shifted version (by adding $c_0 X$ in both size) of equation (2) over each control volume $A_i \in \mathcal{T}$ and using the divergence theorem yields

$$\int_{A_i} X_t(\mathbf{x}, t) d\mathbf{x} - \sum_{\sigma \in \mathcal{E}_i} \int_{\sigma} (\mathbf{D}\nabla X - \mathbf{q}X) \cdot \mathbf{n}_{\sigma} d\sigma + c_0 \int_{A_i} X d\mathbf{x} = \int_{A_i} f(\mathbf{x}, X(\mathbf{x}, t)) d\mathbf{x}.$$
 (98)

For $t \in [0, T]$, $A_i \in \mathcal{T}$ and $\sigma \in \mathcal{E}_i$ using the same notation as in [6], let us set

$$\begin{cases} p_i(t) = X(\mathbf{x}_i, t) - \frac{1}{\operatorname{mes}(A_i)} \int_{A_i} X(\mathbf{x}, t) d\mathbf{x}, \\ \varrho_i(t) = \frac{1}{\operatorname{mes}(A_i)} \int_{A_i} f(\mathbf{x}, X(\mathbf{x}, t)) d\mathbf{x} - f(\mathbf{x}_i, X_i(t)). \end{cases}$$
(99)

As we have assumed that the unique solution X of (2) is the regular, Taylor expansion yields

$$\begin{cases} X_t(\mathbf{x},t) = X_t(\mathbf{x}_i,t) + s_i(\mathbf{x},t), & |s_i(\mathbf{x},t)| \le C_1(X,T) h \\ \int_{A_i} X_t(\mathbf{x},t) d\mathbf{x} = \operatorname{mes}(A_i) X_t(\mathbf{x}_i,t) + S_i, & S_i = \int_{A_i} s_i(\mathbf{x},t) d\mathbf{x}, & |S_i| \le \operatorname{mes}(A_i) C_1(X,T) h. \end{cases}$$
(100)

Using again the regularity of the solution X, we also have

$$X(\mathbf{x},t) = X(\mathbf{x}_i,t) + s'_i(\mathbf{x},t), \qquad |s'_i(\mathbf{x},t)| \le C'_1(X,T)h,$$

therefore

$$|p_i(t)| \le C'_3(X,T) h.$$
(101)

Using the expressions (99) and (100) in (98) yields the following decomposition of our initial continuous problem (98)

$$\operatorname{mes}(A_{i})X_{t}(\mathbf{x}_{i},t) + S_{i} - \sum_{\sigma \in \mathcal{E}_{i}} \operatorname{mes}(\sigma) \left(R_{i,\sigma}(t) + r_{i,\sigma}(t)\right) + \sum_{\sigma \in \mathcal{E}_{i}} \left[\frac{\operatorname{mes}(\sigma)\mu_{\sigma}}{d(i,j)} \left(X(\mathbf{x}_{i},t) - X(\mathbf{x}_{j},t)\right)\right] + \sum_{\sigma \in \mathcal{E}_{i}} \left[q_{i,\sigma}X(\mathbf{x}_{\sigma,+},t)\right] + c_{0}\operatorname{mes}(A_{i}) \left(X(\mathbf{x}_{i},t) - p_{i}(t)\right) = \int_{A_{i}} f(\mathbf{x},X(\mathbf{x},t))d\mathbf{x}.$$
(102)

Let $X_h(t) \in V_h$ solution of (40) ⁵ such that $X_h(t)(\mathbf{x}_i) = X_i(t)$. Subtracting the first equation of (35) from (102) yields

$$\begin{cases} \operatorname{mes}(A_{i})\frac{de_{i}(t)}{dt} + \sum_{\sigma \in \mathcal{E}_{i}} G_{i,\sigma}(t) + W_{i,\sigma}(t) + c_{0}\operatorname{mes}(A_{i})e_{i}(t) \\ = \int_{A_{i}} \left(f(\mathbf{x}, X(\mathbf{x}, t)) - f(\mathbf{x}_{i}, X_{i}(t))\right) d\mathbf{x} \\ + c_{0}\operatorname{mes}(A_{i})p_{i}(t) + \sum_{\sigma \in \mathcal{E}_{i}} \operatorname{mes}(\sigma)(R_{i,\sigma}(t) + r_{i,\sigma}(t)) - S_{i}(t), \quad \forall A_{i} \in \mathcal{T} \end{cases}$$
(103)

 $^{^{5}}$ also solution of (42) or (35)

with

$$e_{i}(t) = X(\mathbf{x}_{i}, t) - X_{i}(t) = X(\mathbf{x}_{i}, t) - X_{h}(t)(\mathbf{x}_{i}), \qquad t \in [0, T],$$

$$G_{i,\sigma}(t) = -\tau_{\sigma}(e_{j}(t) - e_{i}(t)), \qquad \sigma = i|j,$$

$$G_{i,\sigma}(t) = \tau_{\sigma}e_{i}(t), \qquad \sigma \in \mathcal{E}_{i} \cap \partial\Omega,$$

$$W_{i,\sigma}(t) = q_{i,\sigma}(X(\mathbf{x}_{\sigma,+}, t) - X_{\sigma,+}(t)).$$
(104)

Multipling equation (103) by $e_i(t)$ and summing for $A_i \in \mathcal{T}$ yields

$$\int_{A_i \in \mathcal{T}} \left[\frac{\operatorname{mes}(A_i)}{2} \frac{d(e_i^2(t))}{dt} + \sum_{\sigma \in \mathcal{E}_i} e_i(t) (G_{i,\sigma}(t) + W_{i,\sigma}(t)) + c_0 \operatorname{mes}(A_i) e_i^2(t) \right] \\
= \sum_{A_i \in \mathcal{T}} e_i(t) \left[\int_{A_i} \left(f(\mathbf{x}, X(\mathbf{x}, t)) - f(\mathbf{x}_i, X_i(t)) \right) d\mathbf{x} \right] \\
+ \sum_{A_i \in \mathcal{T}} \left[c_0 \operatorname{mes}(A_i) p_i(t) e_i(t) + \sum_{\sigma \in \mathcal{E}_i} \operatorname{mes}(\sigma) e_i(t) (R_{i,\sigma}(t) + r_{i,\sigma}(t)) - e_i(t) S_i(t) \right].$$
(105)

Using the fact that f is differentiable with respect to X and \mathbf{x} , Taylor expansion yields

$$\max(A_i)\varrho_i(t)$$

$$= \int_{A_i} f(\mathbf{x}, X(\mathbf{x}, t)) d\mathbf{x} - \max(A_i) f(\mathbf{x}_i, X_i(t))$$

$$= \int_{A_i} (f(\mathbf{x}, X(\mathbf{x}, t)) - f(\mathbf{x}_i, X_i(t))) d\mathbf{x},$$

$$= \max(A_i) (f(\mathbf{x}_i, X(\mathbf{x}_i, t)) - f(\mathbf{x}_i, X_i(t))) + \int_{A_i} Z_1(\mathbf{x}, t) (X(\mathbf{x}, t) - X(\mathbf{x}_i, t)) d\mathbf{x}$$

$$+ \int_{A_i} Z_2(\mathbf{x}, t) (\mathbf{x} - \mathbf{x}_i) d\mathbf{x}$$

$$= \max(A_i) (f(\mathbf{x}_i, X(\mathbf{x}_i, t)) - f(\mathbf{x}_i, X_i(t))) + \kappa(\mathbf{x}_i, X, f),$$

$$(106)$$

where

$$\kappa(\mathbf{x}_i, X, f) = \int_{A_i} Z_1(\mathbf{x}, t) (X(\mathbf{x}, t) - X(\mathbf{x}_i, t)) d\mathbf{x} + \int_{A_i} Z_2(\mathbf{x}, t) (\mathbf{x} - \mathbf{x}_i) d\mathbf{x}$$
$$Z_1(\mathbf{x}, t) = \int_0^1 \frac{\partial f}{\partial X} \left(\mathbf{x}_i + \tau(\mathbf{x} - \mathbf{x}_i), X(\mathbf{x}_i, t) + \tau(X(\mathbf{x}, t) - X(\mathbf{x}_i, t)) \right) d\tau$$
$$Z_2(\mathbf{x}, t) = \int_0^1 \frac{\partial f}{\partial \mathbf{x}} \left(\mathbf{x}_i + \tau(\mathbf{x} - \mathbf{x}_i), X(\mathbf{x}_i, t) + \tau(X(\mathbf{x}, t) - X(\mathbf{x}_i, t)) \right) d\tau.$$

As we have assumed that the solution $X(t) \in \mathcal{B}$ is differentiable with respect **x** and *f* differentiable with respect to the two variables with derivatives satisfying (66) or (67), one more Taylor expansion yields

$$|\kappa(\mathbf{x}_i, t, X, f)| \le \max(A_i)C_4(\mathcal{B}, \Omega, f, T, X) h.$$

Using (4), (5) or Proposition 2.1, and the fact that $X(t) \in \mathcal{B}$ and $X_h(t) \in \mathcal{B}$ allow to have

$$\operatorname{mes}(A_i)\varrho_i(t) \le \operatorname{mes}(A_i)\left(C_4'(\mathcal{B},\Omega,f,T,X)|X(\mathbf{x}_i,t) - X_i(t)| + C_4(\mathcal{B},\Omega,T,X)h\right).$$
(107)

Let $e_h(t) \in V_h$ a piecewise constant function defined by

$$e_h(t)(\mathbf{x}_i) = e_i(t) = X(\mathbf{x}_i, t) - X_h(t)(\mathbf{x}_i) \quad A_i \in \mathcal{T}, \quad t \in [0, T].$$
 (108)

Since

$$\tau_{\sigma} = \frac{\mathrm{mes}}{d_{\sigma}} \mu_{\sigma}.$$
 (109)

Using (104), reordering the summation and the fact that the transmissibility is symmetric, i.e. $\tau_{i|j} = \tau_{j|i}$, we have

$$\|e_{h}(t)\|_{1,h}^{2} := \sum_{A_{i}\in\mathcal{T}}\sum_{\sigma\in\mathcal{E}_{i}}e_{i}(t)G_{i,\sigma}(t),$$

$$= \sum_{\sigma\in\mathcal{E}}|D_{\sigma}e_{h}(t)|^{2}\frac{\operatorname{mes}(\sigma)\mu_{\sigma}}{d_{\sigma}}.$$
 (110)

Note that μ_{σ} is defined in (30) and (29). As in the proof of Theorem 4.1, using the regularity of the mesh \mathcal{T} ($\zeta_1 h \leq d_{i,\sigma} \leq h$), we have

$$C_5 \le \mu_\sigma \le C'_5, \quad \forall \sigma \in \mathcal{E}.$$
 (111)

 So

$$C_5 \|e_h(t)\|_{1,\mathcal{T}}^2 \le \|e_h(t)\|_{1,h}^2 \le C_5' \|e_h(t)\|_{1,\mathcal{T}}^2.$$
(112)

Note that

$$\begin{cases} |D_{\sigma}e_{h}(t)| = |e_{i}(t) - e_{j}(t)|, & \text{if } \sigma = i|j, \\ |D_{\sigma}e_{h}(t)| = |e_{i}(t)|, & \text{if } \sigma \in \mathcal{E}_{i} \cap \partial\Omega. \end{cases}$$
(113)

Setting $e_{\sigma,+}(t) = X(\mathbf{x}_{\sigma,+}, t) - X_{\sigma,+}(t)$, as in the proof of Theorem 4.1, using the same technique as in bilinear form $b_h^2(.)$ (see (55)-(60)) yields

$$\sum_{A_i \in \mathcal{T}_{\sigma \in \mathcal{E}_i}} \sum_{e_i(t) W_{i,\sigma}(t)} = \sum_{i \in \mathcal{T}_{\sigma \in \mathcal{E}_i}} \sum_{q_{i,\sigma} e_i(t)} (X(\mathbf{x}_{\sigma,+}, t) - X_{\sigma,+}(t))$$
$$= \sum_{A_i \in \mathcal{T}_{\sigma \in \mathcal{E}_i}} \sum_{q_{i,\sigma} e_i(t) e_{\sigma,+}(t)} \sum_{e_i(t) \in \mathcal{E}_i} \sum_{e_i(t) \in \mathcal{E}_i} q_{i,\sigma} e_i(t) e_{\sigma,+}(t) \ge 0.$$
(114)

Using (114) and (107) in the expression (105) yields

$$\begin{cases}
\frac{1}{2} \sum_{A_i \in \mathcal{T}} \operatorname{mes}(A_i) \frac{d(e_i^2(t))}{dt} + \|e_h(t)\|_{1,h}^2 + c_0 \|e_h(t)\|_{0,h}^2 \leqslant C_4'(\mathcal{B}) \|e_h(t)\|_{0,h}^2 \\
+ C_4(\mathcal{B}) h \sum_{A_i \in \mathcal{T}} \operatorname{mes}(A_i) |e_i(t)| + c_0 C_3' h \sum_{A_i \in \mathcal{T}} \operatorname{mes}(A_i) |e_i(t)| \\
+ \sum_{A_i \in \mathcal{T} \sigma \in \mathcal{E}_i} \operatorname{mes}(\sigma) e_i(t) (R_{i,\sigma}(t) + r_{i,\sigma}(t)) + C_1 h \sum_{A_i \in \mathcal{T}} \operatorname{mes}(A_i) |e_i(t)|.
\end{cases}$$
(115)

The continuity of the diffusion and advection flux at each interface yields

$$R_{i,\sigma}(t) = -R_{j,\sigma}(t), \qquad r_{i,\sigma}(t) = -r_{j,\sigma}(t), \qquad \text{for } \sigma = i | j \in \mathcal{E}_{int}$$

Set

$$R_{\sigma}(t) = |R_{i,\sigma}(t)|, \quad r_{\sigma}(t) = |R_{i,\sigma}(t)|, \qquad A_i \in \mathcal{T}, \qquad \sigma \in \mathcal{E}_{int}$$

Using the Cauchy-Schwarz inequality as in [6] for stationary elliptic problems, and reordering the summation over the edges yields

$$\sum_{A_i \in \mathcal{T} \sigma \in \mathcal{E}_i} \sum_{mes(\sigma) e_i(t) (R_{i,\sigma}(t) + r_{i,\sigma}(t))} \\ \leqslant \sum_{\sigma \in \mathcal{E}} \max(\sigma) D_{\sigma} e_h(t) (R_{\sigma}(t) + r_{\sigma}(t)) \\ \leqslant \left(\sum_{\sigma \in \mathcal{E}} \frac{\operatorname{mes}(\sigma)}{d_{\sigma}} (D_{\sigma} e_h(t))^2 \right)^{\frac{1}{2}} \left(\sum_{\sigma \in \mathcal{E}} \operatorname{mes}(\sigma) d_{\sigma} (R_{\sigma} + r_{\sigma})^2 \right)^{\frac{1}{2}},$$

where $d_{\sigma} = d_{i,\sigma}$, for $\sigma \in \mathcal{E}_i$. Using the fact that $\sum_{\sigma \in \mathcal{E}} \operatorname{mes}(\sigma) d_{\sigma} \leq d \operatorname{mes}(\Omega)$, Lemma 4.2 and relation (112) yields

$$\sum_{A_i \in \mathcal{T}_{\sigma \in \mathcal{E}_i}} \sum_{mes(\sigma) e_i(t) (R_{i,\sigma}(t) + r_{i,\sigma}(t))} \\ \leqslant C_3 h (mes(\Omega) d)^{\frac{1}{2}} \|e_h(t)\|_{1,\mathcal{T}} \\ \leqslant (C_5)^{-1} C_3 h (mes(\Omega) d)^{\frac{1}{2}} \|e_h(t)\|_{1,h}.$$
(116)

For an arbitrary constant C > 0, Young's inequality implies that

$$\begin{cases} |C h \sum_{A_i \in \mathcal{T}} \operatorname{mes}(A_i) e_i(t)| = |\sum_{A_i \in \mathcal{T}} (C h \operatorname{mes}(A_i)^{\frac{1}{2}}) (\operatorname{mes}(A_i)^{\frac{1}{2}} e_i(t))| \\ \leqslant \frac{1}{2} ||e_h(t)||_{0,h}^2 + \frac{1}{2} C^2 h^2 \operatorname{mes}(\Omega) \\ C h ||e_h(t)||_{1,h} \leqslant \frac{1}{2} C^2 h^2 + \frac{1}{2} ||e_h(t)||_{1,h}^2. \end{cases}$$
(117)

Using expression (116) and (117) in expression (115) yields

$$\begin{cases}
\frac{1}{2} \left[\sum_{A_i \in \mathcal{T}} \operatorname{mes}(A_i) \frac{d(e_i^2(t))}{dt} + \|e_h(t)\|_{1,h}^2 + 2c_0 \|e_h(t)\|_{0,h}^2 \right] \leq (C_7 \|e_h(t)\|_{0,h}^2 + C_6 h^2 \\
C_6 = C_6(c_0, C_1, C_3, C_3', C_4, C_5), C_7 = C_7(C_4').
\end{cases}$$
(118)

Bounding the left hand side of expression (118) below yields

$$\sum_{A_i \in \mathcal{T}} \max(A_i) \frac{d(e_i^2(s))}{ds} \leqslant 2C_7 \|e_h(s)\|_{0,h}^2 + 2C_6 h^2, \qquad \forall s \in [0,T].$$
(119)

Integrating both sides of expression (119) through interval [0, t], $0 \le t \le T$ yields

$$\|e_h(t)\|_{0,h}^2 \le \|e_h(0)\|_{0,h}^2 + 2C_6 T h^2 + 2C_7 \int_0^t \|e_h(s)\|_{0,h}^2 ds, \qquad \forall t \in [0,T].$$
(120)

Applying the discrete Gronwall yields

$$\|e_{h}(t)\|_{0,h}^{2} \leq C\left(\|e_{h}(0)\|_{0,h}^{2} + h^{2}\right), \qquad (121)$$
$$C = C(\mathcal{B}, \Omega, X, F, \mathbf{D}, \mathbf{q}, T, \zeta_{1}).$$

5. Full discretization and main result

5.1. Exponential Euler method for time discretization

For simplicity we consider a constant time-step $\Delta t > 0$. At time $t_m = m\Delta t \in [0, T]$, the mild solution (65) is given by

$$X_h(t_m) = S_h(t_m)X_{0h} + \int_0^{t_m} S_h(t_m - s)P_hF(X_h(s))ds.$$
 (122)

Then, given the solution X_h at the time t_m , we can construct the corresponding solution at t_{m+1} as

$$X_{h}(t_{m+1}) = S_{h}(\Delta t)X_{h}(t_{m}) + \int_{0}^{\Delta t} S_{h}(\Delta t - s)P_{h}F(X_{h}(t_{m} + s))ds.$$
(123)

Note that the expression in (123) is still an exact form of X_h . The idea behind exponential time differencing is to approximate $P_hF(X_h(t_m+s))$ by a suitable polynomial [14, 15]. We consider the simplest case where $P_hF(X_h(t_m+s))$ is approximated by the constant $P_hF(X_h(t_m))$ and the corresponding scheme (ETD1) is given by

$$X_h^{n+1} = e^{-\Delta t A_h} X_h^n + \Delta t \varphi_1(-\Delta t A_h) P_h F(X_h^m)$$
(124)

where

$$\varphi_1(-\Delta t A_h) = (-\Delta t A_h)^{-1} \left(e^{-\Delta t A_h} - I \right) = \frac{1}{\Delta t} \int_0^{\Delta t} e^{-(\Delta t - s)A_h} ds.$$

Note that the ETD1 scheme in (124) can be rewritten as

$$X_{h}^{m+1} = X_{h}^{m} + \Delta t \varphi_{1}(-\Delta t A_{h})(-A_{h}X_{h}^{m} + P_{h}F(X_{h}^{m})).$$
(125)

This new expression has the advantage that it is computationally more efficient as only one matrix exponential function needs to be evaluated at each step.

5.2. Main result

To achieve the optimal orders of convergence in time and space, the solution X need to be regular.

Theorem 5.1. Let $\mathcal{B} \subset V$ be bounded, consider the solution X of (2) and X_h^m the numerical solution (125) given by combining the finite volume method in space discretization and ETD1 scheme in time integration. Assume that $X(t_k) \in \mathcal{B}$, $X_h^k \in \mathcal{B} \bigcap V_h$ and $X_h(t_k)^6 \in \mathcal{B} \bigcap V_h$ for all $t_k = k\Delta t \leq T \leq t^*$, $k \in \mathbb{N}$. We also assume that the unique mild solution X of (2) is the classical solution (i.e. X is twice continuously differentiable with respect to \mathbf{x} and differentiable with respect to t), Assumption 4.1 is satisfied and the reaction function F satisfies (20). Furthermore assume that $X_0 \in \mathcal{D}(A) \cap \mathcal{B}$, $X_{0h} \in V_h \cap \mathcal{B}$ and $f(\mathbf{x}, u)$ is differentiable respect to \mathbf{x} and u with (66) or (67), then the following estimate holds

$$||X(t_m) - X_h^m||_{0,h} \le C \left(||X_0 - X_{0h}||_{0,h} + \Delta t + h \right),$$

where $C = C(\mathcal{B}, \Omega, X, F, \mathbf{D}, \mathbf{q}, T, \zeta_1).$

Before give the proof, let us provide two important results.

Lemma 5.1. [Norms equivalence]

Consider the discrete $L^2(\Omega)$ norm $\|.\|_{0,h}$ associated to the discrete scalar product (38)or (39) and the discrete norm $\|.\|_{0,H}$ defined with the dual mesh \mathcal{T}_h such that for $v \in C(\Omega)$

$$\|v\|_{0,H} = \sqrt{\sum_{K \in \mathcal{T}_h} h_K^d \sum_{\mathbf{x}_i \in K} v(\mathbf{x}_i)^2}.$$
(126)

Assume that \mathcal{T} is regular (Assumption 4.1 is satisfied, so \mathcal{T}_h is regular according to Remark 4.1), the norms $\|.\|_{0,h}$, $\|.\|_{0,H}$, and the $L^2(\Omega)$ norm $\|.\|$ are equivalent in V_h uniformly with respect to h.

Proof. See [25, Remark 6.16, p. 275] and [25, Theorem 3.43, p. 163].

Proposition 5.1. [Interpolation error]

Let \mathcal{T} be an admissible mesh in the sense of Definition 4.1 and \mathcal{T}_h its dual Delaunay triangulation (remember that $\{\mathbf{x}_i\}$ are vertices of \mathcal{T}_h and centers of \mathcal{T}). Let $I_h : C(\overline{\Omega}) \to V_h$ defined by

$$I_h(u) = \sum_{i \in \mathcal{T}} u(\mathbf{x}_i) \varphi_{\mathbf{x}_i}, \qquad u \in C(\overline{\Omega})$$
(127)

where $\{\varphi_{\mathbf{x}_i}\}_{i\in\mathcal{T}}$ is the nodal basis corresponding to $\{\mathbf{x}_i\}_{i\in\mathcal{T}}$ in the sense of finite element method $(\varphi_{x_i}(x_j) = \delta_{i,j})$. Let X the solution of (2) given by (24). If $X(t) \in H^2(\Omega)$, then there exists a positive constant $C_0 > 0$ independent of X and t such that the following estimate holds

$$||X(t) - I_h(X(t))|| \le C_0 |X(t)|_2 h^2,$$
(128)

⁶This is the solution of (35) or (65)

where $|.|_2$ denotes the semi norm of $H^2(\Omega)$ ⁷. Furthermore, if $X \in C([0,T], H^2(\Omega))$, there exists $C_0 = C_0(X,T)$ such that

$$||X(t) - I_h(X(t))|| \le C_0(X, T)h^2, \qquad \forall t \in [0, T].$$
(129)

Proof. For (128), see [25, Section 3.4, Theorem 3.29, page 138] or [3, Theorem 17.1, page 132] with k = 1 and m = 0. Recall that $C([0,T], H^2(\Omega))$ is the set of continuous functions $v: [0,T] \to H^2(\Omega)$ such that $\sup_{t \in [0,T]} \|v(t)\|_2 < \infty$. To have (129), we obviously have

$$\|X(t) - I_h(X(t))\| \le C_0 \|X(t)\|_2 h^2 \le C_0 \sup_{t \in [0,T]} \|X(t)\|_2 h^2 = C_0(X,T)h^2, \quad \forall t \in [0,T].$$
(130)

As the preparatory results are provided, let us proof our main result (Theorem 5.1).

Proof. We use the equivalence of the norms $\|.\|$ and $\|.\|_{0,h}$ in V_h as we have assumed that the mesh \mathcal{T} is regular (see Lemma 5.1). Using the triangle inequality yields

$$\begin{aligned} \|X(t_m) - X_h^m\|_{0,h} &\leq \|X(t_m) - X_h(t_m)\|_{0,h} + \|X_h(t_m) - X_h^m\|_{0,h} \\ &= I + II. \end{aligned}$$
(131)

As I is already estimated in Theorem 4.2, let us estimate II. From (122) and (124), we have

$$X_h(t_m) = S_h(t_m) X_{0h} + \sum_{k=0}^{m-1} \int_{t_k}^{t_{k+1}} S_h(t_m - s) P_h F(X_h(s)) ds,$$
(132)

and

$$X_h^m = S_h(t_m)X_{0h} + \sum_{k=0}^{m-1} \int_{t_k}^{t_{k+1}} S_h(t_m - s)P_hF(X_h^k)ds.$$
 (133)

The smoothing properties of the semigroup S_h in Proposition 2.2 and the equivalence $\|.\| \equiv \|.\|_{0,h}$ in V_h yields

$$\begin{aligned} \|X_{h}(t_{m}) - X_{h}^{m}\|_{0,h} &\equiv \|X_{h}(t_{m}) - X_{h}^{m}\| \\ &\leq \sum_{k=0}^{m-1} \int_{t_{k}}^{t_{k+1}} \|S_{h}(t_{m} - s)P_{h}\left(F(X_{h}(s)) - F(X_{h}^{k})\right)\|ds \end{aligned}$$

Following [27] we should prove that

$$\|A_h^{-1/2}P_hv\| \le Ch\|v\| + \|v\|_{-1}, \qquad \forall v \in L^2(\Omega).$$
(134)

From [30, (18)] it is well known that

$$|(u,v) - \langle u,v \rangle_{0,h}| \le Ch^{s+p} ||u||_s ||v||_p, \quad \forall u \in H^s(\Omega), \ v \in H^p(\Omega), \qquad s,p \in \{0,1\}.$$
(135)

⁷Note that this semi norm uses only second order derivatives which belong to $L^2(\Omega)$.

Indeed identifying $L^2(\Omega)$ to its dual, as $A_h^{*-1/2}$ is uniformly bounded, using (135) and the definition of the projection P_h yields

$$\begin{split} \|A_{h}^{-1/2}P_{h}v\| &= \sup_{u_{h}\in V_{h}} \frac{|(A_{h}^{-1/2}P_{h}v, u_{h})|}{\|u_{h}\|_{1}} \\ &= \sup_{u_{h}\in V_{h}} \frac{|(P_{h}v, A_{h}^{*-1/2}u_{h})|}{\|u_{h}\|_{1}} \\ &\leq C_{1}h\|v\| + \sup_{u_{h}\in V_{h}} \frac{|\langle P_{h}v, A_{h}^{*-1/2}u_{h}\rangle_{0,h}|}{\|u_{h}\|_{1}} \\ &= C_{1}h\|v\| + \sup_{u_{h}\in V_{h}} \frac{|\langle v, A_{h}^{*-1/2}u_{h}\rangle_{0,h}|}{\|u_{h}\|_{1}} \\ &\leq Ch\|v\| + \sup_{w_{h}\in V_{h}} \frac{|\langle v, w_{h}\rangle|}{\|A_{h}^{*1/2}w_{h}\|_{1}} \\ &\leq Ch\|v\| + \|v\|_{-1}. \end{split}$$
(136)

Comparing with the results in [27], the estimation (134) clearly shows the difference between the finite element method and the finite element method while performing the space discretization of problem of type (1).

Let us back in our main proof. Using (134), Proposition 2.2, the Lipschitz conditions (20) and (21) and the fact that both the full discrete and semi discrete solutions are in \mathcal{B} allows to have

$$\begin{aligned} \|X_{h}(t_{m}) - X_{h}^{m}\|_{0,h} \\ &\leq \sum_{k=0}^{m-1} \int_{t_{k}}^{t_{k+1}} \|S_{h}(t_{m} - s)A_{h}^{1/2}A_{h}^{-1/2}P_{h}\left(F(X_{h}(s)) - F(X_{h}^{k})\right)\| ds, \\ &\leq \sum_{k=0}^{m-1} \int_{t_{k}}^{t_{k+1}} (t_{m} - s)^{-1/2} \left(Ch\|F(X_{h}(s)) - F(X_{h}^{k})\| + \|F(X_{h}(s)) - F(X_{h}^{k})\|_{-1}\right) ds \\ &\leq \sum_{k=0}^{m-1} \int_{t_{k}}^{t_{k+1}} (t_{m} - s)^{-1/2} \left(Ch\|X_{h}(s) - X_{h}^{k}\|_{1} + \|X_{h}(s) - X_{h}^{k}\|\right) ds \\ &\leq C(\mathcal{B})h + \sum_{k=0}^{m-1} \int_{t_{k}}^{t_{k+1}} (t_{m} - s)^{-1/2} \left(\|X_{h}(s) - X_{h}^{k}\|\right) ds \\ &\leq C(\mathcal{B})h + C(\mathcal{B})\sum_{k=0}^{m-1} \int_{t_{k}}^{t_{k+1}} (t_{m} - s)^{-1/2} \left(\|X_{h}(s) - X(s)\|\right) ds \\ &+ C(\mathcal{B})\sum_{k=0}^{m-1} \int_{t_{k}}^{t_{k+1}} (t_{m} - s)^{-1/2} \left(\|X(s) - X_{h}^{k}\|\right) ds \\ &= C(\mathcal{B}) \left(h + H_{1} + H_{2}\right). \end{aligned}$$
(137)

For $s \in [0, T]$, as the solution X is assumed to be regular, Proposition 5.1 yields

$$\begin{aligned} \|X_{h}(s) - X(s)\| &\leq \|X_{h}(s) - I_{h}(X(s)) + I_{h}(X(s)) - X(s)\| \\ &\leq (\|X_{h}(s) - I_{h}(X(s))\| + \|I_{h}(X(s)) - X(s)\|) \\ &\leq (\|X_{h}(s) - I_{h}(X(s))\| + C_{0}(X, T)h^{2}). \end{aligned}$$
(138)

Since $X_h(s) - I_h(X(s)) \in V_h$, the equivalence $\|.\| \equiv \|.\|_{0,h}$ and the uniform estimate of the term I in [0, T] yields

$$\begin{aligned} \|X_{h}(s) - I_{h}(X(s))\| &\equiv \|X_{h}(s) - I_{h}(X(s))\|_{0,h} \\ &= \|X_{h}(s) - X(s)\|_{0,h} \quad \text{(by definition of } \|.\|_{0,h}, \text{ and } I_{h}) \\ &\leq C(\mathcal{B}, \Omega, X, F, \mathbf{D}, \mathbf{q}, \zeta_{1})(\|X_{0h} - X_{0}\|_{0,h} + h), \end{aligned}$$
(139)

which yields

$$II_{1} \leq C(\mathcal{B}) \left(\|X_{0h} - X_{0}\|_{0,h} + h \right) \sum_{k=0}^{m-1} \int_{t_{k}}^{t_{k+1}} (t_{m} - s)^{-1/2} ds + C_{0}(X, T)h^{2}$$
(140)

$$\leq C(\mathcal{B})(\|X_{0h} - X_0\|_{0,h} + h) \left(\sum_{k=0}^{m-1} \int_0^T (t_m - s)^{-1/2} ds\right) + C_0(X, T)h^2 \quad (141)$$

$$\leq C(\mathcal{B}, \Omega, X, F, \mathbf{D}, \mathbf{q}, T, \zeta_1) (\|X_{0h} - X_0\|_{0,h} + h). \quad (142)$$

$$\leq C(\mathcal{B}, \Omega, X, F, \mathbf{D}, \mathbf{q}, T, \zeta_1) \left(\|X_{0h} - X_0\|_{0,h} + h \right).$$
(142)

We also have

$$II_{2} = \sum_{k=0}^{m-1} \int_{t_{k}}^{t_{k+1}} (t_{m} - s)^{-1/2} \|X(s) - X_{h}^{k}\| ds$$

$$\leq \sum_{k=0}^{m-1} \int_{t_{k}}^{t_{k+1}} (t_{m} - s)^{-1/2} \|X(s) - X(t_{k})\| ds$$

$$+ \sum_{k=0}^{m-1} \int_{t_{k}}^{t_{k+1}} (t_{m} - s)^{-1/2} \|X(t_{k}) - X_{h}^{k}\| ds$$

$$= II_{2}^{1} + II_{2}^{2}.$$
(143)

Using Lemma 2.1 yields

$$II_{2}^{1} \leq C(\mathcal{B})\sum_{k=0}^{m-1} \int_{t_{k}}^{t_{k+1}} (t_{m} - s)^{-1/2} (s - t_{k}) ds$$

$$\leq C(\mathcal{B})\Delta t \sum_{k=0}^{m-1} \int_{t_{k}}^{t_{k+1}} (t_{m} - s)^{-1/2} ds$$

$$\leq C(\mathcal{B})\Delta t \int_{0}^{T} (t_{m} - s)^{-1/2} ds$$

$$\leq C(\mathcal{B}, T)\Delta t.$$
(144)

Using the equivalence $\|.\| \equiv \|.\|_{0,h}$ in V_h (Lemma 5.1) allow to have

$$II_{2}^{2} \leq \sum_{k=0}^{m-1} \int_{t_{k}}^{t_{k+1}} (t_{m} - s)^{-1/2} \|X(t_{k}) - I_{h}(X(t_{k})) + I_{h}(X(t_{k})) - X_{h}^{k}\|ds$$
(145)

$$\leq \sum_{k=0}^{m-1} \int_{t_k}^{t_{k+1}} (t_m - s)^{-1/2} \|X(t_k) - I_h(X(t_k))\| + \|I_h(X(t_k)) - X_h^k\|_{0,h} ds \quad (146)$$

$$\leq C(X,T)\left(h^{2} + \sum_{k=0}^{m-1} \int_{t_{k}}^{t_{k+1}} (t_{m} - s)^{-1/2} \|X(t_{k}) - X_{h}^{k}\|_{0,h} ds\right)$$
(147)

Then

$$II \leq C(\mathcal{B})\left((\|X_{0h} - X_0\|_{0,h} + \Delta t + h) + \sum_{k=0}^{m-1} \int_{t_k}^{t_{k+1}} (t_m - s)^{-1/2} \|X(t_k) - X_h^k\|_{0,h} ds \right),$$

Combining estimates I and II yields

$$\|X(t_m) - X_h^m)\|_{0,h} \le C(\mathcal{B}) \left(\|X_{0h} - X_0\|_{0,h} + \Delta t + h + \sum_{k=0}^{m-1} \int_{t_k}^{t_{k+1}} (t_m - s)^{-1/2} \|X(t_k) - X_h^k\|_{0,h} ds \right), \quad (148)$$

where $C(\mathcal{B}) = C(\mathcal{B}, \Omega, X, F, \mathbf{D}, \mathbf{q}, \zeta_1)$. Applying the generalized discrete Gronwall in (148) ends the proof.

Remark 5.1. Using Proposition 5.1, if the initial solution $X_0 \in \mathcal{D}(A) \subset H^2(\Omega) \subset C(\Omega)$, and $X_{0h} = I_h(X_0)$, we obviously have the following estimate

$$\|X(t_m) - X_h^m)\|_{0,h} \le C(\mathcal{B}) \left(\Delta t + h\right).$$
(149)

Remark 5.2. Although optimal orders in time and space are achieved for $X_0 \in \mathcal{D}(A)$, this condition is not enough to ensure the regularity of the solution as stated in Theorem 5.1 and Theorem 4.2, which requires that $X_0 \in C^2(\Omega)$.

6. Numerical simulations

The goal here is to illustre the theoretical result (149) and to compare ETD1 scheme with standard time-stepping methods, implicit Euler and semi implicit schemes. The difference between these schemes is the time integration as the space discretization is performed using finite volume method. So our main focus will be the errors in time. In [4, Section 4, Figure 2], the convergence in space has been studied for linear with exact solution and the optimal order in space (149) has been reached. Here we compare the time errors and the efficiency of the schemes.

Our code was implemented in Matlab 7.7. In the legends of all of our graphs we use the following notation

- "Implicit with Newton" denotes results from the implicit Euler with standard Newton's method.
- "Implicit with Newton V" denotes results from the implicit Euler with a variant of Newton's method where the Jacobian is kept constant [4].
- "Léja ETD1" denotes results from ETD1 with real fast Léja points for matrix exponential.
- "Krylov ETD1" denotes results from ETD1 with Krylov subspace for matrix exponential.
- "Semi implicit" denotes results from the semi-implicit scheme.

We now evaluate the ETD1 method for a non-linear ADR problem where the non-linear reaction term is given by $f(\mathbf{x}, u) = -\theta u^2(1-u)$. We take $\theta = 100$, use a constant velocity of $\underline{\mathbf{q}} = [-0.01, -0.01]^T$, and the dispersion tensor has the entries $D_1 = D_2 = 10^{-4}$. The domain is $\Omega = [0, 1) \times [0, 1)$, which we discretise with $h = \Delta x = \Delta y = 10^{-2}$. We can observe that f satisfies the local Lipschitz condition (5), so existence and uniqueness of the local solution is ensured. Indeed the global solution exists and is given by [29]

$$C(x, y, t) = (1 + \exp(a(x + y - bt) + a(b - 1)))^{-1}$$
(150)

where $a = \sqrt{\theta/(4 \times 10^{-4})}$ and $b = -0.02 + \sqrt{\theta \times 10^{-4}}$. The initial condition and boundary conditions are defined with respect to the exact solution (150).



Figure 2: (a) Convergence of the L^2 norm at T = 1 as a function of Δt . (b) The L^2 norm at T = 1 as a function of CPU time. Both are for the the non-linear ADR in homogeneous porous media (Problem 2).

Figure 2(a) shows the convergence as a function of the chosen time-step Δt , measuring the error at the final time T = 1. The semi-implicit time-stepping method and the ETD1 methods have similar error constants. All schemes have the same rate of convergence $\mathcal{O}(\Delta t)$, which is predicted in our convergence result (149). Figure 2(b) shows the L^2 error as a function of CPU time, which is given in Figure 2(a). ETD1 graphs are also similar to the semi-implicit one. However, all three methods, ETD1 with Leja points and Krylov subspace technique and semi-implicit time-stepping, outperform the implicit time-stepping methods.

Acknowledgements

This work was supported by the Overseas Research Students Awards Scheme (ORSAS) at Heriot Watt University and Robert Bosch Stiftung through the AIMS ARETE chair programme.

References

- [1] M. Afif, & B. Amaziane. Convergence of finite volume schemes for a degenerate convection-diffusion equation arising in flow in porous media. *Comput. Methods Appl. Mech. Engrg*, **191**(2002): 5265–5286.
- [2] D. Henry. Geometric theory of semilinear parabolic equations. Number 840 in Lecture notes in mathematics. Springer, 1981.
- [3] H. Fujita and T. Suzuki. Evolutions problems (part1), in P. G. Ciarlet and J. L. Lions(eds.), Handbook of Numerical Analysis, North-Holland, volume II, 1991, pp. 789–928.
- [4] A. Tambue, G. Lord, and S. Geiger. An exponential integrator for advection-dominated reactive transport in heterogeneous porous media. *Journal of Computational Physics*, 229(10)(2010) 3957–3969.
- [5] A. Tambue. Efficient Numerical Schemes for Porous Media Flow. PhD thesis, Department of Mathematics, Heriot–Watt University, 2010.
- [6] R. Eymard, T. Gallouet, and R. Herbin, Finite volume methods. Updated preprint(2006) of the work appeared in: P. G. Ciarlet, J. L. Lions (Eds.), Handbook of Numerical Analysis Volume 7, North-Holland, Amsterdam, 2000, pp. 713–1020.
- [7] M. Hochbruck and A. Ostermann. Exponential integrators. Acta Numerica, vol. 19 (2010), 209–286.
- [8] B. Amaziane and M. El Ossmani. Convergence Analysis of an Approximation to Miscible Fluid Flows in Porous Media by Combining Mixed Finite Element Finite Volume Methods. *Numer. Method.*. Partial. Differ. E., 24(3)(2008), 799–832.
- [9] R. Eymard, D. Hilhorst, and M. Vohralik. A combined finite volume-finite element scheme for the discretization of strongly nonlinear convection-diffusion-reaction problems on nonmatching grids, *Numer. Method. Partial. Differ. E.*, 26(3)(2010), 612–646.
- [10] A. Tambue, S. Geiger, and G. J Lord. Exponential Time integrators for 3D Reservoir Simulation. In proceedings of the 12th European Conference on the Mathematics of Oil Recovery, Oxford, UK, DOI: 10.3997/2214-4609.20145032,2010.
- [11] S. Geiger, G. L. Lord and Tambue, A. Exponential time integrators for stochastic partial differential equations in 3D reservoir simulation, *Computational Geosciences*, **16**(2) (2012), 323–334.
- [12] A. Martinez, L. Bergamaschi, M. Caliari, and M. Vianello, A massively parallel exponential integrator for advection-diffusion models, J. Comput. Appl. Math. 231(1) (2009) 82–91.
- [13] L. Bergamaschi, M. Caliari, A. Martinez, and M. Vianello, Comparing Léja and Krylov approximations of large scale matrix exponentials, *Comput. Sci. - ICCS*, **3994**(2006) 685–692.
- [14] S. M. Cox and P. C. Matthews, Exponential time differencing for stiff systems, J. Comput. Phys. 176(2) (2002) 430–455.
- [15] A. K. Kassam and L. N Trefethen. Fourth-order time stepping for stiff PDES. SIAM Journal of Computing, 26(4)(2005), 1214–1233.
- [16] J. Baglama, D. Calvetti, and L. Reichel. Fast Leja points. *Electronic Transactions on Numerical Analysis*, 7(1998), 124–140.
- [17] L. Bergamaschi, M. Caliari, and M. Vianello. The RELPM exponential integrator for FE discretizations of advection-diffusion equations. *Computational Science ICCS Proceedings*, **3039**(2004), pp. 434–442.
- [18] G. J Lord and A. Tambue, A modified semi-implict Euler-Maruyama scheme for finite element discretization of SPDEs with Additive Noise. arXiv:1004.1998v1, 2010.
- [19] G. J Lord and A. Tambue. Stochastic Exponential Integrators for finite element discretization of SPDEs with Additive Noise. http://arxiv.org/abs/1005.5315, 2010.
- [20] G. J Lord and A. Tambue. Stochastic exponential integrators for the finite element discretization of SPDEs for multiplicative and additive noise. IMA Journal of Numerical Analysis, 33(2) (2013), 515–543.
- [21] A Pazy. Semigroups of linear operators and applications to partial differential equations. Applied Mathematical Sciences, volume 44, Springer-Verlag, New York, 1983.

- [22] I. Aavatsmark. An introduction to multipoint flux approximations for quadrilateral grids. Comput. Geosci., 6(3-4)(2002):405-432.
- [23] I. Aavatsmark, T. Barkve, Ø Bøe, and T. Mannseth. Discretization on non-orthogonal, quadrilateral grids for inhomogeneous, anisotropic media. J. Comput. Phys., 127(1)(1996) 2–14.
- [24] R. Eymard, T. Gallouet and R. Herbin. A cell-centred finite-volume approximation for anisotropic diffusion operators on unstructured meshes in any space dimension. *IMA Journal of Numerical Analysis*, 26(2006), 326–353.
- [25] P. Knabner, and L. Angermann. Numerical methods for elliptic and parabolic partial differential equations solution. Springer Verlag, Berlin, 2003.
- [26] L. C. EVans Partial Differential Equations. Graduate Studies in Mathematics, Vol. 19, 1997.
- [27] S. Larsson. Nonsmooth data error estimates with applications to the study of the long-time behavior of finite element solutions of semilinear parabolic problems. *Preprint 1992-36, Department of Mathematics, Chalmers University of Technology, available at http://www.math.chalmers.se/~stig/papers/index.html*.
- [28] J. L. Lions. Espaces d'interpolation et domaines du puissances fractionnaires d'oprateurs. J. Math. Soc. Japan, 14 (1962) 233–241.
- [29] M. Caliari, M. Vianello, and L. Bergamaschi. The LEM exponential integrator for advection-diffusionreaction equations. J. Comput. Appl. Math. 210(1-2) (2007) 56–63.
- [30] L. S. Pop, M. Seplveda, F. A. Radu, and O. P. Vera Villagran. Error estimates for the finite volume discretization for the porous medium equation. J. Comput. Appl. Math. 234(7) (2010) 2135–2142.