# Harnack Inequalities of Hitting Distributions of Projections of Planar Symmetric Random Walks on the Lattice Torus

Michael Carlisle<sup>∗</sup> Baruch College, CUNY

October 28, 2021

#### Abstract

We give Harnack inequalities for the hitting distributions of a large family of symmetric random walks on  $\mathbb{Z}^2$ , and their projections onto the lattice torus  $\mathbb{Z}_K^2$ . This extends a framework for the simple random walk in [\[6\]](#page-19-0), and generalizes the results in [\[1\]](#page-19-1) to the toral projection.

## 1 Introduction

Consider a random walk  $S_t = S_0 + \sum_{j=1}^t X_j$ , on  $\mathbb{Z}^2$ , with  $X = \{X_j\}_{j\in\mathbb{N}}$  having the following properties:  $X_1$  is symmetric, has finite covariance matrix equal to a scalar times the identity, i.e.,  $\Gamma := cov(X_1) = cI, c > 0$ , and X is strongly aperiodic.

As usual in the literature, let

$$
p_1(x, y) = p_1(y - x) = P^x(X_1 = y)
$$

<sup>∗</sup>michael.carlisle@baruch.cuny.edu

be the one-step transition probability of X. We say X satisfies **Condition A** if either  $p_1$ has bounded support, or, from any point "just outside" a disc, we will enter the disc with positive probability; *i.e.*, for any  $s \leq n$ , for large enough *n*,

<span id="page-1-0"></span>
$$
\inf_{y:n \le |y| < n+s} \sum_{z \in D(x,n)} p_1(y,z) = \inf_{y \in \partial D(x,n)_s} P^y(X_1 \in D(x,n)) \ge ce^{-\beta s^{1/4}},\tag{1.1}
$$

where the (Euclidean) s-annulus around the disc  $D(x, n)$  is defined as

$$
\partial D(x,n)_s := D(x,n+s) \setminus D(x,n).
$$

In particular, if  $X_1$  has infinite range, then for any  $y \in \partial D(0, n)_s$ , there exists  $x \in D(0, n)$ such that  $p_1(y, x) > 0$ .

Starting at a point  $x \in A^c$ , we define the *hitting distribution* of A to be

$$
H_A(x,y) := P^x(S_{T_A} = y),
$$

where  $T_A$  is the first hitting time of the set A by the random walk:

$$
T_A = \inf\{t \ge 0 : S_t \in A\}.
$$

The *last exit decomposition* of a hitting distribution is based on the Green's function: for A a proper subset of  $\mathbb{Z}^2$ ,  $x \in A^c$ , and  $y \in A$ ,

$$
H_A(x,y) = \sum_{z \in A^c} G_{A^c}(x,z) p_1(z,y), \tag{1.2}
$$

where the (truncated) Green's function, up to escaping a set B, is defined, for  $x, y \in B$ , as the total expected number of visits to  $y$ , starting from  $x$ , before escaping  $B$ :

$$
G_B(x,y) := \mathbb{E}^x \bigg[ \sum_{j=0}^{\infty} 1_{\{S_j = y; j < T_{B^c}\}} \bigg] = \sum_{j=0}^{\infty} P^x(S_j = y; j < T_{B^c}) \tag{1.3}
$$

and 0 if x or  $y \notin B$ . A useful identity relates the Green's function of a set to its expected escape time: if  $x \in B$ ,

<span id="page-1-1"></span>
$$
\mathbb{E}^x(T_{B^c}) = \sum_{z \in B} G_B(x, z). \tag{1.4}
$$

The goal of this paper is to establish Harnack inequalities of hitting distributions of discs and disc complements on the planar and toral lattices. Condition A is sufficient to allow the error terms in our Harnack inequalities induced by the toral projection to drop out of sight, as long as there is a moment condition in place on the size of our jumps: we assume

<span id="page-1-2"></span>
$$
\mathbb{E}|X_1|^M = \sum_{x \in \mathbb{Z}^2} |x|^M p_1(x) < \infty \tag{1.5}
$$

for some  $M > 4$  (and write  $M = 4 + 2\beta$  for some  $\beta > 0$ ). While  $M > 2$  suffices for interior

Harnack inequality results,  $M = 3 + 2\beta$  is needed for results on the plane, and one more moment is used in our arguments in the transition to the torus.

We will switch between the planar and toral representations of the random walk and corresponding stopping times, hitting distributions, etc. Define the projections, for  $x =$  $(x_1, x_2) \in \mathbb{Z}^2$ , by

$$
\pi_K : \mathbb{Z}^2 \to [-K/2, K/2)^2 \cap \mathbb{Z}^2,
$$
  
\n
$$
\pi_K(x) = ((x_1 + \lfloor \frac{K}{2} \rfloor)(\text{mod } K) - \lfloor \frac{K}{2} \rfloor, (x_2 + \lfloor \frac{K}{2} \rfloor)(\text{mod } K) - \lfloor \frac{K}{2} \rfloor);
$$
  
\n
$$
\hat{\pi}_K : \mathbb{Z}^2 \to \mathbb{Z}_K^2, \quad \hat{\pi}_K(x) = (\pi_K x) + (K\mathbb{Z})^2.
$$

(For example, if  $x = (-12, 6)$  and  $K = 11$ , then  $\pi_{11}(\mathbb{Z}^2) = \{-5, \ldots, 5\}^2$ ,  $\pi_{11}(x) =$  $(-1, -5)$ , and  $\hat{\pi}_{11}(x) = (-1, -5) + (11\mathbb{Z})^2$ .

We call the set of lattice points  $\pi_K(\mathbb{Z}^2) = [-K/2, K/2)^2 \cap \mathbb{Z}^2$  the **primary copy** in  $\mathbb{Z}^2$ , and for  $x \in \pi_K(\mathbb{Z}^2)$ ,  $\hat{x} := \hat{\pi}_K x$  is its corresponding element in  $\mathbb{Z}_K^2$ . Any  $z \in \pi_K^{-1} x$ ,  $z \neq \pi_K x$ , is called a **copy** of x. Likewise, for a set  $A \subset \mathbb{Z}^2$ ,  $\hat{A} := \hat{\pi}_K A$  is the toral projection of A, and the set of all copies of A is

$$
\pi_K^{-1}\pi_K A = \hat{\pi}_K^{-1} \hat{A} := \{ z \in \mathbb{Z}^2 : z = x + (iK, jK), \ i, j \in \mathbb{Z}, x \in A \}.
$$

For a given  $\hat{x} \in \mathbb{Z}_K^2$ , we define x to be the (planar) primary copy of that element;  $x :=$  $\pi_K \hat{\pi}_K^{-1} \hat{x}.$ 

While  $X_j$  is the jth step of the planar walk and  $S_j$  its position at time j, we use  $\hat{S}_j := \hat{\pi}_K S_j$ to denote the position of the toral walk at time  $j$ . The distance between two points  $x, y \in \mathbb{Z}^2$  will be the Euclidean distance  $|x - y|$ ; on the torus, the distance between two points  $\hat{x}, \hat{y} \in \mathbb{Z}_K^2$  will be the minimum Euclidean distance  $|\hat{x} - \hat{y}| \le K\sqrt{2}/2$ . To limit the issues regarding this distance, we will restrict any discs on  $\mathbb{Z}_K^2$  to have radius  $n < K/4$ (sometimes written as a diameter constraint:  $2n < K/2$ ).

To bound our functions, we need a precise notion of bounding distance on the lattice torus  $\mathbb{Z}_K^2$ . As in [\[6\]](#page-19-0), a function  $f(x)$  is said to be  $O(x)$  if  $f(x)/x$  is bounded, uniformly in all implicit geometry-related quantities (such as K). That is,  $f(x) = O(x)$  if there exists a universal constant C (not depending on K) such that  $|f(x)| \leq Cx$ . Thus  $x = O(x)$  but Kx is not  $O(x)$ . A similar convention applies to  $o(x)$ .

Next, we will define a few terms describing the distance of a random walk step, relative to a reference disc of radius  $n$  and an s-annulus around the disc. A small jump refers to a step that is short enough to possibly (but not necessarily) stay inside a disc of radius  $n(i.e.,$  $|X_1| < 2n$ . A baby jump refers to a small jump that is too short to hop over an s-annulus from inside a disc (*i.e.*,  $|X_1| < s$ ). A **medium** jump refers to a step that is sufficiently large to hop out of a disc and past an s-annulus, but with magnitude strictly less than K, and cannot land near a toral copy of its launching point  $(i.e., s < |X_1| < K - 2n)$ . large jump is a step which, in the toral setting, would be considered "wrapping around" in one step  $(i.e., |X_1| > K - 2n)$ . A targeted jump is a large jump which lands directly in a copy of the disc or annulus just launched from  $(i.e., j(K-2n) \leq |X_1| \leq j(K+2n)/\sqrt{2}$  for some j). These terms will aid in dealing with differences between planar and toral hitting and escape times.

## 2 Random Walk Preliminaries

In this section we give, without proof, results from [\[3\]](#page-19-2) which are used in our Harnack inequality proofs.

First, for  $x, y \in \mathbb{Z}^2$  such that  $|x| \ll |y|$ , we have, by a Taylor expansion around y,

<span id="page-3-0"></span>
$$
\log|y - x| = \log|y| + O\left(\frac{|x|}{|y|}\right). \tag{2.1}
$$

In particular, if  $x \in D(0, 2r)$  and  $y \in D(0, R/2)^c$ , with  $R = 4mr$ , we have

<span id="page-3-1"></span>
$$
\log|y - x| = \log|y| + O(m^{-1}).\tag{2.2}
$$

Note that [\(2.1\)](#page-3-0) and [\(2.2\)](#page-3-1) hold in the toral case without adjustment.

### 2.1 Expected Hitting Times

Here we state some results about the expected escape and entry times of discs on the plane and torus. Our first generalizes a common disc escape argument, and improves on [\[10,](#page-19-3) Prop. 6.2.6].

**Lemma 2.1.** [\[3,](#page-19-2) Lemma 2.1] Let  $S_t = S_0 + \sum_{j=1}^t X_j$  be a random walk in  $\mathbb{Z}^2$  with  $E|X_1|^2 < \infty$ , and covariance matrix  $\Gamma$  such that  $tr(\Gamma) = \gamma^2 > 0$ . Then, uniformly for  $x \in D(0, n)$ , and for sufficiently large n,

<span id="page-3-3"></span>
$$
\frac{n^2 - |x|^2}{\gamma^2} \le \mathbb{E}^x(T_{D(0,n)^c}) \le \frac{n^2 - |x|^2}{\gamma^2} + 2n + 1.
$$
 (2.3)

Computational bounds on the  $\mathbb{E}^{\hat{x}}(T_{\hat{\pi}_K(D(0,n)_{\tilde{K}}^c)})$ , the expected toral disc escape, have a slight torally-induced error term [\[3,](#page-19-2) (2.29)]:

<span id="page-3-4"></span>
$$
\frac{n^2 - |x|^2}{\gamma^2} \le \mathbb{E}^{\hat{x}}(T_{\hat{\pi}_K(D(0,n)^c_K)}) \le \frac{n^2 - |x|^2}{\gamma^2} + 2n + 1 + O(K^{-M}n^4). \tag{2.4}
$$

#### 2.2 Probability Estimates

By Markov's inequality, large jumps are rare: if  $C_M = \mathbb{E}(|X_1|^M) < \infty$ , then since  $2n <$  $K/2$ ,  $2M<sub>z</sub>$ 

<span id="page-3-2"></span>
$$
P(|X_1| > K - 2n) \le \frac{C_M}{(K - 2n)^M} < \frac{2^M C_M}{K^M} = O(K^{-M}).\tag{2.5}
$$

This leads to the targeted jump probability estimate, i.e., the rare chance that a large jump lands back into the toral disc the walk escaped from in the planar setting [\[3,](#page-19-2) (2.24)]:

$$
P^{x}(T_{\hat{\pi}_{K}(D(0,n)^{c}_{K})} > T_{D(0,n)^{c}}) = \sum_{z \in (\hat{\pi}_{K}^{-1}\hat{\pi}_{K}(D(0,n)) \setminus D(0,n))} \sum_{y \in D(0,n)} G_{D(0,n)}(x,y) p_{1}(y,z)
$$
  

$$
\le cK^{-M} \sum_{y \in D(0,n)} G_{D(0,n)}(x,y) = O(K^{-M}n^{2}). \tag{2.6}
$$

[\[3,](#page-19-2) (2.50)] gives a toral gambler's ruin probability estimate for a radius-ruin, i.e., hitting the center of a disc before escaping it, in  $\mathbb{Z}_K^2$ :

$$
P^{\hat{x}}(T_{\hat{0}} < T_{\hat{\pi}_K(D(0,n)_K^c)}) = \frac{\log(n/|\hat{x}|) + O(|\hat{x}|^{-1/4})}{\log(n)} \left(1 + O((\log n)^{-1})\right) + O(K^{-M}n^2)
$$

$$
= \frac{\log(n/|\hat{x}|) + O(|\hat{x}|^{-1/4})}{\log(n)} \left(1 + O((\log n)^{-1})\right). \tag{2.7}
$$

Arguments similar to the proof of  $(2.7)$  also result in probabilities of "near ruin", *i.e.*, entering a smaller disc (rather than hitting the center), and their "success" counterparts, in both  $\mathbb{Z}^2$  and  $\mathbb{Z}_K^2$ : from [\[3,](#page-19-2) (2.51)-(2.54)], uniformly for  $r < |x| < R$ ,

<span id="page-4-4"></span><span id="page-4-2"></span><span id="page-4-0"></span>
$$
P^{x}(T_{D(0,r)} > T_{D(0,R)^c}) = \frac{\log(|x|/r) + O(r^{-1/4})}{\log(R/r)}
$$
\n(2.8)

<span id="page-4-6"></span><span id="page-4-5"></span><span id="page-4-3"></span>
$$
P^{x}(T_{D(0,r)} < T_{D(0,R)^c}) = \frac{\log(R/|x|) + O(r^{-1/4})}{\log(R/r)}\tag{2.9}
$$

$$
P^{\hat{x}}(T_{\hat{\pi}_K(D(0,r))} > T_{\hat{\pi}_K(D(0,R)_K^c)}) = \frac{\log(|\hat{x}|/r) + O(r^{-1/4})}{\log(R/r)} + O(K^{-M}R^2)
$$
  

$$
= \frac{\log(|\hat{x}|/r) + O(r^{-1/4})}{\log(R/r)} \tag{2.10}
$$
  

$$
P^{\hat{x}}(T_{\hat{\pi}_K(D(0,r))} < T_{\hat{\pi}_K(D(0,R)_K^c)}) = \frac{\log(R/|\hat{x}|) + O(r^{-1/4})}{\log(R/r)} + O(K^{-M}R^2)
$$

<span id="page-4-7"></span><span id="page-4-1"></span>
$$
\log(R/r)
$$
\n
$$
= \frac{\log(R/r)}{\log(R/r)} - \frac{\log(R/r)}{\log(R/r)}.
$$
\n(2.11)

Next, we give bounds from [\[3\]](#page-19-2) on the probabilities of exiting a disc when starting far inside, or entering a disc from far outside it, by jumping over an annulus.

**Lemma 2.2.** [\[3,](#page-19-2) Lemmas 4.1, 4.2] For n sufficiently large,

$$
\sup_{\hat{x}\in\hat{\pi}_K(D(0,n/2))} P^{\hat{x}}(T_{\hat{\pi}_K(\partial D(0,n)_s)} > T_{\hat{\pi}_K(D(0,n+s)^c_K)}) \le c(s^{-M+2} \vee n^{-M+2})\tag{2.12}
$$

$$
\sup_{\hat{x}\in\hat{\pi}_K(D(0,n+s)_K^c)} P^{\hat{x}}(T_{\hat{\pi}_K(D(0,n))} < T_{\hat{\pi}_K(\partial D(0,n)_s)}) \le cn^2 \log(n)^2 (s^{-M} + n^{-M}).\tag{2.13}
$$

#### 2.3 Green's Functions

To calculate Green's functions in  $\mathbb{Z}^2$ , we require the *potential kernel* of X: for our walks in  $\mathbb{Z}^2$ , this function is defined by, for  $x \in \mathbb{Z}^2$ ,

<span id="page-5-2"></span>
$$
a(x) := \lim_{n \to \infty} \sum_{j=0}^{n} [p_j(0) - p_j(x)].
$$
\n(2.14)

For our class of random walks, the potential kernel can be shown to be

$$
a(x) = \frac{2}{\pi_{\Gamma}} \log |x| + C(p_1) + o(|x|^{-1})
$$
\n(2.15)

where  $C(p_1)$  is a constant depending on  $p_1$  but not x, and  $\pi_{\Gamma} = 2\pi$ √ det Γ.

[\[3,](#page-19-2) (2.45)-(2.46)] give computational results for the internal Green's function at zero on  $\mathbb{Z}^2$  and  $\mathbb{Z}_K^2$  before escaping a disc:

<span id="page-5-4"></span>
$$
G_{D(0,n)}(0,0) = \frac{2}{\pi_{\Gamma}} \log n + C' + O(n^{-1/4})
$$
\n(2.16)

$$
\hat{G}_{\hat{\pi}_K(D(0,n))}(\hat{0},\hat{0}) = G_{D(0,n)}(0,0)(1 + O(K^{-M}n^2))
$$
  
= 
$$
\left(\frac{2}{\pi_{\Gamma}}\log n + C' + O(n^{-1/4})\right) (1 + O(K^{-M}n^2))
$$
  
= 
$$
\frac{2}{\pi_{\Gamma}}\log n + C' + O(n^{-1/4}).
$$
 (2.17)

[\[3,](#page-19-2) (2.55)-(2.58)] give calculations and bounds for  $G_{D(0,n)}(x,0), \hat{G}_{\hat{\pi}_K(D(0,n))}(\hat{x},\hat{0}), G_{D(0,n)}(x,z)$ , and  $\hat{G}_{\hat{\pi}_K(D(0,n))}(\hat{x}, \hat{z})$ : for  $x \in D(0, n)$  and  $\hat{x} \in \hat{\pi}_K(D(0, n))$ : for some  $C = C(p_1) < \infty$ ,

<span id="page-5-5"></span><span id="page-5-0"></span>
$$
G_{D(0,n)}(x,0) = P^x(T_0 < T_{D(0,n)^c}) \, G_{D(0,n)}(0,0) \\
= \frac{2}{\pi_\Gamma} \log \left( \frac{n}{|x|} \right) + C + O(|x|^{-1/4}),\n \tag{2.18}
$$

$$
\hat{G}_{\hat{\pi}_K(D(0,n))}(\hat{x},\hat{0}) = \frac{2}{\pi_{\Gamma}} \log \left(\frac{n}{|\hat{x}|}\right) + C + O(|\hat{x}|^{-1/4}) \tag{2.19}
$$

$$
G_{D(0,n)}(x,z) \le G_{D(x,2n)}(0,z-x) \le c \log n. \tag{2.20}
$$

$$
\hat{G}_{\hat{\pi}_K(D(0,n))}(\hat{x}, \hat{z}) = G_{D(0,n)}(x, z) + O(K^{-M}n^2 \log n) \le c \log n.
$$
 (2.21)

[\[3,](#page-19-2) Lemma 2.8] gives that, for any  $0 < \delta < \varepsilon < 1$ , we can find  $0 < c_1 < c_2 < \infty$ , such that for all  $\hat{x} \in \hat{\pi}_K(D(0,n)) \setminus \hat{\pi}_K(D(0,\varepsilon n)), \hat{y} \in \hat{\pi}_K(D(0,\delta n))$  and all n sufficiently large such that  $2n < K/2$ ,

<span id="page-5-3"></span><span id="page-5-1"></span>
$$
c_1 \frac{\rho(\hat{x}) \vee 1}{n} \le \hat{G}_{\hat{\pi}_K(D(0,n))}(\hat{y}, \hat{x}) \le c_2 \frac{\rho(\hat{x}) \vee 1}{n}.
$$
\n(2.22)

Finally, for any  $x, y \in \pi_K(D(0, n)_K^c)$  such that  $|x| \leq |y|$ , there are computational bounds on the external Green's function before entering a disc:

<span id="page-6-4"></span>
$$
G_{D(0,n)^c}(x,y) \le c_j \log |x|,\tag{2.23}
$$

$$
\hat{G}_{\hat{\pi}_K(D(0,n)_K^c)}(\hat{x},\hat{y}) \leq \hat{c}_j \log|\hat{x}|,\tag{2.24}
$$

where  $c_j, \hat{c}_j$  depend on  $j > 2, c_j \geq \hat{c}_j$ , and in the toral case, such that  $|\hat{x}| < (\frac{K}{2})$  $(\frac{K}{2})^{1/j}$  (there is no such restriction on the planar case).

## 3 Interior Harnack inequalities

We call our first Harnack inequality "interior": the starting points are from the interior of a disc, and we examine the probabilities of escaping from a far larger disc around it. We find the planar version first, then move it to the torus.

**Lemma 3.1.** Uniformly for  $1 \leq m \ll r$ , with  $s \ll \frac{r}{4m}$ ,  $x, x' \in D(0, 2r)$ ,  $R = 4mr$ , and  $y \in D(0,R)^c$ ,

$$
H_{D(0,R)^c}(x,y) = (1 + O(m^{-1}))H_{D(0,R)^c}(x',y) + O(R^{-M}\log R),\tag{3.1}
$$

where the error term is completely absorbed, i.e.,

<span id="page-6-3"></span><span id="page-6-2"></span><span id="page-6-1"></span><span id="page-6-0"></span>
$$
H_{D(0,R)^c}(x,y) = (1 + O(m^{-1}))H_{D(0,R)^c}(x',y),
$$
\n(3.2)

if  $s \leq (\log R)^4$  and  $y \in \partial D(0,R)_s$ .

Furthermore, if  $x \in \partial D(0, r)_r$  and  $y \in D(0, R)^c$ ,

$$
P^x \left( S_{T_{D(0,R)^c}} = y, T_{D(0,R)^c} < T_{D\left(0, \frac{r}{4m} + s\right)} \right) \tag{3.3}
$$
\n
$$
= (1 + O(m^{-1})) P^x \left( T_{D(0,R)^c} < T_{D\left(0, \frac{r}{4m} + s\right)} \right) H_{D(0,R)^c}(x, y) + O(R^{-M} \log R),
$$

with a similar loss of the error term if  $y \in \partial D(0,R)_{s}$ .

**Proof** (In this proof, we switch freely between R and  $4mr$ .) First, we decompose  $D(0, R)$ and examine  $H_{D(0,R)^c}(x,y)$ :

$$
H_{D(0,R)^c}(x,y) = \left(\sum_{z \in D(0,2mr)} + \sum_{\substack{z \in D(0,3mr) \\ \langle D(0,2mr) \rangle}} + \sum_{\substack{z \in D(0,4mr) \\ \langle D(0,3mr) \rangle}} \right) G_{D(0,R)}(x,z) p_1(z,y). \tag{3.4}
$$

If X is finite range, then for  $r$  sufficiently large, the first two sums of  $(3.4)$  are zero. Otherwise, we bound the Green's function via [\(2.18\)](#page-5-0) and [\(2.20\)](#page-5-1), and by Markov's inequality,  $\sum_{z \in D(0,2mr)} p_1(z,y) \leq c(mr)^{-M} \leq cR^{-M}$ . Together, these yield, for some  $c < \infty$ ,

<span id="page-7-2"></span>
$$
G_{D(0,R)}(x, z) \le G_{D(0,2R)}(0, z) \le c \log R
$$
  
\n
$$
\implies \sum_{z \in D(0,2mr)} G_{D(0,R)}(x, z) p_1(z, y) \le cR^{-M} \log R.
$$
 (3.5)

By [\(2.15\)](#page-5-2) and [\(2.1\)](#page-3-0), uniformly in  $x \in D(0, 2r)$  and  $y \in D(0, 2mr)^c$ ,

<span id="page-7-0"></span>
$$
a(y - x) = \frac{2}{\pi_{\Gamma}} \log |y - x| + C' + O(|y - x|^{-1})
$$
  
= 
$$
\frac{2}{\pi_{\Gamma}} \log |y| + C' + O(m^{-1}) = a(y) + O(m^{-1}).
$$
 (3.6)

For  $z \in D(0, 4mr) \setminus D(0, 2mr)$ , by the symmetry of the Green's function, the fact that H is a probability,  $[10, (4.28)]$ , and  $(3.6)$ , we have

$$
G_{D(0,R)}(x,z) = G_{D(0,R)}(z,x)
$$
  
= 
$$
\left(\sum_{w \in D(0,R)^c} H_{D(0,R)^c}(z,w)a(w-x)\right) - a(z-x)
$$
(3.7)  
= 
$$
\left(\sum_{w \in D(0,R)^c} H_{D(0,R)^c}(z,w)a(w)\right) - a(z) + O(m^{-1})
$$
  
= 
$$
G_{D(0,4mr)}(z,0) + O(m^{-1}).
$$

By [\(2.18\)](#page-5-0),  $G_{D(0,R)}(z,0) \ge c > 0$  uniformly for  $z \in D(0,3mr) \setminus D(0,2mr)$ , yielding

<span id="page-7-3"></span><span id="page-7-1"></span>
$$
G_{D(0,R)}(x,z) = G_{D(0,R)}(0,z)(1 + O(m^{-1})).
$$
\n(3.8)

For  $z \in D(0, 4mr) \setminus D(0, 3mr)$ , by the strong Markov property at  $T_{D(0, 3mr)}$ ,

$$
G_{D(0,R)}(z,x) = \mathbb{E}^{z}(G_{D(0,R)}(S_{T_{D(0,3mr)}},x);T_{D(0,3mr)} < T_{D(0,4mr)^{c}})
$$
(3.9)  
= 
$$
\mathbb{E}^{z}(G_{D(0,R)}(S_{T_{D(0,3mr)}},x);T_{D(0,3mr)} < T_{D(0,4mr)^{c}};|X_{T_{D(0,3mr)}}| > 2mr)
$$
  
+ 
$$
\mathbb{E}^{z}(G_{D(0,R)}(S_{T_{D(0,3mr)}},x);T_{D(0,3mr)} < T_{D(0,4mr)^{c}};|X_{T_{D(0,3mr)}}| \leq 2mr).
$$

By  $(2.20)$  and  $(2.13)$ , the last term here is bounded, for sufficiently large r, by

$$
c(\log R)P^{z}(|X_{T_{D(0,3mr)}}| \le 2mr) \le c(\log R)P^{z}(T_{D(0,2mr)} < T_{\partial D(0,2mr)_{mr}})
$$
  
\n
$$
\le c(\log R)(2mr)^{2}\log(2mr)^{2}[(mr)^{-M} + (2mr)^{-M}]
$$
  
\n
$$
\le c(\log R)^{3}R^{-M+2} \le cR^{-M+2+\beta}.
$$

Applying [\(3.8\)](#page-7-1) to the first term, then switching it back to its original form, yields, for

 $z \in D(0, 4mr) \setminus D(0, 3mr),$ 

$$
G_{D(0,R)}(z,x) = (1 + O(m^{-1})) G_{D(0,R)}(z,0) + O(R^{-M+2+\beta}).
$$
\n(3.10)

The planar version of  $(2.22)$  gives us  $G_{D(0,R)}(z,x) \geq \frac{c}{m}$  $\frac{c}{mr}$  for  $z \in D(0, 4mr) \setminus D(0, 3mr)$ . This reduces [\(3.10\)](#page-8-0) to

<span id="page-8-0"></span>
$$
G_{D(0,R)}(z,x) = (1 + O(m^{-1})) G_{D(0,R)}(z,0).
$$
\n(3.11)

Combining [\(3.4\)](#page-6-0), [\(3.5\)](#page-7-2), [\(3.8\)](#page-7-1), and [\(3.11\)](#page-8-1) yields [\(3.1\)](#page-6-1).

For [\(3.2\)](#page-6-2), let  $y \in \partial D(0,R)$ . The only thing we need to do here is show that the error terms are absorbed, *i.e.*, for some  $c > 0$ , with  $M = 4 + 2\beta$ ,

<span id="page-8-1"></span>
$$
m^{-1}H_{D(0,R)^c}(x,y) \ge cR^{-M}\log R. \tag{3.12}
$$

Wlog, we can show this for  $x = 0$ . First note that, for  $|z| \leq \frac{R}{100}$ , by [\(2.18\)](#page-5-0) we have

$$
G_{D(0,R)}(z,0) \ge c \log \frac{R}{(R/100)} = c \log 100 \ge c \ge \frac{c}{R}
$$

for some  $c > 0$ , and for  $z \in D(0, R) \setminus D(0, R/100)$ , by the planar version of  $(2.22)$ ,  $G_{D(0,R)}(z,0) \geq \frac{c}{R}$  $\frac{c}{R}$  as well. Hence, by this, a last exit decomposition, and [\(1.1\)](#page-1-0),

$$
m^{-1}H_{D(0,R)^c}(0,y) = m^{-1} \sum_{z \in D(0,R)} G_{D(0,R)}(0,z)p_1(z,y) \ge \frac{c}{mR} \sum_{z \in D(0,R)} p_1(z,y) \quad (3.13)
$$

$$
\ge \frac{c}{mR}e^{-\beta s^{1/4}} = c(mR)^{-1}e^{-\beta \log R} = cm^{-1}R^{-1-\beta} > cR^{-M} \log R.
$$

To show [\(3.3\)](#page-6-3), we start with the decomposition

<span id="page-8-3"></span>
$$
P^x \left( S_{T_{D(0,R)^c}} = y, T_{D(0,R)^c} < T_{D\left(0, \frac{r}{4m} + s\right)} \right) \tag{3.14}
$$
\n
$$
= H_{D(0,R)^c}(x,y) - P^x \left( S_{T_{D(0,R)^c}} = y, T_{D(0,R)^c} > T_{D\left(0, \frac{r}{4m} + s\right)} \right).
$$

By the strong Markov property at  $T_{D\left(0, \frac{r}{4m} + s\right)},$ 

<span id="page-8-2"></span>
$$
P^x \left( S_{T_{D(0,R)^c}} = y, T_{D(0,R)^c} > T_{D\left(0, \frac{r}{4m} + s\right)} \right)
$$
  
=  $\mathbb{E}^x \left[ H_{D(0,R)^c} \left( S_{T_{D(0,\frac{r}{4m} + s)}}, y \right); T_{D(0,R)^c} > T_{D\left(0, \frac{r}{4m} + s\right)} \right].$  (3.15)

By [\(3.1\)](#page-6-1), uniformly in  $w \in D(0, 2r)$ ,

$$
H_{D(0,R)^c}\left(S_{T_{D(0,\frac{r}{4m}+s)}},y\right) = \left(1+O\left(m^{-1}\right)\right)H_{D(0,R)^c}(w,y) + O(R^{-M}\log R).
$$

By [\(2.8\)](#page-4-2) and [\(2.9\)](#page-4-3), with  $m \gg 1$ , uniformly for  $x \in \partial D(0, r)_r$  (say  $|x| = cr$ ,  $1 < c < 2$ ),

 $\exists c', c'' > 0$  such that

<span id="page-9-2"></span>
$$
P^x \left( T_{D(0,R)^c} < T_{D\left(0, \frac{r}{4m} + s\right)} \right) = \frac{\log(c'm) + O\left(\left(\frac{r}{m}\right)^{-1/4}\right)}{\log(c''m^2)} = \frac{1}{2} + o\left(\left(\frac{r}{m}\right)^{-1/4}\right),
$$
\n
$$
P^x \left( T_{D(0,R)^c} > T_{D\left(0, \frac{r}{4m} + s\right)} \right) = \frac{1}{2} + o\left(\left(\frac{r}{m}\right)^{-1/4}\right),\tag{3.16}
$$

so the probabilities are both bounded below by a constant. (The small  $m$  case operates similarly, but due to the small constants involved, the lower bound must be reduced;  $\frac{1}{4}$ for one of them suffices.) Combining these and [\(3.15\)](#page-8-2) into [\(3.14\)](#page-8-3) yields

$$
P^{x}\left(S_{T_{D(0,R)^{c}}}=y, T_{D(0,R)^{c}} < T_{D(0,\frac{r}{4m}+s)}\right)
$$
  
\n
$$
=H_{D(0,R)^{c}}(x,y)-\mathbb{E}^{x}\left[H_{D(0,R)^{c}}\left(S_{T_{D(0,\frac{r}{4m}+s)}},y\right); T_{D(0,R)^{c}} > T_{D(0,\frac{r}{4m}+s)}\right]
$$
  
\n
$$
=H_{D(0,R)^{c}}(x,y)\left[P^{x}\left(T_{D(0,R)^{c}} < T_{D(0,\frac{r}{4m}+s)}\right)+P^{x}\left(T_{D(0,R)^{c}} > T_{D(0,\frac{r}{4m}+s)}\right)\right]
$$
  
\n
$$
-\left(1+O\left(m^{-1}\right)\right)P^{x}\left(T_{D(0,R)^{c}} > T_{D(0,\frac{r}{4m}+s)}\right)+O(R^{-M}\log R)\right]
$$
  
\n
$$
=H_{D(0,R)^{c}}(x,y)\left(1+O\left(m^{-1}\right)\right)P^{x}\left(T_{D(0,R)^{c}} < T_{D(0,\frac{r}{4m}+s)}\right)+O(R^{-M}\log R).
$$

We now move these results to the torus.

**Proposition 3.2.** For large r and  $1 \le m \ll r$  such that  $R = 4mr < K/6$  and  $s \le (\log R)^4$ , uniformly for  $\hat{x}, \hat{x}' \in \hat{\pi}_K(D(0, 2r))$  and  $\hat{y} \in \hat{\pi}_K(D(0, R)^c_K)$ ,

<span id="page-9-1"></span><span id="page-9-0"></span>
$$
\hat{H}_{\hat{\pi}_K(D(0,R)_K^c)}(\hat{x},\hat{y}) = (1+O(m^{-1})) \hat{H}_{\hat{\pi}_K(D(0,R)_K^c)}(\hat{x}',\hat{y}) \n+ O(R^{-M}\log R \vee K^{-M}R^2).
$$
\n(3.17)

Furthermore, uniformly in  $\hat{x} \in \hat{\pi}_K(\partial D(0,r)_r)$  and  $\hat{y} \in \hat{\pi}_K(D(0,R)_K^c)$ ,

$$
P^{\hat{x}}(\hat{S}_{T_{\hat{\pi}_K(D(0,R)^c_K)}}) = \hat{y}, T_{\hat{\pi}_K(D(0,R)^c_K)} < T_{\hat{\pi}_K(D(0,\frac{r}{4m}+s))})
$$
\n
$$
= \left(1 + O\left(m^{-1}\right)\right) P^{\hat{x}}(T_{\hat{\pi}_K(D(0,R)^c_K)} < T_{\hat{\pi}_K(D(0,\frac{r}{4m}+s))}) \hat{H}_{\hat{\pi}_K(D(0,R)^c_K)}(\hat{x},\hat{y})
$$
\n
$$
+ O(R^{-M} \log R \vee K^{-M} R^2). \tag{3.18}
$$

If  $\hat{y} \in \hat{\pi}_K(\partial D(0,R)_s)$ , the error term is absorbed in both of these statements.

**Proof** As before, wlog, we can take  $\hat{x}' = \hat{0}$ . Let s be the size of the annulus for  $\hat{y}$ . For brevity, set

$$
A_p := \{S_{T_{D(0,R)^c}} = y\}, \t d_p := |S_{T_{D(0,R)^c}} - S_{T_{D(0,R)^c} - 1}|,
$$
  
\n
$$
A_t := \{\hat{S}_{T_{\hat{\pi}_K(D(0,R)^c_K)}} = \hat{y}\}, \t d_t := |S_{T_{\hat{\pi}_K(D(0,R)^c_K)}} - S_{T_{\hat{\pi}_K(D(0,R)^c_K)} - 1}|.
$$

Note that x and y are the primary copies of  $\hat{x}$  and  $\hat{y}$ , and so  $|x - y| \leq \frac{K}{\sqrt{2}}$  $\frac{1}{2}$ , but that  $d_p$  is a planar distance using a planar escape time and  $d_t$  is a planar distance using a toral escape time; hence, both can exceed  $\frac{K}{\sqrt{6}}$  $\frac{1}{2}$ , the maximum distance between two points in  $\mathbb{Z}_K^2$ .

To prove  $(3.17)$ , first we re-label  $(3.1)$  as

<span id="page-10-3"></span>
$$
H_{D(0,R)^c}(x,y) = P^x(A_p) = \left(1 + O\left(\frac{r}{R}\right)\right) P^{x'}(A_p) + O\left(R^{-M}\log R\right). \tag{3.19}
$$

We have the decomposition

<span id="page-10-0"></span>
$$
P^{x}(A_{p}) = P^{x}(A_{p}; d_{p} < K - 2R) + P^{x}(A_{p}; d_{p} \ge K - 2R).
$$
 (3.20)

On the plane, the second term of  $(3.20)$  is zero for all but the furthest-away y in the primary copy (*i.e.*,  $K - 2R \leq |x - y| \leq \frac{K}{4}$  $(\frac{1}{2});$  for those y, we have, by  $(2.5), (1.4)$  $(2.5), (1.4)$ , and  $(2.3),$  $(2.3),$ 

$$
P^{x}(A_{p}; d_{p} \ge K - 2R) = \sum_{z \in D(0,R)} G_{D(0,R)}(x, z) P^{z}(|X_{1} - z| > K - 2R)
$$
  

$$
\le cK^{-M}R^{2}.
$$
 (3.21)

The toral version can be written using planar distances as a decomposition, but using the toral disc escape time means a further decomposition comparing the planar and toral escape times a la [\(2.6\)](#page-4-4). We decompose  $\hat{H}_{\hat{\pi}_K(D(0,R)_K^c)}(\hat{x},\hat{y}) = P^{\hat{x}}(A_t)$  as

<span id="page-10-2"></span><span id="page-10-1"></span>
$$
P^{\hat{x}}(A_t) = P^{\hat{x}}(A_t; d_t < K - 2R; T_{D(0,R)^c} = T_{\hat{\pi}_K(D(0,R)^c_K)})
$$
\n
$$
+ P^{\hat{x}}(A_t; d_t \ge K - 2R; T_{D(0,R)^c} = T_{\hat{\pi}_K(D(0,R)^c_K)})
$$
\n
$$
+ P^{\hat{x}}(A_t; T_{D(0,R)^c} < T_{\hat{\pi}_K(D(0,R)^c_K)}).
$$
\n
$$
(3.22)
$$

In the torus, the first term of  $(3.22)$  equals the first term of  $(3.20)$ , plus a large jump error which contains some paths from the second term of  $(3.20)$  (if y is far): by  $(3.21)$ ,

$$
P^{\hat{x}}(A_t; d_t < K - 2R; T_{D(0,R)^c} = T_{\hat{\pi}_K(D(0,R)^c_K)})
$$
\n
$$
= P^x(A_p; d_p < K - 2R) + P^x(A_p; d_p \ge K - 2R; d_t < K - 2R)
$$
\n
$$
= P^x(A_p) + O(K^{-M}R^2).
$$

The second term of  $(3.22)$  only occurs if the final, escaping jump is large: by  $(2.5)$ ,  $(1.4)$ , and [\(2.4\)](#page-3-4), just as in [\(3.21\)](#page-10-2),

$$
P^{\hat{x}}(A_t; d_t \ge K - 2R; T_{D(0,R)^c} = T_{\hat{\pi}_K(D(0,R)^c_K)}) \le P^{\hat{x}}(A_t; d_t \ge K - 2R)
$$
  
= 
$$
\sum_{\hat{z} \in \hat{\pi}_K(D(0,R))} \hat{G}_{\hat{\pi}_K(D(0,R))}(\hat{x}, \hat{z}) P^z(|X_1 - z| > K - 2R) \le cK^{-M}R^2.
$$

The last term of  $(3.22)$  requires a large jump to have occured. Hence, by  $(2.6)$ ,

$$
P^{\hat{x}}(A_t; T_{D(0,R)^c} < T_{\hat{\pi}_K(D(0,R)^c_K)}) \le cK^{-M}R^2.
$$

Therefore, [\(3.22\)](#page-10-1) reduces to

$$
P^{\hat{x}}(A_t) = P^x(A_p) + O(K^{-M}R^2).
$$
\n(3.23)

(Due to targeting, this is generalizable to any planar set  $B \subset D(0,R)$  for  $R \lt K/4$ .) Combining this with  $(3.19)$  gives us  $(3.17)$ :

$$
P^{\hat{x}}(A_t) = P^x(A_p) + O(K^{-M}R^2)
$$
  
=  $(1 + O(\frac{r}{R})) P^{x'}(A_p) + O(K^{-M}R^2) + O(R^{-M}\log R)$   
=  $(1 + O(\frac{r}{R})) P^{\hat{x}'}(A_t) + O(K^{-M}R^2) + O(R^{-M}\log R).$ 

The proof of  $(3.18)$  follows from the Markov property argument for  $(3.3)$ , using the appropriate toral identities:  $(3.17)$  for  $(3.1)$ , and  $(2.10)-(2.11)$  $(2.10)-(2.11)$  for  $(2.8)-(2.9)$  $(2.8)-(2.9)$ .  $\Box$ 

## 4 Exterior Harnack inequality

To aid our construction of an exterior Harnack inequality (moving from outside a disc to a much smaller disc far inside it), we first establish uniform bounds on external Green's functions and probabilities in the torus and plane. Fix  $\delta$  < 1 and use  $r \geq e^n$  for some  $n > 13$ ,  $R = 4mr$  for some  $1 \le m \ll r$ , and  $s \le (\log R)^4$ . First, for  $\hat{x} \in \hat{\pi}_K(\partial D(0,R)_{R/100})$ and  $\hat{y} \in \hat{\pi}_K(D(0,R)_K^c)$ , we show that

<span id="page-11-1"></span><span id="page-11-0"></span>
$$
\hat{G}_{\hat{\pi}_K(D(0,r+s)^\text{c}_K)}(\hat{x},\hat{y}) \ge c > 0. \tag{4.1}
$$

Pick some  $\hat{x}_1 \in \hat{\pi}_K(\partial D(0,R))$ , and, proceeding clockwise, choose points  $\hat{x}_2, ..., \hat{x}_{36} \in$  $\hat{\pi}_K(\partial D(0,R))$  whose rays beginning at  $\hat{0}$  divide  $\hat{\pi}_K(\partial D(0,R))$  into 36 approximately equal arcs. The distance between any two adjacent such  $\hat{x}_j$  is, for sufficiently large R, approximately  $2R\sin(\pi/36) \approx 0.174R$ . Thus, using discs of radius  $R/5$  (so adjacent circles contain their neighbor's centers), and by  $(2.11)$  we have for any  $j = 1, ..., 36$ 

$$
\inf_{\hat{x}\in\hat{\pi}_K(D(x_j,R/5))} P^{\hat{x}}(T_{\hat{\pi}_K(D(x_{j+1},R/5))} < T_{\hat{\pi}_K(D(0,r+s))})\n\geq \inf_{\hat{x}\in\hat{\pi}_K(D(x_{j+1},R/5))} P^{\hat{x}}(T_{\hat{\pi}_K(D(x_{j+1},R/5))} < T_{\hat{\pi}_K(D(\hat{x}_{j+1},R/2)_{K}^c)}) \geq c_1 > 0,
$$
\n
$$
(4.2)
$$

for some  $c_1$  independent of  $n, r, R, n$  large, and  $\hat{x}_{37} = \hat{x}_1$ .

Hence, by the strong Markov property, rotating through the arcs, we have

$$
\inf_{j,k} \inf_{\hat{x} \in \hat{\pi}_K(D(x_j, R/5))} P^{\hat{x}}(T_{\hat{\pi}_K(D(x_k, R/5))} < T_{\hat{\pi}_K(D(0, r+s))}) \ge c_2 := c_1^{36}.\tag{4.3}
$$



**Figure 1:** Possible selections of  $\hat{x}_j$  and  $\hat{x}_{j+1}$ , with associated discs and annuli.

Furthermore, it follows from  $(2.7)$  that for any j,

<span id="page-12-0"></span>
$$
\inf_{\hat{x}, \hat{x}' \in \hat{\pi}_K(D(x_j, R/5))} P^{\hat{x}}(T_{\hat{x}'} < T_{\hat{\pi}_K(D(0, r+s))})
$$
\n
$$
\geq \inf_{\hat{x} \in \hat{\pi}_K(D(x', 2R/5))} P^{\hat{x}}(T_{\hat{x}'} < T_{\hat{\pi}_K(D(x', R/2)_{\hat{K}}^c})) \geq \frac{c_3}{\log R}
$$
\n
$$
(4.4)
$$

for some independent  $c_3 > 0$ .

Since  $\hat{\pi}_K(\partial D(0,R)_{R/100}) \subset \cup_{j=1}^{36} \hat{\pi}_K(D(x_j,R/5)),$  combining [\(4.3\)](#page-11-0) and [\(4.4\)](#page-12-0) we have, for some independent  $c_4 > 0$ ,

<span id="page-12-2"></span><span id="page-12-1"></span>
$$
\inf_{\hat{x}, \hat{x}' \in \hat{\pi}_K(\partial D(0, R)_{R/100})} P^{\hat{x}}(T_{\hat{x}'} < T_{\hat{\pi}_K(D(0, r+s))}) \ge \frac{c_4}{\log R}.\tag{4.5}
$$

It then follows from [\(2.17\)](#page-5-4) that

$$
\inf_{\hat{x},\hat{x}' \in \hat{\pi}_K(\partial D(0,R)_{R/100})} \hat{G}_{\hat{\pi}_K(D(0,r+s)_{K}^c)}(\hat{x},\hat{x}')
$$
\n
$$
= \inf_{\hat{x},\hat{x}' \in \hat{\pi}_K(\partial D(0,R)_{R/100})} P^{\hat{x}}(T_{\hat{x}'} < T_{\hat{\pi}_K(D(0,r+s))}) \hat{G}_{\hat{\pi}_K(D(0,r+s)_{K}^c)}(\hat{x}',\hat{x}')
$$
\n
$$
\geq \frac{c_4}{\log R} \hat{G}_{\hat{\pi}_K(D(x',R/2))}(\hat{x}',\hat{x}') \geq c_5 > 0 \tag{4.6}
$$

for some independent  $c_5 > 0$ . Using the strong Markov property,  $(4.6)$ , and  $(2.13)$ , we see

that

$$
\inf_{\hat{z}\in\hat{\pi}_{K}(D(0,1,01R)^{c}_{K}),\atop\hat{x}\in\hat{\pi}_{K}(\partial D(0,R)_{R/100})}\hat{G}_{\hat{\pi}_{K}(D(0,r+s)^{c}_{K})}(\hat{z},\hat{x})
$$
\n
$$
\geq \mathbb{E}^{\hat{z}}\left(\hat{G}_{\hat{\pi}_{K}(D(0,r+s)^{c}_{K})}(\hat{S}_{T_{\hat{\pi}_{K}(D(0,1,01R))}},\hat{x});\hat{S}_{T_{\hat{\pi}_{K}(D(0,1,01R))}}\in\hat{\pi}_{K}(\partial D(0,R)_{R/100})\right)
$$
\n
$$
\geq c > 0.
$$
\n(4.7)

This gives [\(4.1\)](#page-11-1). Applying the same argument once more,

$$
\inf_{\hat{z}\in\hat{\pi}_{K}(D(0,1.01R)_{K}^{c})}\hat{G}_{\hat{\pi}_{K}(D(0,r+s)_{K}^{c})}(\hat{z},\hat{x})
$$
\n
$$
\hat{z}\in\hat{\pi}_{K}(D(0,R)_{K}^{c})
$$
\n
$$
\geq \mathbb{E}^{\hat{z}}\left(\hat{G}_{\hat{\pi}_{K}(D(0,r+s)_{K}^{c})}(\hat{S}_{T_{\hat{\pi}_{K}(D(0,1.01R))}},\hat{x});\hat{S}_{T_{\hat{\pi}_{K}(D(0,1.01R))}}\in\hat{\pi}_{K}(\partial D(0,R)_{R/100})\right)
$$
\n
$$
\geq c > 0.
$$
\n(4.8)

Hence, for all  $\hat{x}, \hat{y} \in \hat{\pi}_K(D(0,R)_K^c)$ ,

<span id="page-13-2"></span><span id="page-13-0"></span>
$$
\hat{G}_{\hat{\pi}_K(D(0,r+s)_{K}^c)}(\hat{x},\hat{y}) \ge c > 0.
$$
\n(4.9)

Next, we look at the external Green's function near the r-disc: uniformly for  $\hat{x} \in \hat{\pi}_K(D(0,R)^c_K)$ and  $\hat{z} \in \hat{\pi}_K(D(0, 2r)) \setminus \hat{\pi}_K(D(0, 5r/4))$ , we have by [\(4.9\)](#page-13-0) and [\(2.10\)](#page-4-5),

$$
\hat{G}_{\hat{\pi}_K(D(0,r+s)_{K}^c)}(\hat{x}, \hat{z}) = \hat{G}_{\hat{\pi}_K(D(0,r+s)_{K}^c)}(\hat{z}, \hat{x})
$$
\n
$$
= \mathbb{E}^{\hat{z}} \left( \hat{G}_{\hat{\pi}_K(D(0,r+s)_{K}^c)}(\hat{S}_{T_{\hat{\pi}_K(D(0,R)_{K}^c)}}, \hat{x}); T_{\hat{\pi}_K(D(0,R)_{K}^c)} < T_{\hat{\pi}_K(D(0,r+s))} \right)
$$
\n
$$
\geq c P^{\hat{z}} (T_{\hat{\pi}_K(D(0,R)_{K}^c)} < T_{\hat{\pi}_K(D(0,r+s))}) \geq \frac{c}{\log m}.
$$
\n
$$
(4.10)
$$

Getting closer to the disc, for any  $\varepsilon > 0$ , uniformly in  $\hat{x}, \hat{x}' \in \hat{\pi}_K(\partial D(0,R)_{R/100})$  and  $\hat{z} \in \hat{\pi}_K(D(0, 2r)) \setminus \hat{\pi}_K(D(0, r + (1 + \varepsilon)s)),$  we have by the strong Markov property and [\(4.5\)](#page-12-2), for any  $\hat{x}' \in \partial D(0,R)_{R/100}$ ,

$$
\hat{G}_{\hat{\pi}_K(D(0,r+s)_{K}^c)}(\hat{x},\hat{z}) \ge P^{\hat{x}}(T_{\hat{x}'} < T_{\hat{\pi}_K(D(0,r+s))}) \hat{G}_{\hat{\pi}_K(D(0,r+s)_{K}^c)}(\hat{x}',\hat{z})
$$
\n
$$
\ge \frac{c\hat{G}_{\hat{\pi}_K(D(0,r+s)_{K}^c)}(\hat{x}',\hat{z})}{\log R}.\tag{4.11}
$$

In view of  $(2.22)$ , if  $\hat{x}' \in \hat{\pi}_K(\partial D(0,R)_{R/100})$  is chosen as close as possible to the ray from the origin which passes through  $\hat{z}$ , we have

<span id="page-13-1"></span>
$$
\hat{G}_{\hat{\pi}_K(D(0,r+s)_{K}^c)}(\hat{x}',\hat{z}) \ge \hat{G}_{\hat{\pi}_K(D(\hat{x}',|\hat{x}'|-(r+s)))}(\hat{x}',\hat{z}) \ge \frac{c}{R},\tag{4.12}
$$

which, combined with [\(4.11\)](#page-13-1), gives us

<span id="page-14-0"></span>
$$
\inf_{\hat{x}\in\hat{\pi}_K(\partial D(0,R)_R/100)} \hat{G}_{\hat{\pi}_K(D(0,r+s)^c_K)}(\hat{x},\hat{z}) \ge \frac{c}{R\log R}.
$$
\n(4.13)

Using the strong Markov property, [\(4.13\)](#page-14-0), and [\(2.12\)](#page-4-7), we see that

$$
\inf_{\hat{x}\in\hat{\pi}_{K}(D(0,1.01R)_{K}^{c}))}\hat{G}_{\hat{\pi}_{K}(D(0,r+s)_{K}^{c})}(\hat{z},\hat{x})
$$
\n
$$
\geq \mathbb{E}^{\hat{z}}\left(\hat{G}_{\hat{\pi}_{K}(D(0,r+s)_{K}^{c})}(\hat{S}_{T_{\hat{\pi}_{K}(D(0,1.01R)_{K}^{c})}},\hat{x});\hat{S}_{T_{\hat{\pi}_{K}(D(0,1.01R)_{K}^{c})}}\in\hat{\pi}_{K}(\partial D(0,R)_{R/100})\right)
$$
\n
$$
\geq \frac{c}{R\log R}.
$$
\n(4.14)

Hence

<span id="page-14-1"></span>
$$
\inf_{\substack{\hat{x}\in\hat{\pi}_K(D(0,R)^c_{K}))\\\hat{z}\in\hat{\pi}_K(D(0,2r))\setminus\hat{\pi}_K(D(0,r+(1+\varepsilon)s))}}\hat{G}_{\hat{\pi}_K(D(0,r+s)^c_{K})}(\hat{x},\hat{y}) \ge \frac{c}{R\log R}.
$$
\n(4.15)

By removing the (hidden) targeted jump error terms, the entire argument in [\(4.1\)](#page-11-1)-[\(4.15\)](#page-14-1) also applies to the plane. We now find a general planar Harnack inequality for entering a small disc from far outside.

**Proposition 4.1.** Let  $R = 4mr$  with  $1 \le m \ll r$   $(m = o(r^{1/4}))$  and large enough r, and  $s \leq (\log R)^4$ . Then, uniformly for  $x, x' \in D(0, R)^c$  and  $y \in \partial D(0, r)_s$ ,

<span id="page-14-5"></span><span id="page-14-4"></span><span id="page-14-3"></span>
$$
H_{D(0,r+s)}(x,y) = \left(1 + O\left(m^{-1}\log m\right)\right)H_{D(0,r+s)}(x',y). \tag{4.16}
$$

Furthermore, for  $x, x' \in \partial D(0, R)_{\sqrt{R}}$ ,

$$
P^{x}(S_{T_{D(0,r+s)}} = y; T_{D(0,r+s)} < T_{D(0,4mR)^{c}})
$$
\n
$$
= (1+O(m^{-1}\log m)) H_{D(0,r+s)}(x,y) P^{x}(T_{D(0,r+s)} < T_{D(0,4mR)^{c}})
$$
\n
$$
= (1+O(m^{-1}\log m)) P^{x'}(S_{T_{D(0,r+s)}} = y; T_{D(0,r+s)} < T_{D(0,4mR)^{c}}).
$$
\n(4.17)

**Proof** For  $x, x' \in D(0, R)^c$  and  $y \in \partial D(0, r)_s$ , we have the last exit decomposition

$$
H_{D(0,r+s)}(x,y) = \left(\sum_{\substack{z \in D(0,5r/4) \\ \backslash D(0,r+s)}} + \sum_{\substack{z \in D(0,2r) \\ \backslash D(0,5r/4)}} + \sum_{z \in D(0,2r)^c} \right) G_{D(0,r+s)^c}(x,z) p_1(z,y). \tag{4.18}
$$

Let  $x, x' \in \partial D(0,R)_R$  and set  $N \ge 4mR$ . Uniformly for  $z \in D(0, 2r) \cup D(0, N)^c$ , by  $(2.15)$ and [\(2.2\)](#page-3-1),

<span id="page-14-2"></span>
$$
a(x-z) = \frac{2}{\pi_{\Gamma}} \log|x-z| + C' + O(|x-z|^{-1}) = a(x'-z) + O(m^{-1}). \tag{4.19}
$$

Using the same approach as in [\(3.7\)](#page-7-3), [\(4.19\)](#page-14-2) implies that, for  $A(r + s, N) := D(0, N)$  $D(0, r + s),$ 

<span id="page-15-0"></span>
$$
G_{A(r+s,N)}(x,z) = G_{A(r+s,N)}(x',z) + O(m^{-1}),
$$

which, by letting  $N \to \infty$  and applying the dominated convergence theorem,

<span id="page-15-2"></span><span id="page-15-1"></span>
$$
G_{D(0,r+s)^c}(x,z) = G_{D(0,r+s)^c}(x',z) + O(m^{-1}).
$$
\n(4.20)

Applying [\(4.10\)](#page-13-2) to [\(4.20\)](#page-15-0) yields, for  $z \in D(0, 2r) \setminus D(0, 5r/4)$ ,

$$
G_{D(0,r+s)^c}(x,z) = (1 + O(m^{-1}\log m))G_{D(0,r+s)^c}(x',z). \tag{4.21}
$$

Next, by the symmetry of the Green's function, the strong Markov property at  $T_{D(0,5r/4)c}$ , [\(4.21\)](#page-15-1) for  $z \in D(0, 5r/4) \setminus D(0, r + s)$ , and decomposing, we have

$$
G_{D(0,r+s)^c}(x,z) = G_{D(0,r+s)^c}(z,x)
$$
(4.22)  
=  $\mathbb{E}^z(G_{D(0,r+s)^c}(S_{T_{D(0,5r/4)^c}},x); T_{D(0,5r/4)^c} < T_{D(0,r+s)}),$   
=  $\mathbb{E}^z(G_{D(0,r+s)^c}(S_{T_{D(0,5r/4)^c}},x); T_{D(0,5r/4)^c} < T_{D(0,r+s)}, |S_{T_{D(0,5r/4)^c}}| \le 2r)$   
+  $\mathbb{E}^z(G_{D(0,r+s)^c}(S_{T_{D(0,5r/4)^c}},x); T_{D(0,5r/4)^c} < T_{D(0,r+s)}, |S_{T_{D(0,5r/4)^c}}| > 2r).$ 

By  $(4.21)$  on the first term and  $(2.23)$  on the second term,  $(4.22)$  is bounded above:

$$
G_{D(0,r+s)^c}(z,x) \leq (1 + O(m^{-1} \log m))
$$
\n
$$
\mathbb{E}^z(G_{D(0,r+s)^c}(S_{T_{D(0,5r/4)^c}},x'); T_{D(0,5r/4)^c} < T_{D(0,r+s)})
$$
\n
$$
+ c \log(R) P^z(|S_{T_{D(0,5r/4)^c}}| > 2r).
$$
\n
$$
(4.23)
$$

Applying the first two lines of [\(4.22\)](#page-15-2) again, the first term here is

$$
(1 + O(m^{-1} \log m)) \mathbb{E}^{z}(G_{D(0,r+s)^{c}}(S_{T_{D(0,5r/4)^{c}}}, x') ; T_{D(0,5r/4)^{c}} < T_{D(0,r+s)})
$$
  
= 
$$
(1 + O(m^{-1} \log m)) G_{D(0,r+s)^{c}}(z, x') = (1 + O(m^{-1} \log m)) G_{D(0,r+s)^{c}}(x', z).
$$

A last exit decomposition of  $P^z(|S_{T_{D(0,5r/4)^c}}|>2r)$ , then [\(2.19\)](#page-5-5) and [\(1.5\)](#page-1-2) yield

$$
G_{D(0,r+s)^c}(x, z) \leq (1 + O(m^{-1} \log m))G_{D(0,r+s)^c}(x', z)
$$
  
+  $c \log(R) \sum_{\substack{|y| < 5r/4\\2r < |w|}} G_{D(0,5r/4)}(z, y)p_1(y, w)$   

$$
\leq (1 + O(m^{-1} \log m))G_{D(0,r+s)^c}(x', z) + c \log(R) \log(r)r^{-M+2}.
$$

Since this argument is symmetric in x and x', then we have that for  $z \in D(0, 5r/4)$  $D(0, r + s)$ , and  $c \log R = c(\log 4 + \log m + \log r) = O(\log r)$ ,

<span id="page-15-3"></span>
$$
G_{D(0,r+s)^c}(x,z) = (1 + O(m^{-1}\log m))G_{D(0,r+s)^c}(x',z) + O(r^{-M+2}(\log r)^2). \tag{4.24}
$$

Finally, by  $(2.23)$ , for  $z \in D(0, 2r)^c$ ,  $G_{D(0,r+s)^c}(x, z) = O(\log R)$ . Thus, for  $y \in \partial D(0, r)$ , since  $\sum_{z \in D(0,2r)^c} p_1(z,y) \leq O(r^{-M})$  by symmetry and Markov's inequality,

<span id="page-16-0"></span>
$$
\sum_{z \in D(0,2r)^c} G_{D(0,r+s)^c}(x,z) p_1(z,y) = O(r^{-M} \log r). \tag{4.25}
$$

Combining  $(4.21)$ ,  $(4.24)$ , and  $(4.25)$  bounds the sums in  $(4.18)$  to

$$
H_{D(0,r+s)}(x,y) = (1 + O(m^{-1}\log m))H_{D(0,r+s)}(x',y) + O(r^{-M+2}(\log r)^2). \tag{4.26}
$$

To complete the proof of [\(4.16\)](#page-14-4) for  $x, x' \in \partial D(0, R)_R$ , we must show that, uniformly for  $x \in \partial D(0,R)_R$  and  $y \in \partial D(0,r)_s$ ,

<span id="page-16-3"></span><span id="page-16-2"></span><span id="page-16-1"></span>
$$
r^{-M+2}(\log r)^2 \le c(m^{-1}\log m)H_{D(0,r+s)}(x,y). \tag{4.27}
$$

With  $A_r := D(0, 2r) \setminus D(0, r + (1 + \varepsilon)s)$ , using a last exit decomposition and bounding with the planar version of  $(4.15)$ ,

$$
H_{D(0,r+s)}(x,y) = \sum_{z \in D(0,r+s)^c} G_{D(0,r+s)^c}(x,z) p_1(z,y)
$$
(4.28)  

$$
\geq \sum_{z \in A_r} G_{D(0,r+s)^c}(x,z) p_1(z,y) \geq \frac{c''}{R \log R} \sum_{z \in A_r} p_1(z,y)
$$

for any  $\varepsilon > 0$ . Note that the annulus  $A_r$  contains the disc  $D(v, 2(1+\varepsilon)s)$ , where  $v :=$  $(r+3(1+\varepsilon)s)y/|y|$ . Thus,  $2(1+\varepsilon)s \le |y-v| \le 3(1+\varepsilon)s$ , and  $(1.1)$  (where we consider  $y \in \partial D(v, 2(1+\varepsilon)s)_{(1+\varepsilon)s}),$  and with  $s \leq (\log R)^4 \leq c(\log r)^4$ ,

$$
\sum_{z \in A_r} p_1(z, y) \ge \sum_{z \in D(v, 2(1+\varepsilon)s)} p_1(z, y) \ge c e^{-\beta((1+\varepsilon)s)^{1/4}} \ge c r^{-(1+\varepsilon)^{1/4}\beta}.
$$
 (4.29)

Hence, combining [\(4.28\)](#page-16-1) and [\(4.29\)](#page-16-2), and since  $m \leq \sqrt{r}$ ,  $R < r^2$ , some  $\varepsilon' > 0$ , and  $r^{\beta} > (\log r)^3$  for large enough r,

<span id="page-16-4"></span>
$$
c(m^{-1}\log m)H_{D(0,r+s)}(x,y) \ge \frac{c(m^{-1}\log m)r^{-(1+\varepsilon)^{1/4}\beta}}{R\log R}
$$
  
\n
$$
\ge cr^{-(1+\varepsilon)^{1/4}\beta}(m^{-1}\log m)(mr)^{-1}(2\log r)^{-1}
$$
  
\n
$$
\ge cr^{-1-(1+\varepsilon')\beta}m^{-2}(\log m)(\log r)^{-1}
$$
  
\n
$$
\ge cr^{-2-2\beta}(\log m)(\log r)^{2} \ge cr^{-M+2}(\log r)^{2}, \qquad (4.30)
$$

which proves [\(4.27\)](#page-16-3), and hence [\(4.16\)](#page-14-4), for  $x, x' \in \partial D(0, R)_R$ .

Next we show [\(4.16\)](#page-14-4) for  $x \in D(0, 2R)^c$ . Decompose the hitting distribution on whether or

not we enter  $D(0, 2R)$  via the R-annulus: uniformly for  $x \in D(0, 2R)^c$  and  $y \in \partial D(0, r)_s$ ,

$$
H_{D(0,r+s)}(x,y) = P^x(S_{T_{D(0,r+s)}} = y, T_{\partial D(0,R)_R} > T_{D(0,R)})
$$
  
+  $P^x(S_{T_{D(0,r+s)}} = y, T_{\partial D(0,R)_R} < T_{D(0,R)}).$ 

We can bound the first term by  $(2.13)$ :

$$
P^x(S_{T_{D(0,r+s)}} = y, T_{\partial D(0,R)_R} > T_{D(0,R)}) \le P^x(T_{\partial D(0,R)_R} > T_{D(0,R)})
$$
  
\$\le cR^2(\log R)^2(R^{-M} + R^{-M}) \le cR^{-M+2}(\log R)^2 < cr^{-M+2}(\log r)^2.\$

By the strong Markov property at  $T_{\partial D(0,R)_R}$ , the second term can be bounded, uniformly for  $x' \in \partial D(0,R)_R$ , by  $(4.16)$ :

$$
P^x(S_{T_{D(0,r+s)}} = y, T_{\partial D(0,R)_R} < T_{D(0,R)})
$$
  
=  $\mathbb{E}^x(H_{D(0,r+s)}(S_{T_{\partial D(0,R)_R}}, y), T_{\partial D(0,R)_R} < T_{D(0,R)})$   
 $\leq (1 + O(m^{-1} \log m)) H_{D(0,r+s)}(x', y).$ 

Thus, combining the two, we have for  $x \in D(0, 2R)^c$  and  $x' \in \partial D(0, R)_R$ ,

$$
H_{D(0,r+s)}(x,y) = \left(1 + O\left(m^{-1}\log m\right)\right)H_{D(0,r+s)}(x',y) + O(r^{-M+2}(\log r)^2),\tag{4.31}
$$

which gives [\(4.16\)](#page-14-4) for  $x \in D(0, 2R)^c$  and  $x' \in \partial D(0, R)_R$ . Applying (4.16) again for the same x' and  $x'' \in D(0, 2R)^c$  gives  $(4.16)$  for  $x, x'' \in D(0, 2R)^c$ .

To prove  $(4.17)$  for  $x, x' \in \partial D(0,R)_{\sqrt{R}}$ , decompose  $H_{D(0,r+s)}(x,y)$  over the event  $\{T_{D(0,r+s)} >$  $T_{D(0,4mR)^c}\}$  to get

<span id="page-17-0"></span>
$$
P^x(S_{T_{D(0,r+s)}} = y, T_{D(0,r+s)} < T_{D(0,4mR)^c})
$$
\n
$$
= H_{D(0,r+s)}(x,y) - P^x(S_{T_{D(0,r+s)}} = y, T_{D(0,r+s)} > T_{D(0,4mR)^c}).
$$
\n
$$
(4.32)
$$

By the strong Markov property at  $T_{D(0,4mR)^c}$  and [\(4.16\)](#page-14-4), the last term of [\(4.32\)](#page-17-0) can be further decomposed to

$$
P^x(S_{T_{D(0,r+s)}} = y, T_{D(0,r+s)} > T_{D(0,4mR)^c})
$$
\n
$$
= \mathbb{E}^x(H_{D(0,r+s)}(S_{T_{D(0,4mR)^c}}, y); T_{D(0,r+s)} > T_{D(0,4mR)^c})
$$
\n
$$
= (1 + O(m^{-1} \log m))H_{D(0,r+s)}(x, y)P^x(T_{D(0,r+s)} > T_{D(0,4mR)^c}),
$$
\n(4.33)

which gives us the first equality in  $(4.17)$ . The second follows from  $(2.9)$  and  $(2.1)$ , since,

for  $x, x' \in \partial D(0, R)_{\sqrt{R}}$ , if  $|x| = R$  and  $|x'| = R +$ √ R,

$$
\frac{P^{x'}(T_{D(0,r+s)} < T_{D(0,4mR)^c})}{P^{x}(T_{D(0,r+s)} < T_{D(0,4mR)^c})} = \frac{\log\left(\frac{R+\sqrt{R}}{r+s}\right) + O(r^{-1/4})}{\log\left(\frac{R}{r+s}\right) + O(r^{-1/4})}
$$
\n
$$
= 1 + O\left(\frac{\sqrt{R}}{R\log(\frac{R}{r+s})}\right) + o(r^{-1/4}) = 1 + o(m^{-1}\log m). \quad \Box
$$
\n(4.34)

When attempting to move the planar exterior Harnack inequality to the torus, we run into difficulties in dealing with walks that wander and enter far-off copies of  $D(0, r + s)$ instead of the primary copy. We modify the exterior Harnack inequality for the toral case to fit our requirements.

**Proposition 4.2.** Let  $R = 4mr$  with  $1 \le m = o(r^{1/4})$  and large enough r,  $4mR < K/4$ , and  $s \leq (\log R)^4$ . Then, uniformly for  $\hat{x}, \hat{x}' \in \hat{\pi}_K(\partial D(0,R)_{\sqrt{R}})$  and  $\hat{y} \in \hat{\pi}_K(\partial D(0,r)_s)$ ,

$$
P^{\hat{x}}(\hat{S}_{T_{\hat{\pi}_K(D(0,r+s))}} = \hat{y}; T_{\hat{\pi}_K(D(0,r+s))} < T_{\hat{\pi}_K(D(0,4mR)_K^c)} \tag{4.35}
$$
\n
$$
= \left(1 + O\left(m^{-1}\log m\right)\right) P^{\hat{x}'}(\hat{S}_{T_{\hat{\pi}_K(D(0,r+s))}} = \hat{y}; T_{\hat{\pi}_K(D(0,r+s))} < T_{\hat{\pi}_K(D(0,4mR)_K^c)} \tag{4.35}
$$

Proof For brevity, set

<span id="page-18-1"></span>
$$
D^* := D(0, r + s) \cup D(0, 4mR)^c, \qquad A_p := \{S_{T_{D^*}} = y\},
$$
  

$$
\hat{D}^* := \hat{\pi}_K(D(0, r + s)) \cup \hat{\pi}_K(D(0, 4mR)^c_K), \qquad A_t := \{\hat{S}_{T_{\hat{D}^*}} = \hat{y}\}.
$$

We start our walk at the primary copy  $x$ , consider the planar landing at the primary copy y, and decompose  $P^x(A_t) = \hat{H}_{\hat{D}^*}(x, y)$  along the planar large disc escape time  $T_{D(0,4mR)^c}$ and the toral annulus escape time  $T_{\hat{D}^*}$ :

$$
P^{x}(A_{t}) = P^{x}(A_{t}; T_{\hat{D}^{*}} < T_{D(0, 4mR)^{c}}) + P^{x}(A_{t}; T_{\hat{D}^{*}} \geq T_{D(0, 4mR)^{c}}).
$$
(4.36)

Since  $\hat{\pi}_K^{-1} \hat{D}^* \subset D^*$ ,  $T_{D^*} \leq T_{\hat{D}^*}$  a.s. The first term of [\(4.36\)](#page-18-0) happens in the event  $\{T_{D^*} =$  $T_{\hat{D}^*} = T_{D(0,r+s)}$ , so the entirety of its action before the final step is inside the primary copy of  $D(0, 4mR)$ . Hence,

<span id="page-18-0"></span>
$$
P^{x}(A_t; T_{\hat{D}^*} < T_{D(0, 4mR)^c}) = P^{x}(A_t; T_{D^*} = T_{\hat{D}^*}) = P^{x}(A_p).
$$

Note that  $P^x(A_p)$  is [\(4.17\)](#page-14-5). The second term of [\(4.36\)](#page-18-0) only occurs if a targeted jump lands in a non-primary copy of  $D(0, 4mR) \setminus D(0, r + s)$ . Hence, by [\(2.6\)](#page-4-4),

$$
P^x(A_t; T_{\hat{D}^*} \ge T_{D(0, 4mR)^c}) \le P^x(T_{D(0, 4mR)^c} < T_{\hat{\pi}_K(D(0, 4mR)^c_K)})
$$
  
\$\le O(K^{-M}(mR)^2).

[\(4.36\)](#page-18-0) thus reduces to  $P^x(A_t) = P^x(A_p) + O(K^{-M}(mR)^2)$ , which, by [\(4.17\)](#page-14-5), is

$$
P^{x}(A_t) = (1 + O(m^{-1}\log m))P^{x'}(A_t) + O(K^{-M}(mR)^{2}).
$$
\n(4.37)

Since  $M > 4$ , the error term  $O(K^{-M}(mR)^2) = o(K^{-2-\beta})$  is absorbed via [\(4.30\)](#page-16-4) applied to the  $P^x(A_p)$  term above, with [\(3.16\)](#page-9-2) one "level" up  $(D(0, 4mR)^c$  as the outer bound instead of  $D(0, R)^c$ ,  $D(0, r + s)$  instead of  $D(0, \frac{r}{4m} + s)$ , and  $x, x' \in \partial D(0, R)_{\sqrt{R}}$  instead of  $\partial D(0, r)_r$ , which yields [\(4.35\)](#page-18-1).  $\Box$ 

## References

- <span id="page-19-1"></span>[1] Bass, R. and Rosen, J. (2007). Frequent points for random walks in two dimensions. Electronic Journal of Probability 12 1-46.
- [2] Carlisle, Michael (2012). On the Escape of a Symmetric Random Walk From Two Pieces of a Tripartite Set. arXiv:1209.1761 [math.PR].
- <span id="page-19-2"></span>[3] Carlisle, Michael (2012). On Escaping, Entering, and Visiting Discs of Projections of Planar Symmetric Random Walks on the Lattice Torus. arXiv:1209.2383 [math.PR].
- [4] Dembo, A., Peres, Y., Rosen, J., and Zeitouni, O. (2001). Thick points for planar Brownian motion and random walks in two dimensions. Acta Math. 186 239-270.
- [5] Dembo, A., Peres, Y., Rosen, J., and Zeitouni, O. (2004). Cover times for Brownian motion and random walks in two dimensions. Annals of Mathematics 160 433-464.
- <span id="page-19-0"></span>[6] Dembo, A., Peres, Y., Rosen, J., and Zeitouni, O. (2006). Late points for random walks in two dimensions. Annals of Probability 34 219-263.
- [7] Durrett, R. (2005). Probability: Theory and Examples. 3rd Edition. Thomson Brooks/Cole.
- [8] Erdös, P. and Taylor, S. J. (1960). Some problems concerning the structure of random walk paths. Acta Math. Acad. Sci. Hungar. 11, 137-162.
- [9] Lawler, G. (1991). *Intersections of Random Walks*. Birkhäuser, Boston.
- <span id="page-19-3"></span>[10] Lawler, G. and Limic, V. (2010). Random Walk: A Modern Introduction. Cambridge studies in advanced mathematics, 123. Cambridge University Press, New York.
- [11] Lawler, G. and Polaski, T. (1992). Harnack inequalities and difference estimates for random walks with infinite range. Journal of Theoretical Probability 6 781-802.
- [12] Revuz, Daniel and Marc Yor. (2005). Continuous Martingales and Brownian Motion, 3rd Edition. Springer Berlin Heidelberg New York.
- [13] Rosen, J. (2005). A random walk proof of the Erdös-Taylor Conjecture. Periodica Mathematica Hungarica 50 223-245.
- [14] Spitzer, F. (1976). Principles of Random Walk, Second Edition. Springer, Princeton, NJ.