# On statistical indistinguishability of the complete and incomplete markets

Nikolai Dokuchaev

Department of Mathematics & Statistics, Curtin University, GPO Box U1987, Perth, 6845 Western Australia

Submitted: September 21, 2012. Revised: May 30, 2013

#### Abstract

The possibility of statistical evaluation of the market completeness and incompleteness is investigated for continuous time diffusion stock market models. It is known that the market completeness is not a robust property: small random deviations of the coefficients convert a complete market model into a incomplete one. The paper shows that market incompleteness is also non-robust: small deviations can convert an incomplete model into a complete one. More precisely, it is shown that, for any incomplete market from a wide class of models, there exists a complete market model with arbitrarily close paths of the stock prices and the market parameters. This leads to a counterintuitive conclusion that the incomplete markets are indistinguishable from the complete markets in the terms of the market statistics.

**Key words**: stochastic market, diffusion market, completeness, incompleteness, forecasting, price statistics.

JEL classification: C52, C53,C54 C58

MSC2010 classification: 91G70 91G20 91B84 91B26 62P05

# 1 Introduction

The paper studies continuous stock market models and their statistical analysis. The possibility of statistical evaluation is investigated for the market completeness or incompleteness. These concepts are crucial for the modern mathematical finance. The classical Black-Scholes market model with a non-random volatility is complete, meaning that an arbitrarily claim can be replicated with some self-financing strategies and some initial wealth. For incomplete market models, the option replication is not always possible. A market model with the coefficients that depend on some random factor being independent from the driving Brownian motion is usually incomplete (see, e.g, [20]). Typically, the incomplete market models are used to match the statistical properties of the historical or implied volatility. Currently, there are many well developed models for the volatility (see, e.g., [1]-[2], [4]-[9], [15]-[19], [25]-[28]). It is well known that the market completeness is not a robust property: small random deviations can ruin the completeness and convert a complete model into a incomplete one.

In the present paper, we address these problems again. We consider a class of diffusion market models in the setting where the admissible portfolio strategies can use historical observations collected during some time period before the launching time of the replicating strategy. We found that the market incompleteness is non-robust similarly to the market completeness: small deviations can convert an incomplete market model into a complete one. More precisely, it is shown that, for any incomplete market model from a wide class of models, there exists a complete market model with an arbitrarily close paths of the stock prices and the market parameters (Theorems 1-2). This leads to a conclusion that the incomplete markets are indistinguishable from the complete markets in the terms of the market statistics (Corollary 1).

Let us explain why the non-robustness of the incompleteness established in Theorem 1-2) leads to a conclusion that the incomplete markets are indistinguishable from the complete markets in the terms of the market statistics (Corollary 1). In theory, for continuous time models, the volatility can be estimated without error from the historical prices. However, this would require to know the exact continuous time path of the past prices. This is not feasible because the historical prices are given as time series with rational values. Theorems 1-2 imply that arbitrarily small rounding and time discretization errors may lead to different market models with respect to the completeness and incompleteness. This result is counterintuitive: there is a common perception that the case of random volatility leading to the incompleteness can be spotted from the statistics.

This result does not undermine the importance of the incomplete market models. These models reflect the immanent non-predictability of the real world, in particular, unpredictability of the stock price volatility.

Theorems 1-2 have rather theoretical than practical value since they establish some limits for analysis of market structures based solely on econometrics. These results can be considered as one more illustration of possibility of co-existence of different acceptable models based on the same sets of observations, in the framework of the concept from [23]-[24].

It can be noted that our par follows the general approach to non-robustness of certain market

properties introduced by Guasoniy and Rásonyi in [18], where non-robustness of arbitrage opportunities was established. We study the incompleteness which is a different market property: the incompleteness caused by non-hedgeable randomness of coefficients. The properties considered in this paper and in [18] neither exclude nor imply each other. Furthermore, the arbitrage possibility or completeness are some extreme and rare features. The arbitrage possibility is usually caused by abnormally vanishing volatility or fast growing appreciation rate; the completeness is caused by the predictability and the absence of the noise for the volatility. On the other hand, the incompleteness is rather a typical feature. Since it is easier to believe that a noise contamination of a model removes some rare property, the result of the present paper is more counterintuitive then the result in [18].

# 2 The market model

We consider the so-called diffusion market model, where the market dynamic is described by stochastic differential equations (see, e.g., [20]). In these equations, the randomness is presented in two ways: as the white noise being an external input and as the randomness/uncetainty of the coefficients (market parameters) that represents the following features: (i) correlations with the past; (ii) non-Markov properties, and (iii) unpredictability of the future price distributions.

Assume that we are given a probability space  $(\Omega, \mathcal{F}, \mathbf{P})$ , where  $\Omega$  is a set of elementary events,  $\mathcal{F}$  is a complete  $\sigma$ -algebra of events and  $\mathbf{P}$  is a probability measure.

Let  $\delta > 0$  and  $T \in (0, +\infty)$  be given. Let w(t) be a Brownian motion defined on  $t \in [-\delta, T]$ such that  $w(-\delta) = 0$ .

Consider continuous time diffusion model of a securities market consisting of a risk free bond or bank account with the price B(t), and a risky stock with price S(t),  $t \in [-\delta, T]$ . The prices of the stocks evolve as

$$dS(t) = S(t) \left( a(t)dt + \sigma(t)dw(t) \right), \tag{1}$$

where a(t) is an appreciation rate process,  $\sigma(t)$  is a volatility process. The price of the bond evolves as

$$dB(t) = rB(t)dt,$$

where r > 0 is a short rate that is assumed to be constant.

Let  $\mathcal{M}$  be the class of random processes  $\mu(t) = (a(t), \sigma(t)), t \in [-\delta, T]$ , such that the following holds:

- (i) The processes a(t),  $\sigma(t)$ , and  $\sigma(t)^{-1}$  are bounded on  $[-\delta, T] \times \Omega$ .
- (ii)  $\mu(t)$  is independent from  $w(t_2) w(t_1)$  for all  $t, t_1, t_2$  such that  $t_2 > t_1 \ge t \ge -\delta$ .

In this paper, we consider market models with  $\mu = (a, \sigma) \in \mathcal{M}$ .

Let  $\mathcal{F}_t$  be the filtration generated by the process  $(w(t), \mu(t)), t \geq -\delta$ .

We assume that  $S(-\delta)$  and  $B(-\delta)$  are given non-random variables. In this case, equation (1) has an unique solution S(t) that is adapted to  $\mathcal{F}_t$ ,  $t \in [-\delta, T]$ .

By Girsanov Theorem, for any  $\mu = (a, \sigma) \in \mathcal{M}$ , there exists a set  $\mathcal{P}_{\mu} = \{\mathbf{P}_{\mu}\}$  of probability measures  $\mathbf{P}_{\mu}$  such that the process  $\tilde{S}(t) = e^{-rt}S(t)$  is a martingale in  $t \in [0, T]$  under  $\mathbf{P}_{\mu}$ .

#### Strategies for bond-stock-options market

We describe below the rules for the operations of the agents on the market that define the class of admissible strategies that can be used for replication of contingent claims.

Let X(0) > 0 be the initial wealth at time t = 0, and let X(t) be the wealth at time  $t \in [0, T]$ . We assume that the wealth X(t) at time  $t \in [0, T]$  is

$$X(t) = \beta(t)B(t) + \gamma(t)S(t).$$
(2)

Here  $\beta(t)$  is the quantity of the bond portfolio,  $\gamma(t)$  is the quantity of the stock portfolio. The pair of processes  $(\beta(t), \gamma(t))$  describes the state of the bond-stocks securities portfolio at time  $t \in [0, T]$ . Each of these pairs is called a strategy.

The process  $\tilde{X}(t) \stackrel{\Delta}{=} e^{-rt}X(t)$  is called the discounted wealth, and the process  $\tilde{S}(t) \stackrel{\Delta}{=} e^{-rt}S(t)$  is called the discounted stock price, t > 0.

A pair  $(\beta(\cdot), \gamma(\cdot))$  is said to be an admissible self-financing strategy if the following holds.

- (i) The processes  $\beta(t)$  and  $\gamma(t)$  are progressively measurable with respect to the filtration  $\mathcal{F}_t$ ,  $t \in [0, T]$ .
- (ii) There exists  $\mathbf{P}_{\mu} \in \mathcal{P}_{\mu}$  such that

$$\mathbf{E}_{\mu} \int_{0}^{T} \tilde{S}(t)^{2} \gamma(t)^{2} dt < +\infty,$$

where  $\mathbf{E}_{\mu}$  is the expectation with respect to the probability measure  $\mathbf{P}_{\mu}$ .

(iii) The strategy is self-financing, meaning that

$$dX(t) = \beta(t)dB(t) + \gamma(t)dS(t).$$

For this model, the agents applying admissible self-financing strategies are not supposed to know the future; the strategies have to be adapted to the flow of current market information described by  $\mathcal{F}_t$  for  $t \in [0, T]$ .

The property of self-financing is equivalent to

$$d\tilde{X}(t) = \gamma(t)d\tilde{S}(t). \tag{3}$$

(See, e.g., [20], [10]). It follows that the process  $\gamma(t)$  alone defines the strategy.

#### Market completeness

**Definition 2.1** We say that a market model is complete if, for any p > 0, any random variable  $\xi \in L_{2+p}(\Omega, \mathcal{F}_T, \mathbf{P})$  can be replicated. This means that there exists an  $\mathcal{F}_0$ -measurable initial wealth X(0) and an admissible self-financing strategy  $(\beta(t), \gamma(t)), t \in [0, T]$ , such that the corresponding total terminal wealth X(t) is such that  $X(T) = \xi$  a.s.

To avoid technical difficulties, we consider the case where p > 0 only.

It is well known that the model is complete if the process  $\sigma(t)$  is deterministic. It is also known that a market model is incomplete if  $\sigma(\cdot)|_{[0,T]}$  is random and independent from  $w(\cdot)|_{[0,T]}$ . In addition, a model is incomplete if there is an additional Wiener process  $\widehat{w}(\cdot)$  that is independent from  $w(\cdot)$  and such that the process  $\sigma(\cdot)|_{[0,T]}$  is not independent from  $\widehat{w}(\cdot)|_{[0,T]}$ .

# 3 The main result

Let  $\mathcal{M}^{\perp}$  be the set of all  $\mu \in \mathcal{M}$  that are independent from  $w(\cdot)$ .

**Theorem 1** For any  $\mu \in \mathcal{M}^{\perp}$ , for any  $q \geq 1$ , and for any  $\varepsilon > 0$ , there exists  $\mu_{\varepsilon} \in \mathcal{M}^{\perp}$  such that the corresponding market model is complete and

$$\mathbf{E} \int_{-\delta}^{T} |\mu_{\varepsilon}(t) - \mu(t)|^{q} dt + \mathbf{E} \sup_{t \in [-\delta, T]} |S_{\varepsilon}(t) - S(t)|^{q} < \varepsilon.$$
(4)

Here  $S_{\varepsilon}(t)$  is the stock price for the model defined by  $\mu_{\varepsilon}$ , with  $S_{\varepsilon}(-\delta) = S(-\delta)$ .

We denote by  $|\mu_{\varepsilon}(t) - \mu(t)|$  the Euclidian norm of the vector.

**Corollary 1** The incomplete markets are indistinguishable from the complete markets in the terms of the market statistics.

Proof of Theorem 1. It suffices to consider a market model with  $\mu \in \mathcal{M}$  such that the market is incomplete.

Without a loss of generality, we assume that a(t) and  $\sigma(t)$  are defined for all  $t \in \mathbf{R}$ , and that there exists  $\delta_0 > 0$  such that

$$a(t) = 0, \qquad t \notin [-\delta, T + \delta],$$
  

$$\sigma(t) = 0, \qquad t \notin [-\delta - \delta_0, T + \delta_0], \qquad \sigma(t) = 1, \quad t \in [-\delta - \delta_0, -\delta_0) \cup (T, T + \delta_0].$$

Clearly,  $a(\cdot, \omega) \in L_2(\mathbf{R})$  and  $\sigma(\cdot, \omega) \in L_2(\mathbf{R})$  for all  $\omega \in \Omega$ .

Let  $\kappa_{\varepsilon}(t)$  be defined as  $\kappa_{\varepsilon}(t) = \varepsilon^{-1}\kappa_1(t/\varepsilon)$ , where  $\kappa_1(t)$  is the density for the standard normal distribution N(0, 1). Let  $\sigma_{\varepsilon}(t) = \sigma_{\varepsilon}(t, \omega)$  and  $a_{\varepsilon}(t) = a_{\varepsilon}(t, \omega)$  be the convolutions

$$a_{\varepsilon}(t,\omega) = \int_{-\infty}^{\infty} a(s,\omega)\kappa_{\varepsilon}(t-s)ds, \quad \sigma_{\varepsilon}(t,\omega) = \int_{-\infty}^{\infty} \sigma(s,\omega)\kappa_{\varepsilon}(t-s)ds.$$

One may say that  $\mu_{\varepsilon}(t) = (a_{\varepsilon}(t), \sigma_{\varepsilon}(t))$  is the output of a time invariant smoothing Gaussian filter representing averaging with respect to time. It follows that  $\sup_{t,\omega}(|a_{\varepsilon}(t,\omega)| + |\sigma_{\varepsilon}(t,\omega)| + |\sigma_{\varepsilon}(t,\omega)| + |\sigma_{\varepsilon}(t,\omega)^{-1}|)$  is bounded in  $\varepsilon > 0$ . Hence  $\mu_{\varepsilon} \in \mathcal{M}^{\perp}$ . Note that this filter is not a causal filter since the output is calculated using the future values of the process.

For  $x \in L_2(\mathbf{R}) \cup L_1(\mathbf{R})$ , we denote by X = F x the function  $X : \mathbf{R} \to \mathbf{C}$  defined as the Fourier transform of x;

$$X(\nu) = (Fx)(\nu) = \int_{-\infty}^{\infty} e^{-i\nu t} x(t) dt, \quad \nu \in \mathbf{R}$$

If  $x \in L_2(\mathbf{R})$ , then X is defined as an element of  $L_2(\mathbf{R})$ .

Let

$$\widehat{\kappa}_{\varepsilon} = \digamma \kappa_{\varepsilon}, \qquad \widehat{a} = \digamma a, \qquad \widehat{\sigma} = \digamma \sigma, \quad \widehat{a}_{\varepsilon} = \digamma a_{\varepsilon}, \qquad \widehat{\sigma}_{\varepsilon} = \digamma \sigma_{\varepsilon}.$$

By the property of convolution,

$$\widehat{a}_{arepsilon}(
u) = \widehat{\kappa}_{arepsilon}(
u) \widehat{a}(
u), \quad \widehat{\sigma}_{arepsilon}(
u) = \widehat{\kappa}_{arepsilon}(
u) \widehat{\sigma}(
u), \qquad 
u \in \mathbf{R}$$

By the properties of the Fourier transform of  $\kappa_{\varepsilon}$ , for all  $\nu \in \mathbf{R}$ ,  $\hat{\kappa}_{\varepsilon}(\nu) \to 1$  as  $\varepsilon \to 0$  a.s. Since  $\mu(t)$  has a finite support on  $\mathbf{R}$ , we have that  $\hat{a} \in L_2(\mathbf{R}) \cap L_{\infty}(\mathbf{R})$  and  $\hat{\sigma} \in L_2(\mathbf{R}) \cap L_{\infty}(\mathbf{R})$  a.s., and the corresponding norms are bounded in  $\omega$ . By Lebesgue's Dominated Convergence Theorem,

$$\|\widehat{a}_{\varepsilon}(\cdot,\omega) - \widehat{a}(\cdot,\omega)\|_{L_{2}(\mathbf{R})} \to 0, \quad \|\widehat{\sigma}_{\varepsilon}(\cdot,\omega) - \widehat{\sigma}(\cdot,\omega)\|_{L_{2}(\mathbf{R})} \to 0 \quad \text{as} \quad \varepsilon \to 0 \quad \text{a.s.}$$

It follows that

$$\|a_{\varepsilon}(\cdot,\omega) - a(\cdot,\omega)\|_{L_2(\mathbf{R})} \to 0, \quad \|\sigma_{\varepsilon}(\cdot,\omega) - \sigma(\cdot,\omega)\|_{L_2(\mathbf{R})} \to 0 \quad \text{as} \quad \varepsilon \to 0 \quad \text{a.s.}$$

It follows that there exists a subsequence  $\varepsilon = \varepsilon_i \to 0$  such that  $\mu_{\varepsilon}(t,\omega) \to \mu(t,\omega)$  for a.e.  $t,\omega$  as  $\varepsilon \to 0$ . By Lebesgue's Dominated Convergence Theorem, for any  $q \ge 1$ ,

$$\mathbf{E} \int_{-\delta}^{T} |\mu_{\varepsilon}(t) - \mu(t)|^{q} dt \to 0 \quad \text{as} \quad \varepsilon \to 0.$$

By Theorem II.8.1 from [21], it follows that (4) holds for some  $\varepsilon_i = \varepsilon_i(q, \mu)$ .

To complete the proof, it suffices to show that a market model defined by  $\mu_{\varepsilon}(t) = (a_{\varepsilon}(t), \sigma_{\varepsilon}(t))$ with this  $\varepsilon = \varepsilon_i$  is complete in the sense of Definition 2.1.

Let  $\mathcal{A}_t^{\varepsilon}$  be the filtration generated by the process  $\mu_{\varepsilon}(t)$ ,  $t \geq -\delta$ . By Proposition 3 from [11], the process  $\mu_{\varepsilon}(t,\omega)$  is weakly predictable for any  $\omega \in \Omega$  meaning that, for any  $f \in L_{\infty}(0,T)$ , the integrals  $\int_0^T a_{\varepsilon}(t,\omega)f(t)dt$  and  $\int_0^T \sigma_{\varepsilon}(t,\omega)f(t)dt$  can be found with an arbitrarily small error using the values  $\{\mu_{\varepsilon}(\tau,\omega)\}_{\tau\leq 0}$ . Moreover, by Proposition 1 [11], the processes  $a_{\varepsilon}(t,\omega)$  and  $\sigma_{\varepsilon}(t,\omega)$  are analytic functions in t for all  $\omega \in \Omega$ . It follows that  $\mu_{\varepsilon}(t)$  is a  $\mathcal{A}_0^{\varepsilon}$ -measurable random vector for any  $t \in [0,T]$ .

Let  $\mathcal{F}_t^{\varepsilon}$  be the filtration generated by the process  $(w(t), \mu_{\varepsilon}(t)), t \geq -\delta$ . In other words, this filtration is generated by the observations  $\{w(s), \ \mu_{\varepsilon}(s), \ -\delta < s < t\}$ . We have established that the analytic properties of  $\mu_{\varepsilon}(t)$  imply that the same filtration is generated by the process  $(w(t), \mu_{\varepsilon}(t \wedge 0)), t \geq -\delta$ , i.e., this filtration is generated by the observations  $\{w(s), \ \mu_{\varepsilon}(s \wedge 0), \ s \leq t\}$ .

Let  $\theta_{\varepsilon}(t) = \sigma_{\varepsilon}(t)^{-1}(a_{\varepsilon}(t) - r)$  and let  $w_{\varepsilon}(t) = \int_{0}^{t} \theta_{\varepsilon}(s) ds + w(t) - w(0)$ . Let a probability measure  $\mathbf{P}_{\varepsilon}$  be defined such that

$$\frac{d\mathbf{P}_{\varepsilon}}{d\mathbf{P}} = \exp\left(-\frac{1}{2}\int_0^T \theta_{\varepsilon}(t)^2 dt - \int_0^T \theta_{\varepsilon}(t) dw(t)\right).$$

Let  $\mathbf{E}_{\varepsilon}$  be the corresponding expectation. By Girsanov Theorem applied on the conditional probability space given  $\mathcal{F}_0^{\varepsilon}$ , the process  $w_{\varepsilon}(t)$  is a Wiener process conditionally given  $\mathcal{F}_0^{\varepsilon}$  under the conditional probability measure  $\mathbf{P}_{\varepsilon}(\cdot | \mathcal{F}_0^{\varepsilon})$ .

By the assumptions on  $\xi$ ,  $\mathbf{E}\xi^{2+p} < +\infty$  for some p > 0. Hence

$$\mathbf{E}_{\varepsilon}\xi^2 < +\infty, \quad \mathbf{E}_{\varepsilon}\{\xi^2|\mathcal{F}_0^{\varepsilon}\} < +\infty \quad \text{a.s.}.$$

By the Martingale Representation Theorem applied on the conditional probability space given  $\mathcal{F}_0^{\varepsilon}$ , there exists a process  $g_{\varepsilon} = g_{\varepsilon}(t,\omega)$  such that  $g_{\varepsilon}(t)$  is adapted to  $\mathcal{F}_t^{\varepsilon}$  and

$$\mathbf{E}_{\varepsilon} \int_{0}^{T} g_{\varepsilon}(t)^{2} dt < +\infty, \quad \mathbf{E}_{\varepsilon} \left\{ \int_{0}^{T} g_{\varepsilon}(t)^{2} dt \Big| \mathcal{F}_{0}^{\varepsilon} \right\} < +\infty \quad \text{a.s.},$$

and

$$e^{-rT}\xi = e^{-rT}\mathbf{E}_{\varepsilon}\{\xi|\mathcal{F}_{0}^{\varepsilon}\} + \int_{0}^{T}g_{\varepsilon}(t)dw_{\varepsilon}(t).$$

(See, e.g., Theorem 4.2.4 in [22], p.67). It follows that

$$e^{-rT}\xi = e^{-rT}\mathbf{E}_{\varepsilon}\{\xi|\mathcal{F}_{0}^{\varepsilon}\} + \int_{0}^{T}\gamma_{\varepsilon}(t)dS_{\varepsilon}(t),$$

where  $\gamma_{\varepsilon}(t) = g_{\varepsilon}(t)\sigma_{\varepsilon}(t)^{-1}\tilde{S}_{\varepsilon}(t)^{-1}$ , where  $\tilde{S}_{\varepsilon}(t) = e^{-rt}S_{\varepsilon}(t)$ . By (3), it follows that the selffinancing strategy with the initial wealth  $X_{\varepsilon}(0) = e^{-rT}\mathbf{E}_{\varepsilon}\{\xi|\mathcal{F}_{0}^{\varepsilon}\}$  and with the quantity of shares  $\gamma_{\varepsilon}(t)$  is such that the terminal discounted wealth  $\tilde{X}_{\varepsilon}(T)$  is  $e^{-rT}\xi$ . Hence the terminal wealth for this strategy is  $X_{\varepsilon}(T) = \xi$ . This completes the proof.  $\Box$ 

**Remark 1** In our setting, it is essential that the initial wealth  $X_{\varepsilon}(0)$  for the replicating strategy is  $\mathcal{F}_0^{\varepsilon}$ -measurable, where  $\mathcal{F}_t^{\varepsilon}$  is the filtration describing the information flow for  $t \ge -\delta$ , and that  $\mathcal{F}_0^{\varepsilon}$  is a non-trivial. The information about the history before t = 0 is used for the predicting  $\mu_{\varepsilon}(t)$  for  $t \in [0, T]$ . This is what makes the approximating market model complete.

### 4 An economic interpretation

Theorem 1 implies that the selection of a incomplete model cannot be based solely on the market statistics. This does not undermine a practical use of incomplete market models. Selecting these models, we admit the immanent non-predictability of the real world. For instance, we would rather accept a model with the possibility of the unpredictable jumps for the volatility than a model where these jumps can be predicted, even if the statistical data supports both models equally.

Let us discuss the consequences of co-existing of statistically indistinguishable complete and incomplete markets models.

In the proof of Theorem 1, the process  $\mu(t)$  is approximated by an analytic function  $\mu_{\varepsilon}(t)$  that is used to set a new alterative model. For the new model, the future values  $\mu_{\varepsilon}(t)$  are uniquely defined by their values on the time interval  $[\delta, 0]$ . However, since the new and the old models produce arbitrarily close sets of prices, an observer cannot tell apart these models with certainty, i.e., she cannot tell which model generates the observed data. Effectively, the process  $\mu_{\varepsilon}(t)$  in the new model is not observable for an observer from the old model.

It can be further illustrated as the following. Assume that an option trader has collected the marked data  $t \in [-\delta, 0]$  with the purpose to test the following hypotheses  $H_0$  and  $H_A$  about the stock price evolution:

•  $H_0$ : the values  $\mu(t)|_{t \in [0,T]}$  are not  $\mathcal{F}_0$ -measurable for  $t \in [0,T]$  (i.e., the market is incomplete).

•  $H_A$ : the values  $\mu(t)|_{t \in [0,T]}$  are  $\mathcal{F}_0$ -measurable for  $t \in [0,T]$  (i.e., the market is complete).

It can be noted that we can replace the hypothesis  $H_0$  by a hypothesis assuming a particular stochastic volatility model, such as a Markov chain model, Heston model, etc.

The trader has to calculate at time t = 0 the price of an option expiring at time T; different hypothesis lead to different prices. According to Theorem 1, it is impossible to reject  $H_A$ hypothesis based solely on market statistics collected during the time period  $[-\delta, 0]$ .

Due to rounding errors, the statistical indistinguishability leading to this conclusion cannot be fixed via the sample increasing since the statistics for the incomplete market models can be arbitrarily close to the statistics of the alternative complete models.

It can also noted that, unfortunately, the predictability of  $\mu_{\varepsilon}$  cannot be used for option pricing under the "natural" hypothesis  $H_0$ . The stock prices and market parameters under these two hypotheses are pathwise close; however, their properties are quite different with respect to the predicability. The process  $\mu_{\varepsilon}(t)$  is an output of a non-causal smoothing filters, and its calculation would require the future values of  $\mu(t)$  that are unavailable in practice.

## 5 A more general setting

In the previous section, we considered  $\mu \in \mathcal{M}^{\perp}$ , i.e.,  $\mu$  was assumed to be independent from the driving Wiener process. In fact, this assumption was rather technical; analogs of Theorem 1 can be obtained for more general models where  $\mu(\cdot)$  can depend on  $w(\cdot)$  or  $S(\cdot)$ . Let us give an example.

Let  $y(t) = y(t, \omega)$  be a bounded random process with the values at  $\mathbf{R}^N$ ,  $t \in [-\delta, T]$ , such that y is independent from w.

Let  $R(t) = \log S(t)$ . Let  $\delta_0 > 0$  be given, and let  $\tilde{\mathcal{M}}$  be the class of all  $\mu \in \mathcal{M}$  allowing a closed-loop representation

$$\mu(t) = (a(t), \sigma(t))^{\top} = M(y(t), R(t), \bar{w}(\cdot), t).$$
(5)

In (5), M is a measurable bounded function  $M : \mathbf{R}^N \times \mathbf{R} \times C(-\delta - \delta_0, T + \delta_0) \to \mathbf{R}^2, \ \bar{w}(t) = w((-\delta \lor t) \land t)$ . We assume that M is such that the following holds:

- (i)  $M(y,\rho,\xi,t)$  is continuous in  $y \in \mathbf{R}^N$  uniformly in  $(\rho,\xi,t) \in \mathbf{R} \times C(-\delta \delta_0, T + \delta_0) \times [-\delta \delta_0, T + \delta_0]$ .
- (ii)  $M(y, \rho, \xi, t)$  is Lipschitz in  $\rho \in \mathbf{R}$  uniformly in  $(y, \xi, t) \in \mathbf{R}^N \times C(-\delta \delta_0, T + \delta_0) \times [-\delta \delta_0, T + \delta_0]$ .

The choice of y and M defines  $\mu \in \mathcal{M}$ .

Consider equation for the process  $R(t) = \log S(t)$ 

$$dR(t) = a(t)dt - \frac{\sigma(t)^2}{2}dt + \sigma(t)dw(t), \qquad R(0) = \log S(0).$$
(6)

For any  $\mu \in \tilde{\mathcal{M}}$ , the assumptions on M ensure existence of an unique solution of equation (6). This implies solvability of (1) with  $S(t) = e^{R(t)}$ .

We introduce the market model, admissible strategies, and the definition of completeness such as defined above but with the filtration  $\mathcal{F}_t$  redefined as the filtration generated by the process (w(t), y(t)).

**Theorem 2** For any  $\mu \in \tilde{\mathcal{M}}$ , for any  $q \geq 1$ , and for any  $\varepsilon > 0$ , there exists  $\mu_{\varepsilon} \in \tilde{\mathcal{M}}$  such that the corresponding market model is complete and

$$\mathbf{E} \int_{-\delta}^{T} |\mu_{\varepsilon}(t) - \mu(t)|^{q} dt + \mathbf{E} \sup_{t \in [-\delta, T]} |\log S_{\varepsilon}(t) - \log S(t)|^{q} < \varepsilon.$$

Here  $S_{\varepsilon}(t)$  is the stock price for the model defined by  $\mu_{\varepsilon}$  such that  $S_{\varepsilon}(0) = S(0)$ .

Note that Theorem 2 is a generalization of Theorem 1 since  $\mathcal{M}^{\perp} \in \tilde{\mathcal{M}}$ ; the models in Theorem 1 belong to the class  $\tilde{\mathcal{M}}$  with N = 2 and with  $y(t) = \mu(t) = (a(t), \sigma(t))^{\top}$ .

Proof of Theorem 2. Without a loss of generality, we assume that  $y(t,\omega) = 0$  for  $t \notin [-\delta - \delta_0, T + \delta_0]$  for all  $\omega$ . We use approximating models with  $\mu_{\varepsilon}(t) = M(y_{\varepsilon}(t), R_{\varepsilon}(t), w(\cdot), t)$ , where

$$y_{\varepsilon}(t) = \int_{-\infty}^{\infty} y(s)\kappa_{\varepsilon}(t-s)ds$$

is the output of the Gaussian smoothing filter, and where  $R_{\varepsilon}(t) = \log S_{\varepsilon}(t)$  is the solution of the corresponding equation (6) such that  $R_{\varepsilon}(-\delta) = \log S(-\delta)$ . By the definitions,  $\mu_{\varepsilon} \in \tilde{\mathcal{M}}$ . The market model for  $\mu \in \tilde{\mathcal{M}}$  is arbitrage free and complete if y(t) is a bounded deterministic process for  $t \in [0, T]$ . The rest of the proof follows the proof of Theorem 1. The assumption (iii) on Mensures applicability of Theorem II.8.1 from [21] to equations (6) with  $\mu = \mu_{\varepsilon}$ .  $\Box$ 

## 6 Concluding remarks on forecasting and future development

We outline below some possible modifications and future developments.

(i) Theorems 1-2 allow other modifications. For instance, the statement of Theorem 1 holds with  $\mathcal{M}^{\perp}$  replaced by a class  $\hat{\mathcal{M}}$  of all  $\mu \in \tilde{\mathcal{M}}$  such that (5) holds with M such that  $sM(y, \log s, \xi, t)$  is Lipschitz in  $s \in (0, +\infty)$  uniformly in  $(y, \xi, t) \in \mathbb{R}^N \times C(-\delta - \delta_0, T\delta_0) \times$  $[-\delta - \delta_0, T + d_0].$ 

- (ii) The predictability used in the proof of Theorem 1 can be ensured by many different noncausal time invariant smoothing filters. Instead of a Gaussian filter, we can use an ideal low pass filter or a filter with the exponential rate of energy on higher frequencies  $e^{-|\nu|T}$ . The output of a process transferred with these smoothing filters is a process that, at time t = 0, can be predicted on time interval [0, T] (see [11]).
- (iii) Currently, it is unknown if a Gaussian filter can be approximated by causal smoothing filters. It is known that the approximation by causal smoothing filters is impossible for the ideal low pass filters; the distance of the set of the ideal low-pass filters from the set of all causal filters is positive [3]. On the other hand, it is known that a filter with the exponential energy decay allows arbitrarily close approximation by causal filters [12]. This could lead to application of filters with the exponential energy decay on higher frequencies for forecasting of market parameters and approximation of  $\mu_{\varepsilon}$ . It could be interesting to explore this opportunity.
- (iv) It could be interesting to extend the approach of this paper on discrete market models. For this, discrete time predictability criterions from [13]-[14] could be used.

### Acknowledgment

This work was supported by ARC grant of Australia DP120100928 to the author.

# References

- Aït-Sahalia, Y., and Mykland, P. (2004). Estimating diffusions with discretely and possibly randomly spaced data: A general theory. *Annals of Statistics*, 32, 2186-2222.
- [2] Aït-Sahalia, Y., and Yu, J. (2009). High frequency market microstructure noise estimates and liquidity measures. Annals of Applied Statistics 3, pp. 422–457.
- [3] Almira, J.M. and Romero, A.E. (2008). How distant is the ideal filter of being a causal one? Atlantic Electronic Journal of Mathematics 3 (1) 46–55.
- [4] Andersen, T. G. and Bollerslev, T. (1998). Answering the skeptics: Yes, standard volatility models do provide accurate forecasts. *International Economic Review* 39, pp. 885–905.
- [5] Andersen, T.G., Bollerslev, T., Diebold, F.X., and Ebens, H. (2001). The distribution of realized stock return volatility. *Journal of Financial Economics* 61, pp. 4376.

- [6] Andersen, T.G., Bollerslev, T., Diebold, F.X., and Labys, P. (2003). Modeling and forecasting realized volatility. *Econometrica* 71, pp. 579–625
- [7] Barndorff-Nielsen, O.E., Graversen S.E. and Shephard, N. (2003), Power variation & stochastic volatility: a review and some new results. *Journal of Applied Probability* 41A, 133-143.
- [8] Clark, P.K. (1973). A subordinated stochastic process model with finite variance for speculative prices. *Econometrica* 41, 135–155.
- [9] Cvitanic, J., Liptser, R., and Rozovskii, B. (2006). A filtering approach to tracking volatility from prices observed at random times *Ann. Appl. Probab.* 16, Number 3, 1633-1652.
- [10] Dokuchaev N.G. (2002). Dynamic portfolio strategies: quantitative methods and empirical rules for incomplete information. Kluwer Academic Publishers, Boston.
- [11] Dokuchaev, N. (2010). Predictability on finite horizon for processes with exponential decrease of energy on higher frequencies. *Signal processing* 90, iss. 2, 696–701.
- [12] Dokuchaev, N. (2012). On sub-ideal causal smoothing filters. Signal Processing 92, iss. 1, 219-223.
- [13] Dokuchaev, N. (2012). Predictors for discrete time processes with energy decay on higher frequencies. *IEEE Transactions on Signal Processing* 60, No. 11, 6027-6030.
- [14] Dokuchaev, N. (2012). On predictors for band-limited and high-frequency time series. Signal Processing 92, iss. 10, 2571-2575.
- [15] Elliott, R.J., Hunter, W.C., and Jamieson, B.M. (1998). Drift and volatility estimation in discrete time. Jour. of Economic Dynamics & Control 22, 209-218.
- [16] Fouque, J.-P., Papanicolaou, G., and Sircar, R. (2000). Derivatives in Financial Markets with Stochastic Volatility. Cambridge University Press.
- [17] Frey, R. and Runggaldier, W. (2001). A nonlinear filtering approach to volatility estimation with a view towards high frequency data, (2001). International Journal of Theoretical and Applied Finance 4, 199–210.
- [18] Guasoniy, P. and Rásonyi, M. (2012). Fragility of Arbitrage and Bubbles in Diffusion Models. Working paper, http://ssrn.com/abstract=1856223

- [19] Hull, J., and White, A. (1987). The pricing of options on assets with stochastic volatilities. *Journal of Finance* 42, 381–400.
- [20] Karatzas, I., and Shreve, S.E. (1998). Methods of Mathematical Finance. New York: Springer-Verlag.
- [21] Krylov, N.V. (1980). Controlled diffusion processes. Shpringer-Verlag.
- [22] Lambertone D., Lapeyre, B. (1996). Introduction to Stochastic Calculus Applied to Finance, Chapman & Hall, London.
- [23] Madan D.B. (1983). Inconsistent Theories as Scientific Objectives. *Philosophy of Science*, Vol. 50, No. 3, pp. 453–470.
- [24] Madan, D.B., and Eberlein, E. (2012). Dealing with complex realities in financial modeling. *Current science* **103** (6), 647–649.
- [25] Malliavin, P., and Mancino, M.E. (2002). Fourier Series method for measurement of multivariate volatilities. *Finance & Stochastics* 6, 49-62.
- [26] Mandelbrot, B. (1963). The variation of certain speculative prices. Journal of Business 36, 394–419.
- [27] Merton, R.C. (1980). On estimating the expected return on the market. Journal of Financial Economics 8, 323–361.
- [28] Zhang, L., Mykland, P.A., and Aït-Sahalia, Y. (2005). A tale of two time scales: determining integrated volatility with noisy high frequency data. *Journal of the American Statistical Association* 100, pp. 1394–1411.