

Mosco Type Convergence of Bilinear Forms and Weak Convergence of n -Particle Systems

Jörg-Uwe Löbus
 Matematiska institutionen
 Linköpings universitet
 SE-581 83 Linköping
 Sverige

Abstract It is well known that Mosco (type) convergence is a tool in order to verify weak convergence of finite dimensional distributions of sequences of stochastic processes. In the present paper we are concerned with the concept of Mosco type convergence for non-symmetric stochastic processes and, in particular, n -particle systems in order to establish relative compactness.

AMS subject classification (2010) primary 47D07, secondary 60K35, 60J35

Keywords Mosco type convergence, n -particle systems, weak convergence

1 Introduction

To show weak convergence of a sequences of stochastic processes one has to proceed in two basic steps. These are on the one hand, proving relative compactness of the sequences of processes and, on the other hand, showing weak convergence of the finite dimensional distributions of the sequence of stochastic processes.

Mosco convergence has been used in order to prove weak convergence of finite dimensional distributions of sequences of stochastic processes corresponding to symmetric Dirichlet forms, cf. for example [1], [3], [6], [9], [10, 11].

In addition, the paper [21] demonstrates in which way Mosco convergence can be used in order to verify relative compactness of a sequences of stochastic processes. In fact, Mosco convergence and additional properties or appropriate additional conditions on the sequences of processes provide the convergence of a certain sequence of associated capacities of the form $\mathbb{E}e^{-\beta\tau_n}$, $n \in \mathbb{N}$. Here, τ_n is a sequence of certain first exit times and $\beta > 0$. The convergence of the sequence $\mathbb{E}e^{-\beta\tau_n}$, $n \in \mathbb{N}$, is then sufficient for relative compactness. This idea has been adapted to particle systems in [13] and [14] and will also be developed further in Section 3 of the present paper.

We would also like to refer to two more motivations for this paper. During the last decade one may have observed an increasing interest in Mosco convergence relative to Dirichlet forms with changing reference measures or, more general, on sequences of Hilbert spaces, see [5], [7], [8], [20], [23]. Most fundamental in this sense is [12]. In the particular case of sequences of L^2 spaces we would like to refer to [14] which has a documented history beginning 2005.

Initiated by the two established generalizations of Dirichlet forms to the non-symmetric case, namely [17] and [22], also Mosco (type) convergence for non-symmetric Dirichlet forms

has been investigated, cf. [23] and [4]. The paper [14] provides a framework for non-symmetric bilinear forms where neither the Dirichlet property nor closability is necessary. This allows to treat stochastic processes and particle systems without any connection to Dirichlet form theory.

The present paper follows this motivation. However the framework here is more sophisticated. As an application, the particle system considered in [15] doesn't seem to be compatible with [17] or [22]. Moreover the limiting initial distribution is no longer concentrated on one single state as in [14]. However it fits the theory presented in this article.

Looking, for example, at the papers [4], [5], [12], [20], [23] one can conclude that different classes of applications require different generalizations or alternations of Mosco convergence. The present paper together with [15] follows this attitude.

1.1 Basic Definitions and Technical Issues

In order to introduce the basic setting, let ν be a probability measure on a measurable space (E, \mathcal{B}) , and let $(T_t)_{t \geq 0}$ be a strongly continuous contraction semigroup of linear operators on $L^2(E, \nu)$. Suppose that $(T_t)_{t \geq 0}$ is associated with a transition probability function $P(t, x, B)$, $t \geq 0$, $x \in E$, $B \in \mathcal{B}$, i. e., $T_t f = \int f(y) P(t, \cdot, dy)$, $t \geq 0$, $f \in L^2(E, \nu)$. Assume, furthermore, that $P(t, \cdot, E) = 1$ ν -a.e., $t \geq 0$. We mention that contractivity of $(T_t)_{t \geq 0}$ on $L^2(E, \nu)$ gives $\int (P(t, x, B))^2 \nu(dx) \leq \nu(B)$, $B \in \mathcal{B}$, which says that $\nu(B) = 0$ implies $P(t, x, B) = 0$ for ν -a.e. $x \in E$.

If, as in Subsection 2.3, $(T_t)_{t \geq 0}$ is no longer contractive on $L^2(E, \nu)$ we suppose this implication to ensure well-definiteness of $(T_t)_{t \geq 0}$.

Denoting by $(A, D(A))$ the generator of $(T_t)_{t \geq 0}$ and by $\langle \cdot, \cdot \rangle$ the inner product in $L^2(E, \nu)$, we introduce now the class of bilinear forms S we are interested in. Define

$$D(S) := \left\{ u \in L^2(E, \nu) : \lim_{t \rightarrow 0} \left\langle \frac{1}{t}(u - T_t u), v \right\rangle \text{ exists for all } v \in L^2(E, \nu) \right\}$$

and

$$S(u, v) := \lim_{t \rightarrow 0} \left\langle \frac{1}{t}(u - T_t u), v \right\rangle, \quad u \in D(S), \quad v \in L^2(E, \nu).$$

We have $D(A) = D(S)$ according to [19], Section 2.1 and

$$S(u, v) = -\langle Au, v \rangle, \quad u \in D(A), \quad v \in L^2(E, \nu).$$

In this sense we would like to understand the term *bilinear form*. However, as it is customary for Mosco (type) convergence, we also set $S(u, v) := \infty$ if $u \in L^2(E, \nu) \setminus D(S)$ and $v \in L^2(E, \nu)$. We emphasize that this definition of bilinear forms S is adjusted to the Mosco type convergence of non-symmetric forms in Subsection 2.2 and, moreover, of non-positive, non-symmetric forms in Subsection 2.3. The latter situation appears for example in the application of [15].

Let $(G_\beta)_{\beta > 0}$ be the resolvent associated with S , i. e., $G_\beta = (\beta - A)^{-1}$, $\beta > 0$. Using contractivity of the semigroup $(T_t)_{t \geq 0}$ in $L^2(E, \nu)$ and

$$\langle T_t u, u \rangle^2 \leq \langle T_t u, T_t u \rangle \langle u, u \rangle$$

one shows positivity of the form S , that is $S(u, u) \geq 0$ for all $u \in D(S)$. This observation is crucial for the whole concept of Mosco type convergence of sequences S_n of forms on sequences of spaces $L^2(E_n, \nu_n)$ to a limiting form S on $L^2(E, \nu)$ as $n \rightarrow \infty$. However, we also develop a framework of Mosco type convergence of sequences of forms when contractivity is replaced by a technical condition on $A'_n \mathbb{1}$, $n \in \mathbb{N}$, and $A' \mathbb{1}$ where $\mathbb{1}$ is the constant function taking the value one and the $'$ refers to the dual generator.

The first question of interest is on the definition of such a bilinear form. In fact, we construct the weak generator of the semigroup $(T_t)_{t \geq 0}$, the second entry v in $S(u, v)$ is in this sense just a test function. This definition of a bilinear form is far away from classical Dirichlet form theory which includes the notion of Mosco convergence. One more problem arises with the definition of the bilinear form, namely the appropriate notion of Mosco type convergence. We recall that the literature suggest several alternations of the classical one by U. Mosco in [18], cf. [4], [5], and [23].

The results obtained in Sections 6 and 7 of [15] give rise to state that our approach to bilinear forms and Mosco type convergence is beneficial relative to the particle system we investigate there and similar ones. We also want to refer to a discussion on choosing the appropriate definition of bilinear forms relative to the mathematical situation, given in [14], Subsections 2.1 and 2.3.

The next problem one may come across is the existence of $A'_n \mathbb{1}$, $n \in \mathbb{N}$, and $A' \mathbb{1}$ in the case of non-positive bilinear forms. In particular, $A'_n \mathbb{1}$, $n \in \mathbb{N}$, and $A' \mathbb{1}$ should display properties which are useful to show Mosco type convergence, cf. Subsection 2.3 below. This is a purely mathematical issue. In the application of [15], a Fleming-Viot type particle system, this issue restricts the initial configurations of the particles that can be investigated in terms of Mosco type convergence.

We conclude the introduction with a remark on the notation in the paper. The greek letter ν comes always in bold. This letter is exclusively used to denote probability measures over spaces of probability measures, the states of measure valued stochastic processes. Those states of stochastic processes are denoted using the greek letter μ , non-bold.

2 Mosco Type Convergence

In this section, we are extending the framework of [14] in several ways. In Subsection 2.1, we develop the concept of convergence in sequences of L^2 -spaces. This is necessary in order to establish the Mosco type convergence for non-symmetric forms on sequences of L^2 -spaces presented in Subsection 2.2. In Subsection 2.3, we are then able to handle convergence of non-symmetric non-positive forms, the situation we have to face in the application in Section 4.

Again, we want to refer to related research carried out in K. Kuwae, T. Shioya [12] and compared with ours in [14], Subsection 3.2.

2.1 Analysis on Sequences of L^2 -Spaces

Let ν_n , $n \in \mathbb{Z}_+ := \{0, 1, 2, \dots\}$, be mutually orthogonal probability measures on (E, \mathcal{B}) . To ease the notation and to stress its special role, we will mostly use the symbol ν instead of ν_0 . Suppose that ν is a measure with countable base on (E, \mathcal{B}) . In addition, assume that there are mutually exclusive subsets E_n , $n \in \mathbb{Z}_+$, of E such that $\nu_n(E \setminus E_n) = 0$. Let α_n ,

$n \in \mathbb{Z}_+$, be a sequence of positive numbers with $\sum_{n=0}^{\infty} \alpha_n = 1$. Define $\mathbb{M} := \sum_{n=0}^{\infty} \alpha_n \nu_n$. We say that $u \in \bigcap_{n \in \mathbb{Z}_+} L^2(E, \nu_n)$ if u is an equivalence class consisting of all everywhere defined \mathcal{B} -measurable functions satisfying $f_1 = f_2$ \mathbb{M} -a.e. if $f_1, f_2 \in u$ and $\int u^2 d\nu_n < \infty$, $n \in \mathbb{Z}_+$. Let $\langle \cdot, \cdot \rangle_n$ denote the inner product in $L^2(E, \nu_n)$, $n \in \mathbb{N}$, and let $\langle \cdot, \cdot \rangle$ denote the inner product in $L^2(E, \nu)$. Introduce

$$\mathcal{D} := \left\{ \varphi \in \bigcap_{n \in \mathbb{Z}_+} L^2(E, \nu_n) : \langle \varphi, \varphi \rangle_n \xrightarrow{n \rightarrow \infty} \langle \varphi, \varphi \rangle \right\}.$$

Throughout this paper we suppose that there exists a linear subset \mathcal{F} of \mathcal{D} which is dense in $L^2(E, \nu)$ and let \mathcal{C} denote the set of all functions $\varphi \in \mathcal{D}$ satisfying the following conditions:

- (c1) For each $\varphi \in \mathcal{C}$, there exists a *representing sequence* $\varphi_n \in \mathcal{F}$, $n \in \mathbb{N}$, such that $\varphi = \varphi_n$, ν_n -a.e., $n \in \mathbb{N}$.
- (c2) $\langle \varphi, \psi \rangle_n \xrightarrow{n \rightarrow \infty} \langle \varphi, \psi \rangle$ for all $\psi \in \mathcal{F}$.

Introduce

$$\mathcal{V} := \left\{ \psi \in \bigcap_{n \in \mathbb{Z}_+} L^2(E, \nu_n) : \sigma \in \mathcal{D}, \tau \in \mathcal{F} \text{ imply } \sigma\psi \in \mathcal{D}, \tau\psi \in \mathcal{F} \right\}.$$

Lemma 2.1 (a) $\mathcal{F} \subseteq \mathcal{C}$.

(b) The set \mathcal{C} is linear.

(c) The set \mathcal{C} is dense in $L^2(E, \nu)$.

(d) Let $\varphi, \psi \in \mathcal{C}$. We have $\langle \varphi, \psi \rangle_n \xrightarrow{n \rightarrow \infty} \langle \varphi, \psi \rangle$.

(e) Let $\psi \in \mathcal{V}$. Then, for all $\varphi \in \mathcal{C}$, we have $\varphi\psi \in \mathcal{C}$.

Proof. (a) (c1) is trivial and (c2) follows from linearity of \mathcal{F} .

(b) Let $\varepsilon > 0$ and let $\varphi, \psi \in \mathcal{C}$ with representing sequences $\varphi_n \in \mathcal{F}$ and $\psi_n \in \mathcal{F}$, $n \in \mathbb{N}$, cf. condition (c2). Since (c1) and (c2) are obvious for $\varphi + \psi$, it remains to show that $\varphi + \psi \in \mathcal{D}$. We can choose $n_0 \in \mathbb{N}$ and $\tilde{\psi} \in \mathcal{F}$ such that, for all $n > n_0$,

$$\begin{aligned} \langle \psi - \tilde{\psi}, \psi - \tilde{\psi} \rangle &< \varepsilon && \text{since } \mathcal{F} \subseteq L^2(E, \nu), \text{ densely,} \\ |\langle \varphi, \varphi \rangle_n - \langle \varphi, \varphi \rangle| &< \varepsilon && \text{since } \varphi \in \mathcal{C} \subseteq \mathcal{D}, \\ |\langle \psi, \psi \rangle_n - \langle \psi, \psi \rangle| &< \varepsilon && \text{since } \psi \in \mathcal{C} \subseteq \mathcal{D}, \\ |\langle \tilde{\psi}, \tilde{\psi} \rangle_n - \langle \tilde{\psi}, \tilde{\psi} \rangle| &< \varepsilon && \text{since } \tilde{\psi} \in \mathcal{F} \subseteq \mathcal{C} \subseteq \mathcal{D}, \\ |\langle \varphi, \tilde{\psi} \rangle_n - \langle \varphi, \tilde{\psi} \rangle| &< \varepsilon, && \text{cf. (c2),} \\ |\langle \psi, \tilde{\psi} \rangle_n - \langle \psi, \tilde{\psi} \rangle| &< \varepsilon, && \text{cf. (c2).} \end{aligned} \tag{2.1}$$

Therefore, we have

$$\begin{aligned} \langle \psi - \tilde{\psi}, \psi - \tilde{\psi} \rangle_n &\leq \langle \psi - \tilde{\psi}, \psi - \tilde{\psi} \rangle \\ &\quad + |\langle \psi, \psi \rangle_n - \langle \psi, \psi \rangle| + |\langle \tilde{\psi}, \tilde{\psi} \rangle_n - \langle \tilde{\psi}, \tilde{\psi} \rangle| + 2|\langle \psi, \tilde{\psi} \rangle_n - \langle \psi, \tilde{\psi} \rangle| \\ &< 5\varepsilon \end{aligned}$$

which implies

$$\begin{aligned} |\langle \varphi, \psi \rangle_n - \langle \varphi, \tilde{\psi} \rangle_n| &\leq \langle \varphi, \varphi \rangle_n^{1/2} \langle \psi - \tilde{\psi}, \psi - \tilde{\psi} \rangle_n^{1/2} \\ &< \sqrt{5} \sup_{n \in \mathbb{N}} \langle \varphi, \varphi \rangle_n^{1/2} \cdot \varepsilon^{1/2} \end{aligned}$$

and

$$\begin{aligned} |\langle \varphi, \psi \rangle_n - \langle \varphi, \psi \rangle| &\leq |\langle \varphi, \psi \rangle_n - \langle \varphi, \tilde{\psi} \rangle_n| + |\langle \varphi, \tilde{\psi} \rangle_n - \langle \varphi, \tilde{\psi} \rangle| + |\langle \varphi, \tilde{\psi} \rangle - \langle \varphi, \psi \rangle| \\ &< \varepsilon + \left(\sqrt{5} \sup_{n \in \mathbb{N}} \langle \varphi, \varphi \rangle_n^{1/2} + \langle \varphi, \varphi \rangle^{1/2} \right) \cdot \varepsilon^{1/2}. \end{aligned} \quad (2.2)$$

It follows now from (2.1) and (2.2) that

$$\begin{aligned} |\langle \varphi + \psi, \varphi + \psi \rangle_n - \langle \varphi + \psi, \varphi + \psi \rangle| &\leq |\langle \varphi, \varphi \rangle_n - \langle \varphi, \varphi \rangle| + |\langle \psi, \psi \rangle_n - \langle \psi, \psi \rangle| + 2|\langle \varphi, \psi \rangle_n - \langle \varphi, \psi \rangle| \\ &< 4\varepsilon + \left(2\sqrt{5} \sup_{n \in \mathbb{N}} \langle \varphi, \varphi \rangle_n^{1/2} + 2\langle \varphi, \varphi \rangle^{1/2} \right) \cdot \varepsilon^{1/2}. \end{aligned}$$

(c) This is a consequence of (a) and the fact that \mathcal{F} is dense in $L^2(E, \nu)$.

(d) This follows from (2.2).

(e) Let $\varphi \in \mathcal{C}$ and $\psi \in \mathcal{V}$. By hypothesis, we have $\varphi\psi \in \mathcal{D}$. Let $\varphi_n \in \mathcal{F}$, $n \in \mathbb{N}$, be the representing sequence of φ , cf. (c1). Then $\varphi_n\psi \in \mathcal{F}$, $n \in \mathbb{N}$, by hypothesis. In other words, $\varphi_n\psi \in \mathcal{F}$, $n \in \mathbb{N}$, is the representing sequence of $\varphi\psi$, i. e., we have (c1) for $\varphi\psi$. Furthermore, for all $\rho \in \mathcal{F}$, we have $\psi\rho \in \mathcal{F}$ by hypothesis and, by (c2), $\langle \varphi, \psi\rho \rangle_n \xrightarrow{n \rightarrow \infty} \langle \varphi, \psi\rho \rangle$. Thus, $\langle \varphi\psi, \rho \rangle_n \xrightarrow{n \rightarrow \infty} \langle \varphi\psi, \rho \rangle$. Thus, we have (c2) for $\varphi\psi$. \square

Definition 2.2 (a) A sequence $\varphi_n \in \mathcal{C}$, $n \in \mathbb{N}$, is said to be *w-convergent* to $\varphi \in L^2(E, \nu)$ as $n \rightarrow \infty$ (in symbols $\varphi_n \xrightarrow{n \rightarrow \infty} \varphi$) if

(i) $\langle \varphi_n, \psi \rangle_n \xrightarrow{n \rightarrow \infty} \langle \varphi, \psi \rangle$ for all $\psi \in \mathcal{C}$.

(b) A sequence $\psi_n \in \mathcal{C}$, $n \in \mathbb{N}$, is said to be *s-convergent* to $\psi \in L^2(E, \nu)$ as $n \rightarrow \infty$ (in symbols $\psi_n \xrightarrow{n \rightarrow \infty} \psi$) if

(i) ψ_n *w-converges* to ψ as $n \rightarrow \infty$ and

(ii) $\langle \psi_n, \psi_n \rangle_n \xrightarrow{n \rightarrow \infty} \langle \psi, \psi \rangle$.

(c) Speaking of *w-convergence* or *s-convergence* of subsequences $\varphi_{n_k} \in \mathcal{C}$ or $\psi_{n_k} \in \mathcal{C}$, respectively, will mean that in (a) or (b) the index $n \in \mathbb{N}$ is replaced with $n_k \in \mathbb{N}$.

Remarks (1) Let $\psi_n \in \mathcal{C}$, $n \in \mathbb{N}$, *w-converge* to $\psi \in \mathcal{C}$. According to the definition of \mathcal{C} , that ψ_n *s-converges* to $\psi \in \mathcal{C}$ is equivalent to

$$\begin{aligned} \langle \psi_n - \psi, \psi_n - \psi \rangle_n &= \langle \psi_n, \psi_n \rangle_n - 2\langle \psi_n, \psi \rangle_n + \langle \psi, \psi \rangle_n \\ \xrightarrow{n \rightarrow \infty} &0. \end{aligned}$$

(2) Let $\psi \in \mathcal{C}$. It follows from Lemma 2.1 (d), that $\psi_n := \psi$, $n \in \mathbb{N}$, *s-converges* to ψ .

Proposition 2.3 (a) Let $\varphi_n \in \mathcal{C}$, $n \in \mathbb{N}$, be a sequence w -convergent to $\varphi \in L^2(E, \nu)$ as $n \rightarrow \infty$. Then $\langle \varphi_n, \varphi_n \rangle_n$, $n \in \mathbb{N}$, is bounded.

(b) Let $\varphi_n \in \mathcal{C}$, $n \in \mathbb{N}$, be a sequence such that $\langle \varphi_n, \varphi_n \rangle_n$ is bounded. Then there exists a subsequence $\varphi_{n_k} \in \mathcal{C}$, $k \in \mathbb{N}$, w -convergent to some $\varphi \in L^2(E, \nu)$ as $k \rightarrow \infty$.

(c) Let $\varphi_n \in \mathcal{C}$, $n \in \mathbb{N}$, be a sequence w -convergent to $\varphi \in L^2(E, \nu)$ and let $\psi_n \in \mathcal{C}$, $n \in \mathbb{N}$, be a sequence that s -converges to $\psi \in L^2(E, \nu)$ as $n \rightarrow \infty$. Then $\langle \varphi_n, \psi_n \rangle_n \xrightarrow{n \rightarrow \infty} \langle \varphi, \psi \rangle$.

(d) Let $\varphi_n \in \mathcal{C}$, $n \in \mathbb{N}$, be a sequence w -convergent to $\varphi \in \mathcal{C}$ as $n \rightarrow \infty$ and let $\psi_n \in \mathcal{C}$, $n \in \mathbb{N}$, be a sequence w -convergent to $\psi \in \mathcal{C}$ as $n \rightarrow \infty$. Suppose

$$\langle \varphi_n - \varphi, \psi_n - \psi \rangle_n \geq 0, \quad n \in \mathbb{N}. \quad (2.3)$$

Then

$$\liminf_{n \rightarrow \infty} \langle \varphi_n, \psi_n \rangle_n \geq \langle \varphi, \psi \rangle.$$

(e) Let $\varphi_n \in \mathcal{C}$, $n \in \mathbb{N}$, be a sequence w -convergent to $\varphi \in L^2(E, \nu)$. Then we have

$$\liminf_{n \rightarrow \infty} \langle \varphi_n, \varphi_n \rangle_n \geq \langle \varphi, \varphi \rangle.$$

In the following, let $\psi_n \in \mathcal{V}$, $n \in \mathbb{N}$, let $\psi \in L^\infty(E, \nu)$, and assume $\psi_n \rho \xrightarrow{n \rightarrow \infty} \psi \rho$ for all $\rho \in \mathcal{C}$.

(f) Let $\mathcal{C} \ni \varphi_n \xrightarrow{n \rightarrow \infty} \varphi \in L^2(E, \nu)$. Then $\varphi_n \psi_n \xrightarrow{n \rightarrow \infty} \varphi \psi$.

(g) Let $\mathcal{C} \ni \varphi_n \xrightarrow{n \rightarrow \infty} \varphi \in L^2(E, \nu)$. Then $\varphi_n \psi_n \xrightarrow{n \rightarrow \infty} \varphi \psi$.

Proof. For the proofs of (a) through (c), we will refer to [14].

(a) and (c) The situation in Section 3 of [14] is compatible with the setting here. In particular, there it is assumed that E is a metric space and Definition 3.1 yields a linear set \mathcal{C} . Replacing in the statement of [14], Lemma 3.2 (b), $C_b(E)$ by \mathcal{F} and using the above properties of \mathcal{F} the assertion of [14], Lemma 3.2 (b), becomes obvious. The proofs of Proposition 3.3 (a) and (b) can now be followed word for word with \mathcal{F} instead of $C_b(E)$. In this way, we have verified (a) and (c) of the present proposition.

(b) This has been demonstrated in [14], proof of Proposition 2.3 (a).

(d) Because of $\varphi_n \xrightarrow{n \rightarrow \infty} \varphi$, $\psi_n \xrightarrow{n \rightarrow \infty} \psi$, and $\varphi, \psi \in \mathcal{C}$, we have $\langle \varphi_n, \psi \rangle_n \xrightarrow{n \rightarrow \infty} \langle \varphi, \psi \rangle$, $\langle \varphi, \psi_n \rangle_n \xrightarrow{n \rightarrow \infty} \langle \varphi, \psi \rangle$, and $\langle \varphi, \psi \rangle_n \xrightarrow{n \rightarrow \infty} \langle \varphi, \psi \rangle$. The lemma is now a consequence of hypothesis (2.3) and

$$\langle \varphi_n, \psi_n \rangle_n = \langle \varphi_n, \psi \rangle_n + \langle \varphi, \psi_n \rangle_n - \langle \varphi, \psi \rangle_n + \langle \varphi_n - \varphi, \psi_n - \psi \rangle_n.$$

(e) Let $\varepsilon > 0$ and $\tilde{\varphi} \in \mathcal{C}$ such that $\langle \tilde{\varphi} - \varphi, \tilde{\varphi} - \varphi \rangle < \varepsilon$, cf. Lemma 2.1 (c). Since φ_n , $n \in \mathbb{N}$, w -converges to $\varphi \in L^2(E, \nu)$, it follows that

$$\begin{aligned} \langle \varphi_n, \varphi_n \rangle_n - \langle \varphi_n - \tilde{\varphi}, \varphi_n - \tilde{\varphi} \rangle_n &= \langle \varphi_n, \tilde{\varphi} \rangle_n + \langle \tilde{\varphi}, \varphi_n \rangle_n - \langle \tilde{\varphi}, \tilde{\varphi} \rangle_n \\ &\xrightarrow{n \rightarrow \infty} \langle \varphi, \tilde{\varphi} \rangle + \langle \tilde{\varphi}, \varphi \rangle - \langle \tilde{\varphi}, \tilde{\varphi} \rangle \\ &= \langle \varphi, \varphi \rangle - \langle \varphi - \tilde{\varphi}, \varphi - \tilde{\varphi} \rangle. \end{aligned}$$

We have $\langle \varphi - \tilde{\varphi}, \varphi - \tilde{\varphi} \rangle \leq \varepsilon$ from which we get $\liminf_{n \rightarrow \infty} \langle \varphi_n, \varphi_n \rangle_n \geq \langle \varphi, \varphi \rangle$.

(f) Let $\rho \in \mathcal{C}$. According to Lemma 2.1 (e) and hypotheses, we have $\varphi_n \psi_n \in \mathcal{C}$, $n \in \mathbb{N}$, and $\mathcal{C} \ni \psi_n \rho \xrightarrow[n \rightarrow \infty]{s} \psi \rho \in L^2(E, \nu)$. From part (c), we obtain

$$\langle \varphi_n \psi_n, \rho \rangle_n = \langle \varphi_n, \psi_n \rho \rangle_n \xrightarrow[n \rightarrow \infty]{} \langle \varphi, \psi \rho \rangle = \langle \varphi \psi, \rho \rangle.$$

(g) It remains to note that by parts (c) and (f), we have $\varphi_n \psi_n^2 \xrightarrow[n \rightarrow \infty]{w} \varphi \psi^2$ which yields by part (c)

$$\begin{aligned} \langle \varphi_n \psi_n, \varphi_n \psi_n \rangle &= \langle \varphi \psi_n, \varphi_n \psi_n \rangle_n + \langle \varphi_n - \varphi, \varphi_n \psi_n^2 \rangle_n \\ &\xrightarrow[n \rightarrow \infty]{} \langle \varphi \psi, \varphi \psi \rangle. \end{aligned}$$

□

2.2 Mosco Type Convergence of Non-Symmetric Forms on Sequences of L^2 -Spaces

For every $n \in \mathbb{N}$, let $(T_{n,t})_{t \geq 0}$ be a strongly continuous contraction semigroup in $L^2(E, \nu_n)$ and let $(T_t)_{t \geq 0}$ be a strongly continuous contraction semigroup in $L^2(E, \nu)$, all in the sense of Section 1. Denote by $S_n, A_n, (G_{n,\beta})_{\beta > 0}$ the bilinear form in the sense of Section 1, the generator, and the family of resolvents associated with $(T_{n,t})_{t \geq 0}$, $n \in \mathbb{N}$. Similarly, let S, A , and $(G_\beta)_{\beta > 0}$ the bilinear form, the generator, and the family of resolvents associated with $(T_t)_{t \geq 0}$. For $\beta > 0$, $n \in \mathbb{N}$, $\varphi_n \in D(S_n)$, $\psi_n \in L^2(E, \nu_n)$, set $S_{n,\beta}(\varphi_n, \psi_n) := \beta \langle \varphi_n, \psi_n \rangle_n + S_n(\varphi_n, \psi_n)$, and for $\varphi \in D(S)$, $\psi \in L^2(E, \nu)$, define $S_\beta(\varphi, \psi) := \beta \langle \varphi, \psi \rangle + S(\varphi, \psi)$.

Definition 2.4 We say that S_n , $n \in \mathbb{N}$, *pre-converges* to S if

- (i) For every $\varphi \in L^2(E, \nu)$ and every subsequence $\varphi_{n_k} \in D(S_{n_k}) \cap \mathcal{C}$, $k \in \mathbb{N}$, w -converging to φ such that $\sup_{k \in \mathbb{N}} \langle A_{n_k} \varphi_{n_k}, A_{n_k} \varphi_{n_k} \rangle_{n_k} < \infty$, we have

$$S(\varphi, \varphi) \leq \liminf_{k \rightarrow \infty} S_{n_k}(\varphi_{n_k}, \varphi_{n_k}).$$

- (ii) For every $\psi \in D(S)$, there exists a sequence $\psi_n \in D(S_n) \cap \mathcal{C}$, $n \in \mathbb{N}$, s -converging to ψ such that $\sup_{n \in \mathbb{N}} \langle A_n \psi_n, A_n \psi_n \rangle_n < \infty$ and

$$\limsup_{n \rightarrow \infty} S_n(\psi_n, \psi_n) \leq S(\psi, \psi).$$

Lemma 2.5 Let S_n , $n \in \mathbb{N}$, be a sequence of bilinear forms pre-convergent to S . In addition, let $w_n \in D(S_n) \cap \mathcal{C}$, $n \in \mathbb{N}$, be a sequence w -converging to some $w \in L^2(E, \nu)$ and let $v_n \in D(S_n) \cap \mathcal{C}$, $n \in \mathbb{N}$, be a sequence s -converging to $v \in D(S)$ in the sense of condition (ii) of Definition 2.4. Suppose $\sup_{n \in \mathbb{N}} \langle A_n w_n, A_n w_n \rangle_n < \infty$.

(a) The limit $\lim_{n \rightarrow \infty} (S_n(v_n, w_n) + S_n(w_n, v_n))$ exists and we have

$$\lim_{n \rightarrow \infty} (S_n(v_n, w_n) + S_n(w_n, v_n)) = S(v, w) + S(w, v).$$

(b) For $\beta > 0$ and $\limsup_{n \rightarrow \infty} S_{n,\beta}(w_n, w_n) < \infty$ then

$$\lim_{n \rightarrow \infty} (S_{n,\beta}(v_n, w_n) + S_{n,\beta}(w_n, v_n)) = S_\beta(v, w) + S_\beta(w, v).$$

The lemma holds also for subsequences n_k , $k \in \mathbb{N}$, of indices.

Proof. See [14], proof of Lemma 2.5. However note the difference in the definition of pre-convergence, cf. condition (i) in Definition 2.4. Note also that $\sup_{n \in \mathbb{N}} \langle A_n w_n, A_n w_n \rangle_n < \infty$ implies by Proposition 2.3 (a) $\limsup_{n \rightarrow \infty} S_n(w_n, w_n) < \infty$, cf. the corresponding assumptions of Lemma 2.5 in [14].

Also in the present exposition, we observe that the pre-convergence of S_n , $n \in \mathbb{N}$, to S implies the pre-convergence of $S_{n,\beta}$, $n \in \mathbb{N}$, to S_β . Property (i) of Definition 2.4 for $S_{n,\beta}$, $n \in \mathbb{N}$, and S_β follows from Proposition 2.3 (a) and (e). The sequence ψ_n , $n \in \mathbb{N}$, in property (ii) of Definition 2.4 for $S_{n,\beta}$, $n \in \mathbb{N}$, and S_β is the same as that in property (ii) for S_n , $n \in \mathbb{N}$, and S . \square

Let us introduce the following condition.

- (c3) (i) $\mathcal{G} := \{G_\beta g : g \in \mathcal{C}, \beta > 0\} \subseteq \mathcal{C}$ in the sense that for every $g \in \mathcal{C}$ and $\beta > 0$, there is a $u \in \mathcal{C}$ with $G_\beta g = u$ ν -a.e.
- (ii) $\mathcal{G}_n := \{G_{n,\beta} g : g \in \mathcal{C}, \beta > 0\} \subseteq \mathcal{C}$, $n \in \mathbb{N}$, in the sense that for every $g \in \mathcal{C}$, $\beta > 0$, and every $n \in \mathbb{N}$, there exists a $v \in \mathcal{C}$ such that $G_{n,\beta} g = v$ ν_n -a.e.
- (iii) $\mathcal{G}' := \{G'_\beta g : g \in \mathcal{C}, \beta > 0\} \subseteq \mathcal{C}$.
- (iv) $\mathcal{G}'_n := \{G'_{n,\beta} g : g \in \mathcal{C}, \beta > 0\} \subseteq \mathcal{C}$, $n \in \mathbb{N}$.

Remark (3) Imposing condition (i) in Definition 2.4 on S_n , $n \in \mathbb{N}$, and S , we implicitly require that $\varphi \in D(S)$. On the one hand we suppose that $\varphi_{n_k} \in D(S_{n_k}) \cap \mathcal{C}$, $k \in \mathbb{N}$, w -converges to φ . This implies $\sup_{k \in \mathbb{N}} \langle \varphi_{n_k}, \varphi_{n_k} \rangle_{n_k} < \infty$ by Proposition 2.3 (a). On the other hand, we require $\sup_{k \in \mathbb{N}} \langle A_{n_k} \varphi_{n_k}, A_{n_k} \varphi_{n_k} \rangle_{n_k} < \infty$. Thus

$$S(\varphi, \varphi) \leq \liminf_{k \rightarrow \infty} S_{n_k}(\varphi_{n_k}, \varphi_{n_k}) \leq \limsup_{k \rightarrow \infty} \langle -A_{n_k} \varphi_{n_k}, \varphi_{n_k} \rangle_{n_k} < \infty$$

which, by definition, says that $\varphi \in D(S)$.

Conversely, in order to verify condition (i) in Definition 2.4, it makes sense to show that $D(S_{n_k}) \cap \mathcal{C} \ni \varphi_{n_k} \xrightarrow[n \rightarrow \infty]{w} \varphi \in L^2(E, \nu)$ and $\sup_{k \in \mathbb{N}} \langle A_{n_k} \varphi_{n_k}, A_{n_k} \varphi_{n_k} \rangle_{n_k} < \infty$ imply $\varphi \in D(S) = D(A)$. This is how we proceed in the proof of [15], Proposition 6.1, Step 3.

Lemma 2.6 *Let S_n , $n \in \mathbb{N}$, be a sequence of bilinear forms pre-convergent to S . Furthermore, let $\beta > 0$ and let $u_n \in D(S_n) \cap \mathcal{C}$ such that $A_n u_n \in \mathcal{C}$, $n \in \mathbb{N}$, be a w -convergent sequence with $\sup_{n \in \mathbb{N}} \langle A_n u_n, A_n u_n \rangle_n < \infty$. Let $u \in D(S)$. Introduce the following conditions.*

- (iii) *Let u_n , $n \in \mathbb{N}$, and u as above.*

$$\lim_{n \rightarrow \infty} S_{n,\beta}(u_n, \psi_n) = S_\beta(u, \psi) \tag{2.4}$$

for all $\psi \in \mathcal{C}$ and all sequences $\psi_n \in \mathcal{C}$, $n \in \mathbb{N}$, s -convergent to ψ yields

$$\lim_{n \rightarrow \infty} S_{n,\beta}(\psi_n, u_n) = S_\beta(\psi, u) \tag{2.5}$$

for all $\psi \in D(S)$ and all sequences $\psi_n \in D(S_n) \cap \mathcal{C}$, $n \in \mathbb{N}$, s -convergent to ψ in the sense of condition (ii) in Definition 2.4.

- (iv) *Let u_n , $n \in \mathbb{N}$, and u as above. If $\beta u_n - A_n u_n \xrightarrow[n \rightarrow \infty]{w} \beta u - Au$ then $u_n \xrightarrow[n \rightarrow \infty]{w} u$.*

Then (iv) implies (iii).

Remark (4) In (iii), we require that (2.4) holds for all $\psi \in \mathcal{C}$ and all sequences $\psi_n \in \mathcal{C}$, $n \in \mathbb{N}$, s -convergent to ψ . It is equivalent to replace in this sentence $\psi \in \mathcal{C}$ by $\psi \in \mathcal{C}'$ for any dense subset \mathcal{C}' of $L^2(E, \nu)$. For this, recall also Proposition 2.3 (a).

Proof. Let us assume (iv) and (2.4). We verify (2.5). For this, let us specify for a moment $\psi_n := \psi$, $n \in \mathbb{N}$. It follows then from Remark (2) and (2.4) that $\beta u_n - A_n u_n \xrightarrow[n \rightarrow \infty]{w} \beta u - Au$. Condition (iv) implies now that $u_n \xrightarrow[n \rightarrow \infty]{w} u$. From Lemma 2.5 and (2.4), we obtain

$$\begin{aligned} \lim_{n \rightarrow \infty} S_{n,\beta}(\psi_n, u_n) &= \lim_{n \rightarrow \infty} \{S_{n,\beta}(\psi_n, u_n) + S_{n,\beta}(u_n, \psi_n)\} - \lim_{n \rightarrow \infty} S_{n,\beta}(u_n, \psi_n) \\ &= \{S_\beta(\psi, u) + S_\beta(u, \psi)\} - S_\beta(u, \psi) \\ &= S_\beta(\psi, u) \end{aligned}$$

for all $\psi \in D(S)$ and all sequences $\psi_n \in D(S_n) \cap \mathcal{C}$, $n \in \mathbb{N}$, s -convergent to ψ satisfying condition (ii) in Definition 2.4. \square

Definition 2.4 continued Let S_n , $n \in \mathbb{N}$, be a sequence of bilinear forms pre-convergent to S . If, in addition, condition (iii) in Lemma 2.6 is satisfied, then we say that S_n , $n \in \mathbb{N}$, converges to S .

Theorem 2.7 Let $\beta > 0$, suppose that conditions (c1)-(c3) are satisfied, and assume that S_n , $n \in \mathbb{N}$, converges to S in the sense of Definition 2.4.

(a) For all $f \in L^2(E, \nu)$ and all sequences $f_n \in \mathcal{C}$ w -converging to f , $G_{n,\beta} f_n$ w -converges to $G_\beta f$ and $G'_{n,\beta} f_n$ w -converges to $G'_\beta f$ as $n \rightarrow \infty$.

(b) For all $g \in L^2(E, \nu)$ and all sequences $g_n \in \mathcal{C}$ s -converging to g , $G_{n,\beta} g_n$ s -converges to $G_\beta g$ and $G'_{n,\beta} g_n$ s -converges to $G'_\beta g$ as $n \rightarrow \infty$.

Proof. In Step 1 below, we will show that for all $g \in \mathcal{C}$, $G_{n,\beta} g$ s -converges to $G_\beta g$. In Step 2 we will use the ideas of Step 1 to prove that for all $f \in L^2(E, \nu)$ and all sequences $f_n \in \mathcal{C}$ w -converging to f , $G_{n,\beta} f_n$ w -converges to $G_\beta f$. In Step 3, we will demonstrate that for all $g \in L^2(E, \nu)$ and all sequences $g_n \in \mathcal{C}$ s -converging to g , $G_{n,\beta} g_n$ s -converges to $G_\beta g$. Finally, Step 4 will be devoted to the verification of the second part of (a) as a consequence of the first part of (b). A straight forward conclusion will then be the second part of (b).

Step 1 Fix $g \in \mathcal{C}$ and $\beta > 0$. Set $u_n := G_{n,\beta} g$. Because of $\langle u_n, u_n \rangle_n^{1/2} \leq \frac{1}{\beta} \sup_{n' \in \mathbb{N}} \langle g, g \rangle_{n'}^{1/2} < \infty$, $n \in \mathbb{N}$, and Proposition 2.3 (b), there exists a subsequence u_{n_k} , $k \in \mathbb{N}$, w -converging to some $\tilde{u} \in L^2(E, \nu)$. For this, recall also condition (c3). Because of Remark (3), we even may conclude $\tilde{u} \in D(S)$ since

$$\sup_{k \in \mathbb{N}} \langle A_{n_k} u_{n_k}, A_{n_k} u_{n_k} \rangle_{n_k} = \sup_{k \in \mathbb{N}} \langle \beta u_{n_k} - g, \beta u_{n_k} - g \rangle_{n_k} \leq 4 \sup_{k \in \mathbb{N}} \langle g, g \rangle_{n_k} < \infty.$$

Set $u := G_\beta g$ and let $\psi \in D(S)$. We have $\lim_{n \rightarrow \infty} S_{n,\beta}(G_{n,\beta} g, \psi_n) = \lim_{n \rightarrow \infty} \langle g, \psi_n \rangle_n = \langle g, \psi \rangle = S_\beta(G_\beta g, \psi)$ for all sequences $\psi_n \in \mathcal{C}$ s -converging to ψ . Thus, condition (iii) of Definition 2.4, Remark (4), and Lemma 2.5 imply that

$$\begin{aligned} S_\beta(\psi, u) + S_\beta(u, \psi) &= \lim_{k \rightarrow \infty} \{S_{n_k,\beta}(\psi_{n_k}, G_{n_k,\beta} g) + S_{n_k,\beta}(G_{n_k,\beta} g, \psi_{n_k})\} \\ &= S_\beta(\psi, \tilde{u}) + S_\beta(\tilde{u}, \psi) \end{aligned} \tag{2.6}$$

for all $\psi \in D(S)$ and all sequences $\psi_n \in D(S_n) \cap \mathcal{C}$ s -converging to ψ in the sense of condition (ii) of Definition 2.4. Note that, in order to use Lemma 2.5, we verify $\sup_{n \in \mathbb{N}} \langle A_n u_n, A_n u_n \rangle_n < \infty$ as above.

Applying (2.6) to both, $\psi = u$ and $\psi = \tilde{u}$, we conclude $S_\beta(u - \tilde{u}, u - \tilde{u}) = 0$ and thus $u = \tilde{u}$. In other words, $G_{n,\beta}g \xrightarrow[n \rightarrow \infty]{w} G_\beta g$, independent of the above chosen subsequence n_k , $k \in \mathbb{N}$. $G_{n,\beta}g \xrightarrow[n \rightarrow \infty]{s} G_\beta g$ is now a consequence of Proposition 2.3 (e), condition (c3) (ii), and Definition 2.4 (i), which imply

$$\begin{aligned} \beta \langle G_\beta g, G_\beta g \rangle + S(G_\beta g, G_\beta g) &\leq \liminf_{n \rightarrow \infty} \{ \beta \langle G_{n,\beta} g, G_{n,\beta} g \rangle_n + S_n(G_{n,\beta} g, G_{n,\beta} g) \} \\ &= \liminf_{n \rightarrow \infty} \langle g, G_{n,\beta} g \rangle_n \\ &= \langle g, G_\beta g \rangle \\ &= \beta \langle G_\beta g, G_\beta g \rangle + S(G_\beta g, G_\beta g) \end{aligned}$$

and thus $\lim_{n \rightarrow \infty} \langle G_{n,\beta} g, G_{n,\beta} g \rangle_n = \langle G_\beta g, G_\beta g \rangle$.

Step 2 Let $f \in L^2(E, \nu)$, $f_n \in \mathcal{C}$, $n \in \mathbb{N}$, be a sequence w -converging to f . Set $u_n := G_{n,\beta} f_n$. By Proposition 2.3 (a) we have $\langle u_n, u_n \rangle_n^{1/2} \leq \frac{1}{\beta} \sup_{n' \in \mathbb{N}} \langle f_{n'}, f_{n'} \rangle_{n'}^{1/2} < \infty$, $n \in \mathbb{N}$. Because of Proposition 2.3 (b) and Remark (3) there exists a subsequence u_{n_k} w -converging to some $\tilde{u} \in D(S)$ as $k \rightarrow \infty$. Here we have used

$$\sup_{k \in \mathbb{N}} \langle A_{n_k} u_{n_k}, A_{n_k} u_{n_k} \rangle_{n_k} = \sup_{k \in \mathbb{N}} \langle \beta u_{n_k} - f_{n_k}, \beta u_{n_k} - f_{n_k} \rangle_{n_k} \leq 4 \sup_{k \in \mathbb{N}} \langle f_{n_k}, f_{n_k} \rangle_{n_k} < \infty,$$

the latter by Proposition 2.3 (a).

Set $u := G_\beta f$ and let $\psi \in D(S)$. As in Step 1, we have $\lim_{n \rightarrow \infty} S_{n,\beta}(G_{n,\beta} f_n, \psi_n) = \lim_{n \rightarrow \infty} \langle f_n, \psi_n \rangle_n = \langle f, \psi \rangle = S_\beta(G_\beta f, \psi)$ for all sequences $\psi_n \in \mathcal{C}$ s -converging to ψ . Thus, condition (iii) of Definition 2.4, Remark (4), and Lemma 2.5 yield as in Step 1 $S_\beta(\psi, u) + S_\beta(u, \psi) = S_\beta(\psi, \tilde{u}) + S_\beta(\tilde{u}, \psi)$ for all $\psi \in D(S)$. Note again that, in order to use Lemma 2.5, we have $\sup_{k \in \mathbb{N}} \langle A_{n_k} u_{n_k}, A_{n_k} u_{n_k} \rangle_{n_k} < \infty$. Again we may conclude $u = \tilde{u}$. We have thus verified $G_{n,\beta} f_n \xrightarrow[n \rightarrow \infty]{w} G_\beta f$, independent of the above chosen subsequence n_k , $k \in \mathbb{N}$.

Step 3 Now, let $g \in L^2(E, \nu)$ and $g_n \in \mathcal{C}$, $n \in \mathbb{N}$, such that $g_n \xrightarrow[n \rightarrow \infty]{s} g$. Let $\varphi \in \mathcal{C}$, $\varepsilon > 0$, and choose $\tilde{g} \in \mathcal{C}$ with $\langle g - \tilde{g}, g - \tilde{g} \rangle^{1/2} < \varepsilon$. We have

$$\begin{aligned} &|(\langle G_{n,\beta} g_n, \varphi \rangle_n - \langle G_\beta g, \varphi \rangle) - (\langle G_{n,\beta} \tilde{g}, \varphi \rangle_n - \langle G_\beta \tilde{g}, \varphi \rangle)| \\ &\leq |\langle G_{n,\beta}(g_n - \tilde{g}), \varphi \rangle_n| + |\langle G_\beta(g - \tilde{g}), \varphi \rangle| \\ &\leq \frac{1}{\beta} \langle \varphi, \varphi \rangle_n^{1/2} (\langle g_n, g_n \rangle_n - 2\langle g_n, \tilde{g} \rangle_n + \langle \tilde{g}, \tilde{g} \rangle_n)^{1/2} + \frac{\varepsilon}{\beta} \langle \varphi, \varphi \rangle^{1/2} \\ &\xrightarrow[n \rightarrow \infty]{} \frac{1}{\beta} \langle \varphi, \varphi \rangle^{1/2} \langle g - \tilde{g}, g - \tilde{g} \rangle^{1/2} + \frac{\varepsilon}{\beta} \langle \varphi, \varphi \rangle^{1/2} = \frac{2\varepsilon}{\beta} \langle \varphi, \varphi \rangle^{1/2}. \end{aligned}$$

Together with $G_{n,\beta} \tilde{g} \xrightarrow[n \rightarrow \infty]{w} G_\beta \tilde{g}$ (cf. Step 1), this implies $G_{n,\beta} g_n \xrightarrow[n \rightarrow \infty]{w} G_\beta g$. Similarly, we obtain

$$\begin{aligned} &|(\langle G_{n,\beta} g_n, G_{n,\beta} g_n \rangle_n - \langle G_\beta g, G_\beta g \rangle) - (\langle G_{n,\beta} \tilde{g}, G_{n,\beta} \tilde{g} \rangle_n - \langle G_\beta \tilde{g}, G_\beta \tilde{g} \rangle)| \\ &\leq |\langle G_{n,\beta}(g_n + \tilde{g}), G_{n,\beta}(g_n - \tilde{g}) \rangle_n| + |\langle G_\beta(g + \tilde{g}), G_\beta(g - \tilde{g}) \rangle| \\ &\leq \frac{1}{\beta^2} \sup_{n \in \mathbb{N}} \langle g_n + \tilde{g}, g_n + \tilde{g} \rangle_n^{1/2} (\langle g_n, g_n \rangle_n - 2\langle g_n, \tilde{g} \rangle_n + \langle \tilde{g}, \tilde{g} \rangle_n)^{1/2} \\ &\quad + \frac{\varepsilon}{\beta^2} \langle g + \tilde{g}, g + \tilde{g} \rangle^{1/2} \end{aligned}$$

and thus

$$\begin{aligned}
& \limsup_{n \rightarrow \infty} |(\langle G_{n,\beta} g_n, G_{n,\beta} g_n \rangle_n - \langle G_\beta g, G_\beta g \rangle) - (\langle G_{n,\beta} \tilde{g}, G_{n,\beta} \tilde{g} \rangle_n - \langle G_\beta \tilde{g}, G_\beta \tilde{g} \rangle)| \\
& \leq \frac{1}{\beta^2} \sup_{n \in \mathbb{N}} \langle g_n + \tilde{g}, g_n + \tilde{g} \rangle_n^{1/2} \langle g - \tilde{g}, g - \tilde{g} \rangle^{1/2} + \frac{\varepsilon}{\beta^2} \langle g + \tilde{g}, g + \tilde{g} \rangle^{1/2} \\
& = \frac{\varepsilon}{\beta^2} \left(\sup_{n \in \mathbb{N}} \langle g_n + \tilde{g}, g_n + \tilde{g} \rangle_n^{1/2} + \langle g + \tilde{g}, g + \tilde{g} \rangle^{1/2} \right),
\end{aligned}$$

note that $\sup_{n \in \mathbb{N}} \langle g_n + \tilde{g}, g_n + \tilde{g} \rangle_n^{1/2} < \infty$ since $g_n + \tilde{g} \xrightarrow[n \rightarrow \infty]{s} g + \tilde{g}$. Together with $G_{n,\beta} \tilde{g} \xrightarrow[n \rightarrow \infty]{s} G_\beta \tilde{g}$ (cf. Step 1), and $G_{n,\beta} g_n \xrightarrow[n \rightarrow \infty]{w} G_\beta g$, this implies $G_{n,\beta} g_n \xrightarrow[n \rightarrow \infty]{s} G_\beta g$.

Step 4 Let $f \in L^2(E, \nu)$, $f_n \in \mathcal{C}$, $n \in \mathbb{N}$, be a sequence w -converging to f , and let $\varphi \in \mathcal{C}$. By the result of Step 1 and Proposition 2.3 (c), we have

$$\begin{aligned}
\langle G'_{n,\beta} f_n, \varphi \rangle_n & = \langle f_n, G_{n,\beta} \varphi \rangle_n \\
& \xrightarrow[n \rightarrow \infty]{} \langle f, G_\beta \varphi \rangle \\
& = \langle G'_\beta f, \varphi \rangle.
\end{aligned}$$

This means nothing but $G'_{n,\beta} f_n \xrightarrow[n \rightarrow \infty]{w} G'_\beta f$. Moreover, let $g \in L^2(E, \nu)$ and $g_n \in \mathcal{C}$ s -converging to g as $n \rightarrow \infty$. We have just proved that $G'_{n,\beta} g_n \xrightarrow[n \rightarrow \infty]{w} G'_\beta g$ which now implies $G_{n,\beta} G'_{n,\beta} g_n \xrightarrow[n \rightarrow \infty]{w} G_\beta G'_\beta g$. Again by Proposition 2.3 (c),

$$\begin{aligned}
\langle G'_{n,\beta} g_n, G'_{n,\beta} g_n \rangle_n & = \langle g_n, G_{n,\beta} G'_{n,\beta} g_n \rangle_n \\
& \xrightarrow[n \rightarrow \infty]{} \langle g, G_\beta G'_\beta g \rangle \\
& = \langle G'_\beta g, G'_\beta g \rangle.
\end{aligned}$$

Thus, $G'_{n,\beta} g_n$ s -converges to $G'_\beta g$ as $n \rightarrow \infty$. □

Lemma 2.8 *Let S_n , $n \in \mathbb{N}$, be a sequence of bilinear forms pre-convergent to S . Condition (iii) of Lemma 2.6 implies condition (iv).*

Proof. Suppose we have condition (iii). Let us also assume that $\beta u_n - A_n u_n$ w -converges to $\beta u - Au$. We show that u_n w -converges to u .

For all $\psi \in \mathcal{C}$ and all sequences $\psi_n \in \mathcal{C}$, $n \in \mathbb{N}$, s -convergent to ψ , we have because of Proposition 2.3 (c),

$$\begin{aligned}
S_{n,\beta}(u_n, \psi_n) & = \langle \beta u_n - A_n u_n, \psi_n \rangle_n \\
& \xrightarrow[n \rightarrow \infty]{} \langle \beta u - Au, \psi \rangle \\
& = S_\beta(u, \psi).
\end{aligned}$$

From condition (iii), it follows that

$$S_{n,\beta}(\psi_n, u_n) \xrightarrow[n \rightarrow \infty]{} S_\beta(\psi, u).$$

for all $\psi \in D(S)$ and all sequences $\psi_n \in D(S_n) \cap \mathcal{C}$, $n \in \mathbb{N}$, s -convergent to ψ in the sense of condition (ii) in Definition 2.4. In particular, we can take $g \in \mathcal{C}$, $\psi_n := G_{n,\beta} g$, $n \in \mathbb{N}$, and $\psi := G_\beta g$. For this, recall also condition (c3). According to Theorem 2.7, $\psi_n \xrightarrow[n \rightarrow \infty]{s} \psi$.

Furthermore, $\psi_n \in D(S_n) \cap \mathcal{C}$, $n \in \mathbb{N}$, and satisfies condition (ii) in Definition 2.4 since $\sup_{n \in \mathbb{N}} \langle A_n \psi_n, A_n \psi_n \rangle_n = \sup_{n \in \mathbb{N}} \langle \beta \psi_n - g, \beta \psi_n - g \rangle_n < \infty$ and

$$\limsup_{n \rightarrow \infty} S_n(\psi_n, \psi_n) = \lim_{n \rightarrow \infty} \langle g - \beta \psi_n, \psi_n \rangle_n = \langle g - \beta \psi, \psi \rangle = S(\psi, \psi).$$

Thus,

$$\langle g, u_n \rangle_n = S_{n,\beta}(\psi_n, u_n) \xrightarrow{n \rightarrow \infty} S_\beta(\psi, u) = \langle g, u \rangle$$

which means that $u_n \xrightarrow[n \rightarrow \infty]{w} u$. We have verified (iv). \square

Remarks (5) Lemmas 2.6 and 2.8 together show that under (i) and (ii) of Definition 2.4, conditions (iii) and (iv) are equivalent.

(6) By virtue of Remark (1), condition (c3), and Theorem 2.7, for $g \in \mathcal{C}$ and $g_n \in \mathcal{C}$ s -convergent to g it holds that

$$\langle G_{n,\beta} g_n - G_\beta g, G_{n,\beta} g_n - G_\beta g \rangle_n \xrightarrow{n \rightarrow \infty} 0. \quad (2.7)$$

For $f \in L^2(E, \nu)$ and $f_n \in \mathcal{C}$ w -convergent to f , from Theorem 2.7 and Proposition 2.3 (e) it follows that

$$\liminf_{n \rightarrow \infty} \langle G_{n,\beta} f_n, G_{n,\beta} f_n \rangle_n \geq \langle G_\beta f, G_\beta f \rangle$$

as well as

$$\liminf_{n \rightarrow \infty} \langle G'_{n,\beta} f_n, G'_{n,\beta} f_n \rangle_n \geq \langle G'_\beta f, G'_\beta f \rangle.$$

In order to prove convergence of forms S_n , $n \in \mathbb{N}$, to a form S let us introduce one more condition:

(c4) $D(S) \subseteq \mathcal{C}$ in the sense that for every $\varphi \in D(S)$, there is a $u \in \mathcal{C}$ with $\psi = u$ ν -a.e. Moreover, in this sense $\{A\varphi : \varphi \in D(S)\} \subseteq \mathcal{C}$.

In the next proposition we formulate conditions under which (i)-(iii) of Definition 2.4 become necessary. For this recall Remark (3).

Proposition 2.9 *Suppose that conditions (c1) – (c4) are satisfied. Let $(G_\beta)_{\beta \geq 0}$ be the resolvent of a strongly continuous contraction semigroup $(T_t)_{t \geq 0}$ on $L^2(E, \nu)$ and let $(G_{n,\beta})_{\beta \geq 0}$ be the resolvent of a strongly continuous contraction semigroup $(T_{n,t})_{t \geq 0}$ on $L^2(E, \nu_n)$, $n \in \mathbb{N}$. Suppose we have the following.*

(i) $G_{n,\beta} g_n$ s -converges to $G_\beta g$ as $n \rightarrow \infty$ for every $g \in \mathcal{C}$, every sequence $g_n \in D(S_n) \cap \mathcal{C}$ with $g_n \xrightarrow[n \rightarrow \infty]{s} g$, and $\beta > 0$. Furthermore, $G'_{n,\beta} g$ s -converges to $G'_\beta g$ as $n \rightarrow \infty$ for every $g \in \mathcal{C}$ and $\beta > 0$.

(ii) $\psi \in D(S_n) \cap \mathcal{C} = D(A_n) \cap \mathcal{C}$ implies $\psi \in D(A'_n)$.

(iii) Let $D(S_{n_k}) \cap \mathcal{C} \ni \varphi_{n_k} \xrightarrow[n \rightarrow \infty]{w} \varphi \in L^2(E, \nu)$ such that $\sup_{k \in \mathbb{N}} \langle A_{n_k} \varphi_{n_k}, A_{n_k} \varphi_{n_k} \rangle_{n_k} < \infty$. Then $\varphi \in D(S)$ and $\sup_{k \in \mathbb{N}} \langle A'_{n_k} \varphi_{n_k}, A'_{n_k} \varphi_{n_k} \rangle_{n_k} < \infty$.

Then the forms S_n , $n \in \mathbb{N}$, associated with $(T_{n,t})_{t \geq 0}$ converge to the form S associated with $(T_t)_{t \geq 0}$ as $n \rightarrow \infty$ in the sense of Definition 2.4.

Proof. *Step 1* In the first two steps, we verify condition (i) of Definition 2.4. For a clearer presentation, we ignore subsequences. Let us introduce bilinear forms $S^{(\beta)}(w, w) := \beta \langle w - \beta G_\beta w, w \rangle$, $w \in D(S)$, and $S_n^{(\beta)}(w_n, w_n) := \beta \langle w_n - \beta G_{n,\beta} w_n, w_n \rangle_n$, $w_n \in D(S_n) \cap \mathcal{C}$, $n \in \mathbb{N}$, which can be considered a counterpart to the Deny-Yosida approximation in Dirichlet form theory. Classical semigroup theory says that $\beta(w - \beta G_\beta w) \xrightarrow{\beta \rightarrow \infty} -Aw$ in $L^2(E, \nu)$ and therefore

$$S^{(\beta)}(w, w) = \beta \langle w - \beta G_\beta w, w \rangle \xrightarrow{\beta \rightarrow \infty} S(w, w), \quad w \in D(S).$$

and similarly $\lim_{\beta \rightarrow \infty} S_n^{(\beta)}(w_n, w_n) = S_n(w_n, w_n)$, $w_n \in D(S_n) \cap \mathcal{C}$, $n \in \mathbb{N}$. Differentiating with respect to β and applying the resolvent identity we get

$$\begin{aligned} \left| \frac{d}{d\beta} S_n^{(\beta)}(w_n, w_n) \right| &= \left| \frac{d}{d\beta} \beta \langle w_n - \beta G_\beta w_n, w_n \rangle_n \right| \\ &= \left| \langle w_n - 2\beta G_{n,\beta} w_n + \beta^2 G_{n,\beta}^2 w_n, w_n \rangle_n \right| \\ &= \left| \langle (A_n G_{n,\beta})^2 w_n, w_n \rangle_n \right| \\ &= \left| \langle G_{n,\beta} A_n G_{n,\beta} w_n, A'_n w_n \rangle_n \right| \\ &= \left| \langle G_{n,\beta}^2 A_n w_n, A'_n w_n \rangle_n \right| \\ &\leq \frac{1}{\beta^2} \langle A_n w_n, A_n w_n \rangle_n^{1/2} \langle A'_n w_n, A'_n w_n \rangle_n^{1/2}, \quad w_n \in D(S_n) \cap \mathcal{C}, \end{aligned}$$

where, for the last three lines we have applied hypothesis (ii). It follows now from hypothesis (iii) that $\lim_{\beta \rightarrow \infty} S_n^{(\beta)}(w_n, w_n) = S_n(w_n, w_n)$ uniformly in $n \in \mathbb{N}$ whenever $\sup_{n \in \mathbb{N}} \langle A_n \varphi_n, A_n \varphi_n \rangle_n < \infty$ and $D(S_n) \cap \mathcal{C} \ni w_n \xrightarrow[n \rightarrow \infty]{w} w \in L^2(E, \nu)$.

Step 2 Let us complete the verification of condition (i) of Definition 2.4. Let $D(S_n) \cap \mathcal{C} \ni \varphi_n \xrightarrow[n \rightarrow \infty]{w} \varphi \in \mathcal{C}$ and recall (c4). We can decompose

$$\begin{aligned} S_n^{(\beta)}(\varphi_n, \varphi_n) - S^{(\beta)}(\varphi, \varphi) &= \beta \langle \varphi_n - \beta G_{n,\beta} \varphi_n, \varphi_n \rangle_n - \beta \langle \varphi - \beta G_\beta \varphi, \varphi \rangle \\ &= \beta \langle \varphi_n - \beta G_{n,\beta} \varphi, \varphi \rangle_n + \beta \langle \varphi - \beta G_\beta \varphi, \varphi_n \rangle_n \\ &\quad - \beta \langle \varphi - \beta G_\beta \varphi, \varphi \rangle_n - \beta \langle \varphi - \beta G_\beta \varphi, \varphi \rangle \\ &\quad + \beta^2 \langle G_\beta \varphi - G_{n,\beta} \varphi - G'_{n,\beta} \varphi, \varphi_n - \varphi \rangle_n \\ &\quad + \beta \langle \varphi_n - \varphi - \beta G_{n,\beta}(\varphi_n - \varphi), \varphi_n - \varphi \rangle_n. \end{aligned}$$

Let us analyze the items on the right-hand side.

- (1) $\langle \varphi_n - \beta G_{n,\beta} \varphi, \varphi \rangle_n \xrightarrow[n \rightarrow \infty]{w} \langle \varphi - \beta G_\beta \varphi, \varphi \rangle$ because of $\varphi_n - \beta G_{n,\beta} \varphi \xrightarrow[n \rightarrow \infty]{w} \varphi - \beta G_\beta \varphi$, cf. (i) of this proposition,
- (2) $\langle \varphi - \beta G_\beta \varphi, \varphi_n \rangle_n \xrightarrow[n \rightarrow \infty]{w} \langle \varphi - \beta G_\beta \varphi, \varphi \rangle$ because of $\varphi - \beta G_\beta \varphi \in \mathcal{C}$ (cf. condition (c3) and Lemma 2.1 (b)) and $\varphi_n \xrightarrow[n \rightarrow \infty]{w} \varphi$,
- (3) $\langle \varphi - \beta G_\beta \varphi, \varphi \rangle_n \xrightarrow[n \rightarrow \infty]{w} \langle \varphi - \beta G_\beta \varphi, \varphi \rangle$ because of condition (c3), Lemma 2.1 (b) and, thus, $\varphi - \beta G_\beta \varphi \in \mathcal{C}$, and Lemma 2.1 (d),

(4) $\langle G_\beta \varphi - G_{n,\beta} \varphi - G'_{n,\beta} \varphi, \varphi_n - \varphi \rangle_n \xrightarrow{n \rightarrow \infty} 0$ because of $G_\beta \varphi - G_{n,\beta} \varphi - G'_{n,\beta} \varphi \xrightarrow{n \rightarrow \infty} -G'_\beta \varphi$ (cf. (i) of this proposition), $\varphi_n - \varphi \xrightarrow{n \rightarrow \infty} 0$, and Proposition 2.3 (c),

(5) $\langle \varphi_n - \varphi - \beta G_{n,\beta}(\varphi_n - \varphi), \varphi_n - \varphi \rangle_n \geq 0$ because of Lemma 2.1 (b) in [14].

Therefore, $\liminf_{n \rightarrow \infty} S_n^{(\beta)}(\varphi_n, \varphi_n) \geq S^{(\beta)}(\varphi, \varphi)$, $\beta > 0$. With the result of Step 1, we get $\liminf_{n \rightarrow \infty} S_n(\varphi_n, \varphi_n) \geq S(\varphi, \varphi)$ which, by condition (iii) of the present proposition and (c4), is condition (i) of Definition 2.4.

Step 3 Let us verify condition (ii) of Definition 2.4. Among others things, we will use an idea from the proof of [18], Theorem 2.4.1, part (jj). Let $\psi \in D(S)$ and recall that, because of (c4), we have $\psi \in \mathcal{C}$. Moreover let $\Psi := \psi - A\psi$, i. e. $\Psi \in \mathcal{C}$ by (c4) and Lemma 2.1 (b). Furthermore, $\psi = G_1 \Psi$. Let $\tilde{\psi}_n := G_{n,1} \Psi$, i. e. $\tilde{\psi}_n \in \mathcal{C}$, $n \in \mathbb{N}$, by (c3). By condition (i) of the present proposition we have $\psi_n = G_{n,1} \Psi \xrightarrow{n \rightarrow \infty} G_1 \Psi = \psi$ and therefore $G_{n,\beta} \tilde{\psi}_n \xrightarrow{n \rightarrow \infty} G_\beta \psi$. According to Proposition 2.3 (c) this means

$$\begin{aligned} S_n^{(\beta)}(\tilde{\psi}_n, \tilde{\psi}_n) &= \beta \langle \tilde{\psi}_n - \beta G_{n,\beta} \tilde{\psi}_n, \tilde{\psi}_n \rangle_n \\ &\xrightarrow{n \rightarrow \infty} \beta \langle \psi - \beta G_\beta \psi, \psi \rangle = S^{(\beta)}(\psi, \psi), \quad \beta > 0. \end{aligned}$$

On the other hand, it holds that $S^{(\beta)}(\psi, \psi) \xrightarrow{\beta \rightarrow \infty} S(\psi, \psi)$, cf. Step 1. Thus, there exists a sequence β_n , $n \in \mathbb{N}$, with $\beta_n \xrightarrow{n \rightarrow \infty} \infty$ such that

$$\lim_{n \rightarrow \infty} S_n^{(\beta_n)}(\tilde{\psi}_n, \tilde{\psi}_n) = S(\psi, \psi). \quad (2.8)$$

Because of

$$\begin{aligned} S_n^{(\beta_n)}(\tilde{\psi}_n, \tilde{\psi}_n) &= S_n(\beta_n G_{n,\beta_n} \tilde{\psi}_n, \beta_n G_{n,\beta_n} \tilde{\psi}_n) \\ &\quad + \beta_n \langle \tilde{\psi}_n - \beta_n G_{n,\beta_n} \tilde{\psi}_n, \tilde{\psi}_n - \beta_n G_{n,\beta_n} \tilde{\psi}_n \rangle_n \end{aligned} \quad (2.9)$$

and $S_n(\beta_n G_{n,\beta_n} \tilde{\psi}_n, \beta_n G_{n,\beta_n} \tilde{\psi}_n) \geq 0$ it follows from (2.8) that

$$\langle \tilde{\psi}_n - \beta_n G_{n,\beta_n} \tilde{\psi}_n, \tilde{\psi}_n - \beta_n G_{n,\beta_n} \tilde{\psi}_n \rangle_n \xrightarrow{n \rightarrow \infty} 0 \quad (2.10)$$

which implies

$$\begin{aligned} |\langle \tilde{\psi}_n - \beta_n G_{n,\beta_n} \tilde{\psi}_n, \varphi \rangle_n| &\leq \langle \tilde{\psi}_n - \beta_n G_{n,\beta_n} \tilde{\psi}_n, \tilde{\psi}_n - \beta_n G_{n,\beta_n} \tilde{\psi}_n \rangle_n^{1/2} \langle \varphi, \varphi \rangle_n^{1/2} \\ &\xrightarrow{n \rightarrow \infty} 0, \quad \varphi \in \mathcal{C}. \end{aligned}$$

We recall that \mathcal{C} is linear by Lemma 2.1 (b). Thus $\beta_n G_{n,\beta_n} \tilde{\psi}_n \xrightarrow{n \rightarrow \infty} \psi$ and (2.10) provides

$$\beta_n G_{n,\beta_n} \tilde{\psi}_n \xrightarrow{n \rightarrow \infty} \psi.$$

Relations (2.8) and (2.9) imply now

$$\limsup_{n \rightarrow \infty} S_n(\beta_n G_{n,\beta_n} \tilde{\psi}_n, \beta_n G_{n,\beta_n} \tilde{\psi}_n) \leq S(\psi, \psi).$$

In addition, by $\psi_n \in D(A_n)$, $n \in \mathbb{N}$, and $\Psi \in \mathcal{C} \subseteq \mathcal{D}$,

$$\begin{aligned} \sup_{n \in \mathbb{N}} \langle A_n \beta_n G_{n,\beta_n} \tilde{\psi}_n, A_n \beta_n G_{n,\beta_n} \tilde{\psi}_n \rangle_n &= \sup_{n \in \mathbb{N}} \langle \beta_n G_{n,\beta_n} A_n \tilde{\psi}_n, \beta_n G_{n,\beta_n} A_n \tilde{\psi}_n \rangle_n \\ &\leq \sup_{n \in \mathbb{N}} \langle A_n \tilde{\psi}_n, A_n \tilde{\psi}_n \rangle_n = \sup_{n \in \mathbb{N}} \langle \tilde{\psi}_n - \Psi, \tilde{\psi}_n - \Psi \rangle_n \leq 4 \sup_{n \in \mathbb{N}} \langle \Psi, \Psi \rangle_n < \infty. \end{aligned}$$

Choosing $\psi_n := \beta_n G_{n,\beta_n} \tilde{\psi}_n$, $n \in \mathbb{N}$, this verifies condition (ii) of Definition 2.4.

Step 4 In order to verify condition (iii) of Definition 2.4, let $\beta > 0$, $\psi \in \mathcal{C}$, and let $u_n \in D(S_n)$, $n \in \mathbb{N}$, be a w -convergent sequence with the properties mentioned in Lemma 2.6 and $u \in D(S)$. We assume that $\lim_{n \rightarrow \infty} S_{n,\beta}(u_n, \psi_n) = S_\beta(u, \psi)$ for all sequences $\psi_n \in \mathcal{C}$ s -converging to ψ . Recalling hypothesis (i) it follows that for $g \in \mathcal{C}$, $\hat{\psi}_n := G'_{n,\beta} g$, $n \in \mathbb{N}$, and $\hat{\psi} := G'_\beta g$ we have

$$\begin{aligned} \lim_{n \rightarrow \infty} \langle u_n, g \rangle_n &= \lim_{n \rightarrow \infty} S_{n,\beta}(u_n, \hat{\psi}_n) \\ &= S_\beta(u, \hat{\psi}) \\ &= \langle u, g \rangle. \end{aligned}$$

This means that u_n w -converges to u . For this, also note that because of condition (c3), $\hat{\psi}, \hat{\psi}_n \in \mathcal{C}$, $n \in \mathbb{N}$. Let now $\psi \in D(S)$ and $\psi_n \in D(S_n) \cap \mathcal{C}$, $n \in \mathbb{N}$, be a sequence s -convergent to ψ in the sense of Definition 2.4 (ii). Assume $\lim_{n \rightarrow \infty} S_{n,\beta}(u_n, \psi_n) = S_\beta(u, \psi)$. Lemma 2.5 yields $\lim_{n \rightarrow \infty} S_{n,\beta}(\psi_n, u_n) = S_\beta(\psi, u)$. We have verified (iii) of Definition 2.4. \square

We conclude this subsection with the proof of s -convergence of the associated semigroups. For this, let us introduce the following condition.

(c5) $\mathcal{T} := \{T_t g : g \in \mathcal{C}, t > 0\} \subseteq \mathcal{C}$, $\mathcal{T}_n := \{T_{n,t} g : g \in \mathcal{C}, t > 0\} \subseteq \mathcal{C}$, $\mathcal{T}' := \{T'_t g : g \in \mathcal{C}, t > 0\} \subseteq \mathcal{C}$, and $\mathcal{T}'_n := \{T'_{n,t} g : g \in \mathcal{C}, t > 0\} \subseteq \mathcal{C}$, $n \in \mathbb{N}$, in the sense of condition (c3).

Theorem 2.10 *Suppose that (c1), (c2), (c3), (c5) are satisfied. Then, for all $g \in \mathcal{C}$ and $\beta > 0$, $G_{n,\beta} g \xrightarrow[n \rightarrow \infty]{s} G_\beta g$ iff $T_{n,t} g \xrightarrow[n \rightarrow \infty]{s} T_t g$ for all $t > 0$ and $G'_{n,\beta} g \xrightarrow[n \rightarrow \infty]{s} G'_\beta g$ iff $T'_{n,t} g \xrightarrow[n \rightarrow \infty]{s} T'_t g$ for all $t > 0$.*

Proof. In Steps 1 and 2, we demonstrate that $G_{n,\beta} g \xrightarrow[n \rightarrow \infty]{s} G_\beta g$ implies $T_{n,t} g \xrightarrow[n \rightarrow \infty]{s} T_t g$ and that $G'_{n,\beta} g \xrightarrow[n \rightarrow \infty]{s} G'_\beta g$ implies $T'_{n,t} g \xrightarrow[n \rightarrow \infty]{s} T'_t g$. In Step 3 we verify the converse.

Step 1 In this step, let us show that $T_{n,t} f \xrightarrow[n \rightarrow \infty]{w} T_t f$ for all $f \in \mathcal{C}$. For well-definiteness, recall condition (c5). Since both, \mathcal{C} and $D(A^2)$, are dense in $L^2(E, \nu)$ it is sufficient to verify this claim for $f = (G_\beta)^2 h$, $h \in \mathcal{C}$. Set $g = G_\beta h$. With [19], Lemma 4.1 of Chapter 3, we have

$$\begin{aligned} \langle T_{n,t} f - T_t f, \varphi \rangle_n &= \langle T_{n,t} G_\beta g - T_t G_\beta g, \varphi \rangle_n \\ &= \langle T_{n,t} G_\beta g - T_{n,t} G_{n,\beta} g, \varphi \rangle_n + \langle G_{n,\beta} T_{n,t} g - G_{n,\beta} T_t g, \varphi \rangle_n + \langle G_{n,\beta} T_t g - G_\beta T_t g, \varphi \rangle_n \\ &= \langle T_{n,t} G_\beta g - T_{n,t} G_{n,\beta} g, \varphi \rangle_n - \left\langle \int_0^t T_{n,t-s} (G_\beta - G_{n,\beta}) T_s h ds, \varphi \right\rangle_n \\ &\quad + \langle G_{n,\beta} T_t g - G_\beta T_t g, \varphi \rangle_n. \end{aligned}$$

It follows from the Schwarz inequality, (2.7), and contractivity of $T_{n,t}$ in $L^2(E, \nu_n)$ that the first item of the right-hand side tends to zero. That the third item of the right-hand side tends to zero is a consequence of the Schwarz inequality and (2.7). Thus, it remains to demonstrate that $\langle \int_0^t T_{n,t-s} (G_\beta - G_{n,\beta}) T_s h ds, \varphi \rangle_n \xrightarrow[n \rightarrow \infty]{} 0$. But using again Schwarz' inequality, and contractivity of all the $T_{n,t-s}$, this follows from

$$\begin{aligned} &\left| \left\langle \int_0^t T_{n,t-s} (G_\beta - G_{n,\beta}) T_s h ds, \varphi \right\rangle_n \right| \\ &\leq \int_0^t \langle (G_\beta - G_{n,\beta}) T_s h, (G_\beta - G_{n,\beta}) T_s h \rangle_n^{1/2} \langle \varphi, \varphi \rangle_n^{1/2} ds, \end{aligned} \quad (2.11)$$

condition (c5), relation (2.7), and dominated convergence.

Step 2 Now, let us prove that $T_{n,t}f \xrightarrow[n \rightarrow \infty]{s} T_t f$ for all $f \in \mathcal{C}$. Again, we choose $f = (G_\beta)^2 h$ where $h \in \mathcal{C}$, set $g = G_\beta h$, and decompose

$$\begin{aligned} T_{n,t}f - T_t f &= (T_{n,t}G_\beta g - T_{n,t}G_{n,\beta}g) - \left(\int_0^t T_{n,t-s}(G_\beta - G_{n,\beta})T_s h ds \right) + (G_{n,\beta}T_t g - G_\beta T_t g). \end{aligned}$$

That the $L^2(E, \nu_n)$ -norms of the first and the third item tend to zero as $n \rightarrow \infty$ follows from the contractivity of the semigroups and relation (2.7). The arguments used already in Step 1, (2.11), lead to

$$\left\langle \int_0^t T_{n,t-s}(G_\beta - G_{n,\beta})T_s h ds, \int_0^t T_{n,t-s}(G_\beta - G_{n,\beta})T_s h ds \right\rangle_n \xrightarrow[n \rightarrow \infty]{} 0.$$

For the dual operators, we recall that also $(T'_{n,t})_{t \geq 0}$ is a strongly continuous contraction semigroup in $L^2(E, \nu_n)$, $n \in \mathbb{N}$, and $(T'_t)_{t \geq 0}$ is a strongly continuous contraction semigroup in $L^2(E, \nu)$, cf. [19] Subsection 1.10, especially Corollary 10.6. This means we can conclude that $T'_{n,t}f \xrightarrow[n \rightarrow \infty]{s} T'_t f$ for all $f \in \mathcal{C}$ the same way as we did it to show $T_{n,t}f \xrightarrow[n \rightarrow \infty]{s} T_t f$.

Step 3 This follows from $G_\beta f = \int_0^\infty e^{-\beta t} T_t f dt$, $G_{n,\beta} f = \int_0^\infty e^{-\beta t} T_{n,t} f dt$, $n \in \mathbb{N}$, $f \in \mathcal{C}$, the same relations for the dual operators, and dominated convergence. \square

Remark (7) From Theorems 2.7 and 2.10 it follows that for $g \in \mathcal{C}$ it holds that $\langle T_{n,t}g - T_t g, T_{n,t}g - T_t g \rangle_n \xrightarrow[n \rightarrow \infty]{} 0$ and $\langle T'_{n,t}g - T'_t g, T'_{n,t}g - T'_t g \rangle_n \xrightarrow[n \rightarrow \infty]{} 0$. Even more important is the following observation. By the contractivity of the semigroups $(T_{n,t})_{t \geq 0}$ we have for $\mathcal{C} \ni g_n \xrightarrow[n \rightarrow \infty]{s} g \in \mathcal{C}$ the limits $\langle T_{n,t}g_n - T_{n,t}g, T_{n,t}g_n - T_{n,t}g \rangle_n \xrightarrow[n \rightarrow \infty]{} 0$ and $\langle G_{n,\beta}g_n - G_{n,\beta}g, G_{n,\beta}g_n - G_{n,\beta}g \rangle_n \xrightarrow[n \rightarrow \infty]{} 0$.

Under the hypotheses of Theorem 2.10 this says $G_{n,\beta}g_n \xrightarrow[n \rightarrow \infty]{s} G_\beta g$ iff $G_{n,\beta}g \xrightarrow[n \rightarrow \infty]{s} G_\beta g$ iff $T_{n,t}g \xrightarrow[n \rightarrow \infty]{s} T_t g$ iff $T_{n,t}g_n \xrightarrow[n \rightarrow \infty]{s} T_t g$ for all $\beta > 0$ and $t > 0$. The same holds for the dual operators.

2.3 Mosco Type Convergence of Non-Positive Non-Symmetric Forms

For the remainder of this section, let us drop the assumption that the semigroups $(T_{n,t})_{t \geq 0}$, $n \in \mathbb{N}$, and $(T_t)_{t \geq 0}$ are contractive. Anything else for the semigroups remains as introduced in Section 1. As a consequence, we cannot state positivity of the associated bilinear forms. We are interested in substitutes for Theorem 2.7. For this, we collect almost everything still necessary for the remainder of the section in the following condition.

- (c6) (i) $\mathbb{1} \in D(A')$ and $\mathbb{1} \in D(A'_n)$, $n \in \mathbb{N}$, and $\sup_{n \in \mathbb{N}} \|A'_n \mathbb{1}\|_{L^\infty(E, \nu_n)} < \infty$.
- (ii) $T'_t \mathbb{1} \in L^\infty(E, \nu)$ and the limit $A' \mathbb{1} = \lim_{t \rightarrow 0} \frac{1}{t}(T'_t \mathbb{1} - \mathbb{1})$ exists in $L^\infty(E, \nu)$.
- (iii) There exist $N_n \in \mathcal{B}(E_n)$ with $\nu_n(N_n) \xrightarrow[n \rightarrow \infty]{} 0$ such that for $\varphi \in \mathcal{C}$ there is $\Phi_n \equiv \Phi_n(\varphi) \in \mathcal{C}$ with $\Phi_n = A'_n \mathbb{1} \cdot \varphi$ on $E_n \setminus N_n$. Furthermore, $\Phi_n \xrightarrow[n \rightarrow \infty]{s} A' \mathbb{1} \cdot \varphi$. Alternatively assume $A'_n \mathbb{1} \in \mathcal{V}$, $n \in \mathbb{N}$, and $A'_n \mathbb{1} \cdot \varphi \xrightarrow[n \rightarrow \infty]{s} A' \mathbb{1} \cdot \varphi$ for all $\varphi \in \mathcal{C}$.
- (iv) For $n \in \mathbb{N}$, there exists a set $D_n \subseteq D(S_n) \cap L^\infty(E, \nu_n)$ which is dense in $D(S_n)$ with respect to the norm $\|f\|_{D_n} := (\langle f, f \rangle_n + \langle A_n f, A_n f \rangle_n)^{1/2}$.

Define $D(\hat{S}_n) := D(S_n)$, $n \in \mathbb{N}$, $D(\hat{S}) := D(S)$, and

$$\hat{S}_n(u_n, v_n) := S_n(u_n, v_n) + \frac{1}{2} \langle A'_n \mathbf{1} \cdot u_n, v_n \rangle_n, \quad u_n \in D(\hat{S}_n), \quad v_n \in L^2(E, \boldsymbol{\nu}_n), \quad n \in \mathbb{N}, \quad (2.12)$$

and

$$\hat{S}(u, v) := S(u, v) + \frac{1}{2} \langle A' \mathbf{1} \cdot u, v \rangle, \quad u \in D(\hat{S}), \quad v \in L^2(E, \boldsymbol{\nu}). \quad (2.13)$$

Lemma 2.11 *Suppose (c6). We have $\hat{S}_n(u_n, u_n) \geq 0$, $u_n \in D(S_n)$, $n \in \mathbb{N}$, and $\hat{S}(u, u) \geq 0$, $u \in D(S)$.*

Proof. Step 1 We prove the claim for \hat{S} . Note that $T'_t \mathbf{1} \in L^\infty(E, \boldsymbol{\nu})$ (cf. (c6(ii))) and observe

$$\|T_t u\|_{L^1(E, \boldsymbol{\nu})} \leq \|T_t |u|\|_{L^1(E, \boldsymbol{\nu})} = \langle T'_t \mathbf{1}, |u| \rangle, \quad u \in L^2(E, \boldsymbol{\nu}).$$

This implies

$$\langle T_t u, T_t u \rangle \leq \|T_t u\|_{L^1(E, \boldsymbol{\nu})}^2 \leq \int T'_t \mathbf{1} \cdot u^2 d\boldsymbol{\nu}, \quad u \in L^2(E, \boldsymbol{\nu}), \quad t \geq 0,$$

the first inequality in this line by the association with a transition probability function. Consequently,

$$\begin{aligned} \langle T_t u, u \rangle &\leq \left(\int T'_t \mathbf{1} \cdot u^2 d\boldsymbol{\nu} \right)^{1/2} \cdot \langle u, u \rangle^{1/2} \\ &= \langle u, u \rangle + \langle u, u \rangle^{1/2} \left(\left(\int T'_t \mathbf{1} \cdot u^2 d\boldsymbol{\nu} \right)^{1/2} - \langle u, u \rangle^{1/2} \right) \end{aligned}$$

and therefore

$$-\frac{\langle u, u \rangle^{1/2}}{\langle u, u \rangle^{1/2} + \left(\int T'_t \mathbf{1} \cdot u^2 d\boldsymbol{\nu} \right)^{1/2}} \langle \frac{1}{t} (T'_t \mathbf{1} - \mathbf{1}) u, u \rangle \leq \frac{1}{t} \langle u - T_t u, u \rangle, \quad u \in L^2(E, \boldsymbol{\nu}).$$

Recalling $\mathbf{1} \in D(A')$ (cf. (c6(i))) and (c6(ii)), and letting $t \rightarrow 0$, it turns out that

$$-\frac{1}{2} \langle A' \mathbf{1} \cdot u, u \rangle \leq S(u, u), \quad u \in D(S).$$

Step 2 We prove the claim for \hat{S}_n . For this, we choose $u_n \in D_n$ (cf. (c6(iv))) and proceed as in Step 1. We arrive at

$$-\frac{\langle u_n, u_n \rangle_n^{1/2}}{\langle u_n, u_n \rangle_n^{1/2} + \left(\int T'_{n,t} \mathbf{1} \cdot u_n^2 d\boldsymbol{\nu}_n \right)^{1/2}} \langle \frac{1}{t} (T'_{n,t} \mathbf{1} - \mathbf{1}) u_n, u_n \rangle_n \leq \frac{1}{t} \langle u_n - T_{n,t} u_n, u_n \rangle_n.$$

Letting again $t \rightarrow 0$, we obtain $-\frac{1}{2} \langle A'_n \mathbf{1} \cdot u_n, u_n \rangle_n \leq S_n(u_n, u_n)$ for all $u_n \in D_n$. Let us finally mention that, in contrast to Step 1, we do not require $T'_{n,t} \mathbf{1} \in L^\infty(E, \boldsymbol{\nu}_n)$. We compensate this by requiring $u_n \in D_n \subseteq L^\infty(E, \boldsymbol{\nu}_n)$. But with (c6(iv)), we get finally $-\frac{1}{2} \langle A'_n \mathbf{1} \cdot u_n, u_n \rangle_n \leq S_n(u_n, u_n)$ for all $u_n \in D(S_n)$. \square

Let us assume that there are Markov processes associated with the semigroups $(T_{n,t})_{t \geq 0}$, $n \in \mathbb{N}$, and $(T_t)_{t \geq 0}$: For $n \in \mathbb{N}$, let $X^n = ((X_t^n)_{t \geq 0}, (P_\mu^n)_{\mu \in E_n})$ be a process taking values in E_n which corresponds to the semigroup $(T_{n,t})_{t \geq 0}$ and the form S_n . Here, E_n is the subset of E specified in Subsection 2.1. Furthermore, let $X = ((X_t)_{t \geq 0}, (P_\mu)_{\mu \in E})$ be a process associated with the semigroup $(T_t)_{t \geq 0}$ and the form S which takes values in some subset of E . Suppose that the paths of the processes X^n , $n \in \mathbb{N}$, and X are cadlag. For $\beta > 0$, introduce $G_{n,\beta} g_n := \int_0^\infty e^{-\beta t} T_{n,t} g_n dt$, $g_n \in L^\infty(E, \nu_n)$, $n \in \mathbb{N}$, $G_\beta g := \int_0^\infty e^{-\beta t} T_t g dt$, $g \in L^\infty(E, \nu)$. Since the semigroups $(T_{n,t})_{t \geq 0}$, $n \in \mathbb{N}$, and $(T_t)_{t \geq 0}$ are not necessarily contractive, the associated families of resolvents $(G_{n,\beta})_{\beta > 0}$, $n \in \mathbb{N}$, and $(G_\beta)_{\beta > 0}$ may not directly be well-defined on the corresponding L^2 -spaces.

Set $D(\hat{A}_n) := D(A_n)$, $n \in \mathbb{N}$, $D(\hat{A}) := D(A)$, and

$$\hat{A}_n u_n := A_n u_n - \frac{1}{2} A'_n \mathbb{I} \cdot u_n, \quad u_n \in D(\hat{A}_n), \quad n \in \mathbb{N}, \quad (2.14)$$

and

$$\hat{A} u := A u - \frac{1}{2} A' \mathbb{I} \cdot u, \quad u \in D(\hat{A}). \quad (2.15)$$

Let id_n denote the identity operator in $L^2(E, \nu_n)$, $n \in \mathbb{N}$, and let id denote the identity operator in $L^2(E, \nu)$.

Lemma 2.12 *Suppose (c6) and let $C := \frac{1}{2} \|A' \mathbb{I}\|_{L^\infty(E, \nu)} \vee \sup_{n \in \mathbb{N}} \frac{1}{2} \|A'_n \mathbb{I}\|_{L^\infty(E, \nu_n)}$. Then we have the following assertions.*

(a) *The operator \hat{A}_n is the generator of a strongly continuous contraction semigroup $(\hat{T}_{t,n})_{t \geq 0}$ in $L^2(E, \nu_n)$, $n \in \mathbb{N}$, and the operator \hat{A} is the generator of a strongly continuous contraction semigroup $(\hat{T}_t)_{t \geq 0}$ in $L^2(E, \nu)$. In addition, we have $\|\hat{T}_{t,n} g\|_{L^\infty(E, \nu_n)} \leq e^{Ct} \|g\|_{L^\infty(E, \nu_n)}$ if $g \in L^\infty(E, \nu_n)$, $n \in \mathbb{N}$, and $\|\hat{T}_t g\|_{L^\infty(E, \nu)} \leq e^{Ct} \|g\|_{L^\infty(E, \nu)}$ if $g \in L^\infty(E, \nu)$, $t \geq 0$.*

(b) *Let $\alpha \geq C$. The operator $A_n - \alpha id_n$ is the generator of a strongly continuous contraction semigroup $(T_{\alpha,n,t})_{t \geq 0}$ in $L^2(E, \nu_n)$, $n \in \mathbb{N}$, and the operator $A - \alpha id$ is the generator of a strongly continuous contraction semigroup $(T_{\alpha,t})_{t \geq 0}$ in $L^2(E, \nu)$.*

Proof. Adapting the ideas of the proof of Lemma 2.11 we obtain for $u \in L^\infty(E, \nu)$

$$\begin{aligned} & \langle u, u \rangle^{1/2} \cdot \frac{1}{t} \left(\langle T_t u, T_t u \rangle^{1/2} - \langle u, u \rangle^{1/2} \right) \\ & \leq \frac{\langle u, u \rangle^{1/2}}{\left(\int T'_t \mathbb{I} \cdot u^2 d\nu \right)^{1/2} + \langle u, u \rangle^{1/2}} \cdot \frac{1}{t} \left(\int T'_t \mathbb{I} \cdot u^2 d\nu - \langle u, u \rangle \right) \end{aligned}$$

which results for $u = T_s v$, $v \in L^\infty(E, \nu)$, $t \rightarrow 0$ in $\frac{d}{ds} \langle T_s v, T_s v \rangle^{1/2} \leq \frac{1}{2} \|A' \mathbb{I}\| \langle T_s v, T_s v \rangle^{1/2}$, $s \geq 0$. This and approximation of $v \in L^2(E, \nu)$ by $L^\infty(E, \nu)$ -functions implies $\langle T_s v, T_s v \rangle^{1/2} \leq e^{\frac{1}{2} \|A' \mathbb{I}\| s} \langle v, v \rangle^{1/2}$, $v \in L^2(E, \nu)$, $s \geq 0$. Similarly, $\langle T_{n,s} v, T_{n,s} v \rangle^{1/2} \leq e^{\frac{1}{2} \|A'_n \mathbb{I}\| s} \langle v, v \rangle^{1/2}$, $v \in L^2(E, \nu_n)$, $s \geq 0$, $n \in \mathbb{N}$.

(a) We prove the claim for $(\hat{T})_{t \geq 0}$. According to the Phillips-Lumer Theorem (cf. [19], Theorem I.4.3, or [24], Section IX.8) and our Lemma 2.11, it is sufficient to demonstrate that, for some $\hat{\alpha} > \frac{1}{2} \|A' \mathbb{I}\|_{L^\infty(E, \nu)}$ the range of $(\hat{\alpha} id - \hat{A})$ is $L^2(E, \nu)$. But this follows immediately from the Feynman-Kac formula,

$$u = \mathbb{E} \cdot \left(\int_0^\infty \exp \left\{ -\hat{\alpha} t - \int_0^t \frac{1}{2} A' \mathbb{I}(X_s) ds \right\} v(X_t) dt \right) \quad (2.16)$$

(\mathbb{E} stands for the expectation) which represents the solution to

$$\hat{\alpha}u - \hat{A}u = \hat{\alpha}u - Au + \frac{1}{2}A'\mathbb{1} \cdot u = v, \quad v \in L^2(E, \nu).$$

We recall the initial step of the present proof and note that therefore $u = G_{\hat{\alpha}}(v - \frac{1}{2}A'\mathbb{1} \cdot u) \in D(A)$ and that (2.16) corresponds to

$$\hat{T}_t v = \mathbb{E} \left(\exp \left\{ - \int_0^t \frac{1}{2}A'\mathbb{1}(X_s) ds \right\} v(X_t) \right), \quad v \in L^2(E, \nu). \quad (2.17)$$

We get $\|\hat{T}_t g\|_{L^\infty(E, \nu)} \leq e^{Ct} \|g\|_{L^\infty(E, \nu)}$ if $g \in L^\infty(E, \nu)$, $t \geq 0$, from (2.17) and condition (c6(i), (ii)).

(b) Keeping (2.16) and (2.17) in mind, this follows from similar considerations noting that, as a consequence Lemma 2.11, $S_{n, \alpha}$, $n \in \mathbb{N}$, and S_α are non-negative forms. \square

Remark (8) Another consequence of Lemma 2.12 is that, besides the definitions (2.12)-(2.15), \hat{A}_n , \hat{S}_n , $(\hat{T}_{n,t})_{t \geq 0}$, $n \in \mathbb{N}$, and \hat{A} , \hat{S} , $(\hat{T}_t)_{t \geq 0}$ are related as described in Section 1.

In addition, let $(\hat{G}_{n, \beta})_{\beta \geq 0}$, denote the resolvent associated with \hat{A}_n , \hat{S}_n , $(\hat{T}_{n,t})_{t \geq 0}$, $n \in \mathbb{N}$, and let $(\hat{G}_\beta)_{\beta \geq 0}$, denote the resolvent associated with \hat{A} , \hat{S} , $(\hat{T}_t)_{t \geq 0}$.

In order to handle the application in [15], it also seems to be beneficial to consider the following stronger condition in place of (c3) and the related Lemma 2.13.

- (c3') (i) If $\mathcal{C} \subseteq L^\infty(E, \nu)$ then $\{G_\beta g : g \in L^\infty(E, \nu), \beta > 0\} \subseteq \mathcal{C}$ in the sense that for every $g \in L^\infty(E, \nu)$, there is a $u \in \mathcal{C}$ with $G_\beta g = u$ ν -a.e.; otherwise, $D(S) \subseteq \mathcal{C}$.
- (ii) If, for $n \in \mathbb{N}$, $\mathcal{C} \subseteq L^\infty(E, \nu_n)$ then $\{G_{n, \beta} g : g \in L^\infty(E, \nu_n), \beta > 0\} \subseteq \mathcal{C}$ in the sense that for every $g \in L^\infty(E, \nu_n)$, there is a $u \in \mathcal{C}$ with $G_{n, \beta} g = u$ ν_n -a.e.; otherwise, $D(S_n) \subseteq \mathcal{C}$.

Lemma 2.13 *Suppose (c6). (a) If (c3'(i)) then condition (c3(i)) holds for \hat{S} in place of S . (b) If (c3'(ii)) then condition (c3(ii)) holds for \hat{S}_n , in place of S_n , $n \in \mathbb{N}$.*

Proof. We show only (a). *Step 1* Let $\beta > C$ and $g \in \mathcal{C}$. We have

$$\beta G_\beta g - A G_\beta g = g = \beta \hat{G}_\beta g - A \hat{G}_\beta g + \frac{1}{2}A'\mathbb{1} \cdot \hat{G}_\beta g.$$

In addition, we recall that $\hat{G}_\beta g \in L^\infty(E, \nu)$ if $g \in L^\infty(E, \nu)$ which is a consequence of Lemma 2.12 (a) since $\beta > C$. If $g \in L^2(E, \nu)$ then Lemma 2.12 (b) guarantees well-definiteness of $G_\beta g$ for $\beta > C$.

Because of $A'\mathbb{1} \in L^\infty(E, \nu)$ (cf. condition (c6(ii))), we have $g - \frac{1}{2}A'\mathbb{1} \cdot \hat{G}_\beta g \in L^\infty(E, \nu)$ if $\mathcal{C} \subseteq L^\infty(E, \nu)$, or otherwise $g - \frac{1}{2}A'\mathbb{1} \cdot \hat{G}_\beta g \in L^2(E, \nu)$. The above identity implies

$$\hat{G}_\beta g = G_\beta \left(g - \frac{1}{2}A'\mathbb{1} \cdot \hat{G}_\beta g \right).$$

Condition (c3(i)) restricted to $\beta > C$ relative to \hat{S} follows.

Step 2 Let now $\beta \in (0, C]$ and $g \in \mathcal{C}$. The sum $f := \sum_{k=0}^{\infty} (C \hat{G}_{\beta+C})^k g$ converges in $L^2(E, \nu)$ according to Lemma 2.12 (a) and is the unique solution to $f - C \hat{G}_{\beta+C} f = g$. By

$\beta\hat{G}_{\beta+C}f - \hat{A}\hat{G}_{\beta+C}f = f - C\hat{G}_{\beta+C}f = g$ we have $\hat{G}_{\beta}g = \hat{G}_{\beta+C}f$ and as in Step 1 of the present proof

$$\begin{aligned}\hat{G}_{\beta}g &= \hat{G}_{\beta+C}f = G_{\beta+C} \left(f - \frac{1}{2}A'\mathbf{1} \cdot \hat{G}_{\beta+C}f \right) \\ &= G_{\beta+C} \left(g + (C - \frac{1}{2}A'\mathbf{1}) \cdot \hat{G}_{\beta+C}f \right) \\ &= G_{\beta+C} \left(g + (C - \frac{1}{2}A'\mathbf{1}) \cdot \hat{G}_{\beta}g \right).\end{aligned}\tag{2.18}$$

If $\mathcal{C} \subseteq L^\infty(E, \nu)$ then Lemma 2.12 (a) says that $\hat{G}_{\beta}g \in L^\infty(E, \nu)$. In this case, we obtain $g + (C - \frac{1}{2}A'\mathbf{1}) \cdot \hat{G}_{\beta}g \in L^\infty(E, \nu)$ and with the right-hand side of (2.18) and the first part of condition (c3'(i)) we verify $\hat{G}_{\beta}g \in \mathcal{C}$.

If $\mathcal{C} \subseteq L^\infty(E, \nu)$ does not hold then again from (2.18) and condition (c3'(i)) we get $\hat{G}_{\beta}g \in \mathcal{C}$. \square

Theorem 2.14 *Let $C := \frac{1}{2}\|A'\mathbf{1}\|_{L^\infty(E, \nu)} \vee \sup_{n \in \mathbb{N}} \frac{1}{2}\|A'_n\mathbf{1}\|_{L^\infty(E, \nu_n)}$. Suppose (c1), (c2), and (c6). Furthermore, suppose (c3) for \hat{S}_n , $n \in \mathbb{N}$, and \hat{S} in place of S_n , $n \in \mathbb{N}$, and S . Assume that the forms \hat{S}_n , $n \in \mathbb{N}$, converge to the form \hat{S} in the sense of Definition 2.4.*

(a) *Let $\alpha \geq C$. Then the forms $S_{n, \alpha}$, $n \in \mathbb{N}$, converge to the form S_α in the sense of Definition 2.4.*

(b) *For all $f, g \in L^2(E, \nu)$, all sequences $f_n \in \mathcal{C}$ w -converging to f , all sequences $g_n \in \mathcal{C}$ s -converging to g , and all $\beta > 0$, we have $\hat{G}_{n, \beta}f_n \xrightarrow[n \rightarrow \infty]{w} \hat{G}_\beta f$, $\hat{G}'_{n, \beta}f_n \xrightarrow[n \rightarrow \infty]{w} \hat{G}'_\beta f$ and $\hat{G}_{n, \beta}g_n \xrightarrow[n \rightarrow \infty]{s} \hat{G}_\beta g$. Furthermore for all $g \in \mathcal{C}$, we have $\hat{G}'_{n, \beta}g \xrightarrow[n \rightarrow \infty]{s} \hat{G}'_\beta g$.*

(c) *Suppose (c3). For $\beta > C$, the operators $G_{n, \beta}$ and G_β can be continuously extended to operators $G_{n, \beta} : L^2(E, \nu_n) \rightarrow L^2(E, \nu_n)$, $n \in \mathbb{N}$, and $G_\beta : L^2(E, \nu) \rightarrow L^2(E, \nu)$, respectively. For $f, g \in L^2(E, \nu)$, and $f_n, g_n \in \mathcal{C}$, $n \in \mathbb{N}$, as in (b), we have $G_{n, \beta}f_n \xrightarrow[n \rightarrow \infty]{w} G_\beta f$, $G'_{n, \beta}f_n \xrightarrow[n \rightarrow \infty]{w} G'_\beta f$ and $G_{n, \beta}g_n \xrightarrow[n \rightarrow \infty]{s} G_\beta g$, $G'_{n, \beta}g_n \xrightarrow[n \rightarrow \infty]{s} G'_\beta g$.*

Remark (9) Recalling the proof of Theorem 2.7, especially Step 4, it will turn out that it is sufficient to require (c3(i)) as well as (c3(ii)) instead of (c3) (for S_n , $n \in \mathbb{N}$, and S) and (c3(i)) as well as (c3(ii)) instead of (c3) for \hat{S}_n , $n \in \mathbb{N}$, and \hat{S} if we are not interested in the convergence of $G'_{n, \beta}$ or $\hat{G}'_{n, \beta}$.

Proof. In order to show part (a), in Step 1-3 below, we will verify conditions (i)-(iii) of Definition 2.4 for $S_{n, \alpha}$, $n \in \mathbb{N}$, and S_α . In Step 4, we will verify (b) and (c).

Step 1 Let us use the symbols $\varphi_n, \varphi, \psi_n, \psi, u_n, u$, as in Definition 2.4 with S and S_n replaced with S_α and $S_{n, \alpha}$. Introduce $a_n := \alpha - \frac{1}{2}A'_n\mathbf{1}$, $n \in \mathbb{N}$, and $a := \alpha - \frac{1}{2}A'\mathbf{1}$. Since \mathcal{C} is dense in $L^2(E, \nu)$ (cf. Lemma 2.1 (c)), for given $\varepsilon > 0$, there exists $\tilde{\varphi} \in \mathcal{C}$ such that $\langle \varphi - \tilde{\varphi}, \varphi - \tilde{\varphi} \rangle < \varepsilon$. Let such ε and $\tilde{\varphi}$ be given. Condition (i) of Definition 2.4 for $S_{n, \alpha}$, $n \in \mathbb{N}$, and S_α follows from

$$S_{n, \alpha}(\varphi_n, \varphi_n) = \hat{S}_n(\varphi_n, \varphi_n) + \langle a_n^{1/2}\varphi_n, a_n^{1/2}\varphi_n \rangle_n, \quad n \in \mathbb{N},$$

$$S_\alpha(\varphi, \varphi) = \hat{S}(\varphi, \varphi) + \langle a^{1/2}\varphi, a^{1/2}\varphi \rangle,$$

condition (i) for \hat{S}_n , $n \in \mathbb{N}$, and \hat{S} , and

$$\langle a_n^{1/2}\varphi_n, a_n^{1/2}\varphi_n \rangle_n = \langle a_n^{1/2}(\varphi_n - \tilde{\varphi}), a_n^{1/2}(\varphi_n - \tilde{\varphi}) \rangle_n + 2\langle a_n\tilde{\varphi}, \varphi_n \rangle_n - \langle a_n\tilde{\varphi}, \tilde{\varphi} \rangle_n$$

together with condition (c6(i), (ii), (iii)) and Proposition 2.3 (c). For the alternative in (c6(iii)) use Lemma 2.1 (e) and Proposition 2.3 (g).

Step 2 For condition (ii) of Definition 2.4 for $S_{n,\alpha}$, $n \in \mathbb{N}$, and S_α we take the same decomposition as in Step 1 for ψ_n and ψ instead of φ_n and φ , $n \in \mathbb{N}$. It is then an immediate consequence of (ii) for \hat{S}_n , $n \in \mathbb{N}$, and \hat{S} together with condition (c6(i), (ii), (iii)) and Proposition 2.3 (c). For the alternative we use again Lemma 2.1 (e) and Proposition 2.3 (g).

Step 3 In order to verify condition (iii) of Definition 2.4 for $S_{n,\alpha}$, $n \in \mathbb{N}$, and S_α , let $\tilde{\beta} > 0$ and suppose $\sup_{n \in \mathbb{N}} \langle A_n u_n, A_n u_n \rangle_n < \infty$ as well as

$$S_{n,\alpha+\tilde{\beta}}(u_n, \psi_n) \xrightarrow{n \rightarrow \infty} S_{\alpha+\tilde{\beta}}(u, \psi).$$

Introducing $v_n := \hat{G}_{n,\tilde{\beta}}(-A_n u_n + (\alpha + \tilde{\beta})u_n)$, $n \in \mathbb{N}$, and $v := \hat{G}_{\tilde{\beta}}(-Au + (\alpha + \tilde{\beta})u)$, this means nothing but $\sup_{n \in \mathbb{N}} \langle A_n v_n, A_n v_n \rangle_n < \infty$ as well as

$$\hat{S}_{n,\tilde{\beta}}(v_n, \psi_n) \xrightarrow{n \rightarrow \infty} \hat{S}_{\tilde{\beta}}(v, \psi). \quad (2.19)$$

Choosing here $\psi_n = \psi$, $n \in \mathbb{N}$, and recalling Remark (2), we get $-\hat{A}_n v_n + \tilde{\beta} v_n \xrightarrow{n \rightarrow \infty} -\hat{A}v + \tilde{\beta}v$. Lemma 2.8 implies $v_n \xrightarrow{n \rightarrow \infty} v$. Let us conclude $u_n \xrightarrow{n \rightarrow \infty} u$ from this.

Let $\rho \in \mathcal{C}$. We have

$$\begin{aligned} \langle u_n + \hat{G}_{n,\tilde{\beta}}(a_n \cdot u_n), \rho \rangle_n &= \langle \hat{G}_{n,\tilde{\beta}}(-A_n u_n + (\alpha + \tilde{\beta}) \cdot u_n), \rho \rangle_n \\ &= \langle v_n, \rho \rangle_n \\ &\xrightarrow{n \rightarrow \infty} \langle v, \rho \rangle \\ &= \langle u + \hat{G}_{\tilde{\beta}}(a \cdot u), \rho \rangle. \end{aligned} \quad (2.20)$$

On the other hand, let us assume that $u_n \xrightarrow{n \rightarrow \infty} \tilde{u}$ for some $\tilde{u} \in L^2(E, \nu)$. From condition (c6(i), (ii), (iii)) and Proposition 2.3 (c),(f) we obtain the existence of $\tilde{\Psi}_n \in \mathcal{C}$ with $\tilde{\Psi}_n = a_n \cdot u_n$ on $E_n \setminus N_n$ and $\tilde{\Psi}_n \xrightarrow{n \rightarrow \infty} a \cdot \tilde{u}$. It follows from Theorem 2.7 (a) and the hypotheses of the present theorem that

$$\langle u_n + \hat{G}_{n,\tilde{\beta}}(a_n \cdot u_n), \rho \rangle_n \xrightarrow{n \rightarrow \infty} \langle \tilde{u} + \hat{G}_{\tilde{\beta}}(a \cdot \tilde{u}), \rho \rangle, \quad \rho \in \mathcal{C}. \quad (2.21)$$

According to (2.20) and (2.21), it holds that $\tilde{u} - u = -\hat{G}_{\tilde{\beta}}(a \cdot (\tilde{u} - u))$ which implies $\tilde{u} \in D(A)$ since $u \in D(A)$ by hypothesis and $\hat{A}(\tilde{u} - u) - \tilde{\beta}(\tilde{u} - u) = a \cdot (\tilde{u} - u)$. The latter implies

$$\langle -\hat{A}(\tilde{u} - u), \tilde{u} - u \rangle = \langle (\frac{1}{2}A'\mathbb{1} - \alpha - \tilde{\beta})(\tilde{u} - u), \tilde{u} - u \rangle.$$

The left-hand side is non-negative by Lemma 2.11. Since $\alpha \geq \frac{1}{2}\|A'\mathbb{1}\|_{L^\infty(E,\nu)}$ it follows that $u = \tilde{u}$.

Together with the above derived relation $v_n \xrightarrow{n \rightarrow \infty} v$ we now also have $u_n \xrightarrow{n \rightarrow \infty} u$. We recall that from condition (c6(i), (ii), (iii)) and Proposition 2.3 (c), (f) we obtain the existence of $\bar{\Psi}_n \in \mathcal{C}$ with $\bar{\Psi}_n = a_n \cdot u_n$ on $E_n \setminus N_n$ and $\bar{\Psi}_n \xrightarrow{n \rightarrow \infty} a \cdot u$. Together with (2.19) and Proposition 2.3 (c), this leads to

$$\begin{aligned} \hat{S}_{n,\tilde{\beta}}(u_n, \psi_n) &= \hat{S}_{n,\tilde{\beta}}(v_n, \psi_n) - \langle a_n \cdot u_n, \psi_n \rangle_n \\ &\xrightarrow{n \rightarrow \infty} \hat{S}_{\tilde{\beta}}(v, \psi) - \langle a \cdot u, \psi \rangle \\ &= \hat{S}_{\tilde{\beta}}(u, \psi). \end{aligned}$$

Condition (iii) of Definition 2.4 for \hat{S}_n , $n \in \mathbb{N}$, and \hat{S} implies now that

$$\hat{S}_{n,\bar{\beta}}(\psi_n, u_n) \xrightarrow{n \rightarrow \infty} \hat{S}_{\bar{\beta}}(\psi, u). \quad (2.22)$$

Since (2.22) holds for arbitrary $D(S_n) \cap \mathcal{C} \ni \psi_n \xrightarrow{n \rightarrow \infty} \psi \in D(S)$, it is also true for ψ_n replaced with $\psi_n + \hat{G}_{n,\bar{\beta}}(\Psi_n)$ and ψ replaced with $\psi + \hat{G}_{\bar{\beta}}(a \cdot \psi)$ where $\Psi_n \equiv \Psi_n(\psi) \in \mathcal{C}$ with $\Psi_n = A'_n \mathbf{1} \cdot \psi$ on $E_n \setminus N_n$ and E_n is as in (c6(iii)). For $\psi_n + \hat{G}_{n,\bar{\beta}}(\Psi_n) \in \mathcal{C}$, $n \in \mathbb{N}$, recall conditions (c6(iii)) and (c3) for \hat{S}_n as well as \hat{S} , and Lemma 2.1 (e). For $\psi_n + \hat{G}_{n,\bar{\beta}}(\Psi_n) \xrightarrow{n \rightarrow \infty} \psi + \hat{G}_{\bar{\beta}}(a \cdot \psi)$, consult condition (c6(i), (ii), (iii)), Proposition 2.3 (g), and Theorem 2.7. From (2.22), we obtain

$$\begin{aligned} S_{n,\alpha+\bar{\beta}}(\psi_n, u_n) &= \hat{S}_{n,\bar{\beta}}(\psi_n + \hat{G}_{n,\bar{\beta}}(a_n \cdot \psi_n), u_n) \\ &= \hat{S}_{n,\bar{\beta}}(\psi_n + \hat{G}_{n,\bar{\beta}}(\Psi_n), u_n) + \langle a_n \cdot \psi_n - \Psi_n, u_n \rangle_n \\ &\xrightarrow{n \rightarrow \infty} \hat{S}_{\bar{\beta}}(\psi + \hat{G}_{\bar{\beta}}(a \cdot \psi), u) \\ &= S_{\alpha+\bar{\beta}}(\psi, u); \end{aligned}$$

for $\langle a_n \cdot \psi_n - \Psi_n, u_n \rangle_n \xrightarrow{n \rightarrow \infty} 0$ we have already mentioned that $\sup_{n \in \mathbb{N}} \langle u_n, u_n \rangle_n < \infty$. Thus, we have (iii) for $S_{n,\alpha}$, $n \in \mathbb{N}$, and S_α . Part (a) has been verified.

Step 4 Part (b) is a corollary of the Lemmas 2.11 and 2.12 (a) and Theorem 2.7. For part (c), let $\beta > C$ and recall Lemma 2.12 (b). The claim follows from Part (a) of the present theorem and from Theorem 2.7. \square

Remark (10) It follows from $C = \frac{1}{2} \|A' \mathbf{1}\|_{L^\infty(E, \nu)} \vee \sup_{n \in \mathbb{N}} \frac{1}{2} \|A'_n \mathbf{1}\|_{L^\infty(E, \nu_n)} < \infty$, relation (2.17), and the contractivity of the semigroups $(\hat{T}_{t,n})_{t \geq 0}$ in $L^2(E, \nu_n)$, $n \in \mathbb{N}$, and $(\hat{T}_t)_{t \geq 0}$ in $L^2(E, \nu)$, cf. Lemma 2.12 (a), that

$$\langle T_{t,n} v, T_{t,n} v \rangle_n \leq e^{2Ct} \langle v, v \rangle_n, \quad n \in \mathbb{N}, \quad \text{and} \quad \langle T_t v, T_t v \rangle \leq e^{2Ct} \langle v, v \rangle, \quad t \geq 0.$$

Let us assume (c1), (c2), (c3(i), (ii)) and (c5). Keeping Theorem 2.14 (c) and in mind replacing in the proof of Theorem 2.10 and in Remark (7) contractivity by this property, for $\beta > C$ and $\mathcal{C} \ni g_n \xrightarrow{n \rightarrow \infty} g \in \mathcal{C}$ we obtain the following. $G_{n,\beta} g_n \xrightarrow{n \rightarrow \infty} G_\beta g$ iff $G_{n,\beta} g \xrightarrow{n \rightarrow \infty} G_\beta g$ iff $T_{n,t} g \xrightarrow{n \rightarrow \infty} T_t g$ iff $T_{n,t} g_n \xrightarrow{n \rightarrow \infty} T_t g$.

3 Relative Compactness

In this section, we specify the setting of Section 2. For this, let $\mathcal{M}_1(\bar{D})$ denote the space of all probability measures on $(\bar{D}, \mathcal{B}(\bar{D}))$ where D is a bounded d -dimensional domain or, more general, a bounded d -dimensional Riemannian manifold for some $d \in \mathbb{N}$. In addition, let $\mathcal{M}_\partial(\bar{D})$ be the set of all equivalence classes μ such that $m_1, m_2 \in \mu$ implies $m_1|_D = m_2|_D$. In this section we will assume that E is one of the spaces $\mathcal{M}_1(\bar{D})$ or $\mathcal{M}_\partial(\bar{D})$. In the case of $E = \mathcal{M}_\partial(\bar{D})$, we identify all points belonging to ∂D with each other. By $r(x, y) := |x - y| \wedge (\inf_{b \in \partial D} |b - x| + \inf_{b \in \partial D} |b - y|)$ if $x, y \in D$ and $r(x, \partial D) = r(\partial D, x) := \inf_{b \in \partial D} |b - x|$ if $x \in D$, as well as $r(\partial D, \partial D) := 0$ the space $(D \cup \partial D, r)$ becomes a separable, complete, and compact metric space. Furthermore, continuity on D with respect to r coincides with continuity with respect to the Euclidean metric and $\{f \in C(\bar{D}) : f \text{ constant on } \partial D\}$ is the

set of all continuous functions on $(D \cup \partial D, r)$. We would like to refer to similarities of this construction to that in [16].

Let both spaces $\mathcal{M}_1(\overline{D})$ and $\mathcal{M}_\partial(\overline{D})$ be endowed with the Prokhorov metric. We note that in this way $\mathcal{M}_1(\overline{D})$ and $\mathcal{M}_\partial(\overline{D})$ are separable, complete, and compact spaces.

Furthermore, for $n \in \mathbb{N}$, let E'_n be the set of all measures μ in E of the form $\mu = \frac{1}{n} \sum_{i=1}^n \delta_{z_i}$ where $z_1, \dots, z_n \in \overline{D}$ and δ_z denotes the Dirac measure concentrated at z . Furthermore, let $E_1 := E'_1$, $E_{n+1} := E'_{n+1} \setminus \bigcup_{i=1}^n E_i$, $n \in \mathbb{N}$, and $E_0 := E \setminus \bigcup_{n=1}^\infty E_n$. According to the basic setting of Subsection 2.1 E_n and E'_n differ by ν_n -null set, $n \in \mathbb{N}$. It is therefore reasonable to identify $L^p(E, \nu_n)$ with both $L^p(E_n, \nu_n)$ and $L^p(E'_n, \nu_n)$, $1 \leq p \leq \infty$, $n \in \mathbb{N}$.

To be consistent with [14], we will keep on writing $C_b(E)$ for $\mathcal{C}(E)$. Choose $\mathcal{F} := C_b(E)$ and note that therefore \mathcal{C} is now the space of all functions $\varphi \in \mathcal{D}$ satisfying the following.

(c1') φ is bounded and continuous on E_n , $n \in \mathbb{N}$.

(c2') $\langle \varphi, \psi \rangle_n \xrightarrow{n \rightarrow \infty} \langle \varphi, \psi \rangle$ for all $\psi \in C_b(E)$.

Obviously, \mathcal{F} is dense in $L^2(E, \nu)$. Under the above choice of \mathcal{F} , compatibility with Section 2 and well-definiteness of \mathcal{C} is subject to the subsequent.

Proposition 3.1 $\mathcal{F} = C_b(E)$ is a subset of \mathcal{D} if and only if the measures ν_n , $n \in \mathbb{N}$, are weakly convergent to ν as $n \rightarrow \infty$; in symbols $\nu_n \xrightarrow{n \rightarrow \infty} \nu$.

Let us assume that $\nu_n \xrightarrow{n \rightarrow \infty} \nu$ and note that (c1') and (c2') are now the defining properties of $\mathcal{C} \subseteq \mathcal{D}$.

As in Section 2, let us assume that there are Markov processes X^n and X associated with the semigroups $(T_{n,t})_{t \geq 0}$, $n \in \mathbb{N}$, and $(T_t)_{t \geq 0}$. Suppose that the paths of the processes X^n , $n \in \mathbb{N}$, are cadlag.

Define the measures $P_{\nu_n} := \int_E P_\mu^n \nu_n(d\mu)$, $n \in \mathbb{N}$, and $P_\nu := \int_E P_\mu \nu(d\mu)$, and introduce the processes $\mathbf{X}^n = ((X_t^n)_{t \geq 0}, P_{\nu_n})$ and $\mathbf{X} = ((X_t)_{t \geq 0}, P_\nu)$. Moreover, let \mathbb{E}_μ^n be the expectation corresponding to P_μ^n , $\mu \in E_n$, and let \mathbb{E}_{ν_n} be the expectation corresponding to P_{ν_n} , $n \in \mathbb{N}$. Let us introduce the set of test functions we are going to work with in this section. Suppose the following.

(c7) There exists an algebra $\tilde{C}_b(E) \subseteq C_b(E)$ of everywhere on E defined functions with $\tilde{C}_b(E) \subseteq \mathcal{G}$ in the sense that, for every $f \in \tilde{C}_b(E)$, there is a $g \equiv g(f) \in \mathcal{C}$ and a $\beta > 0$ with $f = \beta G_\beta g$ ν -a.e. $\tilde{C}_b(E)$ contains the constant functions and separates points in E .

Remark (1) Note that, for $f \in \tilde{C}_b(E)$, the existence of one $g \equiv g(f) \in \mathcal{C}$ and one $\beta > 0$ such that $f = \beta G_\beta g$ ν -a.e. implies that, for all $\beta' > 0$, there is a $g' \equiv g'(f, \beta') \in \mathcal{C}$ such that $f = \beta' G_{\beta'} g'$ ν -a.e. This follows from $Af = \beta f - \beta g \in \mathcal{C}$ and $g' := g + (1/\beta - 1/\beta')Af = (g + (Af - \beta f)/\beta) - (Af - \beta' f)/\beta' = -(Af - \beta' f)/\beta'$.

For $f \in \tilde{C}_b(E)$, $g = g(f) \in \mathcal{C}$, and a given sequence $\varepsilon_n > 0$, $n \in \mathbb{N}$, introduce

$$B := \bigcup_{n=1}^{\infty} \{\mu \in E_n : |\beta G_{n,\beta} g(\mu) - f(\mu)| \geq \varepsilon_n \|g\|\} . \quad (3.1)$$

Furthermore, let $\tau_{B^c} \equiv \tau_{B^c}^n(g)$ denote the first exit time of \mathbf{X}^n from the set $B^c \cap E_n$, $n \in \mathbb{N}$. Let $T > 0$ and set

$$\gamma_n \equiv \gamma_n(f) := \sup_{s \in [0, T+1]} |\beta G_{n,\beta} g(X_s^n) - f(X_s^n)| , \quad n \in \mathbb{N}.$$

In order to prove relative compactness of the families of processes $f(\mathbf{X}^n) = ((f(X_t^n))_{t \geq 0}, P_{\nu_n} \circ f^{-1})$, $n \in \mathbb{N}$, we need one more technical condition. In particular, we specify the sequence $\varepsilon_n > 0$, $n \in \mathbb{N}$.

(c8) There is a sequence $\varepsilon_n > 0$, $n \in \mathbb{N}$, with $\varepsilon_n \xrightarrow{n \rightarrow \infty} 0$ such that with $B \equiv B((\varepsilon_n)_{n \in \mathbb{N}})$ defined in (3.1)

$$\mathbb{E}_{\nu_n} (e^{-\beta \tau_{B^c}}) \xrightarrow{n \rightarrow \infty} 0$$

whenever $\langle f - \beta G_{n,\beta} g, f - \beta G_{n,\beta} g \rangle_n \xrightarrow{n \rightarrow \infty} 0$.

Theorem 3.2 (a) *Let the following be satisfied:*

(i) *Conditions (c3), (c7), and (c8) hold.*

(ii) *We have the hypotheses of Theorem 2.14, namely*

(c3) *for \hat{S}_n , $n \in \mathbb{N}$, and \hat{S} in place of S_n , $n \in \mathbb{N}$, and S ,*

(c6),

the forms \hat{S}_n , $n \in \mathbb{N}$, converge to the form \hat{S} in the sense of Definition 2.4.

Then, for $f \in \tilde{C}_b(E)$, the family of processes $f(\mathbf{X}^n) = ((f(X_t^n))_{t \geq 0}, P_{\nu_n} \circ f^{-1})$, $n \in \mathbb{N}$, is relatively compact with respect to the topology of weak convergence of probability measures over the Skorokhod space $D_{[-\|f\|, \|f\|]}([0, \infty))$.

(b) *The family of processes $\mathbf{X}^n = ((X_t^n)_{t \geq 0}, P_{\nu_n})$, $n \in \mathbb{N}$, is relatively compact with respect to the topology of weak convergence of probability measures over the Skorokhod space $D_E([0, \infty))$.*

Proof. (a) We will apply Theorem 3.8.6 of S.N. Ethier, T. Kurtz [2]. For $n \in \mathbb{N}$ and $t \geq 0$, let \mathcal{F}_t^n denote the σ -algebra generated by the family $(X_s^n)_{0 \leq s \leq t}$. In Steps 1 and 2 below, we will keep $n \in \mathbb{N}$ fixed. In Step 3, we will then pass to the limit as $n \rightarrow \infty$.

Step 1 Let $f \in \tilde{C}_b(E)$ and $\beta > 0$. Because of (c7) and Remark (1), there exist $g_1, g_2 \in \mathcal{C}$ with $f^2 = \beta G_{\beta} g_1$ ν -a.e. and $f = \beta G_{\beta} g_2$ ν -a.e. For $0 < \delta < 1$ and $0 \leq t \leq T$, $0 \leq u \leq \delta$, and $\beta > 0$, we have

$$\begin{aligned} & \mathbb{E}_{\nu_n} \left[(f(X_{t+u}^n) - f(X_t^n))^2 \mid \mathcal{F}_t^n \right] \\ &= \mathbb{E}_{\nu_n} \left[(f(X_{t+u}^n))^2 - (f(X_t^n))^2 \mid \mathcal{F}_t^n \right] - 2f(X_t^n) \mathbb{E}_{\nu_n} [f(X_{t+u}^n) - f(X_t^n) \mid \mathcal{F}_t^n] \\ &\leq \left| \mathbb{E}_{\nu_n} [f^2(X_{t+u}^n) - \beta G_{n,\beta} g_1(X_{t+u}^n) \mid \mathcal{F}_t^n] \right| + \left| \mathbb{E}_{\nu_n} [\beta G_{n,\beta} g_1(X_{t+u}^n) \right. \\ &\quad \left. - \beta G_{n,\beta} g_1(X_t^n) \mid \mathcal{F}_t^n] \right| + \left| \mathbb{E}_{\nu_n} [\beta G_{n,\beta} g_1(X_t^n) - f^2(X_t^n) \mid \mathcal{F}_t^n] \right| \\ &\quad + 2\|f\| \left| \mathbb{E}_{\nu_n} [f(X_{t+u}^n) - \beta G_{n,\beta} g_2(X_{t+u}^n) \mid \mathcal{F}_t^n] \right| + 2\|f\| \left| \mathbb{E}_{\nu_n} [\beta G_{n,\beta} g_2(X_{t+u}^n) \right. \\ &\quad \left. - \beta G_{n,\beta} g_2(X_t^n) \mid \mathcal{F}_t^n] \right| + 2\|f\| \left| \mathbb{E}_{\nu_n} [\beta G_{n,\beta} g_2(X_t^n) - f(X_t^n) \mid \mathcal{F}_t^n] \right| \end{aligned}$$

$$\begin{aligned}
&\leq 2 \mathbb{E}_{\nu_n} \left[\sup_{s \in [0, T+1]} \left| \beta G_{n, \beta} g_1(X_s^n) - f^2(X_s^n) \right| \middle| \mathcal{F}_t^n \right] \\
&\quad + \mathbb{E}_{\nu_n} \left[\sup_{r \in [0, T]} \int_r^{r+\delta} |A_n(\beta G_{n, \beta} g_1)(X_s^n)| ds \middle| \mathcal{F}_t^n \right] \\
&\quad + 4 \|f\| \mathbb{E}_{\nu_n} \left[\sup_{s \in [0, T+1]} \left| \beta G_{n, \beta} g_2(X_s^n) - f(X_s^n) \right| \middle| \mathcal{F}_t^n \right] \\
&\quad + 2 \|f\| \mathbb{E}_{\nu_n} \left[\sup_{r \in [0, T]} \int_r^{r+\delta} |A_n(\beta G_{n, \beta} g_2)(X_s^n)| ds \middle| \mathcal{F}_t^n \right] \\
&= 2 \mathbb{E}_{\nu_n} [\gamma_n(f^2) | \mathcal{F}_t^n] + \mathbb{E}_{\nu_n} \left[\sup_{r \in [0, T]} \beta \int_r^{r+\delta} |g_1(X_s^n) - \beta G_{n, \beta} g_1(X_s^n)| ds \middle| \mathcal{F}_t^n \right] \\
&\quad + 4 \|f\| \mathbb{E}_{\nu_n} [\gamma_n(f) | \mathcal{F}_t^n] + 2 \|f\| \mathbb{E}_{\nu_n} \left[\sup_{r \in [0, T]} \beta \int_r^{r+\delta} |g_2(X_s^n) - \beta G_{n, \beta} g_2(X_s^n)| ds \middle| \mathcal{F}_t^n \right] \\
&\leq 2 \mathbb{E}_{\nu_n} [\gamma_n(f^2) | \mathcal{F}_t^n] + 2\beta\delta \|g_1\| + 4 \|f\| \mathbb{E}_{\nu_n} [\gamma_n(f) | \mathcal{F}_t^n] + 4\beta\delta \|f\| \|g_2\|. \tag{3.2}
\end{aligned}$$

Step 2 For some arbitrary sequence $\varepsilon_n > 0$, $n \in \mathbb{N}$, satisfying $\varepsilon_n \xrightarrow{n \rightarrow \infty} 0$ and $\tau_{B^c} \equiv \tau_{B^c}^n(g_2)$ we have

$$\begin{aligned}
\mathbb{E}_{\nu_n} \gamma_n(f) &= \mathbb{E}_{\nu_n} (\chi_{\{\gamma_n \leq \varepsilon_n \|g_2\|\}} \gamma_n(f)) + \mathbb{E}_{\nu_n} (\chi_{\{\gamma_n > \varepsilon_n \|g_2\|\}} \gamma_n(f)) \\
&\leq \varepsilon_n \|g_2\| + (\|f\| + \|g_2\|) \mathbb{E}_{\nu_n} (\chi_{\{\gamma_n > \varepsilon_n \|g_2\|\}}) \\
&= \varepsilon_n \|g_2\| + (\|f\| + \|g_2\|) e^{\beta(T+1)} \mathbb{E}_{\nu_n} (e^{-\beta(T+1)} \chi_{\{\gamma_n > \varepsilon_n \|g_2\|\}}) \\
&\leq \varepsilon_n \|g_2\| + (\|f\| + \|g_2\|) e^{\beta(T+1)} \mathbb{E}_{\nu_n} (e^{-\beta\tau_{B^c}} \chi_{\{\gamma_n > \varepsilon_n \|g_2\|\}}). \tag{3.3}
\end{aligned}$$

It follows now from $f = \beta G_{\beta} g_2$ ν -a.e. and $f - \beta G_{\beta} g_2 \in \mathcal{C}$ (cf. (c3) and (c7)) that $\langle f - \beta G_{\beta} g_2, f - \beta G_{\beta} g_2 \rangle_n \xrightarrow{n \rightarrow \infty} 0$. Together with $G_{n, \beta} g_2 \xrightarrow{n \rightarrow \infty} G_{\beta} g_2$ (cf. Theorem 2.14 (c)), this leads to

$$\langle f - \beta G_{n, \beta} g_2, f - \beta G_{n, \beta} g_2 \rangle_n \xrightarrow{n \rightarrow \infty} 0.$$

Relation (3.3) and condition (c8) imply now

$$\mathbb{E}_{\nu_n} \gamma_n(f) \xrightarrow{n \rightarrow \infty} 0 \quad \text{and similarly} \quad \mathbb{E}_{\nu_n} \gamma_n(f^2) \xrightarrow{n \rightarrow \infty} 0. \tag{3.4}$$

Step 3 Setting

$$\gamma_n(\delta) := 2\gamma_n(f^2) + 4\|f\| \gamma_n(f) + 2\beta\delta \|g_1\| + 4\beta\delta \|f\| \|g_2\|, \quad n \in \mathbb{N},$$

and taking into consideration (3.2), we observe that

$$\mathbb{E}_{\nu_n} \left[(f(X_{t+u}^n) - f(X_t^n))^2 | \mathcal{F}_t^n \right] \leq \mathbb{E}_{\nu_n} [\gamma_n(\delta) | \mathcal{F}_t^n], \quad f \in \tilde{C}_b(E), \quad n \in \mathbb{N},$$

and from (3.4), we obtain

$$\lim_{\delta \rightarrow 0} \lim_{n \rightarrow 0} \mathbb{E}_{\nu_n}(\delta) = 0.$$

Relative compactness of the family $f(\mathbf{X}^n)$, $n \in \mathbb{N}$, follows now from [2], Theorems 3.7.2 and 3.8.6, Remark 3.8.7, and the fact that the processes $f(\mathbf{X}^n)$, $n \in \mathbb{N}$, take values in the compact interval $[\inf f, \sup f]$.

(b) Let us recall that E is compact. According to condition (c7), the Stone-Weierstrass Theorem, implies that $\tilde{C}_b(E)$ is dense in $C_b(E)$ with respect to the sup-norm. Now the claim is a consequence of part (a) and [2], Theorem 3.9.1. \square

References

- [1] F. CONRAD, M. GROTHAUS, J. LIERL, O. WITTICH, Convergence of Brownian motion with a scaled Dirac delta potential. *Proc. Edinb. Math. Soc. (Series 2)* **55** No. 2 (2012), 403-427.
- [2] S. N. ETHIER, T. KURTZ, *Markov processes, Characterization and convergence*, New York Chichester Brisbane Toronto Singapore: John Wiley 1986.
- [3] M. GROTHAUS, Y. G. KONDRATIEV, M. RÖCKNER N/V -limit for stochastic dynamics in continuous particle systems. *Probab. Theory Related Fields* **137** No. 1-2 (2007), 121-160.
- [4] M. HINO, Convergence of non-symmetric forms. *J. Math. Kyoto Univ.* **38** No. 2 (1998), 329-341.
- [5] M. HINZ Approximation of jump processes on fractals. *Osaka J. Math.* **46** No. 1 (2009), 141-171.
- [6] P. KIM Weak convergence of censored and reflected stable processes. *Stochastic Process. Appl.* **116** No. 12 (2006), 1792-1814.
- [7] A. V. KOLESNIKOV, Convergence of Dirichlet forms with changing speed measures on \mathbb{R}^d . *Forum Math.* **17** No. 2 (2005), 225-259.
- [8] A. V. KOLESNIKOV, Mosco convergence of Dirichlet forms in infinite dimensions with changing reference measures. *J. Funct. Anal.* **230** No. 2 (2006), 382-418.
- [9] A. V. KOLESNIKOV, Weak convergence of diffusion processes on Wiener space. *Probab. Theory Related Fields* **140** No. 1-2 (2008), 1-17.
- [10] K. KUWAE, T. UEMURA Weak convergence of symmetric diffusion processes. *Probab. Theory Related Fields* **109** No. 2 (1997), 159-182.
- [11] K. KUWAE, T. UEMURA Weak convergence of symmetric diffusion processes II. *Probability theory and mathematical statistics (Tokyo, 1995)*, River Edge, NJ: World Sci. Publ., (1996), 266-275.
- [12] K. KUWAE, T. SHIOYA, Convergence of spectral structures: a functional analytic theory and its applications to spectral geometry. *Comm. Anal. Geom.* **11** No. 4 (2003), 599-673.
- [13] J.-U. LÖBUS, A stationary Fleming-Viot type Brownian particle system. *Math. Z.* **263** No. 3 (2009), 541-581.
- [14] J.-U. LÖBUS, Weak convergence of n -particle systems using bilinear forms. *Milan J. Math.* **81** No. 1 (2013), 37-77.

- [15] J.-U. LÖBUS, Mosco type convergence and weak convergence for a Fleming-Viot type particle system, Preprint (2013), <http://www.mai.liu.se/~julob/Lo14-3.pdf>
- [16] MANDELKERN, Metrization of the one-point compactification. *Proc. Amer. Math. Soc.* **107** No. 4 (1989), 1111-1115.
- [17] Z.-M. MA AND M. RÖCKNER, *Introduction to the Theory of (Non-symmetric) Dirichlet Forms*, Berlin: Springer 1992.
- [18] U. MOSCO, Composite media and asymptotic Dirichlet forms. *J. Funct. Anal.* **123** No. 2 (1994), 368-421.
- [19] A. PAZY, *Semigroups of linear operators and applications to partial differential equations*, New York: Springer 1983.
- [20] O. V. PUGACHEV, On the closability and convergence of Dirichlet forms. *Proc. Steklov Inst. Math.*, **270** (2010), 216-221.
- [21] W. SUN, Weak convergence of Dirichlet processes. *Sci. China Ser. A* **41** No. 1 (1998), 8-21.
- [22] W. STANNAT, The theory of generalized Dirichlet forms and its applications in analysis and stochastics. *Mem. Amer. Math. Soc.* **142** No. 678 (1999).
- [23] J. M. TÖLLE Convergence of non-symmetric forms with changing reference measures, Diploma thesis, Faculty of Mathematics, University of Bielefeld, 2006.
- [24] K. YOSIDA, *Functional Analysis*, Berlin Heidelberg New York: Springer 1980.