The Number of Spanning Trees of an Infinite Family of Outerplanar, Small-World and Self-Similar Graphs

Francesc Comellas, Alícia Miralles

Dep. Matemàtica Aplicada IV, EPSC, Universitat Politècnica de Catalunya, c/ Esteve Terradas 5, Castelldefels (Barcelona), Catalonia, Spain (comellas@ma4.upc.edu).

Hongxiao Liu, Zhongzhi Zhang

School of Computer Science and Shanghai Key Lab of Intelligent Information Processing, Fudan University, Shanghai 200433, China (zhangzz@fudan.edu.cn, 10210240077@fudan.edu.cn).

Abstract

In this paper we give an exact analytical expression for the number of spanning trees of an infinite family of outerplanar, small-world and self-similar graphs. This number is an important graph invariant related to different topological and dynamic properties of the graph, such as its reliability, synchronization capability and diffusion properties. The calculation of the number of spanning trees is a demanding and difficult task, in particular for large graphs, and thus there is much interest in obtaining closed expressions for relevant infinite graph families. We have also calculated the spanning tree entropy of the graphs which we have compared with those for graphs with the same average degree.

Key words: spanning trees, tree entropy, complex networks, self-similarity

1. Introduction

Finding the number of spanning trees of graph is an old problem relevant in areas as diverse as mathematics, chemistry, physics, and computer science. This graph invariant is a parameter related, for example, to the reliability of a network [2] its synchronization [14] and the study of random walks [8]. The number of spanning trees of any finite graph can be computed from the well known Kirchhoff's matrix-tree theorem as the product of all nonzero eigenvalues of the Laplacian matrix of the graph [5], however this is a demanding and difficult task, in particular for large graphs. Thus, there has been much interest in finding other methods to produce exact expressions for the number of spanning trees of relevant graph families such as grids [11], lattices [13] and Sierpinski gaskets [1, 15].

In this paper we give an exact analytical expression for the number of spanning trees of an infinite family of outerplanar, small-world and selfsimilar graphs which was introduced by two of the authors in [3]. This family also has clustering zero and an exponential degree distribution. The combination of modularity, small clustering coefficient, small-world properties and exponential degree distribution can be found in some networks associated with several social and technological systems [9, 10]. Moreover, outer planarity is a relevant property as it is known that many algorithms that are NP-complete for general graphs perform polynomial in outerplanar graphs [12].

2. The M(t) graph family: construction and properties

This family of graphs M(t) was introduced in [3] and can be constructed following a recursive-modular method, see Fig. 1.

For t = 0, M(0) has two vertices and one edge.

For t = 1, M(1) is obtained from two graphs M(0) connected by two new edges.

For $t \geq 2$, M(t) is obtained from two graphs M(t-1) by connecting them with two new edges. In each M(t-1) the two vertices chosen are adjacent with maximum degree and have also been used at step t-1 to connect two M(t-2).



Figure 1: Graphs M(t) produced at iterations t = 0, 1, 2, 3 and t.

The number of vertices and edges of $M(t) = (V_t, E_t)$ are, respectively, $|V_t| = 2^{t+1}$ and $|E_t| = 3 \cdot 2^t - 2$.

The graph M(t) is outerplanar, that is, the graph is planar and has an embedding in the plane such that all vertices lie always in the exterior face of the graph while the edges never cross. This family of graphs has the degrees of the endvertices positively correlated –the graphs are assortative–while their degree distribution follows an exponential law $P_{\rm cum}(k) = 2^{2-k}$. See [3, 4] for more details and properties.

For large t, the diameter and average distance of M(t) scale logarithmically with the order of the graph, meaning that the graph is small-world, although its clustering coefficient is zero.

3. The number of spanning trees of M(t)

In this section we make use of the construction method of M(t) described in the previous section to obtain a set of recursive equations for the number of spanning trees and spanning subgraphs of the graph, which then can be solved by induction to find the exact number of spanning trees of M(t) at any iteration step t. For this calculation, we adapt the decimation technique given in [6], which has been also considered to find the number of spanning trees of the Sierpinski gasket, the pseudofractal web, and several fractal lattices, see [1, 18, 19].

As shown in the previous section, M(t) can be obtained from two M(t-1) graphs interconnected by adding two new edges. As the four endvertices of these new edges are particularly relevant when counting the number of spanning trees of M(t), we denote them as a_t , b_t , c_t and d_t and we will refer to them as *hubs*, see Fig. 1.

We recall here that a spanning subgraph of M(t) is a subgraph with the same vertex set as M(t) and a number of edges $|E'_t|$ such that $|E'_t| \leq |E_t|$. A spanning tree of M(t) is a spanning subgraph which is a tree and thus $|E'_t| = |V_t| - 1$.

Let s(t) denote the total number of spanning trees of M(t) and let g(t) be the number of spanning subgraphs of M(t) which consist of two trees such that one of the hub vertices used to form M(t+1) belongs to one tree and the other is in the second tree (note that one of the trees might be just one hub).

To find s(t), we classify the different cases that contribute to s(t) taking into account the connections between hub vertices of the two copies of M(t-1) that form M(t).

We denote the existence of an edge between hubs a_t , c_t by (a_t, c_t) and we represent that there is no edge between them by $\overline{(a_t, c_t)}$. We use the same notation for hubs b_t and d_t .

- 1. If $\overline{(a_t, c_t)}$ and $\overline{(b_t, d_t)}$ then there is no contribution to s(t). 2. If (a_t, c_t) and $\overline{(b_t, d_t)}$, the number of spanning trees that contribute to s(t) is

$$s(t-1)^2$$
. (1)

3. If $\overline{(a_t, c_t)}$ and (b_t, d_t) , like in the previous case, the number of spanning trees is

$$s(t-1)^2$$
. (2)

4. If (a_t, c_t) and (b_t, d_t) . In this case the number of spanning trees contributing to s(t) is s(t-1)g(t-1), i.e. the product of the number of spanning trees of the copy M(t-1) containing hubs a_t and b_t with g(t-1), which counts the number of spanning subgraphs of M(t-1)such that each contains the hubs c_t and d_t but does not have a path connecting them because they belong to different trees of the spanning graph. We have also to consider the symmetrical case with respect to the sets $\{a_t, b_t\}$ and $\{c_t, d_t\}$. Thus, the total number of spanning trees that corresponds to this case is

$$2s(t-1)g(t-1).$$
 (3)

By adding (1), (2) and (3) we find that the total number of spanning trees of M(t) is

$$s(t) = 2s(t-1)^{2} + 2s(t-1)g(t-1)$$
(4)

Now we need to obtain a recursive relation for g(t). A similar reasoning than above leads to the following equation

$$g(t) = s(t-1)^{2} + 2s(t-1)g(t-1).$$
(5)

Subtracting equation (5) from equation (4) we obtain

$$s(t) - g(t) = s(t-1)^2$$

and thus

$$g(t-1) = s(t-1) - s(t-2)^2.$$
 (6)

Introducing (6) into (4) we have:

$$s(t) = 4s(t-1)^2 - 2s(t-1)s(t-2)^2$$
 with $s(0) = 1$ and $s(1) = 4$.

To solve this equation, we rewrite it as

$$\frac{s(t)}{s(t-1)^2} = 4 - 2\frac{1}{\frac{s(t+1)}{s(t+2)^2}}$$
(7)

and, if we introduce

$$q(t) = s(t)/s(t-1)^2, \quad t \ge 2,$$

$$q(1) = 4,$$
(8)

equation (7) becomes

$$q(t) = 4 - \frac{2}{q(t-1)}, \quad \text{with} \quad t \ge 2,$$
 (9)

which can be solved to obtain

$$q(t) = 2 - \sqrt{2} + \frac{2\sqrt{2}}{1 - (3 - 2\sqrt{2})^t}.$$
(10)

From equation (8) we have:

$$s(t) = q(t)s(t-1)^{2} = q(t)q(t-1)^{2}s(t-2)^{4} = \cdots$$

= $q(t)q(t-1)^{2}q(t-2)^{2^{2}}q(t-3)^{2^{3}}\cdots q(t-i)^{2^{i}}\cdots q(1)^{2^{t-1}},$

and thus, using (10), we can write:

Theorem 3.1. The number of spanning trees of M(t), $t \ge 1$, is

$$s(t) = \prod_{i=1}^{t} \left[2 - \sqrt{2} + \frac{2\sqrt{2}}{1 - (3 - 2\sqrt{2})^i} \right]^{2^{t-i}}$$

4. Spanning tree entropy of M(t)

After having an exact expression for the number of spanning trees of M(t), we can calculate its spanning tree entropy, which is defined as in [7, 17]:

$$h = \lim_{V_t \to \infty} \frac{\ln s(t)}{V_t}.$$

and we obtain

 $h \simeq 0.657.$

We can now compare this asymptotic value of the entropy of the spanning trees of M(t) with those of other graph families with the same average degree: As an example, the value for the honeycomb lattice is 0.807 [17] and the 4-8-8 (bathroom tile) and 3-12-12 lattices have entropy values 0.787 and 0.721, respectively [13] while the spanning tree entropy for Hanoi graphs is 0.677 [20]. Thus, the asymptotic value of the entropy of the spanning trees of M(t) is the lowest known for graphs with average degree 3. This means that the number of spanning trees in M(t), although growing exponentially, do it at a lower rate than graphs with the same average degree. This result suggests that the family of graphs M(t) would be less reliable to a random removal of edges than the graphs mentioned above (as they have more spanning trees). However, the degree distribution also affects the reliability of a graph and it is well known that graphs with a scale-free distribution of degrees are much more resilient than homogeneous graphs (like regular and exponential graphs, where all vertices are statistically identical), see [16]. Thus, the large inhomogeneity in the degree distribution of the graphs M(t) might increase their robustness and it would be of interest to study, for these and other relevant graph families, the connections among spanning tree entropy, degree distribution and other topological parameters like degree correlation, in relation to their dynamic properties (reliability, synchronizability, etc.)

5. Conclusion

In this paper we find the number of spanning trees of the M(t) graph family by using a method, based on their self-similar structure, which allows us to obtain an analytical exact expression valid for any number of iterations. From the number of spanning trees we find their spanning tree entropy which is lower than in other graph families with the same average degree.

Acknowledgments

Z.Z. and H.L. are supported by the National Natural Science Foundation of China under grant No. 61074119. F.C. and A.M. are supported by the Ministerio de Economia y Competitividad, Spain, and the European Regional Development Fund under project MTM2011-28800-C02-01 and partially supported by the Catalan Research Council under grant 2009SGR1387.

References

- S.-C. Chang, L.-C. Chen, W.-S. Yang, Spanning trees on the Sierpinski gasket, J. Stat. Phys. 126 (2007) 649–667.
- [2] C.J. Colbourn, The Combinatorics of Network Reliability, New York, Oxford University Press 1987.

- [3] F. Comellas, A. Miralles, Modeling complex networks with self-similar outerplanar unclustered graphs, Phys. A, 388 (2009) 2227-2233.
- [4] F. Comellas, A. Miralles, Vertex labeling and routing in self-similar outerplanar unclustered graphs modeling complex networks J. Phys. A: Math. Gen. 42 (2009) 425001.
- [5] C. Godsil, G. Royle, Algebraic Graph Theory, Graduate Texts in Mathematics 207, Springer, New York 2001.
- [6] M. Knezevic, J. Vannimenus, Large-scale properties and collapse transition of branched polymers: Exact results on fractal lattices. Phys. Rev. Lett. 56 (1986) 1591.
- [7] R. Lyons, Asymptotic enumeration of spanning trees, Combin. Probab. Comput. 14 (2005) 491–522.
- [8] P. Marchal, Loop-erased random walks, spanning trees and hamiltonian cycles, Elect. Comm. in Probab. 5 (2000) 39-50.
- [9] M.E.J. Newman, The structure and function of complex networks, SIAM Review 45 (2003) 167–256.
- [10] R. Ferrer i Cancho, C. Janssen, R.V. Solé, Topology of technology graphs: Small world patterns in electronic circuits, Phys. Rev. E 64 (2001) 046119
- [11] S. D. Nikolopoulos, C. Papadopoulos, The number of spanning trees in Kncomplements of quasi-threshold graphs, Graphs Combin. 20 (2004) 383–394.
- [12] A. Brandstaedt, V.B. Le, P.J. Spinrad, Graph Classes: A Survey, SIAM Monographs on Discrete Mathematics and Applications. Philadelphia, PA, 1999.
- [13] R. Shrock and F. Y. Wu, Spanning trees on graphs and lattices in d dimensions, J. Phys. A: Math. Gen. 33 (2000) 3881.
- [14] N. Takashi, A. E. Motter, Synchronization is optimal in nondiagonalizable networks. Phys. Rev. E, 73 (2006) 065106(R).
- [15] E. Teufl, S. Wagner, Resistance Scaling and the Number of Spanning Trees in Self-Similar Lattices, J. Stat. Phys., 142 (2011) 879-897.
- [16] Y. Tu, How robust is the Internet?, Nature 406 (2000) 353-354.
- [17] F. Y. Wu, Number of spanning trees on a lattice, J. Phys. A: Math. Gen. 10 (1977) 113-115.
- [18] Z. Zhang, H. Liu, B. Wu, S. Zhou, Enumeration of spanning trees in a pseudofractal scale-free web, Europhys. Lett. 90 (2010) 68002.

- [19] Z. Zhang, H. Liu, B. Wu, T. Zou, Spanning trees in a fractal scale-free lattice, Phys. Rev. E 83 (2011) 016116.
- [20] Z. Zhang, M. Li, S. Wu, F. Comellas, The number and structure of spanning trees in the Hanoi graph, submitted.