Symmetry and Self-Duality in Categories of Probabilistic Models

Alexander Wilce

This note adds to the recent spate of derivations of the probabilistic apparatus of finite-dimensional quantum theory from various axiomatic packages. We offer two different axiomatic packages that lead easily to the Jordan algebraic structure of finite-dimensional quantum theory. The derivation relies on the Koecher-Vinberg Theorem, which sets up an equivalence between order-unit spaces having homogeneous, self-dual cones, and formally real Jordan algebras.

1 Introduction and Overview

The last several years have seen a spate of derivations of the probabilistic apparatus of finite-dimensional quantum theory from various axiomatic packages, many having an information-theoretic motivation [7, 8, 10, 12, 13]. This note (which in part echoes, but greatly improves upon [19]) adds to the flow. I offer two different, though overlapping, axiomatic packages, both stressing symmetry principles, that lead quickly and easily to the Jordan algebraic structure of finite-dimensional quantum theory. Quickly and easily, at any rate, if one is familiar with the Koecher-Vinberg Theorem [11, 15], which sets up an equivalence between order-unit spaces having homogeneous, self-dual cones, and formally real Jordan algebras.

A probabilistic system can be described, in a standard way, in terms of an order-unit space A, the positive elements of which are scalar multiples of "effects". The strategy, then, is to show that certain strong, but not unreasonable, assumptions force the positive cone A_+ to be homogeneous and self-dual, and hence, isomorphic to the cone of squares of such a Jordan algebra. In [3], several conditions are adduced that lead to a homogeneous and *weakly* self-dual cone — that is, a homogeneous cone that is *order-isomorphic* to its dual cone in A^* . However, proper self-duality is a much more stringent condition, requiring that the isomorphism be mediated by an inner product.

The line of attack here is to assume that systems individually have a great deal of symmetry, and collectively, can be organized into a symmetric monoidal category [1, 2, 14]. Here is a sketch. Further details can be found in the longer paper [20].

2 Probabilistic Models

A *test space* [16] is a pair (X, \mathfrak{A}) where X is a set of *outcomes* and \mathfrak{A} is a covering of X by non-empty (for our purposes here, finite) subsets called *tests*, each understood as the set of possible outcomes of some measurement, experiment, etc. Two outcomes $x, y \in X$ are *distinguishable* iff they belong to a common test. In this case, I write $x \perp y$. Notice that there is, as yet, no linear structure in view, let alone an inner product; so this notation is promissory.

A *state* on a test space (X,\mathfrak{A}) is a function $\alpha: X \to [0,1]$, summing independently to unity on each test. A *symmetry* of (X,\mathfrak{A}) is a mapping $g: X \to X$ such that $g(E), g^{-1}(E) \in \mathfrak{A}$ for every $E \in \mathfrak{A}$. By a *probabilistic model*, I mean a structure $(X,\mathfrak{A},\Omega,G)$ where (X,\mathfrak{A}) is a test space, Ω is a compact convex set of states on (X,\mathfrak{A}) , and G is a group acting on (X,\mathfrak{A}) by symmetries, and leaving Ω invariant.

For illustration, if **H** is a finite-dimensional Hilbert space (real or Complex), the corresponding *quantum model* is $(X(\mathbf{H}), \mathfrak{A}(\mathbf{H}), \Omega(\mathbf{H}), U(\mathbf{H}))$, where $X = X(\mathbf{H})$ is the set of rank-one projection operators on \mathbf{H} , $\mathscr{A} = \mathfrak{A}(\mathbf{H})$ is set of (projective) *frames*, i.e., maximal pairwise orthogonal sets of projections, $\Omega(\mathbf{H})$ is the convex set of density operators on \mathbf{H} , and $U(\mathbf{H})$ is the group of unitary operators on \mathbf{H} , acting on $X(\mathbf{H})$ by conjugation.

Categories of Models. I will be interested in categories of models. A *morphism* from a model $(X, \mathfrak{A}, \Omega, G)$ to a model $(Y, \mathfrak{B}, \Gamma, H)$ is a pair (ϕ, ψ) , where

- (i) $\phi: X \to Y$ with $\phi(\mathfrak{A}) \subseteq \mathfrak{B}$, $\phi^*(\Gamma) \subseteq \Omega$
- (ii) $\psi \in \text{Hom}(G, H)$;
- (iii) $\phi(gx) = \psi(g)\phi(x)$ for all $x \in X, g \in G$.

In what follows, \mathscr{C} is a symmetric monoidal category of probabilistic models $A = (X(A), \mathfrak{A}(A), \Omega(A), G(A))$, with morphisms as above. I shall make two further assumptions:

- (1) For every $A \in \mathcal{C}$, $G(A) \subseteq \mathcal{C}(A,A)$.
- (2) The model $A \otimes B \in \mathscr{C}$ is a *composite* of the models $A, B \in \mathscr{C}$, in the sense of [5]. This means, in particular, that there are canonical injections $\otimes : X(A) \times X(B) \to X(A \otimes B)$ and $\otimes : \Omega(A) \times \Omega(B) \to \Omega(A \otimes B)$, with

$$E \otimes F = \{x \otimes y | x \in E, y \in F\} \in \mathfrak{A}(A \otimes B)$$

for every $E \in \mathfrak{A}(A), F \in \mathfrak{A}(B)$, and

$$(\alpha \otimes \beta)(x \otimes y) = \alpha(x)\beta(y)$$

for all $\alpha \in \Omega(A)$, $\beta \in \Omega(B)$, $x \in X(A)$ and $y \in X(B)$. A *bipartite state* between $A, B \in \mathcal{C}$ is a state ω in $\Omega(A \otimes B)$. It is also part of the definition that the *un-normalized conditional state* $\hat{\omega}(x) := \omega(x, \cdot)$ belong to $\Omega(B)$ for every $x \in X$, and similarly with A and B reversed.

Models Linearized. Every model $A = (X(A), \mathfrak{A}(A), \Omega(A), G(A)) \in \mathscr{C}$ generates, in a standard and quite canonical way, an order-unit space $\mathbf{E}(A)$. To be precise, $\mathbf{E}(A)$ is the span in \mathbb{R}^{Ω} of the evaluation functionals associated with measurement outcomes $x \in X$.) In the case of a quantum model $A(\mathbf{H}) = (X(\mathbf{H}), \mathfrak{A}(\mathbf{H}), \Omega(\mathbf{H}), U(\mathbf{H}))$, one has $\mathbf{E}(A) \simeq \mathscr{L}(\mathbf{H})$, the space of Hermitian operators on \mathbf{H} , with the usual ordering and $u_A = \mathbf{1}_{\mathbf{H}}$.

The construction $A \mapsto \mathbf{E}(A)$ is functorial, so we obtain from $\mathscr C$ a category $\mathbf{E}(\mathscr C)$ of order-unit spaces and positive linear mappings. It is natural to enlarge this to a category $\mathscr E$ in which each hom-set $\mathscr E(A,B)$ is an ordered linear space, and in which, e.g., $\mathscr E(I,A) \simeq \mathbf{E}(A)$. In what follows, I assume that such a "linearized" category $\mathscr E$ has been fixed.

3 Bi-Symmetric Models

To tighten this structure further, I now ask that every $A \in \mathscr{C}$ enjoy a property I call bi-symmetry.

Definition. A model $A \in \mathcal{C}$ is *bi-symmetric* iff

- (i) G(A) acts transitively on the pure states (that is, extreme points) of $\Omega(A)$,
- (ii) G(A) acts transitively on \mathfrak{A} , and on pairs (x,y) of outcomes with $x \perp y$.

Alexander Wilce 277

If, in place of (ii), we require that arbitrary bijections $f: E \to F, E, F \in \mathfrak{A}$, extend to elements of G, then A is *fully bi-symmetric*.

If A is bi-symmetric, then G acts transitively. Clearly, the quantum model discussed above is fully bi-symmetric. Bi-symmetry and full bi-symmetry, are very natural conditions. (See [17] for further discussion and motivation of the latter.)

Definition. A *SPIN form*¹ for the model *A* is a real bilinear form *B* on $\mathbf{E}(A)$ that is *symmetric*, *positive* in the sense that $B(a,b) \geq 0$ for all $a,b \in \mathbf{E}(A)_+$, *invariant*, in the sense that B(ga,gb) = B(a,b) for all $g \in G(A)$, and *normalized*, in the sense that $B(u_A,u_A) = 1$. A SPIN form is *orthogonalizing* iff B(x,y) = 0 for all distinguishable measurement outcomes $x,y \in X(A)$.

An example is the usual tracial inner product on $\mathcal{L}(\mathbf{H})$. Call $\mathbf{E}(A)$ irreducible iff (with respect to any SPIN form B), the subspace $u^{\perp} = \{a \in \mathbf{E}(A) | B(a,u) = 0\}$ (this is independent of B) is irreducible under the group G(A). Quantum models are irreducible in this sense.

Theorem 1. If $\mathbf{E}(A)$ is irreducible, it supports at most one orthogonalizing SPIN form, which — if it exists — is an inner product.

4 Conjugates and Daggers

At this point, the aim is to find sufficient conditions for the existence of an orthogonalizing SPIN form on $\mathbf{E}(A)$. I shall provide two.

Definition. By a *conjugate* for a model A, I mean a structure $(\overline{A}, \gamma_A, \eta_A)$, where $\gamma_A : A \mapsto \overline{A}$ is an isomorphism of models, and η_A is a bipartite state on $A \times \overline{A}$ such that $\eta(x, \gamma_A(x)) = 1/n$ (where n is the rank of A) for every $x \in X(A)$.

In the case of a quantum-mechanical model $A = A(\mathbf{H})$ associated with a Hilbert space \mathbf{H} , the obvious conjugate model is just that associated with the conjugate Hilbert space $\overline{\mathbf{H}}$, with γ_A taking the rank-one projection x to the corresponding projection \overline{x} on $\overline{\mathbf{H}}$, and with η_A the pure state associated with the unit vector $\frac{1}{\sqrt{n}}\sum_i e_i \otimes \overline{e}_i$, $\{e_i\}$ any basis for \mathbf{H} (note that this is basis-independent).

Returning to the general case, note that by averaging over G(A), we can choose η_A to be invariant, in the sense that $\eta_A(gx, \gamma_A(gy)) = \eta_A(gx, gy)$ for all $g \in G(A)$. This gives us an invariant SPIN form on $\mathbf{E}(A)$, defined on outcomes by $B(x,y) := \eta(x, \gamma_A(y))$. Applying Theorem 1, we have

Theorem 2. Let $\mathbf{E}(A)$ be irreducible, and suppose A has a conjugate. Then $\mathbf{E}(A)$ carries a canonical orthogonalizing inner product.

Under some mild auxiliary hypotheses, the existence of a conjugate for every $A \in \mathscr{C}$ (with γ_A and η_A appropriately belonging to \mathscr{C}) can be used to construct a dagger on the category $\mathscr{E} \supseteq \mathbf{E}(\mathscr{C})$ discussed above. In fact, however, the mere existence of a reasonable dagger-monoidal structure on \mathscr{E} is enough to obtain much the same result.

Theorem 3. Suppose \mathscr{E} supports a dagger-monoidal structure, such that for every $g \in G(A)$, $g^{\dagger} = g^{-1}$ (i.e., $g \in G(A)$ is "unitary"). If $\mathbf{E}(A)$ is irreducible, then it carries an orthogonalizing inner product.

In order to obtain the self-duality of $\mathbf{E}(A)_+$ for an irreducible model A, it now suffices to assume either of two simple further conditions:

Theorem 4. Suppose that either

¹This is probably not the best choice of terminology.

- (a) In the context of Theorem 2, A has a conjugate such that the state η_A is an isomorphism state or
- (b) In the context of Theorem 3, A is sharp, meaning that every outcome has probability one in a unique state on $\mathbf{E}(A)$.

Then $\mathbf{E}(A)_+$ is self-dual.

The homogeneity of $\mathbf{E}(A)_+$ can now be enforced by any of several conditions discussed in [4, 19]. Applying the Koecher-Vinberg Theorem, we can conclude that $\mathbf{E}(A)$ carries a unique Jordan product making it a formally real Jordan algebra.

One of these conditions is so simple it's worth pausing to describe it. Any bipartite state ω between $A, B \in \mathscr{C}$ gives rise to a natural positive linear mapping $\hat{\omega} : \mathbf{E}(A) \to \mathbf{E}(B)^*$, uniquely defined by $\hat{\omega}(x)(y) = \omega(x,y)$. Where $\hat{\omega}$ is an *order-isomorphism* — that is, where $\hat{\omega}$ is an order-isomorphism (that is, invertible and having a positive inverse), we call ω an *isomorphism state*. A basic observation from [4], translated into the present context, is that if every state in the interior of $\Omega(B)$ is the marginal of an isomorphism state, then the cone in $\mathbf{E}(B)$ generated by $\Omega(B)$ is homogeneous.

5 Image-Closure

In order to extend these results to possibly reducible systems, I impose one further constraint on \mathscr{C} . Call a morphism $(\phi, \psi): (X, \mathfrak{A}, \Omega, G) \to (Y, \mathfrak{B}, \Gamma, H)$ is *surjective* iff $\phi(X) = Y$, $\mathfrak{B} \subseteq \phi(\mathfrak{A})$, $H = \psi(G)$, and $\Gamma = \{\beta \in \Omega(Y, \mathfrak{B}) | \phi^*(\beta) \in \Omega\}$. In this case, we call $(Y, \mathfrak{B}, \Gamma, H)$ the *image* of $(X, \mathfrak{A}, \Omega, G)$ under (ϕ, ψ) . Call \mathscr{C} *image-closed* iff, for any $A \in \mathscr{C}$ and any surjective morphism $(\phi, \psi): (X_A, \mathfrak{A}_A, \Omega_A, G_A) \to (Y, \mathfrak{B}, \Gamma, H)$, (i) the model $B := (Y, \mathfrak{B}, \Gamma, H)$ belongs to \mathscr{C} , and (ii) $(\phi, \psi) \in \mathscr{C}(A, B)$. in \mathscr{C} , again belong to \mathscr{C} .

Theorem 5. Let \mathscr{C} be an image-closed category of bi-symmetric probabilistic models, and let \mathscr{E} be the corresponding linearized category as discussed in Section 1. If either

- (a) every $A \in \mathcal{C}$ has a conjugate $\overline{A} \in \mathcal{C}$, with η_A an isomorphism state, or
- (b) \mathscr{E} has a dagger-monoidal structure making every $g \in G(A)$ unitary for all $A \in \mathscr{E}$, and every $A \in \mathscr{E}$ is sharp, then for every $A \in \mathscr{E}$, $\mathbf{E}(A)_+$ is self-dual.

Again, adding any of the sufficient conditions for homogeneity from [4, 17] — or simply assuming it outright — will yield a category of formally real Jordan algebras.

Operationally, it is reasonable to suppose that any image $\phi(A)$ of a model $A \in \mathscr{C}$ can be *simulated* by means of the model A. Hence, if we wish to think of \mathscr{C} as closed under operationally reasonable constructions, it is not far-fetched that $\phi(A)$ should belong to \mathscr{C} . In fact, the image of a bi-symmetric model is 2-symmetric, so one can simply "close up" \mathscr{C} without sacrificing this assumption. (To suppose that this closure continues to support, e.g., a symmetric-monoidal structure, or conjugate systems, is a sharper constraint, of course.) Categories of finite-dimensional quantum models turn out to be image-closed for the simple reason that a quantum model *has no* non-trivial images.

6 Conclusion

These results raise any number of interesting questions. For one thing, it is possible that the assumptions are stronger than advertised, singling out a narrower class than formally real Jordan algebras. It is noteworthy that I have not had to assume that \mathscr{C} 's monoidal product is locally tomographic. In fact, in a forthcoming paper with Howard Barnum [6], we show (using a result of Hanche-Olsen) that if \mathscr{C} is

Alexander Wilce 279

a dagger-monoidal category of finite-dimensional order-unit spaces with homogeneous self-dual cones, then local tomography, plus the existence in $\mathscr C$ of a system having the structure of a qubit, implies that every $A \in \mathscr C$ is isomorphic to the Hermitian part of a finite-dimensional complex C^* algebra.

References

- [1] S. Abramsky and B. Coecke, *A categorical semantics of quantum protocols*. Proceedings of the 19th IEEE conference on Logic in Computer Science (LiCS'04). IEEE Computer Science Press (2004). doi:10.1109/LICS.2004.1319636.
- [2] J. Baez, Quantum quandaries: a category-theoretic perspective. arXiv:quant-ph/0404040 (2004).
- [3] H. Barnum, R. Duncan and A. Wilce, *Symmetry, compact closure, and dagger-compactness for categories of convex operational models.* arXiv:1004.2920 (2010), to appear in J. Phil. Logic.
- [4] H. Barnum, P. Gaebler and A. Wilce, *Ensemble Steering, weak self-duality and the structure of probabilistic theories*. arXiv:0912.5532v2 (2009).
- [5] H. Barnum and A. Wilce, *Information processing in convex operational theories*. ENTCS 270 (2011), 3–15. doi:10.1016/j.entcs.2011.01.002.
- [6] H. Barnum and A. Wilce, Local tomography, homogeneous self-dual cones, and the structure of quantum theory. To appear.
- [7] G. Chiribella, G. M. D'Ariano, and P. Perinotti, *Probabilistic theories with purification*. Phys. Rev. A. **81** (2010), arXiv:0908.1583v5 (2009), and Informational derivation of quantum theory, arXiv:1011.6451 (2010).
- [8] B. Dakic and C. Brukner, Quantum theory and beyond: is entanglement special? arXiv:0911.0695 (2009).
- [9] J. Faraut and A. Koranyi, Analysis on Symmetric Cones. Oxford, 1994.
- [10] L. Hardy, Quantum theory from five reasonable axioms. arXiv:quant-ph/0101012 (2001)
- [11] M. Koecher, *Die Geodätischen von Positivitätsbereichen*. Math. Ann. **135** (1958), 384–432. doi:10.1007/BF01351796.
- [12] L. Masanes, M. Müller, A derivation of quantum theory from physical requirements. New J. Phys., 13 (2011). doi:10.1088/1367-2630/13/6/063001. arXiv:1004.1483, 2011.
- [13] J. Rau, *On quantum vs. classical probability*. Annals of Physics 324 (2009), 2622–2637. doi:10.1016/j.aop.2009.09.013. arXiv:0710.2119, 2007.
- [14] P. Selinger, *Dagger compact closed categories and completely positive maps*. ENTCS 2007. doi:10.1016/j.entcs.2006.12.018.
- [15] E. Vinberg, *Homogeneous cones*. Sov. Math. Dokl. **1** (1960), 787–790.
- [16] A. Wilce, *Test Spaces*. In D. Gabbay, K. Engesser and D. Lehman (eds.), *The Handbook of Quantum Logic*, volume II, North-Holland, 2010.
- [17] A. Wilce, Symmetry and composition in probabilistic theories. ENTCS 270 (2011), 191–207. doi:10.1016/j.entcs.2011.01.031. arXiv:0910.1527, 2009.
- [18] A. Wilce, *Formalism and interpretation in quantum theory*. Foundations of Physics 40(4): 434–462, 2010. doi:10.1007/s10701-010-9410-x. philsci:3794, 2008.
- [19] A. Wilce, Four and a half axioms for finite-dimensional quantum theory. arXiv:0912.5530 (2009), to appear in a volume in memory of Itamar Pitowsky.
- [20] A. Wilce, Symmetry, Self-Duality and the Jordan Structure of Quantum Mechanics. arXiv:1110.6607v1 (2011).