Early Warning Signals and the Prosecutor's Fallacy

Carl Boettiger^{a,*}, Alan Hastings^b

^aCenter for Population Biology, 1 Shields Avenue, University of California, Davis, CA, 95616 United States. ^bDepartment of Environmental Science and Policy, University of California, Davis

Abstract

1

2

3

4

5

Early warning signals have been proposed to forecast the possibility of a critical transition, 6 such as the eutrophication of a lake, the collapse of a coral reef, or the end of a glacial period. 7 Because such transitions often unfold on temporal and spatial scales that can be difficult to 8 approach by experimental manipulation, research has often relied on historical observations as a 9 source of natural experiments. Here we examine a critical difference between selecting systems for study based on the fact that we have observed a critical transition and those systems for 11 which we wish to forecast the approach of a transition. This difference arises by conditionally selecting systems known to experience a transition of some sort and failing to account for the bias this introduces – a statistical error often known as the Prosecutor's Fallacy. By analysing simulated systems that have experienced transitions purely by chance, we reveal an elevated rate 15 of false positives in common warning signal statistics. We further demonstrate a model-based 16 approach that is less subject to this bias than these more commonly used summary statistics. 17 We note that experimental studies with replicates avoid this pitfall entirely.

Keywords: early warning signals, tipping point, alternative stable states, likelihood methods

20 1. Introduction

21

22

Mathematics ... while assisting the trier of fact in the search of truth, must not cast a spell over him. – California Supreme court, 1968.

In the case of *People v. Collins* 1968, California Supreme Court considered the evidence of an expert witness described by the court as "an instructor of mathematics at a state college", which concluded that the probability that a randomly selected individual would match the description given by the victim would be less than 1 in 12 million (Supreme Court, 1968). The prosecution

*Corresponding author. Email address: cboettig@ucdavis.edu (Carl Boettiger) had produced an individual matching the prosecutor's detailed description, and convinced by
the mathematics, the lower courts had found him guilty.

The prosecution has only observed that the probability of seeing the evidence (E) they produced given a random innocent individual (I), P(E|I) is very small. From this one cannot conclude that the individual is indeed guilty, that is, that the probability the individual is innocent given the evidence P(I|E) is also very small. In a city with millions of people, there might be several individuals who match the description of the evidence. Mathematically P(E|I)need not equal P(I|E), instead, these expressions are related by Bayes theorem,

$$P(E|I) = P(I|E)\frac{P(E)}{P(I)},$$
(1)

³⁵ $P(E) \ll 1$ and $P(I) \approx 1$, so $P(E|I) \approx P(I|E)P(E)$, and consequently we cannot conclude ³⁶ that $P(I|E) \ll 1$ from $P(E|I) \ll 1$. Realizing this mistake, the California Supreme Court ³⁷ reversed the decision, and the case became a widely recognized example of the Prosecutor's ³⁸ Fallacy (Thompson and Schumann, 1987). Here we explore how a similar misconception can ³⁹ arise from the use of historical data to evaluate methods for detecting early warning signals of ⁴⁰ critical transitions.

Catastrophic transitions or tipping points, where a complex system shifts suddenly from one 41 state to another, have been implicated in a wide array of ecological and global climate systems 42 such as lake ecosystems (Carpenter, 2011), coral reefs (Mumby et al., 2007), savannah (Kéfi 43 et al., 2007), fisheries (Berkes et al., 2006), and tropical forests (Hirota et al., 2011). Re-44 cent research has begun to identify statistical patterns commonly associated with these sudden 45 catastrophic transitions which could be used as an *early warning sign* to identify an approaching 46 tipping point, which might provide managers time to react to and avert an undesirable state 47 shift (Scheffer et al., 2009; Lenton, 2011). An array of statistical patterns associated with tip-48 ping point phenomena has been suggested for the detection of early warning signals associated 49 with such sudden transitions. Two of the most commonly used are a pattern of increasing 50 variance (Carpenter and Brock, 2006) and a pattern of increasing autocorrelation (van Nes and 51 Scheffer, 2007), which have been tested in both experimental manipulation (Drake and Griffen, 52 2010; Carpenter, 2011; Veraart et al., 2011; Dai et al., 2012) and historical observations (Livina 53 and Lenton, 2007; Dakos et al., 2008; Lenton et al., 2012; Ditlevsen and Johnsen, 2010; Guttal 54

and Jayaprakash, 2008; Thompson and Sieber, 2010).

⁵⁶ Testing patterns on historical data

Historical examples of sudden transitions taken from the paleo-climate record provide an 57 important way to test and evaluate potential leading indicator methods, and have been widely 58 used for this purpose (Livina and Lenton, 2007; Dakos et al., 2008; Lenton et al., 2012; Ditlevsen 59 and Johnsen, 2010; Guttal and Jayaprakash, 2008; Thompson and Sieber, 2010). Similarly, it 60 has been suggested that data gathered from ecological systems such as lakes that were monitored 61 before they experienced sudden eutrophication, or grasslands subjected to overgrazing, could 62 contain data that could help reveal when similar systems are approaching a tipping point (Car-63 penter, 2011). 64

However, testing methods for early warning signals against historical examples of transitions 65 is susceptible to statistical mistakes that arise from selecting data conditional on that data 66 having already exhibited a sudden transition. A central tenant of early warning theory is that 67 the system in question is slowly approaching a tipping point that lies some unknown distance 68 away. If nothing is done to remedy the situation, this slow change will inevitably carry the 69 system beyond the tipping point, which introduces a sudden, rapid transition into an undesirable 70 state (Scheffer et al., 2009). This process can be described mathematically as a *bifurcation*, in 71 which a slowly changing parameter reaches a critical value that causes the system stability to 72 change. 73

Not all sudden transitions are caused by some "guilty" process slowly driving the system over 74 a tipping point – the kind of process that early warning signals are designed to detect. Some 75 systems may experience such transitions purely by chance, leaving a stable state on an extremely 76 unlikely excursion that happens to stray to far from the stable attractor (e.q.) Ditlevsen and 77 Johnsen, 2010; Lenton, 2011, consider this possibility in transitions that arise from analyzing 78 historical climate record). Like the evidence presented before the California Supreme Court in 79 1968, the chance of observing such an "innocent" transition a priori may be very small, but when 80 selected from a historical record of many possible transitions, this possibility can no longer be 81 ignored. 82

Figure 1 shows a schematic illustrating critical transitions under each of these scenarios. In

the left panel, the system experiences a bifurcation and should contain an early warning signal. In the right panel, a similar-looking trajectory emerges from a simulation of a stable system which should not contain a warning signal. While the simulation of the bifurcation scenario shown on the left produces a similar transition every time, the transition shown on the right is somewhat less likely, occurring in only 1% of simulations.

[Figure 1 about here.]

⁹⁰ 2. Methods and Results

89

To investigate if early warning signals are vulnerable to this fallacy, we simulate a system that 91 is not driven towards a bifurcation such as in Fig reffig:1(b). This simulation approach allows 92 us to determine whether examining historical events is a valid way to test the utility of these 93 indicators. We simulated 20,000 replicates of a stochastic individual-based birth-death process 94 with an Allee threshold (Courchamp et al., 2008), which arises from positive fitness effects at 95 low densities. Above the Allee threshold the population returns to a positive equilibrium size, 96 whereas below the threshold the population decreases to zero. The model can be represented as 97 a continuous time birth-death process where births and deaths are Poisson events which depend 98 on the current density with rates given by 99

$$b(n) = \frac{Kn^2}{n^2 + h^2},$$
(2)

$$d(n) = en + a, (3)$$

a model with a linear death rate and density-dependent birth rate that drives the Allee 100 effect at low densities and limits growth at high densities. In this model n indicates the discrete 101 number of individuals in the population, K indicates a carrying capacity as set by a limiting 102 resource, e a per-capita death rate (the e scaling term in the birth equation allows the carrying 103 capacity K to correspond to a positive equilibrium point), a an additional mortality imposed 104 on the population such as harvest, h is a parameter controlling at what population size the 105 addition of more individuals switches from conferring a positive benefit on growth from Allee 106 interactions n < h to a negative impact on growth due to increased competition, n > h. The 107

key feature of this model is the alternate stable states introduced by this effect; other functional forms for Eq. (2) could serve equally well for these simulations (see *e.g.* Scheffer et al., 2001). Though this system can be forced through a bifurcation by increasing the death rate, in these simulations all parameters are held constant and no bifurcation occurs. Consequently we do not anticipate an early warning signal of an approaching bifurcation.

The simulation starts from the positive equilibrium population size. Though the chance of a transition across the Allee threshold in any given time step is small, given enough time this system will eventually experience such a rare event driving the population extinct. We ran each replicate over 50,000 time units, sampling the system every 50 time units. In this time window 266 of the 1,000 replicates experience population collapse. To keep the examples of comparable sample size, we focus on a section of the data 500 time points prior to the system approaching the transition.

To test whether selecting systems that have experienced spontaneous transitions could bias 120 the analysis towards false positive detection of early warning signals, (the Prosecutor's Fallacy) 121 we selected replicates conditional on having collapsed in the simulations. We then selected a 122 window around each system that ended just before the collapse, while the population values 123 were still above the Allee threshold. For each replicate, we calculated the most common early 124 warning indicators, variance and autocorrelation (e.g. Carpenter and Brock, 2006; Dakos et al., 125 2008; Scheffer et al., 2009), around a moving window equal to half the length of that time series. 126 To test for the presence of a warning signal in these indicators we computed values of 127 Kendall's τ for both indicators for each of the 266 replicates. Kendall's τ is a non-parametric 128 measure of rank correlation frequently used to identify an increasing trend ($\tau > 0$) in early 129 warning signals (Dakos et al., 2008, 2011), defined as $\tau \frac{1}{2}n(n-1)$ in n observations. ¹ τ takes 130 values in (-1, 1). The distribution of τ values observed across these replicates is shown in Fig-131 ure 2. We compare the distribution of τ from all the simulations to the distribution conditioned 132 on experiencing a chance transition to the alternative stable state. To avoid an effect of sample 133 size the time series are all chosen to be the same length. 134

¹A pair of observations (x_i, y_i) and (x_j, y_j) are concordant if $x_i > x_j$ and $y_i > y_j$ or $x_i < x_j$ and $y_i < y_j$ and discordant otherwise; equalities excepted.

To demonstrate the effect we observe is not unique to models with Allee effects, we provide an example of the effect arising in a discrete-time model with two non-zero stable states adapted from (May, 1977),

$$X_{t+1} = X_t \exp\left(r\left(1 - \frac{X_t}{K}\right) - \frac{a * X_t^{Q-1}}{X_t^Q + H^Q}\right).$$
(4)

which combines a logistic growth model with a saturating predator response (See May (1977)
for detailed discussion), shown in Figure 3. Code to replicate the analysis can be found at
https://github.com/cboettig/earlywarning/tree/prosecutor/.

[Figure 3 about here.]

For each of these replicates we also take a model-based approach, estimating parameters for an approximate linear model of the system approaching a saddle node bifurcation, as described by Boettiger and Hastings (2012),

$$dX = \sqrt{r_t}(\phi(r_t) - X_t)dt + \sigma\sqrt{\phi(r_t)}dB_t$$
(5)

In this model, the parameter m describes the approach towards the saddle-node bifurcation. Estimates m < 0 are expected in systems approaching a bifurcation, while for stable systems mshould be approximately zero. None of the estimates across the 266 simulations differed from zero in our study, hence the model-based estimation shows no evidence of bias on data that has been selected conditional on collapse.

¹⁵¹ 3. Discussion

The attempts to detect early warning signs for critical transitions are based on the concept of a deteriorating environment as embodied in a changing parameter Scheffer et al. (2009), which is a different kind of transition than one which is driven instead by stochasticity in an environment which is otherwise constant and exhibiting no directional change. When trying to use historical data to understand critical transitions we often do not know which category, changing environment or simply chance, an observed large change falls into.

We have shown here that systems which undergo rare sudden transitions due to chance look 158 statistically different from their counterparts that do not, even though they are driven by the 159 same stochastic process. In particular, such conditionally selected examples are more likely to 160 show signs associated with an early warning of an approaching tipping point, such as increasing 161 variance or increasing autocorrelation, as measured by Kendall's τ . This increases the risk of 162 false positives – cases in which a warning signal being tested appears to have successfully detected 163 an underlying change in the system leading to a tipping point, when in fact the example comes 164 instead from a stable system with no underlying change in parameters. Figure 2 shows that 165 many of the chance crashes show values of τ that are significantly larger than those observed in 166 the otherwise identical replicates that did not experience a chance transition, thus "detecting" 167 an underlying change in the system dynamics that is not in fact present. 168

¹⁶⁹ 3.1. Chance transitions are false positives for early warning signals

It seems tempting to argue that this bias towards positive detection in historical examples 170 is not problematic – each of these systems did indeed collapse, so the increased probability of 171 exhibiting warning signals could be taken as a successful detection. Unfortunately this is not 172 the case. At the moment the forecast is made, these systems are not likely to transition, since 173 they experience a strong pull towards the original stable state. A closer look at the patterns 174 involved shows why common indicators such as autocorrelation and variance can be misleading. 175 As the system gets farther from its stable point, it it more likely to draw a random step that 176 returns it towards the stable point. Despite this, there is always some probability that it will 177 move further still, so systems that do cross the tipping point must do so rather quickly by a 178 string of events. This pattern, clearly visible before the crashes in each of the examples in Figure 179 1, produces a string of observations that appear more highly autocorrelated (if we are sampling 180 the system frequently enough to catch the excursion at all) than we observe in the rest of the 181 fluctuations around the equilibrium. Yet this autocorrelation comes from a chance trajectory 182 moving quickly away from the stable state, not from the critical slowing down pattern in the 183 return times to the stable state which precede a saddle-node bifurcation and motivate the early 184 warning signal. 185

186

This longer than expected excursion results in a higher than expected variance in that window

as well. Both variance and autocorrelation are calculated using a moving window over the time-187 series, which allows the method to pick out a pattern of change as the window moves along the 188 sequence. If this chance excursion that precedes the crash happens to fill a significant part of 189 the moving window, the resulting pattern will tend to show an increase in autocorrelation or 190 variance. If the chance excursion is relatively rapid compared to the frequency at which the 191 system is observed (spacing of the data) or the width of the moving window, the excursion may 192 not significantly alter the general pattern. In this way, some of the events in which a crash is 193 observed will appear to present these statistical patterns of increased variance or autocorrelation 194 without being harbingers of approaching critical transitions. 195

196 3.2. The truncation of observations

If we had a complete knowledge of the system dynamics, then we could eliminate the bias 197 we observe here since the bias arises from the transient branch of the trajectory that crosses 198 the threshold, and if the system were truncated at the minimum of the potential then the 199 effect we emphasize here would not appear. But, it is not possible to truncate the system in 200 any practical application. The precise location of the minimum of the potential which is the 201 location of the deterministic equilibrium is unknown. Moreover, under the hypothesis that the 202 system is approaching a critical transition, the location of the minimum potential moves so it 203 cannot easily be estimated by previous observations, (see Figure 1c where the equilibrium point 204 moves in the direction of the transition). Thus it is neither practical nor desirable to suggest 205 that historical time series can be used by following a simple truncation rule that avoids the 206 branch of a trajectory crossing the threshold to another basin of attraction. Exactly where a 207 particular study will choose to truncate such a trajectory will necessarily be arbitrary without an 208 underlying model of the process. Frequently this is done by removing the very steep, monotonic 209 branch of the trajectory expected once the system crosses the unstable threshold. Such an 210 approach corresponds with our choice of termination and produces the bias we discuss here. 211

The examples of Figure 1, though only single replicates, may be useful in illustrating these issues. Figure 1c, top panel shows a sample trajectory of a system with a parameter shift, while 1b shows a trajectory without a shift. Both trajectories become more highly autocorrelated and higher variance near the end of the time series (time increases on the y axis in Figure 1). The part of the time series following the critical transition shows a fast and monotonic trajectory to the unstable trajectory, and would usually be excluded by an analysis for warning signals in advance of the transition. No such clear pattern exists prior to the transition in Figure 1b. An alternative proposal to terminate the trajectory in panel B earlier would also risk decreasing the signal seen in panel c, and would be inconsistent with the application of warning signals in the forecasting context, where there would be no such truncation.

222 3.3. Comparing to the model-based method

In our numerical experiment, the model-based estimate of early warning signals appears more 223 robust than the summary statistics, producing the same estimates on both the conditionally 224 selected replicates as on a random sample of the replicates. This is a consequence of the more 225 rigid specifications that come with a model-based approach – the pattern expected is less general 226 than any increase in variance or autocorrelation, but instead must be one that matches its 227 approximation of the saddle-node bifurcation. This observation highlights the difference between 228 the pattern driving the false positive trends in increasing variance and increasing autocorrelation 229 and the pattern anticipated in the saddle-node model. This should not however be taken as 230 evidence that the model-based approach is immune to the bias of the Prosecutor's Fallacy. 231

232 3.4. Importance of experimental approaches

The problem we highlight ultimately stems from the difficulty of having only a single realization with which to examine a complex problem. The only way to deal with this problem embodied is through replication, as can be done in an experimental system in laboratory manipulations such as Drake and Griffen (2010); Veraart et al. (2011); Dai et al. (2012) and at the scale of whole lake ecosystems in Carpenter (2011). Experimental procedures avoid the hazard of the Prosecutor's fallacy by generating a complete sample of replicates, rather than selecting a subset of cases from some larger historical sample.

240 4. Acknowledgments

This research was supported by funding from NSF Grant EF 0742674 to AH and a Computa tional Sciences Graduate Fellowship from the Department of Energy grant DE-FG02-97ER25308

and NERSC Supercomputing grant DE-AC02-05CH11231 to CB. The authors thank M. Baskett, T.A. Perkins and N. Ross for helpful comments on earlier drafts of the manuscript, and
also P. Ditlevsen and an anonymous reviewer for their comments.

246 **References**

- 247 Berkes, F., Hughes, T. P., Steneck, R. S., Wilson, J. A., Bellwood, D. R., Crona, B., Folke, C.,
- Gunderson, L. H., Leslie, H. M., Norberg, J., Nyström, M., Olsson, P., Osterblom, H., Scheffer,
 M., Worm, B., Mar. 2006. Ecology. Globalization, roving bandits, and marine resources.
 Science (New York, N.Y.) 311 (5767), 1557–8.
- Boettiger, C., Hastings, A., May 2012. Quantifying limits to detection of early warning for
 critical transitions. Journal of The Royal Society Interface 9 (75), 2527 2539.
- ²⁵³ Carpenter, J., Feb. 2011. May the Best Analyst Win. Science 331 (6018), 698–699.
- Carpenter, S. R., Brock, W. A., 2006. Rising variance: a leading indicator of ecological transition.
 Ecology letters 9 (3), 311–8.
- ²⁵⁶ Courchamp, F., Berec, L., Gascoigne, J., 2008. Allee Effects in Ecology and Conservation.
 ²⁵⁷ Oxford University Press, USA.
- Dai, L., Vorselen, D., Korolev, K. S., Gore, J., May 2012. Generic Indicators for Loss of Resilience
 Before a Tipping Point Leading to Population Collapse. Science 336 (6085), 1175–1177.
- Dakos, V., Kéfi, S., Rietkerk, M., Nes, E. H. V., Scheffer, M., 2011. Slowing Down in Spatially
 Patterned Ecosystems at the Brink of Collapse. The American Naturalist.
- Dakos, V., Scheffer, M., van Nes, E. H., Brovkin, V., Petoukhov, V., Held, H., Sep. 2008. Slowing
 down as an early warning signal for abrupt climate change. Proceedings of the National
 Academy of Sciences 105 (38), 14308–12.
- Ditlevsen, P. D., Johnsen, S. J., Oct. 2010. Tipping points: Early warning and wishful thinking.
 Geophysical Research Letters 37 (19), 2–5.

- Drake, J. M., Griffen, B. D., Sep. 2010. Early warning signals of extinction in deteriorating
 environments. Nature 467 (7314), 456–459.
- Guttal, V., Jayaprakash, C., Dec. 2008. Spatial variance and spatial skewness: leading indicators
 of regime shifts in spatial ecological systems. Theoretical Ecology 2 (1), 3–12.
- ²⁷¹ Hirota, M., Holmgren, M., Van Nes, E. H., Scheffer, M., Oct. 2011. Global Resilience of Tropical
 ²⁷² Forest and Savanna to Critical Transitions. Science 334 (6053), 232–235.
- ²⁷³ Kéfi, S., Rietkerk, M., Alados, C. L., Pueyo, Y., Papanastasis, V. P., Elaich, A., de Ruiter,
 ²⁷⁴ P. C., Sep. 2007. Spatial vegetation patterns and imminent desertification in Mediterranean
 ²⁷⁵ arid ecosystems. Nature 449 (7159), 213–7.
- Lenton, T. M., Jun. 2011. Early warning of climate tipping points. Nature Climate Change 1 (4),
 201–209.
- Lenton, T. M., Livina, V., Dakos, V., 2012. Early warning of climate tipping points from critical
 slowing down: comparing methods to improve robustness. Philosophical Transactions of The
 Royal Society A (in press).
- Livina, V. N., Lenton, T. M., Feb. 2007. A modified method for detecting incipient bifurcations
 in a dynamical system. Geophysical Research Letters 34 (3), 1–5.
- May, R. M., Oct. 1977. Thresholds and breakpoints in ecosystems with a multiplicity of stable
 states. Nature 269 (5628), 471–477.
- Mumby, P. J., Hastings, A., Edwards, H. J., Nov. 2007. Thresholds and the resilience of
 Caribbean coral reefs. Nature 450 (7166), 98–101.
- Scheffer, M., Bascompte, J., Brock, W. A., Brovkin, V., Carpenter, S. R., Dakos, V., Held, H.,
 van Nes, E. H., Rietkerk, M., Sugihara, G., 2009. Early-warning signals for critical transitions.
 Nature 461 (7260), 53–9.
- Scheffer, M., Carpenter, S. R., Foley, J. A., Folke, C., Walker, B., Oct. 2001. Catastrophic shifts
 in ecosystems. Nature 413 (6856), 591–6.

- ²⁹² Supreme Court, C., 1968. People v. Collins.
- Thompson, J. M. T., Sieber, J., Dec. 2010. Climate tipping as a noisy bifurcation: a predictive
 technique. IMA Journal of Applied Mathematics 76 (1), 27–46.
- Thompson, W. C., Schumann, E. L., 1987. Interpretation of statistical evidence in criminal
 trials: The prosecutor's fallacy and the defense attorney's fallacy. Law and Human Behavior
 11 (3), 167–187.
- van Nes, E. H., Scheffer, M., Jun. 2007. Slow recovery from perturbations as a generic indicator
 of a nearby catastrophic shift. The American naturalist 169 (6), 738–47.
- Veraart, A. J., Faassen, E. J., Dakos, V., van Nes, E. H., Lürling, M., Scheffer, M., Dec. 2011.
- Recovery rates reflect distance to a tipping point in a living system. Nature, 2–5.



Figure 1: The Prosecutor's Fallacy. (a) Plot of the model functions shown in Eq (2) with parameters a = 180, K = 500, e = .5, and h = 200. When the death rate is higher than the birth rate, the system dynamics drive the state (population size) to smaller values. When birth rate is higher, the system moves right, as indicated by the arrows.(b) The potential energy is given by the negative integral of b(n) - d(n), shown in the lower plot. The potential function gives an intuitive picture of the stability of a system by imagining the curve as a surface on which a ball is free to bounce across – wells correspond to stable points and peaks to unstable points. While most trajectories remain near the stable well, some transition out merely by chance. An example of such a trajectory is shown in the top panel, in which time increases along the vertical axis. Though initially oscillating around the stable state, a chance excursion carries it beyond the Allee threshold (vertical dotted line). Such chance trajectories can produce the statistical patterns as observed in true critical transitions seen in panel (c): Early warning signals are aimed at detecting systems which are slowly moving towards a tipping point or bifurcation, illustrated in the successive curves (deteriorating and critical). Top panel: An example trajectory from a simulation under this process shows the state of the system as the potential moves towards the bifurcation point. The original position of the Allee threshold is shown by the vertical dotted line (though it moves slightly as the parameter changes).



Figure 2: The distribution of the correlation statistic τ for two early warning indicators (variance, autocorrelation) on replicates conditionally selected for having collapsed by chance in simulations is shown in grey bars. Solid lines indicate the estimated density of the statistic from a random sample of the simulations (not conditional on observing a transition). Positive values of τ correspond to a pattern of an indicator increasing with time; typically taken as evidence that a system is approaching a critical transition. In these simulations, the pattern arises instead from the Prosecutor's fallacy of conditional selection.



Figure 3: The identical analysis from Figure 2 is shown for the model in Eq (4) using parameters r = 0.75, K = 10, a = 1.7, Q = 3, and H = 1. A similar statistical bias, particularly towards positive values of *tau* occurs in this model as well.