

Note on divisible difference sets from Galois rings $GR(9, n)$

Koji Momihara*

Abstract

In this note, we give a new construction of divisible difference sets in \mathbb{Z}_9^n using Galois ring $GR(3^2, n)$ under the assumption of the existence of skew Hadamard difference sets in \mathbb{F}_{3^n} .

Keywords: divisible difference set; skew Hadamard difference set

1 Introduction

We assume that the reader is familiar with the basic theory of difference sets as can be found in [2].

Let G be a group of order $v = mn$. A k -subset of G is called a *divisible difference set* with parameters (m, n, k, λ, μ) if the list of differences $\{xy^{-1} \mid x, y \in D; x \neq y\}$ covers every element in $G \setminus N$ exactly μ times and the elements in $N \setminus \{e\}$ exactly λ times, where N is a specified subgroup of G of order n . For detailed descriptions of divisible difference sets, see [13]. Also, for known constructions of divisible difference sets, see [3, 7, 9, 8, 10].

The definition of divisible difference sets immediately yields the group ring equation

$$DD^{(-1)} = k + \lambda(N - e) + \mu(G - N),$$

where $D = \sum_{x \in D} x \in \mathbb{Z}[G]$ and $D^{(-1)} = \sum_{x \in D} x^{-1}$. We now restrict our attention to abelian groups. In this case, it is known [14] that D is a divisible difference set in G with parameters (m, n, k, λ, μ) if and only if for any nontrivial character χ of G

$$|\chi(D)| = \begin{cases} \sqrt{k - \lambda} & \text{if } \chi \text{ is nontrivial on } N; \\ \sqrt{k - \lambda + (\lambda - \mu)n} & \text{if } \chi \text{ is trivial on } N \text{ but nontrivial on } G. \end{cases}$$

A difference set D in a group G is called *skew Hadamard* if G is the disjoint union of D , $D^{(-1)}$, and $\{e\}$. The primary example (and for many years, the only known example in abelian groups) of skew Hadamard difference sets is the classical Paley (quadratic residue) difference set in $(\mathbb{F}_q, +)$ consisting of the nonzero squares of \mathbb{F}_q , where \mathbb{F}_q is the finite field

*Department of Mathematics, Faculty of Education, Kumamoto University, 2-40-1 Kurokami, Kumamoto 860-8555, Japan, (momihara@educ.kumamoto-u.ac.jp).

of order q , a prime power congruent to 3 modulo 4. In the case where G is an (additively written) abelian group of order v , it is known [14] that a subset D such that $D \cup -D \cup \{0\}$ in G is a skew Hadamard difference set if and only if

$$\chi(D) = \frac{-1 \pm \sqrt{-v}}{2}$$

for any nontrivial character χ of G .

Skew Hadamard difference sets are currently under intensive study, see [4, 5, 6, 11, 12, 16, 17]. There was a major conjecture in this area: Up to equivalence the Paley difference sets mentioned above are the only skew Hadamard difference sets in abelian groups. This conjecture turned out to be false: Ding and Yuan [4] constructed a family of skew Hadamard difference sets in $(\mathbb{F}_{3^m}, +)$, where $m \geq 3$ is odd, and showed that two examples in the family are inequivalent to the Paley difference sets. Very recently, Muzychuk [11] constructed infinitely many inequivalent skew Hadamard difference sets in an elementary abelian group of order q^3 . Also, the author gave a construction of inequivalent skew Hadamard difference sets in \mathbb{F}_q using cyclotomic strongly regular graphs [12].

In this note, we give a new construction of divisible difference sets in the 3-groups \mathbb{Z}_9^n using Galois ring $GR(3^2, n)$ under the assumption of the existence of skew Hadamard difference sets in \mathbb{F}_{3^n} .

2 Group characters of $GR(p^2, n)$

Let $\mu_1 : \mathbb{Z}/p^2\mathbb{Z} \rightarrow \mathbb{Z}/p\mathbb{Z}$ be the reduction map modulo p . We can extend μ_1 to $(\mathbb{Z}/p^2\mathbb{Z})[x]$ in the natural way. Let $f(x)$ be a primitive polynomial of degree n over \mathbb{F}_p and $g(x) \in \mathbb{Z}/p^2\mathbb{Z}$ be preimage of $f(x)$ under μ_1 . There is a unique $g(x)$ whose root ξ satisfies $\xi^{p^n-1} = 1$. The ring $\mathbb{R}_n = (\mathbb{Z}/p^2\mathbb{Z})(\xi)$ is an algebraic extension of $\mathbb{Z}/p^2\mathbb{Z}$ of degree n , which is called a *Galois ring* and denoted by $GR(p^2, n)$. The ring $GR(p^2, n)$ has the unique maximal ideal $p\mathbb{R}_n = \{p\xi^i \mid 0 \leq i \leq p^n - 1\} \cup \{0\}$ and the residue class field $\mathbb{R}_n/p\mathbb{R}_n$ is isomorphic to \mathbb{F}_{p^n} . We can take Teichmüller system $\mathbb{T}_n = \{0, 1, \xi, \dots, \xi^{p^n-2}\}$ as a set of representatives of $\mathbb{R}_n/p\mathbb{R}_n$. An arbitrary element $\alpha \in \mathbb{R}_n$ is uniquely determined by $\alpha = \alpha_0 + p\alpha_1$, $\alpha_0, \alpha_1 \in \mathbb{T}_n$. We denote the set of all invertible elements in \mathbb{R}_n by \mathbb{R}_n^* . Every element of \mathbb{R}_n^* is uniquely represented as $\alpha_0(1 + p\alpha_1)$, $\alpha_0 \in \mathbb{T}_n \setminus \{0\}$, $\alpha_1 \in \mathbb{T}_n$.

The map $\tau(\alpha) = \alpha_0$ is a homomorphism of \mathbb{R}_n into the cyclic group $\mathbb{T}_n \setminus \{0\}$ and $(\mathbb{P}_n :=) Ker(\tau)$ is the group of principal units, namely elements of the form $1 + p\alpha_1 \in \mathbb{R}_n^*$. The τ is precisely given by $\tau = \alpha^{p^n}$. It is obvious that $\overline{x + y} = \overline{x} + \overline{y} = \tau(x + y)$ and $\overline{x \cdot y} = \overline{x} \cdot \overline{y} = \tau(xy)$ for any $x, y \in \mathbb{R}_n$, where \overline{x} means the residue class of $\mathbb{R}_n/p\mathbb{R}_n$ containing x . The set of principal units has the structure of an elementary abelian group of order p^n and is isomorphic to the additive group of $\mathbb{R}_n/p\mathbb{R}_n (\simeq \mathbb{F}_{p^n})$ via the map $1 + p\alpha_1 \mapsto \overline{\alpha_1}$. Thus, \mathbb{R}_n^* is a direct product of a cyclic group of order $p^n - 1$ and an elementary abelian group of order p^n . Hereafter, we identify $\mathbb{R}_n/p\mathbb{R}_n$ with \mathbb{F}_q and denote each element of $\mathbb{R}_n/p\mathbb{R}_n$ and \mathbb{F}_q by \overline{a} in common.

The Frobenius map f from \mathbb{R}_n to \mathbb{R}_n is the ring automorphism $f : \alpha_0 + p\alpha_1 \rightarrow \alpha_0^p + p\alpha_1^p$, $\alpha_0, \alpha_1 \in \mathbb{T}_n$. The set of elements of \mathbb{R}_n invariant under f is identical with $\mathbb{Z}/p^2\mathbb{Z}$. For $\alpha \in \mathbb{R}_n$,

the trace function from \mathbb{R}_n to $\mathbb{Z}/p^2\mathbb{Z}$ is defined by

$$\mathrm{Tr}_{\mathbb{R}_n}(\alpha) = \sum_{i=0}^{n-1} \alpha^{f^i}.$$

This is a linear function of \mathbb{R}_n into $\mathbb{Z}/p^2\mathbb{Z}$. The additive characters ϕ_β of \mathbb{R}_n can be described by the trace function as follows:

$$\phi_\beta(\alpha) = \zeta_{p^2}^{\mathrm{Tr}_{\mathbb{R}_n}(\beta\alpha)} \text{ for any } \alpha \in \mathbb{R}_n,$$

where $\zeta_k = \exp(\frac{2\pi i}{k})$. The ϕ_β 's for $\beta \in \mathbb{R}_n$ give all additive characters of \mathbb{R}_n , see [15] for the proof.

A multiplicative character ψ of \mathbb{R}_n^* is defined by

$$\psi(\alpha\beta) = \psi(\alpha)\psi(\beta),$$

and each value of $\psi(\alpha)$ is a $p^n(p^n - 1)$ th root of unity. Let ψ_P denote a multiplicative character of \mathbb{R}_n trivial on $\mathbb{T}_n \setminus \{0\}$, which induces a character of \mathbb{P}_n . Also, let ψ_T denote a multiplicative character of \mathbb{R}_n trivial on \mathbb{P}_n , which induces a character of $\mathbb{T}_n \setminus \{0\}$. It is clear that any multiplicative character of \mathbb{R}_n is uniquely determined as a product $\psi_P\psi_T$. Put $\alpha = a(1 + pb) \in \mathbb{R}_n^*$. Obviously, the multiplicative character $\psi_P = \psi_{P,\ell}$ for $\ell \in \mathbb{T}_n$ of \mathbb{R}_n is given by

$$\psi_{P,\ell}(\alpha) = \psi_{P,\ell}(1 + pb) = \chi_{\bar{\ell}}(\bar{b}),$$

where $\chi_{\bar{\ell}}$ is an additive character of \mathbb{F}_{p^n} defined by

$$\chi_{\bar{\ell}}: \mathbb{F}_q \rightarrow \mathbb{C}^*, \quad \chi_{\bar{\ell}}(\bar{x}) = \zeta_p^{\mathrm{Tr}_{\mathbb{F}_q/\mathbb{F}_p}(\bar{\ell}x)},$$

where $\mathrm{Tr}_{q/p}$ is the trace from \mathbb{F}_q to \mathbb{F}_p . Also, the multiplicative character $\psi_T = \psi_{T,m}$ for $0 \leq m \leq p^n - 2$ of \mathbb{R}_n is given by

$$\psi_{T,m}(\alpha) = \psi_{T,m}(a) = \lambda_{q-1}^m(\bar{a}),$$

where λ_{q-1} is the multiplicative character of \mathbb{F}_{p^n} of order $q - 1$ define by

$$\lambda_{q-1}: \mathbb{F}_q^* \rightarrow \mathbb{C}^*, \quad \lambda_{q-1}^m(\bar{\xi}^s) = \zeta_{q-1}^{sm}.$$

We extend the domain of ψ to all elements of \mathbb{R}_n by defining as $\psi_{P,\ell}(\alpha) := 0$ for any $\alpha \in p\mathbb{R}_n$ and any $\ell \in \mathbb{T}_n$, and as $\psi_{T,m}(\alpha) := 0$ for any $\alpha \in p\mathbb{R}_n$ and any $1 \leq m \leq p^n - 2$, and as $\psi_{T,0}(\alpha) := 1$ for any $\alpha \in p\mathbb{R}_n$.

Hereafter, we use the following notations.

- \mathbb{R}_n : the Galois ring $GR(p^2, n)$
- \mathbb{T}_n : the Teichmüller system of \mathbb{R}_n
- ϕ_1 : the canonical additive character of \mathbb{R}_n
- ψ : a multiplicative character of \mathbb{R}_n such that $\psi = \psi_P\psi_T$
- $\chi_{\bar{1}}$: the canonical additive character of \mathbb{F}_{p^n}
- λ : a multiplicative character of \mathbb{F}_{p^n}

Similar to Gauss sums of $\mathbb{F}_q [1]$, we can define Gauss sums of \mathbb{R}_n . For a multiplicative character $\psi = \psi_P \psi_T$ and a nontrivial additive character ϕ_β of \mathbb{R}_n , the sum

$$G(\phi_\beta, \psi) = \sum_{\alpha \in \mathbb{R}_n} \phi_\beta(\alpha) \psi(\alpha)$$

is called a *Gauss sum* of $GR(p^2, n)$. Note that $G(\phi_\beta, \psi) = \sum_{\alpha \in \mathbb{R}_n^*} \phi_\beta(\alpha) \psi(\alpha)$ by the definition of ψ .

3 Divisible difference sets over $GR(9, n)$

In the following, we compute the exact value of $G(\phi_1, \psi)$.

Lemma 3.1. *Let $\psi = \psi_{P,\ell} \psi_{T,m}$. Then it holds that*

$$G(\phi_1, \psi) = \begin{cases} 0 & \text{if } \psi_{P,\ell} \text{ is trivial;} \\ p^n \phi_1(c) \psi_{T,m}(c) & \text{otherwise,} \end{cases}$$

where $c \in \mathbb{T}_n \setminus \{0\}$ is determined by $\bar{c} = -\bar{\ell}$ in \mathbb{F}_{p^n} .

Proof: By the definition of $G(\phi_1, \psi)$, it holds that

$$\begin{aligned} G(\phi_1, \psi) &= \sum_{\alpha = a(1+pb) \in \mathbb{R}_n^*} \phi_1(\alpha) \psi(\alpha) \\ &= \sum_{a \in \mathbb{T}_n \setminus \{0\}} \sum_{b \in \mathbb{T}_n} \phi_1(a(1+pb)) \psi_{P,\ell}(1+pb) \psi_{T,m}(a). \end{aligned}$$

Since

$$\phi_1(a(1+pb)) = \zeta_{p^2}^{\text{Tr}_{\mathbb{R}_n}(a+pub)} = \zeta_{p^2}^{\text{Tr}_{\mathbb{R}_n}(a)} \zeta_{p^2}^{p \text{Tr}_{\mathbb{F}_{p^n}}(\bar{a}b)} = \phi_1(a) \chi_{\bar{\Gamma}}(\bar{a}b),$$

we have

$$\begin{aligned} G(\phi_1, \psi) &= \sum_{a \in \mathbb{T}_n \setminus \{0\}} \sum_{b \in \mathbb{T}_n} \phi_1(a) \chi_{\bar{\Gamma}}(\bar{a}b) \chi_{\bar{\Gamma}}(\bar{b}\ell) \psi_{T,m}(a) \\ &= \sum_{a \in \mathbb{T}_n \setminus \{0\}} \phi_1(a) \psi_{T,m}(a) \sum_{\bar{b} \in \mathbb{F}_{p^n}} \chi_{\bar{\Gamma}}((\bar{a} + \ell)\bar{b}) \\ &= \begin{cases} 0 & \text{if } \ell = 0; \\ p^n \phi_1(c) \psi_{T,m}(c) & \text{otherwise,} \end{cases} \end{aligned}$$

where $c \in \mathbb{T}_n$ is determined by $\bar{c} = -\bar{\ell}$ in \mathbb{F}_{p^n} . □

Theorem 3.2. *Let $p = 3$ and n be an odd integer. Furthermore, let $b \in \mathbb{T}_n \setminus \{0\}$ such that $\text{Tr}_{\mathbb{F}_q/\mathbb{F}_p}(\bar{b}) \neq 0$ and C be a skew Hadamard difference set in \mathbb{F}_q . Then, the set $D = \{x(1+py) \mid \bar{x} \in C, y \in T_{b,0}\} \leq \mathbb{R}_n^*$ forms a $(3^n, 3^n, \frac{3^{n-1}(3^n-1)}{2}, \frac{3^{n-1}(3^{n-1}-1)}{2}, \frac{3^{n-1}(3^{n-1}-1)}{4})$ divisible difference set in \mathbb{R}_n relative to $p\mathbb{R}_n$, where*

$$T_{b,i} = \{x \in \mathbb{T}_n \mid \text{Tr}_{\mathbb{F}_q/\mathbb{F}_p}(\bar{b}x) = i\}.$$

Proof: It is clear that the parameters $v = |\mathbb{R}_n|$, $n = |p\mathbb{R}_n|$, $k = \frac{3^{n-1}(3^n-1)}{2}$, $\mu = \frac{3^{n-1}(3^{n-1}-1)}{2}$, $\lambda = \frac{3^{n-1}(3^{n-1}-1)}{4}$ satisfy the equation

$$(v - n)\lambda + (n - 1)\mu = k(k - 1).$$

We prove that

$$|\phi_1(aD)| = \begin{cases} \frac{3^{n-1}\sqrt{1+3^n}}{2} & \text{if } a \in p\mathbb{R}_n^*; \\ 3^{n-1} & \text{if } a \in \mathbb{R}_n^*. \end{cases}$$

(i) For $a \in p\mathbb{R}_n^*$, write $a = p\xi^r$. Note that $\phi_1(p\xi^r) = \chi_{\overline{1}}(\overline{\xi^r})$ and $\sum_{\overline{x} \in C} \chi_{\overline{1}}(\overline{\xi^r x}) = \frac{-1 \pm \sqrt{-3^n}}{2}$ by the definition of C . Hence, we have

$$\phi_1(aD) = |T_{b,0}| \cdot \sum_{\overline{x} \in C} \chi_{\overline{1}}(\overline{\xi^r x}) = \frac{3^{n-1}(-1 \pm \sqrt{-3^n})}{2},$$

which implies that $|\phi_1(aD)| = \frac{3^{n-1}\sqrt{1+3^n}}{2}$.

(ii) For $a \in \mathbb{R}_n^*$, write $a = \xi^s(1 + p\xi^r)$. By the orthogonality of characters,

$$\begin{aligned} \phi_1(aD) &= \frac{1}{q(q-1)} \sum_{\psi \in \widehat{\mathbb{R}_n^*}} G(\phi_1, \psi^{-1}) \psi(aD) \\ &= \frac{1}{3(q-1)} \sum_{\psi \in H} \sum_{x \in D} G(\phi_1, \psi^{-1}) \psi(ax), \end{aligned}$$

where H is the subgroup of $\widehat{\mathbb{R}_n^*}$, the character group of \mathbb{R}_n^* , such that each of H is trivial on $\{a(1 + pb) \mid a \in \mathbb{T}_n \setminus \{0\}, b \in T_{b,0}\}$. The $\psi(ax)$ for $\psi \in H$ and $x = \xi^y(1 + p\xi^z) \in D$ can be written as $\lambda_{q-1}^m(\overline{\xi^s \xi^y}) \chi_{\overline{b}}(\overline{c \xi^r})$ for some $m \in \{0, 1, \dots, q-2\}$ and $\overline{c} \in \mathbb{F}_3$. Hence, by Lemma 3.1, we have

$$|\phi_1(aD)| = \frac{3^{n-1}}{q-1} \cdot \left| \phi_1(1) \chi_{\overline{b}}(\overline{\xi^r}) \left(\sum_{m=0}^{q-2} \lambda_{q-1}^m(\overline{\xi^s C}) \right) + \phi_1(\xi^{\frac{3^n-1}{2}}) \chi_{\overline{b}}(\overline{-\xi^r}) \left(\sum_{m=0}^{q-2} \lambda_{q-1}^m(\overline{-\xi^s C}) \right) \right|. \quad (1)$$

Since C is skew Hadamard, either one of $\sum_{m=0}^{q-2} \lambda_{q-1}^m(\overline{\xi^s C})$ or $\sum_{m=0}^{q-2} \lambda_{q-1}^m(\overline{-\xi^s C})$ is zero and another is $q-1$, the (1) is equal to 3^{n-1} , which completes the proof. \square

As described in Introduction, a lot of inequivalent skew Hadamard difference sets in \mathbb{F}_q were found recently. One may expect that we obtain many inequivalent divisible difference sets in \mathbb{Z}_q^n applying the construction of Theorem 3.2 to such inequivalent skew Hadamard difference sets. We close this note providing the following open problem.

Problem 3.3. *Determine whether or not divisible difference sets constructed in Theorem 3.2 are equivalent if the assumed skew Hadamard difference sets are inequivalent.*

Acknowledgments

This work was partially supported by JSPS under Grant-in-Aid for Research Activity start-up 23840032.

References

- [1] B. Berndt, R. Evans, K. S. Williams, *Gauss and Jacobi Sums*, Wiley, (1997).
- [2] T. Beth, D. Jungnickel, H. Lenz, *Design Theory*, vol. I, 2nd ed., Encyclopedia of Mathematics and its Applications 78, Cambridge University Press, Cambridge, (1999).
- [3] J. A. Davis, New semiregular divisible difference sets, *Discr. Math.*, **188** (1998), 99-109.
- [4] C. Ding, J. Yuan, A family of skew Hadamard difference sets, *J. Combin. Theory, Ser. A*, **113** (2006), 1526–1535.
- [5] C. Ding, Z. Wang, Q. Xiang, Skew Hadamard difference sets from the Ree-Tits slice symplectic spreads in $PG(3, 3^{2h+1})$, *J. Combin. Theory, Ser. A*, **114** (2007), 867–887.
- [6] T. Feng, Q. Xiang, Cyclotomic constructions of skew Hadamard difference sets, *J. Combin. Theory, Ser. A*, **119** (2012), 245–256.
- [7] Y. J. Ionin, A technique for constructing divisible difference sets, *J. Geom.*, **67** (2000), 164–172.
- [8] K. H. Leung, S. Ling, S. L. Ma, K. B. Tay, Almost perfect sequences with $\theta = 2$, *Arch. Math.*, **70** (1998), 128–131.
- [9] K. Liu, *Divisible difference sets and difference sets from cyclotomy*, Ph.D. Thesis, The Ohio State University, (1999).
- [10] S. L. Ma, S. K. Sehgal, A family of semi-regular divisible difference sets, *Des. Codes Cryptogr.*, **11** (1997), 73–78.
- [11] M. E. Muzychuk, On skew Hadamard difference sets, arXiv:1012.2089.
- [12] K. Momihara, Skew Hadamard difference sets from cyclotomic strongly regular graphs, arXiv:1207.2197.
- [13] A. Pott, *Finite geometry and character theory*, Springer, New York, Heidelberg, Berlin, (1995).
- [14] R. J. Turyn, Character sums and difference sets, *Pacific J. Math.*, **15** (1965), 319–346.
- [15] K. Yamamoto, M. Yamada, Hadamard difference sets over an extension of $Z/4Z$, *Util. Math.*, **34** (1988), 169–178.
- [16] G. B. Weng, L. Hu, Some results on skew Hadamard difference sets, *Des. Codes Cryptogr.*, **50** (2009), 93–105.
- [17] G. B. Weng, W. S. Qiu, Z. Wang, Q. Xiang, Pseudo-Paley graphs and skew Hadamard difference sets from presemifields, *Des. Codes Cryptogr.*, **44** (2007), 49–62.