1	A phylogeny of birds based on over 1,500 loci collected by target enrichment and
2	high-throughput sequencing
3	
4	John E. McCormack,* ¹ Michael G. Harvey, ^{1,2} Brant C. Faircloth, ³ Nicholas G. Crawford, ⁴ Travis
5	C. Glenn ⁵ , & Robb T. Brumfield ^{1,2}
6	
7	¹ Museum of Natural Science, Louisiana State University, Baton Rouge, LA 70803, USA
8	² Department of Biological Sciences, Louisiana State University, Baton Rouge, LA 70803, USA
9	³ Department of Ecology and Evolutionary Biology, University of California, Los Angeles, CA
10	90095, USA
11	⁴ Department of Biology, Boston University, Boston, MA 02215, USA
12	⁵ Department of Environmental Health Science, University of Georgia, Athens, GA 30602, USA
13	
14	
15	
16	
17	
18	
19	
20	
21	*Corresponding author: John McCormack, Moore Laboratory of Zoology, Occidental College,
22	1600 Campus Rd., Los Angeles, CA 90041 Phone: 323-259-1352 E-mail: mccormack@oxy.edu
23	

24 ABSTRACT

Evolutionary relationships among birds in Neoaves, the clade comprising the vast majority of 25 avian diversity, have vexed systematists due to the ancient, rapid radiation of numerous lineages. 26 We applied a new phylogenomic approach to resolve relationships in Neoaves using target 27 enrichment (sequence capture) and high-throughput sequencing of ultraconserved elements 28 (UCEs) in avian genomes. We collected sequence data from UCE loci for 32 members of 29 Neoaves and one outgroup (chicken) and analyzed data sets that differed in their amount of 30 missing data. An alignment of 1,541 loci that allowed missing data was 87% complete and 31 32 resulted in a highly resolved phylogeny with broad agreement between the Bayesian and maximum-likelihood (ML) trees. Although results from the 100% complete matrix of 416 UCE 33 loci were similar, the Bayesian and ML trees differed to a greater extent in this analysis, 34 suggesting that increasing from 416 to 1,541 loci led to increased stability and resolution of the 35 tree. Novel results of our study include surprisingly close relationships between phenotypically 36 divergent bird families, such as tropicbirds (Phaethontidae) and the sunbittern (Eurypygidae) as 37 well as between bustards (Otididae) and turacos (Musophagidae). This phylogeny bolsters 38 support for monophyletic waterbird and landbird clades and also strongly supports controversial 39 40 results from previous studies, including the sister relationship between passerines and parrots and the non-monophyly of raptorial birds in the hawk and falcon families. Although significant 41 challenges remain to fully resolving some of the deep relationships in Neoaves, especially among 42 43 lineages outside the waterbirds and landbirds, this study suggests that increased data will yield an increasingly resolved avian phylogeny. 44

45 The diversification of modern birds occurred extremely rapidly, with all major orders and most families becoming distinct within a short window of 0.5 to 5 million years around the 46 Cretaceous-Tertiary boundary [1-4]. As with other cases of ancient, rapid radiation, resolving 47 deep evolutionary relationships in birds has posed a significant challenge. Some authors have 48 hypothesized that the initial splits within Neoaves might be a hard polytomy that will remain 49 irresolvable even with expanded data sets (reviewed in [5]). However, several recent studies 50 have suggested that expanded genomic and taxonomic coverage will lead to an increasingly 51 resolved avian tree of life [2,6,7]. 52

Using DNA sequence data to reconstruct rapid radiations like the Neoaves phylogeny 53 presents a practical challenge on several fronts. First, short speciation intervals provide little time 54 for substitutions to accrue on internal branches, reducing the phylogenetic signal for rapid 55 56 speciation events. Traditionally, the solution to this problem has been to collect additional sequence data, preferably from a rapidly evolving molecular marker such as mitochondrial DNA 57 [8]. However, rapidly evolving markers introduce a new set of problems to the inference of 58 ancient radiations: through time, substitutions across rapidly evolving markers overwrite older 59 substitutions, resulting in signal saturation and homoplasy [9]. To address this challenge, some 60 researchers have inferred ancient phylogeny using rare genomic changes, like retroposon 61 insertions and indels, because rare changes are unlikely to occur in the same way multiple times, 62 thereby minimizing homoplasy [10,11]. Though successful in some cases [12], retroposons are 63 64 often insufficiently numerous to fully resolve relationships between taxa that rapidly radiated [13], and although often billed as being homoplasy-free, we now know that shared retroposon 65 insertions can be due to independent events [14]. 66

67 A second challenge to reconstructing ancient, rapid radiations is the randomness inherent to the process of gene sorting (i.e., coalescent stochasticity), which occurs even when gene 68 histories are estimated with 100% accuracy [15]. The amount of conflict among gene-tree 69 topologies due to coalescent stochasticity increases as speciation intervals get shorter [16]. 70 Hemiplasy refers to gene-tree discord deep in phylogenies resulting from stochastic sorting 71 processes that occurred long ago, but where the alleles are now fully sorted [17]. Accounting for 72 hemiplasy requires increasing the number of loci interrogated and analyzing the resulting 73 sequence data using species-tree methods that accommodate discordant gene histories [18-20]. 74 75 Despite these challenges, our understanding of Neoaves phylogeny has steadily improved as genomic coverage and taxonomic coverage have increased [21]. Hackett et al. [6] – based on 76 169 species and 19 loci – provided a more resolved phylogeny of all birds than ever before. 77 Combined with other studies during the previous decade, we now have a resolved backbone for 78 the avian tree of life, including three well-supported clades: Neoaves, Palaeognathae (e.g., 79 ostrich, emu, tinamous) and Galloanserae (e.g., ducks and chickens) [2,6,22-25]. Nonetheless, 80 many relationships within Neoaves remain challenging to resolve despite the application of 81 molecular tools such as whole mitochondrial genomes [26-28] and rare genomic changes [12-82 14,29]. Specifically, many of the basal nodes and the evolutionary affinities of enigmatic 83 84 lineages (e.g., tropicbirds, hoatzin, sunbittern/kagu) within Neoaves continue to be poorly supported even when addressed with large data sets comprising a variety of molecular markers. 85 86 This raises the question: Are there certain relationships deep in the Neoaves phylogeny that cannot be resolved regardless of the scope of the data collected? 87 Here, we apply a new method for collecting large amounts of DNA sequence data to 88

89 address evolutionary relationships in Neoaves. This method, which involves simultaneous

90 capture and high-throughput sequencing of hundreds of loci, addresses the main challenges of resolving ancient, rapid radiations - and is applicable throughout the tree of life. The markers we 91 target are anchored by ultraconserved elements (UCEs), which are short stretches of highly 92 93 conserved DNA. UCEs were originally discovered in mammals [30], but are also found in a wide range of other organisms [31-33]. UCEs allow for the convenient isolation and capture of 94 independent loci among taxonomically distant species while providing phylogenetic signal in 95 flanking regions [33,34]. Because variation in the flanks increases with distance from the core 96 UCE, these markers display a balance between having a high enough substitution rate while 97 minimizing saturation, providing information for estimating phylogenies at multiple evolutionary 98 timescales [33,35]. UCEs are rarely found in duplicated genomic regions [36], making the 99 determination of orthology more straightforward than in other markers (e.g., exons) or whole 100 101 genomes, and UCEs are numerous among distantly related taxa, facilitating their use as discrete loci in species-tree analysis [33,35]. We employed sequence capture (i.e., bait-capture or target 102 enrichment) to collect UCE sequence data from genomic DNA of 32 non-model bird species 103 (Fig. 1) and used outgroup UCE data from the chicken genome to reconstruct evolutionary 104 relationships in Neoaves. 105

106

107 METHODS

108 We extracted DNA from tissue samples of 32 vouchered museum specimens (Table 1; Fig. 1),

109 each from a different family within the traditional Neoaves group [37], using a phenol-

110 chloroform protocol [38]. All samples for this project were loaned by, and used with permission

111 of, the Louisiana State University Museum of Natural Science. We prepared sequencing libraries

112 from purified DNA using Nextera library preparation kits (Epicentre Biotechnologies, Inc.),

113 incorporating modifications to the protocol outlined in Faircloth et al. [33]. Briefly, following limited-cycle (16-19 cycles) PCR to amplify libraries for enrichment and concentration of 114 amplified libraries to 147 ng/ μ L using a Speed-Vac, we individually enriched libraries for 2,386 115 UCE loci using 2,560 synthetic RNA capture probes (MyBaits, Mycroarray, Inc.). We designed 116 capture probes targeting UCE loci that had high sequence identity between lizards and birds 117 because previous work indicated that UCE loci from this set were useful for deep-level avian 118 phylogenetics [33]. Following enrichment, we incorporated a custom set of indexed, Nextera 119 adapters to each library [39] using enriched product as template in a limited-cycle PCR (16 120 121 cycles), and we sequenced equimolar pools of enriched, indexed libraries using $1\frac{1}{2}$ lanes of single-end, 100 bp sequencing on an Illumina Genome Analyzer IIx (LSU Genomics Facility). 122 The LSU Genomics Facility demultiplexed pooled reads following the standard Illumina 123 124 pipeline, and we combined demultiplexed reads from each run for each taxon prior to adapter trimming, quality filtering, and contig assembly. 125 We filtered reads for adapter contamination, low-quality ends, and ambiguous bases 126 using an automated pipeline (https://github.com/faircloth-lab/illumiprocessor) that incorporates 127 Scythe (https://github.com/vsbuffalo/scythe) and Sickle (https://github.com/najoshi/sickle). We 128

assembled reads for each taxon using Velvet v1.1.04 [40] and VelvetOptimiser v2.1.7 (S

130 Gladman; http://bioinformatics.net.au/software.shtml), and we computed coverage across UCEs

using tools from the AMOS package, as described in [33]. We used the PHYLUCE software

132 package (<u>https://github.com/faircloth-lab/phyluce</u>; version m1.0-final) to align assembled contigs

back to their associated UCE loci, remove duplicate matches, create a taxon-specific database of

134 contig-to-UCE matches, and include UCE loci from the chicken (Gallus gallus) genome as

135 outgroup sequences. We then generated two alignments across all taxa: one containing no

missing data (i.e., all loci required to be present in all taxa) and one allowing up to 50% of the 136 species to have data missing for a given locus. We built alignments using MUSCLE [41]. The 137 steps specific to this analysis are available from https://gist.github.com/47e03463db0573c4252f. 138 For both alignments (missing data and no missing data), we prepared a concatenated 139 alignment for MrBayes v3.1.2 [42] by estimating the most-likely finite-sites substitution model 140 for individual UCE loci. Using a parallel implementation of MrAIC from the PHYLUCE 141 package, we selected the best-fitting substitution model for all loci using AICc, and we grouped 142 loci having the same substitution model into partitions. We assigned the parent substitution 143 144 model to each partition, for a total of 20 partitions, and we analyzed these alignments using two independent MrBayes runs (4 chains) of 10M iterations each (thinning=100). We sampled 145 50,000 trees from the posterior distribution (burn-in=50%) after convergence by ensuring the 146 average standard deviation of split frequencies was < 0.00001 and the potential scale reduction 147 factor for estimated parameters was approximately 1.0. We confirmed convergence with 148 Effective Sample Size values >200 in TRACER [43] and by assessing the variance in tree 149 topology with AWTY [44]. We also prepared a concatenated alignment in PHYLIP format with 150 a single partition containing all sequence data, and we analyzed this alignment using the fast-151 approximation, maximum likelihood (ML) algorithm in RaXML (raxmlHPC-MPI-SSE3; v. 152 (7.3.0) with 1,000 bootstrap replicates [45,46]. 153 For the data set with no missing data, we also estimated a species tree on 250 nodes of a 154 155 Hadoop cluster (Amazon Elastic Map Reduce) using a map-reduce implementation

156 (https://github.com/ngcrawford/CloudForest) of a workflow combining MrAIC to estimate and

select the most-appropriate finite-sites substitution model. We used PhyML 3.0 [47] to estimate

gene trees, and PHYBASE to estimate species trees from gene trees using the STAR (Species

Trees from Average Ranks of Coalescences) method [48]. We performed 1,000 multi-locus, nonparametric bootstrap replicates for the STAR tree by resampling nucleotides within loci as well as resampling loci within the data set [49]. We only performed the species tree analysis on the alignment with no missing data due to concerns about how missing loci might affect a coalescent analysis.

To assess phylogenetically informative indels, we scanned alignments by eye in Geneious 5.4 (Biomatters Ltd, Aukland, New Zealand), recording indels that were 2 bp or more in length and shared between two or more ingroup taxa. We then mapped informative indels onto the resolved 416-locus Bayesian phylogeny.

168

169 RESULTS

170 We provide summary statistics for sequencing and alignment in Table 1. We obtained an average of 2.6 million reads per sample (range = 1.1 - 4.9 million). These reads assembled into an 171 average of 1,830 contigs per sample (range = 742 - 2,418). An average (per sample) of 1,412 of 172 these contigs matched the UCE loci from which we designed target capture probes (range = 694173 -1,681). The average length of UCE-matching contigs was 429 base pairs (bp) (range = 244 -174 598), and the average coverage of UCE-matching contigs was 71 times (range = 44 - 138). The 175 percentage of original sequencing reads that were "on target" (i.e., helped build UCE-matching 176 contigs) averaged 24% across samples (range = 15% - 35%). 177

When we selected loci allowing 50% of species for a given locus to have missing data, the final data set contained 1,541 UCE loci and produced a concatenated alignment that was 87% complete across 32 Neoaves species and the chicken outgroup. The average length of these 1,541 loci was 350 bp (min=90, max=621), and the total concatenated alignment length was 539,526 182 characters (including indels) with 24,703 informative sites.

Generally, the Bayesian and ML phylogenies for the 1,541 locus alignment were similar 183 in their topology and amount of resolution (Fig. 2a; see Fig. S1 for fully resolved trees). Of the 184 31 nodes, 27 (87%) were highly supported in the Bayesian tree (>0.95 PP), whereas a subset of 185 20 of those nodes (65%) were also highly supported in the ML tree (>75% bootstrap score). An 186 additional 7 nodes (23%) appeared in both the Bayesian and ML trees, but support in the ML tree 187 was low (bisected nodes in Fig. 2a). Four nodes (16%) had either low support in both trees (and 188 thus are collapsed in Fig. 2a) or had high support in the Bayesian tree, but did not appear in the 189 190 ML tree (white nodes in Fig. 2a). A phylogram for the 1,541 locus Bayesian tree (Fig. S2) showed long terminal branches and short internodes near the base of the tree, consistent with 191 previous studies suggesting an ancient, rapid radiation of Neoaves. 192

For the data set requiring no missing data, we recovered 416 UCE loci across 29 Neoaves 193 species and the chicken outgroup. Enrichments for three species performed relatively poorly 194 (Table 1; *Micrastur*, *Trogon*, and *Vidua*), and we excluded these samples to boost the number of 195 loci recovered. The average length of these 416 loci was 397 bp, and the total concatenated 196 alignment length was 165,163 characters (including indels) with 7,600 informative sites. 197 198 Bayesian and ML trees differed more in their topology and resolution than was observed for the 1,541 locus trees above (Fig. 2b; see Fig. S3 for fully resolved trees). Of the 28 nodes, 24 (86%) 199 were highly supported in the Bayesian tree (>0.95 PP), whereas only a subset of 14 (50%) was 200 201 highly supported in the ML tree (>75% bootstrap score). We recovered an additional three nodes (11%) in both the Bayesian and ML trees, but support for these nodes in the ML tree was low 202 (bisected nodes in Fig. 2b). Twelve nodes (43%) disagreed between the Bayesian and ML trees, 203 204 a frequency much higher than the 16% disagreement we observed from the 1,541 locus analysis.

The STAR species tree from the 416 locus data set (Fig. 3; Fig. S3c) was much less resolved and had lower support values than either the Bayesian or ML tree estimated for these data. There has been little study on what constitutes high bootstrap support for a species tree analysis, but only 11 nodes (39%) had over 50% support. Despite the differences in resolution between the Bayesian, ML, and STAR species tree for the 416 locus analysis, when we collapsed weakly supported nodes (PP < 0.90, ML bootstrap < 70%, species-tree bootstrap < 40%), there very few strongly supported contradictions among the three trees.

We identified 44 indels greater than two bp in length that were shared among two or more ingroup taxa (Table S1). Only 13 of these indels validated clades found in the phylogenetic trees generated from nucleotide data. The four clades supported by the 13 indels represented four of the six longest internal branches of the phylogeny (Fig. 4).

216

217 DISCUSSION

Containing 1,541 loci and 32 species, our study is among the largest comparative avian 218 phylogenomics data sets assembled for the purpose of elucidating avian evolutionary 219 relationships. By strengthening support for controversial relationships and resolving several new 220 221 parts of the avian tree (discussed below), our results suggest that increasing sequence data will lead to an increasingly resolved bird tree of life, with some caveats. Our sampling strategy 222 sought to balance the number of taxa included with the number of loci interrogated. We sampled 223 224 the genome much more broadly than the 19 loci of Hackett et al. [6], but with reduced taxonomic sampling (32 species compared to 169 species). Additionally, compared to Hackett et al. [6], our 225 loci were shorter (350 bp vs. 1,400 bp), meaning that although our 1,541 locus data set contained 226 227 roughly 80 times the number of loci, our total alignment length was only about 17 times larger.

Another recent avian phylogenomic study [50] included 1,995 loci, producing a concatenated alignment roughly 1.5 times larger than ours, but this study included only 9 Neoaves species, 5 of which were passerines, which limited the potential of that study for phylogenetic inference.

232 Increasing data increases resolution of the avian tree of life

One striking result of our study is that Bayesian and ML trees based on 1,541 loci were in much stronger agreement with one another than Bayesian and ML trees estimated from 416 loci (Fig. 2). The stronger agreement was driven primarily by increased resolution and support of the 1,541 locus ML tree (i.e., it became more similar to the Bayesian tree). In contrast, although the 416locus Bayesian tree was highly resolved, its ML counterpart was much less so and conflicted in topology with the Bayesian tree to a greater degree.

Combined with results of other studies, this suggests that increasing loci leads to 239 increasing support and stability of the avian tree. In discussing our results below, we rely 240 primarily on relationships found in the 1,541 locus tree due to the stronger congruence among 241 analytical methods, as well as recent research suggesting that analyses of incomplete data 242 matrices may be beneficial for studies with highly incomplete taxonomic sampling [51]. Most 243 simulation studies assessing the effect of missing data found that a common negative effect of 244 missing data was erosion of support values rather than an artificial increase in support [52]. We 245 did not observe lower support values in the tree with more missing data, and, in fact, we 246 247 observed the opposite, suggesting minimal negative effects of missing data. This is perhaps unsurprising given that the threshold amount of missing data producing negative effects in 248 simulation studies was often much higher than our level of missing data (many studies assessing 249 250 50-90% missing data, whereas we had 13%). Where relevant, we compare the 416 locus tree and

species tree to the 1,541 locus tree, and we discuss a few results from the 416 locus tree that areparticularly well supported or interesting.

253

254 Low support for the species tree and differences between Bayesian and ML trees

The low support for many nodes in the species tree (Fig. 3) is understandable given the length of 255 individual UCE loci. We estimated the species tree using methods that take gene trees as input, 256 rather than those that jointly estimating both gene trees and species trees [53], which is too 257 computationally intensive for large data sets. Therefore, the resolution of the species tree is 258 259 entirely dependent on the quality and resolution of the individual gene trees. Because we assembled relatively short UCE loci (397 bp for the 416 locus data set) from enriched reads, each 260 locus, considered individually, is not likely to contain much signal informing basal relationships. 261 262 Concatenation effectively masks this reduction in signal by joining all loci, maximizing the information content on short internal branches, and helping to resolve relationships when 263 speciation intervals are short. Of course, this benefit of concatenation comes with the cost of 264 ignoring the independent histories of genes and potentially inflating support values for nodes 265 affected by substantial coalescent stochasticity [54,55], especially when using Bayesian methods. 266 While the low information content of shorter UCE loci clearly posed a problem for 267 inferring the species tree, this is a methodological limitation of this study rather than a general 268 limitation of the UCE enrichment approach. For this study, we sequenced single-end, 100 bp 269 270 reads on an Illumina GAIIx. However, it is now possible to obtain paired-end reads as long as 250 bp from the Illumina platform, which will facilitate assembly of longer loci from fewer reads 271 than we obtained during this study. Tighter control on the average size of DNA fragments used 272 273 for enrichment (i.e., using fragments of the maximum size allowed by the sequencing platform)

274 and increased sequencing depth can also increase the size of recovered loci to 600-700 bp (B. Faircloth, unpublished data). Using UCE loci that averaged ~750 bp, we did not observe poorly 275 resolved species trees in a study of rapid radiation of mammals [35]. Thus, increasing the length 276 277 of loci recovered is clearly an important step towards addressing the dual problems of low information content and coalescent stochasticity in resolving the avian tree of life, although it 278 remains to be seen how denser taxon sampling will interact with these problems and affect future 279 analyses. In any event, given our results and those of prior studies, the more exigent problem in 280 this case appears to be low information content. 281

282 Although there were very few contradictory relationships in highly supported parts of the trees, there was an obvious difference in resolution between the Bayesian and ML trees for the 283 416 locus alignment, and to a lesser degree, for the 1,541 locus alignment. One possible 284 explanation for the lower resolution of the ML trees is that bootstrapping may not be the best 285 way to assess confidence with UCE data, given the expected skewed distribution of phylogenetic 286 information across sites (i.e., more toward the flanks) [33]. Also, it is common to observe higher 287 support values for trees estimated by Bayesian methods, and in some cases PPs can be 288 deceptively high [56,57]. There is also current debate concerning whether Bayesian methods 289 might suffer from a "star tree paradox", where a simultaneous divergence of three or more 290 lineages nonetheless appears resolved in bifurcating fashion with high PP [58,59]. Bayesian 291 methods also might be more prone to long-branch attraction [60]. Research on these concerns is 292 293 ongoing and salient to our results, in which the Bayesian trees tended to group several basally diverging lineages with long branches together into clades with high PP that were not supported 294 by the ML trees. On the other hand, ML bootstraps can underestimate support compared to 295 296 Bayesian methods [61,62] – an effect suggested by our observation that many weakly supported

nodes in the 416 locus ML tree, for which Bayesian analysis showed high PP, became well
supported in the ML tree when we increased the size of the data matrix to 1,541 loci.

299

300 Defining a backbone for the Neoaves phylogeny

We found strong congruence across data sets and analytical methods for previously 301 hypothesized, but still tenuously supported, waterbird (Aequornithes; [63]) and landbird clades 302 [2,6] that diverge deep in the Neoaves phylogeny (Fig. 2). We address relationships within 303 landbirds and waterbirds below, but their position as sister clades in three of four trees contrasts 304 305 with previous studies that placed a number of additional taxa close to the waterbirds [2,6,23]. Both Bayesian trees supported a third clade – including families as diverse as hummingbirds, 306 flamingos, cuckoos, trumpeters, bustards, and turacos - bearing some resemblance to the 307 Metaves clade recovered in earlier molecular studies [2,6,23], but differing by including 308 bustards, trumpeters, and turacos, which have not typically been considered part of Metaves. 309 However, this clade did not appear in either ML tree or the species tree, suggesting that the 310 grouping of these taxa could be an artifact resulting from long-branch attraction, as discussed 311 above. Although we uncovered novel, well-supported sister relationships between some of these 312 species toward the tips of the tree (see below), their deeper evolutionary affinities will need to be 313 explored with increased taxonomic sampling to break up long branches and provide further 314 information on state changes deep in the tree. Our study thus suggests that resolving the avian 315 316 tree outside of waterbirds and landbirds is the final frontier in deep-level bird systematics.

317

318

319 The surprising relationship between tropicbirds and the sunbittern

320 This study adds to the overwhelming evidence for a sister relationship between the phenotypically divergent flamingo and grebe families [2,5,6,64-66]. Our results also suggest 321 another surprisingly close affinity between morphologically disparate groups – tropicbirds and 322 323 the sunbittern. Three of four analyses lent strong support to this relationship, for which ML support increased sharply (43% to 96%) when genomic sampling increased from 416 to 1,541 324 loci (Fig. 2; Fig. S1 & S2). A close relationship between the sunbittern and tropicbirds is 325 surprising because of dissimilarities in appearance, habitat, and geography. Tropicbirds are 326 pelagic seabirds with mostly white plumage, elongated central tail feathers, and short legs that 327 328 make walking difficult. Meanwhile, the sunbittern is a cryptic resident of lowland and foothill Neotropical forests that spends much of its time foraging on the ground in and near freshwater 329 streams and rivers. The kagu, a highly terrestrial bird restricted to the island of New Caledonia 330 (not sampled in our study), is the sister species of the sunbittern [6,22,23] and may superficially 331 bear some similarity to tropicbirds. These results should spark further research into shared 332 morphological characteristics of tropicbirds, the sunbittern, and the kagu. 333

334

335 A sister relationship between bustards and turacos?

Another surprising sister relationship uncovered in our study is that between turacos and bustards
(Fig. 2a). Turacos are largely fruit-eating arboreal birds of sub-Saharan Africa, whereas bustards
are large, omnivorous, terrestrial birds widely distributed in the Old World. Despite some
overlap in their biogeography, the two families have little in common and have, to our
knowledge, never been hypothesized to be closely related based on phenotypic characteristics.
Previous molecular studies have placed members of these two families near one another
evolutionarily [2,6], but never as sister taxa. Our study did not include a member of the cuckoo

family, which has often been considered a close relative of the turacos and thus might be its true
sister taxon. An additional note of caution is that a turaco-bustard relationship was not supported
outside the 1,541 locus tree, but neither was it contradicted. Thus, although confirming results
are needed, our study provides some support for the idea that turacos and bustards are much
more closely related than previously thought, if not actually sister families.

348

349 Further clarity for waterbird relationships

We found consistent support across all analyses for relationships among the six sampled families within the waterbirds (Figs. 2 and 3). Prior to the availability of molecular data, the relationships within this clade were difficult to resolve due to the extreme morphological diversity of its members and the scarcity of apomorphic morphological characters [63]. The topology we recovered within this portion of the tree is identical to that of Hackett et al. [6]. For example, in both studies loons are the outgroup to all other waterbirds, and the morphologically divergent penguins are sister to tube-nosed seabirds in the family Procellariidae.

357

358 *Hoatzin: still a riddle wrapped in a mystery...*

Hoatzin (*Opisthicomus hoazin*), the only extant member of Opisthocomidae, is arguably the most
enigmatic living bird species due to its unique morphology, folivorous diet, and confusion
relative to its evolutionary affinities across numerous molecular phylogenies. One phylogenetic
study found no support for a sister relationship between hoatzin and the Galloanserae, nor with
turacos, cuckoos, falcons, trogons, or mousebirds in Neoaves; the study found some, albeit weak,
support for a sister relationship between hoatzin and doves [67]. The 416 locus Bayesian tree
placed the hoatzin sister to a shorebird (Fig. 2b) with high support, but we did not observe this

366 relationship in either the ML tree or the species tree. Furthermore, support for any definitive placement of the hoatzin eroded in the 1,541 locus tree (Fig. 2a). A close relationship of hoatzin 367 to shorebirds would be extremely surprising and in stark contrast to any prior hypotheses [68]. 368 Our results raise the question of whether or not more data will eventually lead to a definitive 369 conclusion on the phylogenetic position of the hoatzin. Given the phylogenetic distinctiveness of 370 the hoatzin, better taxonomic sampling may be as beneficial as further genomic sampling in the 371 search for shared, derived characters deep in the tree. Thus, we present a link between the 372 hoatzin and shorebirds, a large family whose members are found in diverse terrestrial and aquatic 373 374 habitats, as an intriguing phylogenetic hypothesis.

375

376 An early divergence for pigeons and doves?

Another place where our 416 locus trees showed support for a relationship not found in the 1,541 377 locus trees was in the placement of the pigeon and dove family (Columbidae). Most prior studies 378 either placed pigeons and doves in an unresolved position [6] or sister to sandgrouse 379 (Pteroclididae) within Metaves [2]. However, amino acid sequences of feather beta-keratins have 380 suggested a basal position of Columbidae within Neoaves [69]. We found complete support in 381 the 416-locus Bayesian tree for a sister relationship between Columbidae and the rest of Neoaves 382 (Fig. 2b). We also recovered this relationship in the 416-locus ML tree and species tree, although 383 with weak support (Fig. S2). However, the 1,541 locus trees disagreed by placing pigeons and 384 385 doves in a more conventional position sister to sandgrouse and instead placing trumpeters sister to the rest of Neoaves (Fig. 2a). 386

387

388 Support for controversial relationships within the landbirds

389 One of the biggest challenges to conventional thought on bird phylogeny contained in Hackett et al. [6] was in the relationships among landbirds. Their finding that parrots were the sister family 390 to passerines is still viewed as controversial (bootstrap support for parrots + passerines from 391 Hackett et al. [6] was 77%), despite corroborating evidence from rare genomic changes encoded 392 in retroposons [12] and expanded data sets [7]. Our results across all analyses strongly support 393 the sister relationship between passerines (in this study represented by a suboscine Pitta and an 394 oscine Vidua) and parrots (perfect support in all Bayesian and ML trees; 85% support in the 395 species tree). 396

Our results also support another controversial finding from Hackett et al. [6]: the absence of a sister relationship between raptorial birds in the hawk (Accipitridae) and falcon (Falconidae) families. Both ML and Bayesian trees from the 1,541 locus analysis provided perfect support for falcons sister to the parrot + passerine clade, whereas the representative of the hawk family was sister to the vultures with high support, improving upon the weak support for hawks + vultures from Hackett et al. [6].

Finally, the larger 1,541 analysis helped resolve deeper relationships within the landbirds among four main clades: (i) passerines + parrots + falcons, (ii) hawks + vultures, (iii) the group sometimes called the "near passerines" (e.g., barbet, woodpecker, woodhoopoe, motmot, and trogon, also known as the CPBT clade in [7] because it includes the families Coraciiformes, Piciformes, Bucerotiformes, and Trogoniformes), and (iv) owls (Fig. 2a). The Bayesian tree placed owls sister to the "near passerines" and then hawks + vultures sister to owls + "near passerines", a topology that also appeared in the ML tree with weak support.

410 Meanwhile, the evolutionary affinities of mousebirds, whose position in prior studies has
411 been uncertain [6,7], remain equivocal. The 416 locus trees positioned mousebirds sister to the

412 "near passerines", but the 1,541 locus trees placed mousebirds sister to passerines. Wang et al.

413 [7] also found mousebirds moving between these two clades depending on the analysis. Other

relationships within the "near passerines" were consistent with previous results [2,6] except that

the positions of trogons and motmots switched between the 416 and 1,541 locus trees.

416

417 A scarcity of indels on short internal branches

Our finding that informative indels were generally scarce (found only on four of the longest 418 internal branches in the phylogeny; Fig. 4) corroborates previous work on rare genomic changes 419 in retroposons, which also found little evidence for shared events deep in the bird phylogeny 420 [12,13]. The low prevalence of informative indels may be exacerbated by the lack of major 421 structural changes in and around UCE loci, although this has not been well studied. Previous 422 423 work on nuclear introns has identified a handful of indels supporting major subdivisions deep in avian phylogeny [23,70,71]. However, lessons from coalescence theory caution that, when 424 drawing phylogenetic inferences from rare genomic changes, numerous loci supporting 425 particular subdivisions are required to account for the expected high variance in gene histories 426 [35]. The study of bird phylogeny awaits a genome-scale analysis of many hundreds of rare 427 428 genomic events including indels, retroposons, and microRNAs.

429

430 *Conclusions*

Our results, combined with other recent studies [2,6], demonstrate that increasing sequence data
leads to improved resolution of the bird tree of life. Major challenges clearly remain in
corroborating results across analytical methods and data types. One of these challenges is a
species tree for birds. While we have focused here on the seemingly more pressing problem of

obtaining phylogenetic signal and high support values from concatenated data sets, we
acknowledge that a proper accounting of the ultra-rapid radiation of avian lineages will require
methods that reconcile discordant gene trees, which could lead to different results. Nevertheless,
the incremental progress of resolving the bird tree of life is a major turnaround from more
pessimistic attitudes that predated the decreased sequencing costs of the last decade and the
advent of high-throughput sequencing technologies [72].

The framework we outline here, sequence capture using UCEs, is a powerful approach 441 that can scale to hundreds of taxa, thousands of loci, and include longer flanking sequences with 442 443 different library preparation and sequencing regimes. Because UCEs occur in many organisms, the method is broadly applicable across the tree of life [32,33]. Data from sequence capture 444 approaches can also be mixed, in hybrid fashion, with UCEs excised from whole genome 445 assemblies [33,34,73] or other types of molecular markers, providing a powerful method for 446 collecting and analyzing phylogenomic data from non-model species to elucidate their 447 evolutionary histories. 448

449

450 LITERATURE CITED

451 1. Feduccia A (1999) The origin and evolution of birds. New Haven, Connecticut: Yale
452 University Press.

453 2. Ericson PGP, Anderson CL, Britton T, Elzanowski A, Johansson US, et al. (2006)

454 Diversification of Neoaves: integration of molecular sequence data and fossils. Biol Lett
455 2: 543-547.

3. Brown JW, Rest J, García-Moreno J, Sorenson M, Mindell D (2008) Strong mitochondrial
 DNA support for a Cretaceous origin of modern avian lineages. BMC Biol 6: 6.

4. Chojnowski JL, Kimball RT, Braun EL (2008) Introns outperform exons in analyses of basal
avian phylogeny using clathrin heavy chain genes. Gene 410: 89-96.

- 460 5. Cracraft J, Barker FK, Braun M, Harshman J, Dyke GJ, et al. (2004) Phylogenetic
- relationships among modern birds (Neornithes): toward an avian tree of life. In: Cracraft
 J, Donoghue M, editors. Assembling the tree of life. New York, NY: Oxford University
- 463 Press. pp. 468-489.
- 464 6. Hackett SJ, Kimball RT, Reddy S, Bowie RCK, Braun EL, et al. (2008) A phylogenomic
 465 study of birds reveals their evolutionary history. Science 320: 1763-1768.
- 466 7. Wang N, Braun EL, Kimball RT (2012) Testing hypotheses about the sister group of the
 467 Passeriformes using an independent 30-locus data set. Mol Biol Evol 29: 737-750.
- 8. Zink RM, Barrowclough GF (2008) Mitochondrial DNA under siege in avian
 phylogeography. Mol Ecol 17: 2107-2121.
- 470 9. Whitfield JB, Lockhart PJ (2007) Deciphering ancient rapid radiations. Trends Ecol Evol 22:
 471 258-265.
- 472 10. Rokas A, Holland PWH (2000) Rare genomic changes as a tool for phylogenetics. Trends
 473 Ecol Evol 15: 454-459.
- 474 11. Shedlock AM, Takahashi K, Okada N (2004) SINEs of speciation: tracking lineages with
 475 retroposons. Trends Ecol Evol 19: 545-553.
- 476 12. Suh A, Paus M, Kiefmann M, Churakov G, Franke FA, et al. (2011) Mesozoic retroposons
 477 reveal parrots as the closest living relatives of passerine birds. Nature Comm 2: 443.
- 478 13. Matzke A, Churakov G, Berkes P, Arms EM, Kelsey D, et al. (2012) Retroposon insertion
 479 patterns of neoavian birds: strong evidence for an extensive incomplete lineage sorting
 480 era. Mol Biol Evol 29: 1497-1501.
- 481 14. Han KL, Braun EL, Kimball RT, Reddy S, Bowie RCK, et al. (2011) Are transposable
 482 element insertions homoplasy free?: an examination using the avian tree of life. Syst Biol
 483 60: 375-386.
- 484 15. Maddison WP (1997) Gene trees in species trees. Syst Biol 46: 523-536.
- 16. Degnan JH, Salter LA (2005) Gene tree distributions under the coalescent process. Evolution
 59: 24-37.
- 487 17. Avise JC, Robinson TJ (2008) Hemiplasy: a new term in the lexicon of phylogenetics. Syst
 488 Biol 57: 503-507.
- 18. Knowles L (2009) Estimating species trees: methods of phylogenetic analysis when there is
 incongruence across genes. Syst Biol 58: 463-467.

- 491 19. Edwards SV (2008) Is a new and general theory of molecular systematics emerging?
 492 Evolution 63: 1-19.
- 20. Liu L, Yu L, Kubatko LS, Pearl DK, Edwards SV (2009) Coalescent methods for estimating
 phylogenetic trees. Mol Phylogenet Evol 53: 320-328.
- 495 21. Edwards SV, Jennings WB, Shedlock AM (2005) Phylogenetics of modern birds in the era of
 496 genomics. Proc R Soc B 272: 979.
- 497 22. Livezey BC, Zusi RL (2007) Higher-order phylogeny of modern birds (Theropoda, Aves:
 498 Neornithes) based on comparative anatomy. II. Analysis and discussion. Zool J Linn Soc
 499 149: 1-95.
- 500 23. Fain MG, Houde P (2004) Parallel radiations in the primary clades of birds. Evolution 58:
 501 2558-2573.
- 502 24. Groth JG, Barrowclough GF (1999) Basal divergences in birds and the phylogenetic utility of
 503 the nuclear RAG-1 gene. Mol Phylogenet Evol 12: 115-123.
- 504 25. Haddrath O, Baker AJ (2012) Multiple nuclear genes and retroposons support vicariance and
 505 dispersal of the palaeognaths, and an Early Cretaceous origin of modern birds. Proc Roy
 506 Soc B In press.
- 507 26. Gibb GC, Penny D (2010) Two aspects along the continuum of pigeon evolution: A South 508 Pacific radiation and the relationship of pigeons within Neoaves. Mol Phylogenet Evol
 509 56: 698-706.
- 27. Pratt RC, Gibb GC, Morgan-Richards M, Phillips MJ, Hendy MD, et al. (2009) Toward
 resolving deep Neoaves phylogeny: data, signal enhancement, and priors. Mol Biol Evol
 26: 313-326.
- 28. Pacheco MA, Battistuzzi FU, Lentino M, Aguilar RF, Kumar S, et al. (2011) Evolution of
 modern birds revealed by mitogenomics: timing the radiation and origin of major orders.
 Mol Biol Evol 28: 1927-1942.
- 516 29. Braun E, Kimball R, Han KL, Iuhasz-Velez N, Bonilla A, et al. (2011) Homoplastic
 517 microinversions and the avian tree of life. BMC Evol Biol 11: 141.
- 30. Bejerano G, Pheasant M, Makunin I, Stephen S, Kent WJ, et al. (2004) Ultraconserved
 elements in the human genome. Science 304: 1321-1325.

- 31. Janes DE, Chapus C, Gondo Y, Clayton DF, Sinha S, et al. (2011) Reptiles and mammals
 have differentially retained long conserved noncoding sequences from the Amniote
 ancestor. Genome Biol Evol 3: 102-113.
- 32. Siepel A, Bejerano G, Pedersen JS, Hinrichs AS, Hou M, et al. (2005) Evolutionarily
 conserved elements in vertebrate, insect, worm, and yeast genomes. Genome Res 15:
 1034-1050.
- 526 33. Faircloth BC, McCormack JE, Crawford NG, Harvey MG, Brumfield RT, et al. (2012)
 527 Ultraconserved elements anchor thousands of genetic markers for target enrichment
 528 spanning multiple evolutionary timescales. Syst Biol 61: 717-726.
- 34. Crawford NG, Faircloth BC, McCormack JE, Brumfield RT, Winker K, et al. (2012) More
 than 1000 ultraconserved elements provide evidence that turtles are the sister group of
 archosaurs. Biol Lett 8: 783-786.
- 532 35. McCormack JE, Faircloth BC, Crawford NG, Gowaty PA, Brumfield RT, et al. (2012)
- 533 Ultraconserved elements are novel phylogenomic markers that resolve placental mammal 534 phylogeny when combined with species-tree analysis. Genome Res 22: 746-754.
- 36. Derti A, Roth FP, Church GM, Wu C-t (2006) Mammalian ultraconserved elements are
 strongly depleted among segmental duplications and copy number variants. Nat Genet
 38: 1216-1220.
- 37. Sibley CG, Monroe BL (1990) Distribution and taxonomy of birds of the world. New Haven:
 Yale University Press.
- 540 38. Sambrook J, Russell DW (2001) Molecular cloning: a laboratory manual. Cold Spring
 541 Harbor: CSHL Press.
- 542 39. Faircloth BC, Glenn TC (2012) Not all sequence tags are created equal: designing and
 543 validating sequence identification tags robust to indels. PLoS One 7: e42543.
- 40. Zerbino DR, Birney E (2008) Velvet: Algorithms for de novo short read assembly using de
 Bruijn graphs. Genome Res 18: 821-829.
- 546 41. Edgar RC (2004) MUSCLE: a multiple sequence alignment method with reduced time and
 547 space complexity. BMC Bioinformatics 5: 113-119.
- 42. Ronquist F, Huelsenbeck JP (2003) MrBayes 3: Bayesian phylogenetic inference under
 mixed models. Bioinformatics 19: 1572-1574.

- 43. Rambaut A, Drummond AJ (2007) Tracer MCMC Trace Analysis Tool, v1.4. Available
 from: <<u>http://beastbioedacuk/Tracer/></u>.
- 44. Nylander JAA, Wilgenbusch JC, Warren DL, Swofford DL (2008) AWTY (are we there
 yet?): a system for graphical exploration of MCMC convergence in Bayesian
 phylogenetics. Bioinformatics 24: 581-583.
- 45. Stamatakis A (2006) RAxML-VI-HPC: maximum likelihood-based phylogenetic analyses
 with thousands of taxa and mixed models. Bioinformatics 22: 2688-2690.
- 46. Stamatakis A, Hoover P, Rougemont J (2008) A rapid bootstrap algorithm for the RAxML
 Web servers. Syst Biol 57: 758-771.
- 47. Guindon S, Dufayard JF, Lefort V, Anisimova M, Hordijk W, et al. (2010) New algorithms
 and methods to estimate maximum-likelihood phylogenies: assessing the performance of
 PhyML 3.0. Syst Biol 59: 307-321.
- 48. Liu L, Yu L, Pearl DK, Edwards SV (2009) Estimating species phylogenies using
 coalescence times among sequences. Syst Biol 58: 468-477.
- 49. Seo TK (2008) Calculating bootstrap probabilities of phylogeny using multilocus sequence
 data. Mol Biol Evol 25: 960-971.
- 566 50. Nabholz B, Kunstner A, Wang R, Jarvis ED, Ellegren H (2011) Dynamic evolution of base
 567 composition: causes and consequences in avian phylogenomics. Mol Biol Evol 28: 2197568 2210.
- 569 51. Wiens JJ, Tiu J (2012) Highly incomplete taxa can rescue phylogenetic analyses from the
 570 negative impacts of limited taxon sampling. PLoS One 7: e42925.
- 571 52. Wiens JJ, Morrill MC (2011) Missing data in phylogenetic analysis: reconciling results from
 572 simulations and empirical data. Syst Biol 60: 719-731.
- 573 53. Edwards SV, Liu L, Pearl DK (2007) High-resolution species trees without concatenation.
 574 Proc Natl Acad Sci USA 104: 5936-5841.
- 575 54. Kubatko L, Degnan J (2007) Inconsistency of phylogenetic estimates from concatenated data
 576 under coalescence. Syst Biol 56: 17-24.
- 577 55. Mossel E, Vigoda E (2005) Phylogenetic MCMC algorithms are misleading on mixtures of
 578 trees. Science 309: 2207-2209.
- 56. Suzuki Y, Glazko GV, Nei M (2002) Overcredibility of molecular phylogenies obtained by
 Bayesian phylogenetics. Proc Natl Acad Sci USA 25: 16138-16143.

- 581 57. Douady CJ, Delsuc F, Boucher Y, Doolittle WF, Douzery EJP (2003) Comparison of
- 582 Bayesian and maximum likelihood bootstrap measures of phylogenetic reliability. Mol
 583 Biol Evol 20: 248-254.
- 58. Kolaczkowski B, Thornton JW (2006) Is there a star tree paradox? Mol Biol Evol 23: 18191823.
- 586 59. Lewis PO, Holder MT, Holsinger KE (2005) Polytomies and Bayesian phylogenetic
 587 inference. Syst Biol 54: 241-253.
- 60. Kolaczkowski B, Thornton JW (2009) Long-branch attraction bias and inconsistency in
 Bayesian phylogenetics. PLoS One 4: e7891.
- 590 61. Erixon P, Svennblad B, Britton T, Oxelman B (2003) Reliability of Bayesian posterior
 591 probabilities and bootstrap frequencies in phylogenetics. Syst Biol 52: 665-673.
- 592 62. Huelsenbeck JP, Larget B, Miller RE, Ronquist F (2002) Potential applications and pitfalls of
 593 Bayesian inference of phylogeny. Syst Biol 51: 673-688.
- 63. Mayr G (2011) Metaves, Mirandornithes, Strisores and other novelties: a critical review of
 the higher-level phylogeny of neornithine birds. J Zool Syst Evol Res 49: 58-76.
- 64. Chubb AL (2004) New nuclear evidence for the oldest divergence among neognath birds: the
 phylogenetic utility of ZENK (i). Mol Phylogenet Evol 30: 140-151.
- 598 65. van Tuinen M, Butvill DB, Kirsch JAW, Hedges SB (2001) Convergence and divergence in
 599 the evolution of aquatic birds. Proc R Soc Lond B 268: 1345-1350.
- 600 66. Morgan-Richards M, Trewick S, Bartosch-Härlid A, Kardailsky O, Phillips M, et al. (2008)
 601 Bird evolution: testing the Metaves clade with six new mitochondrial genomes. BMC
 602 Evol Biol 8: 20.
- 603 67. Sorenson MD, Oneal E, Garcia-Moreno J, Mindell DP (2003) More taxa, more characters:
 604 the hoatzin problem is still unresolved. Mol Biol Evol 20: 1484-1498.
- 605 68. Thomas B (1996) Family Opisthocomidae (hoatzins). In: del Hoyo J, Jordi A, Sargatal C,
 606 editors. Handbook of the birds of the world, volume 3, hoatzins to auks. Barcelona: Lynx
 607 Ediciones. pp. 24-32.
- 608 69. Glenn TC, French JO, Heincelman TJ, Jones KL, Sawyer RH (2008) Evolutionary
- relationships among copies of feather beta (β) keratin genes from several avian orders.
 Integr Comp Biol 48: 463-475.

611	70. Pasko L, Ericson PGP, Elzanowski A (2011) Phylogenetic utility and evolution of indels: A
612	study in neognathous birds. Mol Phylogenet Evol 61: 760-771.
613	71. Prychitko TM, Moore WS (2003) Alignment and phylogenetic analysis of β -fibrinogen
614	intron 7 sequences among avian orders reveal conserved regions within the intron. Mol
615	Biol Evol 20: 762-771.
616	72. Poe S, Chubb AL (2004) Birds in a bush: five genes indicate explosive evolution of avian
617	orders. Evolution 58: 404-415.
618	73. Haussler D, O'Brien SJ, Ryder OA, Barker FK, Clamp M, et al. (2009) Genome 10K: a
619	proposal to obtain whole-genome sequence for 10,000 vertebrate species. J Hered 100:
620	659-674.
621	
622	Acknowledgments We thank Scott Herke and the LSU Genomics Facility for assistance with
623	sequencing. Donna Dittmann (LSUMZ) assisted with tissue loans. Illustrations for Figure
624	1 are artistic interpretations based on photos used with permission or under Creative
625	Commons license. Photo credits for Fig. 1: (1) Prin Pattawaro; (2) Alan Manson; (3)
626	Farelli; (4) Unknown; (5) Enoch Joseph Wetsy; (6) Vijay Cavale; (7) Nancy Wyman; (8)
627	Lip Kee; (9) Steve Turner; (10) Keith Murdock; (11) Unknown; (12) Eduardo Lopez;
628	(13) Jan Sevcik; (14) Srihari Kulkarni; (15) Arthur Grosset; (16) Tom Tarrant; (17)
629	Fir0002/Flagstaffotos; (18) Mark Hannaford; (19) Unknown; (20) Jose Garcia; (21) "The
630	Lilac Breasted Roller"; (22) Paul Baker; (23) Unknown; (24) Linda De Volder; (25) Utz
631	Klingenböck; (26) Pixxl (Lisa M); (27) Bobby K; (28) Tarique Sani; (29) Lee Harding;
632	(30) Doug Backlund; (31) Doug Pratt; (32) Jeff Whitlock. M. Alfaro and two anonymous
633	reviewers provided comments on the manuscript.
634	
635	Availability Assembled contigs, alignments, and gene trees for both data sets are available from
636	Dryad (doi: 10.5061/dryad.sd080). All source code used for UCE data processing is available
637	from https://github.com/faircloth-lab/phyluce under BSD and Creative Commons licenses.

- 638 Version controlled, reference probe sets and outgroup data are available from
- 639 <u>https://github.com/faircloth-lab/uce-probe-sets</u>. UCE contigs used in analyses are available from

- 640 Genbank (accessions: JQ328245 JQ335930, KC358654 KC403881). Protocols for UCE
- 641 enrichment, probe design, and additional information regarding techniques are available from
- 642 <u>http://ultraconserved.org</u>.

		_			All contigs			Contigs Aligned to UCE loci							
Family	Scientific name	Common Name	Museum tissue no.	Number of trimmed reads	Count	Avg. size	Avg. coverage	Reads in contigs	Count	Avg. size	Avg. coverage	Reads in contigs	Contigs match >1 locus ¹	Contigs "on- target" ²	Reads "on- target" ³
Pittidae (1)	Pitta guajana	Banded Pitta	LSUMZ B36368	2,723,264	2369	386	63.1	914,414	1572	457.4	71.3	719,095	32	0.66	0.26
Viduidae (2)	Vidua macroura	Pin-tailed Whydah	LSUMZ B16749	1,098,154	1203	240	41.3	288,210	959	244.2	43.5	234,214	2	0.80	0.21
Psittacidae (3)	Psittacula alexandri	Red-breasted Parakeet	LSUMZ B36554	2,745,979	2312	421	55	974,441	1487	508.1	62.7	752,493	42	0.64	0.27
Falconidae (4)	Micrastur	Collared Forest Falcon	LSUMZ B11298	1,405,847	742	309	49.9	229,417	694	309.8	51.1	214,967	8	0.94	0.15
Coliidae (5)	Urocolius indicus	Red-faced Mousebird	LSUMZ B34225	2,822,685	2208	398	73.9	877,590	1495	465.3	84.0	695,586	43	0.68	0.25
Megalaimidae (6)	Megalaima virens	Great Barbet	LSUMZ B20788	2,302,531	1370	341	58.6	466,552	1174	351.1	62.7	412,208	10	0.86	0.18
Picidae (7)	Sphyrapicus varius	Yellow-bellied Sapsucker	FLMNH 43569	2,693,567	1952	388	61.2	757,975	1542	416.5	65.9	642,192	46	0.79	0.24
Phoeniculidae (8)	Rhinopomastus	Common Scimitarbill	LSUMZ B34262	1,829,285	1742	382	55.9	665,679	1425	411.1	59.3	585,753	24	0.82	0.32
Momotidae (9)	Momotus momota	Blue-crowned Motmot	LSUMZ B927	2,694,269	2195	383	51.9	840,829	1587	430.7	57.3	682,265	45	0.72	0.25
Trogonidae (10)	Trogon personata	Masked Trogon	LSUMZ B7644	2,371,840	1263	316	80.8	399,423	1117	315.1	84.6	351,958	13	0.88	0.15
Tytonidae (11)	Tyto alba	Barn Owl	LSUMZ B19295	3,543,135	1833	338	60.7	620,375	1464	360.9	67.0	528,413	22	0.80	0.15
Accipitridae (12)	Gampsonyx swainsonii	Pearl Kite	LSUMZ B15046	2,605,257	1588	525	64.6	833,617	1351	557.6	67.2	753,293	8	0.85	0.29
Cathartidae (13)	Cathartes aura	Turkey Vulture	LSUMZ B17242	2,837,787	2166	462	69.4	1,001,122	1551	528.9	76.6	820,238	27	0.72	0.29
Phalacrocoracidae (14)	Phalacrocorax carbo	Great Cormorant	LSUMZ B45740	4,892,448	1601	521	133.8	834,275	1384	554.1	137.9	766,906	10	0.86	0.16
Scopidae (15)	Scopus umbretta	Hamerkop	LSUMZ B28330	3,322,061	2024	533	75	1,079,622	1580	598.1	78.7	944,999	46	0.78	0.28
Balaenicipitidae (16)	Balaeniceps rex	Shoebill	LSUMZ B13372	1,906,136	1784	420	52.8	749,552	1485	448.9	55.2	666,057	19	0.83	0.35
Spheniscidae (17)	Eudyptula minor	Little Penguin	LSUMZ B36558	3,009,607	2418	434	66.6	1,049,164	1681	507.5	73.5	852,753	42	0.70	0.28
Hydrobatidae (18)	Oceanites oceanicus	Wilson's Storm Petrel	LSUMZ B37197	2,519,648	1930	488	73.4	942,397	1574	535.6	76.9	842,403	18	0.82	0.33
Gaviidae (19)	Gavia immer	Common Loon	LSUMZ B7923	2,947,546	2132	386	48.4	821,803	1492	431.7	55.3	644,027	17	0.70	0.22
Nyctibiidae (20)	Nyctibius grandis	Great Potoo	LSUMZ B15415	4,224,329	2060	377	95	776,650	1474	421.0	105.2	620,400	78	0.72	0.15
Trochilidae (21)	Colibri coruscans	Sparkling Violetear	LSUMZ B5574	2,496,109	1881	384	64.4	723,418	1435	425.8	70.4	608,046	25	0.76	0.24
Phaethontidae (22)	Phaethon rubicauda	Red-tailed Tropicbird	LSUMZ B35135	2,956,951	1875	423	71.2	792,485	1450	460.9	77.8	668,317	36	0.77	0.23
Eurypygidae (23)	Eurypyga helias	Sunbittern	LSUMZ B1980	3,181,048	1988	416	78.8	827,124	1585	450.2	85.1	713,511	16	0.80	0.22
Opisthocomidae (24)	Opisthocomus hoazin	Hoatzin	LSUMZ B9660	1,848,363	1427	307	57.9	438,153	1257	309.4	61.7	388,853	8	0.88	0.21
Otididae (25)	Ardeotis kori	Kori Bustard	FLMNH 44254	2,058,864	2000	389	52.1	777,365	1489	436.0	57.0	649,136	54	0.74	0.32
Musophagidae (26)	Tauraco erythrolophus	Red-crested Turaco	LSUMZ B5354	3,031,838	2134	402	70	858,470	1571	447.8	78.4	702,976	37	0.74	0.23
Columbidae (27)	Treron vernans	Pink-necked Green Pigeon	LSUMZ B47229	1,949,899	1771	370	46.4	655,866	1337	409.7	48.5	547,817	47	0.75	0.28
Pteroclididae (28)	Pterocles exustus	Chestnut-bellied Sandgrouse	LSUMZ B20765	2,167,890	1303	341	71.7	444,614	1130	351.0	75.5	396,601	30	0.87	0.18
Phoenicopteridae (29)	Phoenicopterus	Chilean Flamingo	LSUMZ B37257	2,826,576	1878	371	68.4	696,317	1486	400.5	73.9	595,072	56	0.79	0.21
Podicipedidae (30)	Podiceps auritus	Horned Grebe	LSUMZ B19296	2,929,983	1502	391	77.4	587,752	1296	402.1	79.7	521,175	2	0.86	0.18
Charadriidae (31)	Phegornis mitchelli	Diademed Sandpiper-plover	LSUMZ B103926	2,488,988	1892	355	65.5	671,797	1518	381.9	70.3	579,714	49	0.80	0.23
Psophiidae (32)	Psophia crepitans	Grey-winged Trumpeter	LSUMZ B7513	2,224,282	2010	368	64.9	739,996	1550	401.9	70.2	622,967	26	0.77	0.28

643 Table 1. Summary of descriptive statistics for samples, Illumina sequencing, and UCE loci.

¹ Potential paralogs that were removed from the data set

² The number of contigs aligned to UCE loci / the total number of contigs

³ The number of reads aligning to UCE loci / total reads

645

Figure 1. Neoaves species used in this study. Species are listed in Table 1. Numbers match
those in table and on the tips of phylogenies. Illustrations are based on photos (see
Acknowledgments).

649

Figure 2. Relationships in Neoaves. A. Phylogeny based on 1,541 loci from 32 species and an 650 alignment that was 87% complete. B. Phylogeny based on 416 loci in 29 species and an 651 alignment that was 100% complete. A, B. Branch lengths are not shown to permit easier 652 interpretation of the topology (see Fig. 4 for phylogram of 416-locus tree and Fig. S2 for 653 phylogram of 1,541-locus tree). Bayesian trees are shown (nodes < 0.90 PP collapsed) with 654 655 circles on nodes indicating level of support for each node and congruence with the ML trees (see legend in figure). Support is shown for nodes that have less than 1.0 PP or less than 100% ML 656 bootstrap support (PP | ML). If only a bootstrap score is shown (e.g., 46), then PP for that node = 657 1.0. NP = node not present in ML tree. Thus, white nodes with no values indicate $1.0 \mid NP$. 658 659 Figure 3. Species tree estimated from 416 individual UCE gene trees. We collapsed nodes 660 receiving less than 40% bootstrap support. 661 662 663 Figure 4. Indels on the phylogram of the 416-locus Bayesian tree. Hash marks indicate the phylogenetic position of the 13 indels that supported clades found in the DNA sequence data 664

trees. The number of indels supporting each clade is shown.

667 SUPPORTING INFORMATION LEGENDS

668

- **Table S1. Indels greater than 1 bp. Informative indels (n=13) that corroborate Bayesian**
- 670 phylogeny are indicated with bold names.

671

- Figure S1. Fully resolved trees from the 1,541 locus analysis with support values. A.
- 673 Bayesian tree. **B.** Maximum-likelihood tree.

674

Figure S2. Phylogram of the 1,541 locus Bayesian tree.

- 677 Figure S3. Fully resolved trees from the 416 locus analysis with support values. A. Bayesian
- 678 tree. **B.** Maximum-likelihood tree. **C.** Species tree.







Θ >0.95 PP, <70% BS

○ >0.90 PP, not in ML tree



Pittidae Psittacidae Megalaimidae Picidae Momotidae Phoeniculidae Coliidae Tytonidae Cathartidae Accipitridae Charadriidae Phalacrocoracidae Balaenicipitidae Scopidae Hydrobatidae Spheniscidae Gaviidae Otididae Psophiidae Musophagidae Trochilidae Nyctibiidae Opisthocomidae Pteroclididae Eurypygidae Phaethontidae Podicipedidae Phoenicopteridae Columbidae Phasianidae



Table S1. Indels greater than 1 bp. Informative indels (n=13) that corroborate Bayesian

phylogeny are indicated with bold names.

UCE ¹	size ²	type ³	Species (informative indels in bold)
chr8 4091	2	deletion	Rhinopomastus, Sphyrapicus
chr1 32309	3	insertion	Pitta, Rhinopomastus, Psittacula, Momotus, Podiceps,
chr3 5661	2	insertion	Rhinopomastus, Sphyrapicus
chr3 5661	3	deletion	Eurypyga, Opisthocomus
chr13 707	6	deletion	Eurypyga, Treron
chr9 3551	4	deletion	Colibri, Rhinopomastus, Treron, Eurypyga
chr9_3551	7	deletion	Megalaima, Sphyrapicus
chr9 3551	3	deletion	Psittacula, Ardeotis
chr2 21162	4	deletion	Opisthocomus, Treron, Phoenicopterus, Podiceps
chr13_2902	3	insertion	Gampsonyx, Phalacrocorax
chr7_6244	5	insertion	Balaeniceps, Phalacrocorax
chr2_3317	4	deletion	Scopus, Balaeniceps
chr15_3386	4	deletion	Psittacula, Gampsonyx
chr15_3386	4	deletion	Urocolius, Scopus
chr1_32247	4	deletion	Momotus, Urocolius
chr1_32247	4	deletion	Phoenicopterus, Podiceps
chr3_5522	10	deletion	Sphyrapicus, Phaethon
chr5_10912	2	deletion	Megalaima, Sphyrapicus
chr2_23600	5	insertion	Megalaima, Sphyrapicus
chr7_10289	2	deletion	Momotus, Sphyrapicus
chr8_5177	6	deletion	Megalaima, Urocolius
chr1_32424	2	deletion	Colibri, Ardeotis
chr6_4126	6	insertion	Colibri, Pterocles, Rhinopomastus, Gampsonyx, Podiceps, Psophia
chr6_4126	4	insertion	Pitta, Gampsonyx
chr12_1611	4	deletion	Momotus, Sphyrapicus, Megalaima
chr2_12990	4	deletion	Megalaima, Sphyrapicus
chr3_19997	2	deletion	Rhinopomastus, Urocolius, Psophia
chr7_10443	3	deletion	Megalaima, Treron, Sphyrapicus
chr8_4221	3	deletion	Rhinopomastus, Motmotus, Sphyrapicus
chr1_15632	3	deletion	Sphyrapicus, Megalaima, Opisthocomus
chr11_3419	3	deletion	Balaeniceps, Motmotus, Gampsonyx
chr7_10549	4	deletion	Tauraco, Phalacrocorax
chr15_2007	2	deletion	Sphyrapicus, Megalaima, Psittacula, Tauraco, Podiceps
chr9_3633	6	deletion	Scopus, Balaeniceps
chr2_18663	2	deletion	Rhinopomastus, Eurypyga
chr6_8088	4	deletion	Nyctibius, Psittacula, Oceanites
chr1_28710	3	deletion	Sphyrapicus, Eudyptyla
chr1_28710	3	deletion	Sphyrapicus, Megalaima
chr11_4777	3	deletion	Phoenicopterus, Podiceps
chr5_14389	2	deletion	Megalaima, Sphyrapicus
chr1_5427	2	deletion	Balaeniceps, Scopus
chr5_2017	2	deletion	Megalaima, Sphyrapicus
chr2_18589	2	deletion	Cathartes, Psophia
chr2_18589	2	deletion	Rhinopomastus, Psittacula, Ardeotis

1 Location relative to chicken genome

2 in base pairs

3 relative to chicken outgroup

Figure S1. Fully resolved trees from the 1,541 locus analysis with support values. A. Bayesian tree. **B.** maximum-likelihood tree.





Figure S2. Phylogram of the 1,541 locus Bayesian tree.

0.004 substitutions/site

Figure S3. Fully resolved trees from the 416 locus analysis with support values. A. Bayesian

tree. B. Maximum-likelihood tree. C. Species tree.

