# The perturbation of the group inverse under the stable perturbation in a unital ring

Fapeng Du<sup>\*</sup>

School of Mathematical & Physical Sciences, Xuzhou Institute of Technology Xuzhou 221008, Jiangsu Province, P.R. China

Yifeng Xue<sup>†</sup>

Department of mathematics and Research Center for Operator Algebras East China Normal University, Shanghai 200241, P.R. China

#### Abstract

Let  $\mathfrak{R}$  be a ring with unit 1 and  $a \in \mathfrak{R}$ ,  $\bar{a} = a + \delta a \in \mathfrak{R}$  such that  $a^{\#}$  exists. In this paper, we mainly investigate the perturbation of the group inverse  $a^{\#}$  on  $\mathfrak{R}$ . Under the stable perturbation, we obtain the explicit expressions of  $\bar{a}^{\#}$ . The results extend the main results in [23, 24] and some related results in [22]. As an application, we give the representation of the group inverse of the

matrix  $\begin{bmatrix} d & b \\ c & 0 \end{bmatrix}$  on the ring  $\mathfrak{R}$  for certain  $d, b, c \in \mathfrak{R}$ .

#### Introduction 1

Let  $\mathfrak{R}$  be a ring with unit and  $a \in \mathfrak{R}$ . we consider an element  $b \in \mathfrak{R}$  and the following equations:

(1) aba = a, (2) bab = b, (3)  $a^k ba = a^k$ , (4) ab = ba.

If b satisfies (1), then b is called a pseudo-inverse or 1-inverse of a. In this case, a is called regular. The set of all 1-inverse of a is denoted by  $a^{\{1\}}$ ; If b satisfies (2), then b is called a 2-inverse of a, and a is called anti-regular. The set of all 2-inverse of a is denoted by  $a^{\{2\}}$ ; If b satisfies (1) and (2), then b is called the generalized inverse of a, denoted by  $a^+$ ; If b satisfies (2), (3) and (4), then b is called the Drazin inverse of a, denoted by  $a^{D}$ . The smallest integer k is called the index of a, denoted by ind(a). If ind(a) = 1, we say a is group invertible and b is the group inverse of a, denoted by  $a^{\#}$ .

The notation so-called stable perturbation of an operator on Hilbert spaces and Banach spaces is introduced by G. Chen and Y. Xue in [4, 6]. Later the notation is generalized to Banach Algebra by Y. Xue in [23] and to Hilbert  $C^*$ -modules by Xu, Wei and Gu in [21]. The stable perturbation of linear operator was widely investigated by many authors. For examples, in [5], G. Chen and Y. Xue study the perturbation for Moore–Penrose inverse of an operator on Hilbert spaces; Q. Xu, C. Song and Y. Wei studied the stable perturbation of the Drazin inverse of the square matrices when  $I - A^{\pi} - B^{\pi}$ is nonsingular in [20] and Q. Huang and W. Zhai worked over the perturbation of closed operators in [16, 17], etc.. Some further results can be found in [10, 11, 12, 13].

The Drazin inverse has many applications in matrix theory, difference equations, differential equations and iterative methods. In 1979, Campbell and Meyer proposed an open problem: how to find an explicit expression for the Drazin inverse of the matrix  $\begin{bmatrix} A & B \\ C & D \end{bmatrix}$  in terms of its sub-block in [1]? The representation of the Drazin inverse of a triangular matrix  $\begin{bmatrix} A & B \\ 0 & D \end{bmatrix}$  has been given in [3, 9, 15]. In [8], Deng and Wei studied the Drazin inverse of the anti-triangular matrix  $\begin{bmatrix} A & B \\ C & 0 \end{bmatrix}$  and given its

representation under some conditions.

<sup>\*</sup>E-mail: jsdfp@163.com

<sup>&</sup>lt;sup>†</sup>Corresponding author, E-mail: yfxue@math.ecnu.edu.cn

In this paper, we investigate the stable perturbation for the group inverse of an element in a ring. Assume that  $1 - a^{\pi} - \bar{a}^{\pi}$  is invertible, we present the expression of  $\bar{a}^{\#}$  and  $\bar{a}^{D}$ . This extends the related results in [22, 24]. As an applications, we study the representation for the group inverse of the anti-triangular matrix  $\begin{bmatrix} d & b \\ c & 0 \end{bmatrix}$  on the ring.

### 2 Some Lemmas

Throughout the paper,  $\mathfrak{R}$  is always a ring with the unit 1. In this section, we give some lemmas:

**Lemma 2.1.** Let  $a, b \in \mathfrak{R}$ . Then 1 + ab is invertible if and only if 1 + ba is invertible. In this case,  $(1 + ab)^{-1} = 1 - a(1 + ba)^{-1}b$  and

 $(1+ab)^{-1}a = a(1+ba)^{-1}, \ b(1+ab)^{-1} = (1+ba)^{-1}b.$ 

**Lemma 2.2.** Let  $a, b \in \mathfrak{R}$ . If 1 + ab is left invertible, then so is 1 + ba.

**Proof.** Let  $c \in \mathfrak{R}$  such that c(1 + ab) = 1. Then

$$1 + ba = 1 + bc(1 + ab)a = 1 + bca(1 + ba).$$

Therefore, (1 - bca)(1 + ba) = 1.

**Lemma 2.3.** Let a be a nonzero element in  $\mathfrak{R}$  such that  $a^+$  exists. If  $s = a^+a + aa^+ - 1$  is invertible in  $\mathfrak{R}$ , then  $a^{\#}$  exists and  $a^{\#} = a^+s^{-1} + (1 - a^+a)s^{-1}a^+s^{-1}$ .

**Proof.** According to [18] or [22, Theorem 4.5.9],  $a^{\#}$  exists. We now give the expression of  $a^{\#}$  as follows.

Put  $p = a^+a$ ,  $q = aa^+$ . Then we have

$$ps = pq = sq, \ qs = qp = sp, \ sa = a^+a^2.$$
 (2.1)

Set  $y = a^+ s^{-1}$ . Then by (2.1),

$$yp = a^+s^{-1}a^+a = a^+aa^+s^{-1} = y = py,$$
  
 $pay = a^+aaa^+s^{-1} = pqs^{-1} = p,$   
 $ypa = a^+s^{-1}a^+aa = a^+a = p.$ 

Put  $a_1 = pap = pa$ ,  $a_2 = (1 - p)ap = (1 - p)a$ . Then  $a = a_1 + a_2$  and it is easy to check that  $a^{\#} = y + a_2 y^2$ . Using (2.1), we can get that  $a^{\#} = a^+ s^{-1} + (1 - a^+ a)a(a^+ s^{-1})^2 = a^+ s^{-1} + (1 - a^+ a)s^{-1}a^+ s^{-1}$ . Let  $M_2(\mathfrak{R})$  denote the matrix ring of all  $2 \times 2$  matrices over  $\mathfrak{R}$  and let  $1_2$  denote the unit of  $M_2(\mathfrak{R})$ .

**Corollary 2.4.** Let  $b, c \in \Re$  have group inverse  $b^{\#}$  and  $c^{\#}$  respectively. Assume that  $k = b^{\#}b + c^{\#}c - 1$  is invertible in  $\Re$ . Then  $\begin{bmatrix} 0 & b \\ c & 0 \end{bmatrix}^{\#}$  exists with  $\begin{bmatrix} 0 & b \\ c & 0 \end{bmatrix}^{\#} = \begin{bmatrix} 0 & k^{-1}c^{\#}k^{-1} \\ k^{-1}b^{\#}k^{-1} & 0 \end{bmatrix}$ . In particular, when  $b^{\#}bc^{\#}c = b^{\#}b$  and  $c^{\#}cb^{\#}b = c^{\#}c$ ,  $\begin{bmatrix} 0 & b \\ c & 0 \end{bmatrix}^{\#} = \begin{bmatrix} 0 & b^{\#}bc^{\#} \\ c^{\#}cb^{\#} & 0 \end{bmatrix}$ .

**Proof.** Set 
$$a = \begin{bmatrix} 0 & b \\ c & 0 \end{bmatrix}$$
. Then  $a^+ = \begin{bmatrix} 0 & c^\# \\ b^\# & 0 \end{bmatrix}$  and  
 $a^+a + aa^+ - 1_2 = \begin{bmatrix} b^\# b + c^\# c - 1 & 0 \\ 0 & b^\# b + c^\# c - 1 \end{bmatrix} = \begin{bmatrix} k \\ k \end{bmatrix}$ 

is invertible in  $M_2(\mathfrak{R})$ . Noting that  $bb^{\#}k^{-1} = k^{-1}cc^{\#}$ . Thus, by Lemma 2.3,

$$\begin{aligned} a^{\#} &= a^{+} \begin{bmatrix} k^{-1} \\ k^{-1} \end{bmatrix} + (1_{2} - a^{+}a) \begin{bmatrix} k^{-1} \\ k^{-1} \end{bmatrix} a^{+} \begin{bmatrix} k^{-1} \\ k^{-1} \end{bmatrix} \\ &= \begin{bmatrix} 0 & c^{\#}k^{-1} + (1 - c^{\#}c)k^{-1}c^{\#}k^{-1} \\ b^{\#}k^{-1} + (1 - b^{\#}b)k^{-1}b^{\#}k^{-1} & 0 \end{bmatrix} \\ &= \begin{bmatrix} 0 & k^{-1}c^{\#}k^{-1} \\ k^{-1}b^{\#}k^{-1} & 0 \end{bmatrix}. \end{aligned}$$

When  $b^{\#}bc^{\#}c = b^{\#}b$  and  $c^{\#}cb^{\#}b = c^{\#}c$ ,  $k^{-1} = k$ . In this case,  $\begin{bmatrix} 0 & b \\ c & 0 \end{bmatrix}^{\#} = \begin{bmatrix} 0 & b^{\#}bc^{\#} \\ c^{\#}cb^{\#} & 0 \end{bmatrix}$ .

**Lemma 2.5.** Let  $a, b \in \mathfrak{R}$  and p be a non-trivial idempotent element in  $\mathfrak{R}$ , i.e.,  $p \neq 0, 1$ . Put x = pap + pb(1-p).

- (1) If pap is group invertible and  $(pap)(pap)^{\#}b(1-p) = pb(1-p)$ , then x is group invertible too and  $x^{\#} = (pap)^{\#} + [(pap)^{\#}]^2 pb(1-p).$
- (2) If x is group invertible, then so is the pap and  $(pap)(pap)^{\#}b(1-p) = pb(1-p)$ .

**Proof.** (1) It is easy to check that  $p(pap)^{\#} = (pap)^{\#}p = (pap)^{\#}$ . Put  $y = (pap)^{\#} + [(pap)^{\#}]^2 pb(1-p)$ .

Then xyx = x, yxy = y and xy = yx, i.e.,  $y = x^{\#}$ . (2) Set  $y_1 = px^{\#}p$ ,  $y_2 = px^{\#}(1-p)$ ,  $y_3 = (1-p)x^{\#}p$  and  $y_4 = (1-p)x^{\#}(1-p)$ . Then  $x^{\#} = y_1 + y_2 + y_3 + y_4$ . From  $xx^{\#}x = x$ ,  $x^{\#}xx^{\#} = x^{\#}$  and  $xx^{\#} = x^{\#}x$ , we can obtain that  $y_3 = y_4 = 0$  and

 $(pap)y_1(pap) = pap, \quad y_1(pap)y_1 = y_1, \quad y_1(pap) = (pap)y_1, \quad y_1(pap)pb(1-p) = pb(1-p),$ 

that is,  $(pap)^{\#} = y_1$  and  $(pap)(pap)^{\#}b(1-p) = pb(1-p)$ .

At the end of this section, we will introduce the notation of stable perturbation of an element in a ring.

Let  $\mathcal{A}$  be a unital Banach algebra and  $a \in \mathcal{A}$  such that  $a^+$  exists. Let  $\bar{a} = a + \delta a \in \mathcal{A}$ . Recall from [23] that  $\bar{a}$  is a stable perturbation of a if  $\bar{a}\mathcal{A} \cap (1 - aa^+)\mathcal{A} = \{0\}$ . This notation can be easily extended to the case of ring as follows.

**Definition 2.6.** Let  $a \in \mathfrak{R}$  such that  $a^+$  exists and let  $\bar{a} = a + \delta a \in R$ . We say  $\bar{a}$  is a stable perturbation of a if  $\bar{a}\mathfrak{R} \cap (1 - aa^+)\mathfrak{R} = \{0\}.$ 

Using the same methods as appeared in the proofs of [23, Proposition 2.2] and [22, Theorem 2.4.7], we can obtain:

**Proposition 2.7.** Let  $a \in \mathfrak{R}$  and  $\bar{a} = a + \delta a \in \mathfrak{R}$  such that  $a^+$  exists and  $1 + a^+ \delta a$  is invertible in  $\mathfrak{R}$ . Then the following statements are equivalent:

- (1)  $\bar{a}^+$  exists and  $\bar{a}^+ = (1 + a^+ \delta a)^{-1} a^+$ .
- (2)  $\bar{a}\mathfrak{R} \cap (1 aa^+)\mathfrak{R} = \{0\}$  (that is,  $\bar{a}$  is a stable perturbation if a).
- (3)  $\bar{a}(1+a^+\delta a)^{-1}(1-a^+a)=0.$
- (4)  $(1 aa^+)(1 + \delta aa^+)^{-1}\bar{a} = 0.$
- (5)  $(1 aa^+)\delta a(1 a^+a) = (1 aa^+)\delta a(1 + a^+\delta a)^{-1}a^+\delta a(1 a^+a).$
- (6)  $\Re \bar{a} \cap \Re (1 a^+ a) = \{0\}.$

#### Main results 3

In this section, we investigate the stable perturbation for group inverse and Drazin inverse of an element a in  $\mathfrak{R}$ .

Let  $a \in \mathfrak{R}$  and  $\bar{a} = a + \delta a \in \mathfrak{R}$  such that  $a^{\#}$  exists and  $1 + a^{\#} \delta a$  is invertible in  $\mathfrak{R}$ . Put  $a^{\pi} = (1 - a^{\#} a)$ ,  $\Phi(a) = 1 + \delta a a^{\pi} \delta a [(1 + a^{\#} \delta a)^{-1} a^{\#}]^2$  and  $B = \Phi(a)(1 + \delta a a^{\#}), C(a) = a^{\pi} \delta a (1 + a^{\#} \delta a)^{-1} a^{\#}.$  These symbols will be used frequently in this section.

**Lemma 3.1.** Let  $a \in \mathfrak{R}$  and  $\bar{a} = a + \delta a \in \mathfrak{R}$  such that  $a^{\#}$  exists and  $1 + a^{\#}\delta a$  is invertible in  $\mathfrak{R}$ . Suppose that  $\Phi(a)$  is invertible, then  $(Ba)^{\#} = Baa^{\#}B^{-1}a^{\#}B^{-1}$ .

**Proof.** Put  $P = aa^{\#}$ . Noting that  $\Phi(a)(1-P) = 1-P$ , we have  $P\Phi(a)P = P\Phi(a)$ ,  $\Phi^{-1}(a)(1-P) = P\Phi(a)$ . (1-P) and PBP = PB,  $B^{-1}(1-P) = (1+\delta aa^{\#})^{-1}(1-P)$ ,  $a^{\#}B^{-1}(1-P) = 0$ , i.e.,  $a^{\#}B^{-1} = a^{\#}B^{-1}P$ . Thus,  $BPB^{-1}Ba = Ba$  and

$$(Ba)(Baa^{\#}B^{-1}a^{\#}B^{-1}) = BPB^{-1} = (Baa^{\#}B^{-1}a^{\#}B^{-1})(Ba), \\ (Baa^{\#}B^{-1}a^{\#}B^{-1})(BPB^{-1}) = Baa^{\#}B^{-1}a^{\#}B^{-1}.$$

These indicate  $(Ba)^{\#} = Baa^{\#}B^{-1}a^{\#}B^{-1}$ .

**Theorem 3.2.** Let  $a \in \mathfrak{R}$  such that  $a^{\#}$  exists. Let  $\bar{a} = a + \delta a \in \mathfrak{R}$  with  $1 + a^{\#}\delta a$  invertible in  $\mathfrak{R}$ . Suppose that  $\Phi(a)$  is invertible and  $\bar{a}\mathfrak{R} \cap (1 - aa^{\#})\mathfrak{R} = \{0\}$ . Put  $D(a) = (1 + a^{\#}\delta a)^{-1}a^{\#}\Phi^{-1}(a)$ . Then  $\bar{a}^{\#}$  exists with

$$\bar{a}^{\#} = (1 + C(a))(D(a) + D^2(a)\delta aa^{\pi})(1 - C(a))$$

*Proof.* Put  $P = aa^{\#}$ . By Proposition 2.7 (3), we have  $a^{\pi}(1 + \delta aa^{\#})^{-1}\bar{a} = 0$  and

$$P\bar{a}(1+a^{\#}\delta a)^{-1} = a(aa^{\#}+a^{\#}\delta a)(1+a^{\#}\delta a)^{-1} = a(1+a^{\#}\delta a-a^{\pi})(1+a^{\#}\delta a)^{-1} = a(aa^{\#}+a^{\#}\delta a)(1+a^{\#}\delta a)(1+a^{\#}\delta a)(1+a^{\#}\delta a)^{-1} = a(aa^{\#}+a^{\#}\delta a)(1+a^{\#}\delta a)(1+a$$

Thus, we have

$$\begin{split} &(1-C(a))\bar{a}(1+C(a)) \\ &= [P+a^{\pi}(1+\delta aa^{\#})^{-1}]\bar{a}[1+a^{\pi}\delta a(1+a^{\#}\delta a)^{-1}a^{\#}] \\ &= P\bar{a}[1+a^{\pi}\delta a(1+a^{\#}\delta a)^{-1}a^{\#}] \\ &= P\bar{a}+P\bar{a}a^{\pi}\delta a(1+a^{\#}\delta a)^{-1}a^{\#} \\ &= P\bar{a}+P\delta aa^{\pi}\delta a(1+a^{\#}\delta a)^{-1}a^{\#}] \\ &= P\delta a+P[a+\delta aa^{\pi}\delta a(1+a^{\#}\delta a)^{-1}a^{\#}] \\ &= P\delta a(1-P)+P\delta aP+P[a+\delta aa^{\pi}\delta a(1+a^{\#}\delta a)^{-1}a^{\#}]P \\ &= P\delta a(1-P)+P[\bar{a}+a+\delta aa^{\pi}\delta a(1+a^{\#}\delta a)^{-1}a^{\#}]P \\ &= P\delta a(1-P)+P[\bar{a}(1+a^{\#}\delta a)^{-1}+\delta aa^{\pi}\delta a(1+a^{\#}\delta a)^{-1}a^{\#}(1+a^{\#}\delta a)^{-1}](1+a^{\#}\delta a)P \\ &= P\delta a(1-P)+P[a+\delta aa^{\pi}\delta a(1+a^{\#}\delta a)^{-1}a^{\#}(1+a^{\#}\delta a)^{-1}](1+a^{\#}\delta a)P \\ &= P\delta a(1-P)+P[a+\delta aa^{\pi}\delta a(1+a^{\#}\delta a)^{-1}a^{\#}(1+a^{\#}\delta a)^{-1}]a^{\#}(1+\delta aa^{\#})a \\ &= P\delta a(1-P)+P[a+\delta aa^{\pi}\delta a(1+a^{\#}\delta a)^{-1}a^{\#}(1+a^{\#}\delta a)^{-1}a^{\#}](1+\delta aa^{\#})a \\ &= P\delta a(1-P)+P[aa^{\#}+\delta aa^{\pi}\delta a(1+a^{\#}\delta a)^{-1}a^{\#})^{2}](1+\delta aa^{\#})a \\ &= P\delta a(1-P)+P[1+\delta aa^{\pi}\delta a((1+a^{\#}\delta a)^{-1}a^{\#})^{2}](1+\delta aa^{\#})a \\ &= P\delta a(1-P)+P\Phi(a)(1+\delta aa^{\#})aP. \end{split}$$

By Lemma 3.1, we have

$$P(Ba)^{\#}P = PBaa^{\#}B^{-1}a^{\#}B^{-1}P = PBPB^{-1}a^{\#}B^{-1}P = a^{\#}B^{-1} = P(Ba)^{\#}$$

and  $PBa(Ba)^{\#}\delta a = P\delta a$ . So  $P(Ba)^{\#}P(Ba)P = P(Ba)^{\#}(Ba)$  and

$$P(Ba)PP(Ba)^{\#}P = P(Ba)(Ba)^{\#}P = P(Ba)^{\#}(Ba)P = P(Ba)^{\#}P(Ba)P$$
$$P(Ba)^{\#}P(Ba)P(Ba)^{\#}P = P(Ba)^{\#}(Ba)(Ba)^{\#}P = P(Ba)^{\#}P,$$
$$P(Ba)P(Ba)^{\#}P(Ba)P = P(Ba)P(Ba)^{\#}(Ba)P = P(Ba)P,$$

i.e.,  $(P(Ba)P)^{\#} = P(Ba)^{\#} = a^{\#}B^{-1}$ . So  $P(Ba)P(P(Ba)P)^{\#} = P$  and hence, we have by Lemma 2.5 (1),

$$[(1 - C(a))\bar{a}(1 + C(a))]^{\#} = a^{\#}B^{-1} + [a^{\#}B^{-1}]^{2}\delta a(1 - P).$$

Therefore,

$$\begin{split} \bar{a}^{\#} &= (1+C(a))[(1-C(a))\bar{a}(1+C(a))]^{\#}(1-C(a)) \\ &= (1+C(a))(D(a)+D^{2}(a)\delta aa^{\pi})(1-C(a)) \\ &= (1+a^{\pi}\delta a(1+a^{\#}\delta a)^{-1}a^{\#})(1+a^{\#}\delta a)^{-1}a^{\#}\left[1+\delta aa^{\pi}\delta a[(1+a^{\#}\delta a)^{-1}a^{\#}]^{2}\right]^{-1} \\ &\times \left[1+(1+a^{\#}\delta a)^{-1}a^{\#}\left[1+\delta aa^{\pi}\delta a[(1+a^{\#}\delta a)^{-1}a^{\#}]^{2}\right]^{-1}\delta aa^{\pi}\right](1-a^{\pi}\delta a(1+a^{\#}\delta a)^{-1}a^{\#}). \quad \Box \end{split}$$

Now we consider the case when  $a \in \mathfrak{R}$  and  $\bar{a} = a + \delta a \in \mathfrak{R}$  such that  $a^{\#}$ ,  $\bar{a}^{\#}$  exist. Firstly, we have **Proposition 3.3.** Let  $a \in \mathfrak{R}$ ,  $\bar{a} = a + \delta a \in \mathfrak{R}$  such that  $a^{\#}$ ,  $\bar{a}^{\#}$  exist. Then the following statements are equivalent:

(1) 
$$\mathfrak{R} = \bar{a}\mathfrak{R} \dotplus (1 - aa^{\#})\mathfrak{R} = a\mathfrak{R} \dotplus (1 - \bar{a}\bar{a}^{\#})\mathfrak{R} = \mathfrak{R}\bar{a} \dotplus \mathfrak{R}(1 - aa^{\#}) = \mathfrak{R}a \dotplus \mathfrak{R}(1 - \bar{a}\bar{a}^{\#}).$$

- (2)  $K = K(a, \bar{a}) = \bar{a}\bar{a}^{\#} + aa^{\#} 1$  is invertible.
- (3)  $\bar{a}\mathfrak{R} \cap (1 aa^{\#})\mathfrak{R} = \{0\}, \ \mathfrak{R}\bar{a} \cap \mathfrak{R}(1 aa^{\#}) = \{0\} \ and \ 1 + \delta aa^{\#} \ is \ invertible.$

**Proof.** (1)  $\Rightarrow$  (2): Since  $\Re = \bar{a}\Re + (1 - aa^{\#})\Re = a\Re + (1 - \bar{a}\bar{a}^{\#})\Re$ , we have for any  $y \in \Re$ , there are  $y_1 \in \mathfrak{R}, y_2 \in \mathfrak{R}$  such that

$$(1 - \bar{a}\bar{a}^{\#})y = (1 - \bar{a}\bar{a}^{\#})(1 - aa^{\#})y_1, \ \bar{a}\bar{a}^{\#}y = \bar{a}\bar{a}^{\#}aa^{\#}y_2$$

Put  $z = aa^{\#}y_2 - (1 - aa^{\#})y_1$ . Then

$$K(a,\bar{a})z = (\bar{a}\bar{a}^{\#} + aa^{\#} - 1)(aa^{\#}y_2 - (1 - aa^{\#})y_1) = y.$$

Since  $\Re = \Re \bar{a} + \Re (1 - aa^{\#}) = \Re \bar{a} + \Re (1 - \bar{a}\bar{a}^{\#})$ , we have for any  $y \in \Re$ , there are  $y_1, y_2 \in \Re$  such that

$$y(1 - \bar{a}\bar{a}^{\#}) = y_1(1 - aa^{\#})(1 - \bar{a}\bar{a}^{\#}), y\bar{a}\bar{a}^{\#} = y_2aa^{\#}\bar{a}\bar{a}^{\#}.$$

Put  $z = y_2 a a^{\#} - y_1 (1 - a a^{\#})$ . Then

$$zK(a,\bar{a}) = (y_2aa^{\#} - y_1(1 - aa^{\#}))(\bar{a}\bar{a}^{\#} + aa^{\#} - 1) = y.$$

The above indicates  $K(a, \bar{a})$  is invertible when we take y = 1.

 $(2) \Rightarrow (3)$ : Let  $y \in \bar{a} \Re \cap (1 - a a^{\#}) \Re$ . Then  $\bar{a} \bar{a}^{\#} y = y, a^{\#} y = 0$ . Thus  $K(a, \bar{a}) y = 0$  and hence y = 0, that is,  $\bar{a}\mathfrak{R} \cap (1 - aa^{\#})\mathfrak{R} = \{0\}$ . Similarly, we have  $\mathfrak{R}\bar{a} \cap \mathfrak{R}(1 - aa^{\#}) = \{0\}$ . Let  $T = aK^{-1}\bar{a}^{\#} - a^{\pi}$ . Since  $\bar{a}a^{\#}aK^{-1} = \bar{a}$ , we have  $(1 + \delta aa^{\#})T = K$ , that is,  $(1 + \delta aa^{\#})$  has

right inverse  $TK^{-1}$ .

Since  $K^{-1}aa^{\#}\bar{a} = \bar{a}$ , we have  $(\bar{a}^{\#}K^{-1}a - a^{\pi})(1 + a^{\#}\delta a) = K$ , that is,  $1 + a^{\#}\delta a$  has left inverse  $K^{-1}(\bar{a}^{\#}K^{-1}a - a^{\pi})$ . This indicates that  $1 + \delta a a^{\#}$  has left inverse  $1 - \delta a K^{-1}(\bar{a}^{\#}K^{-1}a - a^{\pi})a^{\#}$  by Lemma 2.2. Finally,  $1 + \delta a a^{\#}$  is invertible.

 $(3) \Rightarrow (1)$ : By Lemma 2.1,  $1 + a^{\#} \delta a$  is also invertible. So from

$$1 + \delta a a^{\#} = \bar{a} a^{\#} + (1 - a a^{\#}), \ 1 + a^{\#} \delta a = a^{\#} \bar{a} + (1 - a a^{\#})$$

and Lemma 2.7, we get that

$$\mathfrak{R} = \bar{a}\mathfrak{R} \dotplus (1 - aa^{\#})\mathfrak{R} = \mathfrak{R}\bar{a} \dotplus \mathfrak{R}(1 - aa^{\#}).$$

We now prove that

$$\mathfrak{R} = a\mathfrak{R} + (1 - \bar{a}\bar{a}^{\#})\mathfrak{R} = \mathfrak{R}a + \mathfrak{R}(1 - \bar{a}\bar{a}^{\#}), \ a\mathfrak{R} \cap (1 - \bar{a}\bar{a}^{\#})\mathfrak{R} = \mathfrak{R}a \cap \mathfrak{R}(1 - \bar{a}\bar{a}^{\#}) = \{0\}.$$

For any  $y \in a\mathfrak{R} \cap (1 - \bar{a}\bar{a}^{\#})\mathfrak{R}$ , we have  $aa^{\#}y = y$ ,  $\bar{a}y = 0$ . So  $(1 + a^{\#}\delta a)y = (1 - a^{\#}a)y = 0$  and hence y = 0.

Similarly, we have  $\Re a \cap \Re(1 - \bar{a}\bar{a}^{\#}) = \{0\}.$ 

By Lemma 2.7,  $\bar{a}^+ = (1 + a^{\#} \delta a)^{-1} a^{\#}$  and  $\bar{a}^+ \bar{a} = (1 + a^{\#} \delta a)^{-1} a^{\#} a (1 + a^{\#} \delta a)$ . So  $(1 - \bar{a}^+ \bar{a}) \Re = (1 + a^{\#} \delta a)^{-1} a^{\#} a (1 + a^{\#} \delta a)$ .  $(1+a^{\#}\delta a)^{-1}(1-a^{\#}a)\mathfrak{R}$ . From  $(1-\bar{a}^{\#}\bar{a})(1-\bar{a}^{+}\bar{a})=1-\bar{a}^{+}\bar{a}$ , we get that  $(1-\bar{a}^{+}\bar{a})\mathfrak{R} \subset (1-\bar{a}^{\#}\bar{a})\mathfrak{R}$ . Note that  $a\mathfrak{R} = a^{\#}\mathfrak{R}$  and  $(1 + a^{\#}\delta a)a\mathfrak{R} = a^{\#}(1 + \delta aa^{\#})\mathfrak{R} = a^{\#}a\mathfrak{R}$ . So

$$\mathfrak{R} \supset a\mathfrak{R} + (1 - \bar{a}^{\#}\bar{a})\mathfrak{R} \supset (1 + a^{\#}\delta a)^{-1}a^{\#}a\mathfrak{R} + (1 + a^{\#}\delta a)^{-1}(1 - a^{\#}a)\mathfrak{R} = \mathfrak{R}.$$

Similarly, we can get  $\Re a + \Re(1 - \bar{a}^{\#}\bar{a}) = \Re$ .

Now we present a theorem which can be viewed as the inverse of Theorem 3.2 as follows:

**Theorem 3.4.** Let  $a \in \mathfrak{R}$  and  $\bar{a} = a + \delta a \in \mathfrak{R}$  such that  $a^{\#}$ ,  $\bar{a}^{\#}$  exist. If  $K(a, \bar{a})$  is invertible, then  $\Phi(a)$  is invertible.

**Proof.** Since  $K(a, \bar{a})$  is invertible, we have  $\bar{a}\mathfrak{R} \cap (1 - aa^{\#})\mathfrak{R} = \{0\}$  and  $1 + \delta aa^{\#}$  is invertible in  $\mathfrak{R}$  by Propositon 3.3. Thus, from the proof of Theorem 3.2, we have

$$(1 - C(a))\bar{a}(1 + C(a)) = P\delta a(1 - P) + P\Phi(a)(1 + \delta aa^{\#})aP$$
  
= PBaP + P\delta a(1 - P).

Since  $(1 - C(a))\bar{a}(1 + C(a))$  is group invertible, it follows from Lemma 2.5 (2) that PBaP is group invertible and  $PBa(PBa)^{\#}\delta a(I - P) = P\delta a(I - P)$ . Consequently,

$$[(1 - C(a))\bar{a}(1 + C(a))]^{\#} = (1 - C(a))\bar{a}^{\#}(1 + C(a))$$
$$= (PBa)^{\#}P + ((PBa)^{\#})^{2}\delta a(1 - P).$$

Thus,

$$(1 - C(a))\bar{a}\bar{a}^{\#}(1 + C(a)) = [PBaP + P\delta a(1 - P)][(PBa)^{\#}P + ((PBa)^{\#})^{2}\delta a(1 - P)]$$
  
$$= PBa(PBa)^{\#}P + (PBa)^{\#}\delta a(1 - P)$$
  
$$(1 - C(a))K(a,\bar{a})(1 + C(a)) = (1 - C(a))\bar{a}\bar{a}^{\#}(1 + C(a)) - (1 - C(a))a^{\pi}(1 + C(a))$$
  
$$= (1 - C(a))\bar{a}\bar{a}^{\#}(1 + C(a)) - a^{\pi}(1 + C(a))$$
  
$$= PBa(PBa)^{\#}P + P(PBa)^{\#}\delta a(1 - P) - (1 - P)C(a)P - (1 - P)C(a)$$

Since  $(1 - C(a))K(a, \bar{a})(1 + C(a))$  is invertible, we get that

$$\rho(a) = (PBa)^{\#} PBa - (PBa)^{\#} \delta aC(a) = PBa[(PBa)^{\#}]^2 (PBa - \delta aC(a))$$

is invertible in  $P\Re P$ . So we have  $P = PBa[(Ba)^{\#}]^2(Ba - \delta aC(a))\rho^{-1}(a)$  and that  $\Phi(a)$  has right inverse.

Set  $E(a) = a^{\#}(1 + \delta a a^{\#})^{-1} \delta a a^{\pi}$ . Then  $1 - E(a) = P + (1 + a^{\#} \delta a)^{-1} a^{\pi}$  and  $(1 - E(a))^{-1} = 1 + E(a)$ . From Lemma 2.7, we have  $\bar{a}(1 + a^{\#} \delta a)^{-1} a^{\pi} = 0$  and

$$a^{\#}(1+\delta aa^{\#})^{-1}\bar{a} = (1+a^{\#}\delta a)^{-1}a^{\#}\bar{a} = (1+a^{\#}\delta a)^{-1}(1+a^{\#}\delta a-a^{\pi})$$
$$= 1-(1+a^{\#}\delta a)^{-1}a^{\pi}.$$

Put  $\psi(a) = 1 + [(1 + a^{\#} \delta a)^{-1} a^{\#}]^2 \delta a a^{\pi} \delta a$  and  $R = (1 + a^{\#} \delta a) \psi(a)$ . Then  $(1 - P)\psi(a) = 1 - P$ ,  $P\psi(a)P = \psi(a)P$  and

$$\begin{split} &(1+E(a))\bar{a}(1-E(a))\\ &= [1+a^{\#}(1+\delta aa^{\#})^{-1}\delta aa^{\pi}]\bar{a}[P+(1+a^{\#}\delta a)^{-1}a^{\pi}]\\ &= [1+a^{\#}(1+\delta aa^{\#})^{-1}\delta aa^{\pi}]\bar{a}P\\ &= \bar{a}P+a^{\#}(1+\delta aa^{\#})^{-1}\delta aa^{\pi}\bar{a}P\\ &= \bar{a}P+a^{\#}(1+\delta aa^{\#})^{-1}\delta aa^{\pi}\delta aP\\ &= aP+\delta aP+a^{\#}(1+\delta aa^{\#})^{-1}\delta aa^{\pi}\delta aP\\ &= (1-P)\delta aP+P\delta aP+aP+a^{\#}(1+\delta aa^{\#})^{-1}\delta aa^{\pi}\delta aP\\ &= (1-P)\delta aP+P[\bar{a}+a^{\#}(1+\delta aa^{\#})^{-1}\delta aa^{\pi}\delta a]P\\ &= (1-P)\delta aP+P(1+\delta aa^{\#})[(1+\delta aa^{\#})^{-1}\bar{a}+(1+\delta aa^{\#})^{-1}a^{\#}(1+\delta aa^{\#})^{-1}\delta aa^{\pi}\delta a]P\\ &= (1-P)\delta aP+Pa(1+a^{\#}\delta a)[a^{\#}(1+\delta aa^{\#})^{-1}\bar{a}+a^{\#}(1+\delta aa^{\#})^{-1}a^{\#}(1+\delta aa^{\#})^{-1}\delta aa^{\pi}\delta a]P\\ &= (1-P)\delta aP+Pa(1+a^{\#}\delta a)[a^{\#}(1+\delta aa^{\#})^{-1}\bar{a}+[(1+a^{\#}\delta a)^{-1}a^{\#}]^{2}\delta aa^{\pi}\delta a]P\\ &= (1-P)\delta aP+Pa(1+a^{\#}\delta a)[1+[(1+a^{\#}\delta a)^{-1}a^{\#}]^{2}\delta aa^{\pi}\delta a]P\\ &= (1-P)\delta aP+PaRP. \end{split}$$

Since  $(1 + E(a))\bar{a}(1 - E(a))$  is group invertible, we can deduce that aRP is group invertible and

$$[(1 + E(a))\bar{a}(1 - E(a))]^{\#} = (1 + E(a))\bar{a}^{\#}(1 - E(a))$$
$$= P(aRP)^{\#} + (1 - P)\delta a((aRP)^{\#})^{2}$$

and

$$(1 + E(a))\bar{a}\bar{a}^{\#}(1 - E(a)) = [(1 - P)\delta aP + PaRP][P(aRP)^{\#} + (1 - P)\delta a((aRP)^{\#})^{2}]$$
  
=  $PaR(aRP)^{\#} + (1 - P)\delta a(aRP)^{\#}.$ 

Thus, from the invertibility of  $K(a, \bar{a})$ , we get that

$$(1+E(a))K(a,\bar{a})(1-E(a))$$
  
=  $(1+E(a))\bar{a}\bar{a}^{\#}(1-E(a)) - (1+E(a))a^{\pi}(1-E(a))$   
=  $PaRP(aRP)^{\#} + (1-P)\delta aP(aRP)^{\#} - (1+E(a))a^{\pi}$   
=  $PaRP(aRP)^{\#} + (1-P)\delta a(aRP)^{\#} - PE(a)(1-P) - (1-P)$ 

is invertible in  $\mathfrak R$  and hence

$$\eta(a) = aRP(aRP)^{\#} - E(a)\delta a(aRP)^{\#} = [aRP - E(a)\delta a][(aRP)^{\#}]^2 aRP$$
$$= [aRP - E(a)\delta a][(aRP)^{\#}]^2 a(1 + a^{\#}\delta a)\psi(a)P$$

is invertible in  $P\Re P$ . So  $P\psi(a)P$  is left invertible and  $\psi(a)$  is left invertible and hence  $\Phi(a)$  is left invertible by Lemma 2.2. Therefore,  $\Phi(a)$  is invertible.

Let  $a \in \mathfrak{R}$  such that  $a^D$  exists and ind(a) = s. As we know if  $a^D$  exists, then  $a^l$  has group inverse  $(a^l)^{\#}$  and  $a^D = (a^l)^{\#} a^{l-1}$  for any  $l \ge s$ .

From Theorem 3.2 and Theorem 3.4, we have the following corollary:

**Corollary 3.5.** Let a and b be nonzero elements in R such that  $a^D$  and  $b^D$  exist. Put s = ind(a) and t = ind(b). Suppose that  $K(a, b) = bb^D + aa^D - 1$  is invertible in  $\mathfrak{R}$ . Then for any  $l \ge s$  and  $k \ge t$ , we have

- (1)  $1 + (a^D)^l (b^k a^l)$  is invertible in  $\mathfrak{R}$  and  $b^k \mathfrak{R} \cap (1 a^D a) \mathfrak{R} = \{0\}.$
- (2)  $W_{k,l} = 1 + E_{k,l} Z_{k,l} (1 + (a^D)^l E_{k,l})^{-1} (a^D)^l$  is invertible in  $\mathfrak{R}$ , here  $E_{k,l} = b^k a^l$  and  $Z_{k,l} = a^{\pi} E_{k,l} (a^D)^l (1 + E_{k,l} (a^D)^l)^{-1}$ .
- (3)  $b^D = (1 + Z_{k,l})[H_{k,l} + H_{k,l}^2 E_{k,l} a^{\pi}](1 Z_{k,l})b^{k-1}$ , where  $H_{k,l} = (1 + (a^D)^l E_{k,l})^{-1}(a^D)^l W_{k,l}^{-1}$ .

**Proof.** Noting that  $(a^D)^l = (a^l)^{\#}$ ,  $aa^D = a^l(a^l)^{\#}$ ,  $bb^D = b^k(b^k)^{\#}$ ,  $l \ge s$ ,  $k \ge t$ , we have

$$K(a,b) = b^{k}(b^{k})^{\#} + a^{l}(a^{l})^{\#} - 1, \quad 1 + (a^{D})^{l}(b^{k} - a^{l}) = 1 + (a^{l})^{\#}(b^{k} - a^{l}).$$

Applying Theorem 3.2 and Theorem 3.4 to  $b^k$  and  $a^l$ , we can get the assertions.

## 4 The representation of the group inverse of certain matrix on $\Re$

As an application of Theorem 3.2 and Theorem 3.4, we study the representation of the group inverse of  $\begin{bmatrix} d & b \\ c & 0 \end{bmatrix}$  on the ring  $\Re$ .

**Proposition 4.1.** Let  $b, c, d \in \mathfrak{R}$ . Suppose that  $b^{\#}$  and  $c^{\#}$  exist and  $k = b^{\#}b + c^{\#}c - 1$  is invertible. If  $b^{\pi}d = 0$  or  $dc^{\pi} = 0$ , then  $\begin{bmatrix} d & b \\ c & 0 \end{bmatrix}^{\#}$  exists and

$$\begin{bmatrix} d & b \\ c & 0 \end{bmatrix}^{\#} = \begin{bmatrix} -k^{-1}c^{\#}k^{-1}b^{\#}k^{-1}dc^{\pi}k^{-1} & k^{-1}c^{\#}k^{-1} \\ k^{-1}b^{\#}k^{-1}(1+dk^{-1}c^{\#}k^{-1}b^{\#}k^{-1}dc^{\pi}k^{-1}) & -k^{-1}b^{\#}k^{-1}dk^{-1}c^{\#}k^{-1} \end{bmatrix}$$

if  $b^{\pi}d = 0$ . When  $dc^{\pi} = 0$ , we have

$$\begin{bmatrix} d & b \\ c & 0 \end{bmatrix}^{\#} = \begin{bmatrix} -k^{-1}b^{\pi}dk^{-1}c^{\#}k^{-1}b^{\#}k^{-1} & (1+k^{-1}b^{\pi}dk^{-1}c^{\#}k^{-1}b^{\#}d)k^{-1}c^{\#}k^{-1} \\ k^{-1}b^{\#}k^{-1} & -k^{-1}b^{\#}k^{-1}dk^{-1}c^{\#}k^{-1} \end{bmatrix}.$$

*Proof.* Set  $a = \begin{bmatrix} 0 & b \\ c & 0 \end{bmatrix}$ ,  $\delta a = \begin{bmatrix} d & 0 \\ 0 & 0 \end{bmatrix}$  and  $\bar{a} = \begin{bmatrix} d & b \\ c & 0 \end{bmatrix}$ . Since  $b^{\#}bk = kc^{\#}c$ ,  $c^{\#}ck = kb^{\#}b$ , it follows from Corollary 2.4 that

$$\begin{split} 1_{2} + a^{\#} \delta a &= 1_{2} + \begin{bmatrix} 0 & b^{\#} b k^{-1} c^{\#} k^{-1} \\ c^{\#} c k^{-1} b^{\#} k^{-1} & 0 \end{bmatrix} \begin{bmatrix} d & 0 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ k^{-1} b^{\#} k^{-1} d & 1 \end{bmatrix} \\ (1_{2} + a^{\#} \delta a)^{-1} a^{\#} &= \begin{bmatrix} 1 & 0 \\ -k^{-1} b^{\#} k^{-1} d & 1 \end{bmatrix} \begin{bmatrix} 0 & k^{-1} c^{\#} k^{-1} \\ 0 & k^{-1} c^{\#} k^{-1} \end{bmatrix} \\ &= \begin{bmatrix} 0 & k^{-1} c^{\#} k^{-1} \\ k^{-1} b^{\#} k^{-1} & -k^{-1} b^{\#} k^{-1} d k^{-1} c^{\#} k^{-1} \end{bmatrix} \\ &= \begin{bmatrix} 0 & b \\ k^{-1} b^{\#} k^{-1} & -k^{-1} b^{\#} k^{-1} d k^{-1} c^{\#} k^{-1} \end{bmatrix} \\ &= \begin{bmatrix} b c^{\#} c k^{-1} b^{\#} k^{-1} & 0 \\ 0 & c b^{\#} b k^{-1} c^{\#} k^{-1} \end{bmatrix} \\ &= \begin{bmatrix} b b^{\#} k^{-1} & 0 \\ 0 & c c^{\#} k^{-1} \end{bmatrix} \\ &= \begin{bmatrix} b b^{\#} k^{-1} & 0 \\ 0 & c c^{\#} k^{-1} \end{bmatrix} \\ &a^{\pi} = 1 - a a^{\#} = \begin{bmatrix} -c^{\pi} k^{-1} & 0 \\ 0 & -b^{\pi} k^{-1} \end{bmatrix} \\ &= \begin{bmatrix} d & b \\ c & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ -c^{\#} c k^{-1} b^{\#} k^{-1} d & 1 \end{bmatrix} \begin{bmatrix} -c^{\pi} k^{-1} & 0 \\ 0 & -b^{\pi} k^{-1} \end{bmatrix} \\ &= \begin{bmatrix} d - b c^{\#} c k^{-1} b^{\#} k^{-1} d & b \\ c & 0 \end{bmatrix} \begin{bmatrix} -c^{\pi} k^{-1} & 0 \\ 0 & -b^{\pi} k^{-1} \end{bmatrix} \\ &= \begin{bmatrix} d - b c^{\#} c k^{-1} b^{\#} k^{-1} d & b \\ c & 0 \end{bmatrix} \begin{bmatrix} -c^{\pi} k^{-1} & 0 \\ 0 & -b^{\pi} k^{-1} \end{bmatrix} \\ &= \begin{bmatrix} -c^{\pi} k^{-1} d & b \\ c & 0 \end{bmatrix} \begin{bmatrix} -c^{\pi} k^{-1} & 0 \\ 0 & -b^{\pi} k^{-1} \end{bmatrix} = \begin{bmatrix} k^{-1} b^{\pi} d c^{\pi} k^{-1} & 0 \\ 0 & 0 \end{bmatrix} \end{split}$$

and

$$\delta a a^{\pi} \delta a = \begin{bmatrix} d & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} -c^{\pi} k^{-1} & 0 \\ 0 & -b^{\pi} k^{-1} \end{bmatrix} \begin{bmatrix} d & 0 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} -dc^{\pi} k^{-1} d & 0 \\ 0 & 0 \end{bmatrix}$$
$$a^{\pi} \delta a = \begin{bmatrix} -k^{-1} b^{\pi} d & 0 \\ 0 & 0 \end{bmatrix}, \quad \delta a a^{\pi} = \begin{bmatrix} -dc^{\pi} k^{-1} & 0 \\ 0 & 0 \end{bmatrix}.$$

If  $b^{\pi}d = 0$  or  $dc^{\pi} = 0$ , then  $\bar{a}(1 + a^{\#}\delta a)^{-1}a^{\pi} = 0$  and  $\delta aa^{\pi}\delta a = 0$ . Thus,  $\Phi(a) = 1_2$  and  $D(a) = 1_2$  $(1 + a^{\#} \delta a)^{-1} a^{\#} \Phi^{-1}(a) = (1 + a^{\#} \delta a)^{-1} a^{\#}.$ When  $b^{\pi} d = 0$ ,  $C(a) = a^{\pi} \delta a (1 + a^{\#} \delta a)^{-1} a^{\#} = 0$  and

$$D(a)\delta aa^{\pi} = \begin{bmatrix} 0 & k^{-1}c^{\#}k^{-1} \\ k^{-1}b^{\#}k^{-1} & -k^{-1}b^{\#}k^{-1}dk^{-1}c^{\#}k^{-1} \end{bmatrix} \begin{bmatrix} -dc^{\pi}k^{-1} & 0 \\ 0 & 0 \end{bmatrix}$$
$$= \begin{bmatrix} 0 & 0 \\ -k^{-1}b^{\#}k^{-1}dc^{\pi}k^{-1} & 0 \end{bmatrix}.$$

By Theorem 3.2, we have

$$\begin{split} \bar{a}^{\#} &= (1_2 + C(a))(D(a) + D^2(a)\delta a a^{\pi})(1_2 - C(a)) \\ &= D(a)(1_2 + D(a)\delta a a^{\pi}) \\ &= \begin{bmatrix} 0 & k^{-1}c^{\#}k^{-1} \\ k^{-1}b^{\#}k^{-1} & -k^{-1}b^{\#}k^{-1}dk^{-1}c^{\#}k^{-1} \end{bmatrix} \begin{bmatrix} 1 & 0 \\ -k^{-1}b^{\#}k^{-1}dc^{\pi}k^{-1} & 1 \end{bmatrix} \\ &= \begin{bmatrix} a_1 & k^{-1}c^{\#}k^{-1} \\ a_2 & -k^{-1}b^{\#}k^{-1}dk^{-1}c^{\#}k^{-1} \end{bmatrix}, \end{split}$$

where  $a_1 = -k^{-1}c^{\#}k^{-2}b^{\#}k^{-1}dc^{\pi}k^{-1}$ ,  $a_2 = k^{-1}b^{\#}k^{-1} + k^{-1}b^{\#}k^{-1}dk^{-1}c^{\#}k^{-2}b^{\#}k^{-1}dc^{\pi}k^{-1}$ . Since  $cc^{\#}b^{\#} = kb^{\#}$ , it follows that

$$c^{\#}k^{-2}b^{\#} = c^{\#}(c^{\#}c)k^{-1}k^{-1}b^{\#} = c^{\#}k^{-1}k^{-1}c^{\#}cb^{\#} = c^{\#}k^{-1}b^{\#}$$

So  $a_1 = -k^{-1}c^{\#}k^{-1}b^{\#}k^{-1}dc^{\pi}k^{-1}, a_2 = k^{-1}b^{\#}k^{-1}(1 + dk^{-1}c^{\#}k^{-1}b^{\#}k^{-1}dc^{\pi}k^{-1}).$ 

When  $dc^{\pi} = 0$ , we have by Theorem 3.2,

$$\begin{split} \bar{a}^{\#} &= (1_2 + C(a))D(a)(1_2 - C(a)) = (1_2 + C(a))D(a) \\ &= \begin{bmatrix} 1 & -k^{-1}b^{\pi}dk^{-1}c^{\#}k^{-1} \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 0 & k^{-1}c^{\#}k^{-1} \\ k^{-1}b^{\#}k^{-1} & -k^{-1}b^{\#}k^{-1}dk^{-1}c^{\#}k^{-1} \end{bmatrix} \\ &= \begin{bmatrix} -k^{-1}b^{\pi}dk^{-1}c^{\#}k^{-1}b^{\#}k^{-1} & (1+k^{-1}b^{\pi}dk^{-1}c^{\#}k^{-1}dk^{-1}c^{\#}k^{-1} \\ k^{-1}b^{\#}k^{-1} & -k^{-1}b^{\#}k^{-1}dk^{-1}c^{\#}k^{-1} \end{bmatrix}. \quad \Box$$

Combining Proposition 4.1 with Corollary 2.4, we have

**Corollary 4.2.** Let  $b, c, d \in \mathfrak{R}$ . Assume that  $b^{\#}$  and  $c^{\#}$  exist and satisfy conditions:  $b^{\#}bc^{\#}c = b^{\#}b$ ,  $c^{\#}cb^{\#}b = c^{\#}c$ .

(1) If 
$$b^{\pi}d = 0$$
, then  $\begin{bmatrix} d & b \\ c & 0 \end{bmatrix}^{\#} = \begin{bmatrix} b^{\#}bc^{\#}b^{\#}db^{\pi} & b^{\#}bc^{\#} \\ c^{\#}cb^{\#}(1 - db^{\#}bc^{\#}b^{\#}db^{\pi}) & -c^{\#}cb^{\#}db^{\#}bc^{\#} \end{bmatrix}$ 

(2) If 
$$dc^{\pi} = 0$$
, then  $\begin{bmatrix} d & b \\ c & 0 \end{bmatrix}^{\#} = \begin{bmatrix} b^{\pi}db^{\#}bc^{\#}b^{\#} & (1 - b^{\pi}db^{\#}bc^{\#}b^{\#}d)b^{\#}bc^{\#} \\ c^{\#}cb^{\#} & -c^{\#}cb^{\#}db^{\#}bc^{\#} \end{bmatrix}$ 

Recall from [14] that an involution \* on  $\mathfrak{R}$  is an involutory anti–automorphism, that is,

$$(a^*)^* = a, \ (a+b)^* = a^* + b^*, \ (ab)^* = b^*a^*, \ a^* = 0$$
 if and only if  $a = 0$ 

and  $\mathfrak{R}$  is called the \*-ring if  $\mathfrak{R}$  has an involution.

**Corollary 4.3.** Let  $\mathfrak{R}$  be a \*-ring with unit 1 and let p be a nonzero idempotent element in  $\mathfrak{R}$ . Then

$$\begin{bmatrix} pp^* & p \\ p & 0 \end{bmatrix}^{\#} = \begin{bmatrix} pp^*(1-p) & p \\ p - (pp^*)^2(1-p) & -pp^*p \end{bmatrix}, \quad \begin{bmatrix} p^*p & p \\ p & 0 \end{bmatrix}^{\#} = \begin{bmatrix} (1-p)p^*p & p - (1-p)(p^*p)^2 \\ p & -pp^*p \end{bmatrix}.$$

**Proof.** Since  $p^{\#} = p$ , we can get the assertions easily by using Corollary 4.2.

**Remark 4.4.** (1) If  $\mathfrak{R}$  is a skew field and b = c in Proposition 4.1, the conclusion of Proposition 4.1 is contained in [26].

(2) Let p be an idempotent matrix. The group inverse of  $\begin{bmatrix} a & b \\ c & 0 \end{bmatrix}$  is given in [2] for some  $a, b, c \in \{pp^*, p, p^*\}$ . The group inverse of this type of matrices is also discussed in [7].

Acknowledgement. The authors thank to the referee for his (or her) helpful comments and suggestions.

### References

- S.L. Campbell, C.D. Meyer, Generalized inverses of linear transformations. Dover, New York, 1991 (Originally published: Pitman, London, 1979).
- [2] C. Cao and X. Tang, Representations of the group inverse of some 2 × 2 block matrices, International Math. Forum, 1 (2006), 1511–1517.
- [3] N. Castro-Gonzlez, J.J. Koliha, New additive results for the g-Drazin inverse, Proc. Roy. Soc. Edinburgh, 134A (2004), 1085–1097.
- [4] G. Chen and Y. Xue, Perturbation analysis for the operator equation Tx = b in Banach spaces, J. Math. Anal. Appl., **212** (1997), 107–125.
- [5] G. Chen and Y. Xue, The expression of generalized inverse of the perturbed operators under type I perturbation in Hilbert spaces, Linear Algebra Appl. 285 (1998), 1–6.
- [6] G. Chen, Y. Wei and Y. Xue, Perturbation analysis of the least square solution in Hilbert spaces, Linear Algebra Appl. 244 (1996), 69–80.

- [7] C. Deng, A comment on some recent results concerning the Drazin inverse of an anti-triangular block matrix, Filomat, 26(2) (2012), 135–145.
- [8] C. Deng and Y. Wei. A note on the Drazin inverse of an anti-triangular matrix. Linear Algebra Appl., 431 (2009), 1910-1922.
- D.S. Djordjević and P.S. Stanimirović, On the generalized Drazin inverse and generalized resolvent, Czechoslovak Math. J. 51(126)(2001), 617-634.
- [10] F. Du and Y. Xue, Note on stable perturbation of bounded linear operators on Hilbert spaces, Funct. Anal. Appr. Comput., 3(2) (2011), 47–56.
- [11] F. Du and Y. Xue, Perturbation analysis of  $A_{T,S}^{(2)}$  on banach spaces, Electronic J. Linear Algebra, 23 (2012), 586-598.
- [12] F. Du and Y. Xue, The characterizations of the stable perturbation of a closed operator by a linear operator in Banach spaces, Linear Algebra Appl., (accepted).
- [13] F. Du and Y. Xue, Perturbation analysis of  $A_{T,S}^{(2)}$  on Hilbert spaces, (preprint).
- [14] R.E. Hartwig, Block generalized inverse. Archive for Rational Mechanics and Analysis, 61(3) (1976), 197-251.
- [15] R.E. Hartwig, G. Wang and Y. Wei, Some additive results on Drazin inverse, Linear Algebra Appl. 322 (2001), 207–217.
- [16] Q. Huang, On perturbations for oblique projection generalized inverses of closed linear operators in Banach spaces, Linear Algebra Appl., 434 (2011), 2468–2474.
- [17] Q. Huang and W. Zhai, Perturbation and expressions for generalized inverses in Banach spaces and Moore–penrose inverses in Hilbert spaces of closed operators, Linear Algebra Appl., 435 (2011), 117–127.
- [18] P. Patrício and A.D. Costa, On the Drazin index of reular elements, Central Euro. J. Math., 7(2) (2009), 200–205.
- [19] Y. Wang and H. Zhang, Perturbation analysis for oblique projection generalized inverses of closed operators in Banach spaces, Linear Algebra Appl. 426 (2007), 1–11.
- [20] Q. Xu, C. Song and Y. Wei, The Stable perturbation of the Drazin inverse of the square matrices. SIAM J. Matrix Anal. & Appl., 31(3) (2010), 1507–1520.
- [21] Q. Xu, Y. Wei and Y. Gu, Sharp norm-estimation for Moore-Penrose inverses of stable perturbations of Hilbert C<sup>\*</sup>-module operators, SIAM J. Numer. Anal., 47(6) (2010), 4735–4758.
- [22] Y. Xue, Stable perturbations of operators and related topics, World Scientific, 2012.
- [23] Y. Xue, Stable perturbation in Banach spaces, J. Aust. Math. Soc., 83 (2007), 1–14.
- [24] Y. Xue and G. Chen Perturbation analysis for the Drazin inverse under satable perturbation in Banach space. Missouri J. Math. Sci. 19(2) (2007), 106–120.
- [25] Y. Xue and G. Chen, Some equivalent conditions of stable perturbation of operators in Hilbert spaces, Applied Math. Comput. 147 (2004), 765–772.
- [26] J. Zhao and C. Bu, Group inverse for the block matrix with two identical subblocks over skew fields, Electronic J. Linear Algebra, 21 (2010), 63–75.