

**Systematically Dissecting the Global Mechanism of miRNA Functions in
Pluripotent Stem Cells**

Anyou Wang^{1*}, et al

*Corresponding author

Anyou Wang

anyou.wang@alumni.ucr.edu

Running title: Global mechanisms of miRNA roles in stem cells

Key words: miRNAs; global mechanism; stem cells; systems network; pluripotency; functions.

Abstract:

MicroRNAs (miRNAs) critically modulate stem cell properties like pluripotency, but the fundamental mechanism remains largely unknown. This study systematically analyzes multiple-omics data and builds a systems physical network including genome-wide interactions between miRNAs and their targets to reveal the systems mechanism of miRNA functions in mouse pluripotent stem cells. Globally, miRNAs directly repress the pluripotent core factors during differentiation state. Surprisingly, during pluripotent state, the top important miRNAs do not directly regulate the pluripotent core factors as thought, but they only directly target the pluripotent signal pathways and directly repress developmental processes. Furthermore, at pluripotent state miRNAs predominately repress DNA methyltransferases, the core enzymes for DNA methylation. The decreasing methylation repressed by miRNAs in turn activates the top miRNAs and pluripotent core factors, creating an active circuit system to modulate pluripotency. MiRNAs vary their functions with different stem cell states. While miRNAs directly repress pluripotent core factors to facilitate the differentiation during cell differentiation, they also help stem cells to maintain pluripotency by activating pluripotent cores through directly repressing DNA methylation systems and primarily inhibiting development.

Introduction

MicroRNAs (miRNAs), short (~22 nts) conserved endogenous non-coding RNAs, inhibit messenger RNA targets by repressing translation or reducing mRNA stability [1]. MiRNAs critically modulate many cellular events, including the balance between proliferation and differentiation during organ development [1]. In pluripotent stem cells (including induced pluripotent stem cells and embryonic stem cells, referred to as stem cells hereafter), miRNAs play important roles in regulating stem cell bioprocesses [2-6].

miRNAs modulate stem cell pluripotency and differentiation [2-4]. Knocking out the key miRNA processing enzymes Dicer [2-3] or DGCR8 [4] causes stem cells to lose their pluripotency. MiRNA-290 cluster has been proposed to regulate the core pluripotency factors like POU5F1 [7-9]. MiRNA-302/367 cluster has also been used to induce pluripotency [10]. On the other hand, miRNAs like let-7 induce stem cell differentiation [11]. However, these recent studies have mostly focused on individual gene functions in stem cells although genome-wide data might be employed, and the conclusions drawn from these current studies are unavoidably biased on genes selected by these studies. Therefore, these studies only provide partial mechanisms of miRNA functions in stem cells, and the overall systems mechanisms of how miRNAs regulate stem cell processes remain largely elusive.

MiRNAs generally do not work alone to perform their functions [12]. One miRNA might

target more than 100 genes [13-14], and one gene can be repressed by multiple miRNAs in a sequence-specific fashion [12-13, 15]. In turn, proteins can physically bind to the promoters and enhancers of miRNAs to regulate miRNA activations [16]. These binary interactions between miRNAs and proteins would form a complex systematic network. This complexity of miRNA interaction network make it challenged for conventional approaches like gene-knockout to unbiasedly capture the real mechanisms of miRNA functions in stem cells.

This present study employed systems physical network approaches [17] and constructed a comprehensive and unbiased map of genome-wide interactions between miRNAs and their targets to investigate the global basis of miRNA roles in pluripotent stem cells. Results of the present study lay a conceptual framework for future studies and applications of miRNAs in stem cells.

Results

Physical network of miRNA and protein interactions in stem cells

To systematically reveal the roles of miRNAs in stem cells, this study first constructed a systems network [17] of interactions between miRNAs and proteins. These interactions contain binary interactions from two directions, from miRNAs to mRNAs coding for proteins, and from proteins to miRNA promoters and enhancers. The miRNA-targets were inferred from CLIP-seq data, which simultaneously identify miRNAs-mRNA interactions by measuring miRNA-Argonaute complexes [18-19] (materials and

methods). Protein-targets were inferred from ChIP-seq, which measures protein interactions with DNA [20] (Figure 1, Table S1 and materials and methods). The CLIP-seq and ChIP-seq provide data of physical binding interactions. The systems interaction network constructed here includes physical interactions of both miRNA-targets and protein-targets.

The entire network contains ~10,000 nodes and ~233,000 interactions (Figure 1C) and is accessible online ([.....](#)). Both ChIP-seq and CLIP-seq measures genome-wide targets and thus this network provides a global map of miRNA targets in stem cells. For example, the genome-wide interactions between miRNAs and the pluripotent core factors (NANOG, POU5F1 and SOX2) could be extracted from this network (Figure 1D). Due to its natural interactions generated from experimental data, this network would provide accurate interactions between miRNA and their targets, and results generated from this network would be reliable.

The primary role of miRNAs in stem cells

To understand the primary role of all miRNAs activated (up- or down-regulated when compared with somatic cells) at pluripotent state in stem cells, we searched for the biological functions of the network activated by miRNAs in stem cells. To avoid the dataset biases, we included different datasets of miRNAs and genes coding for proteins and selected miRNAs and genes that are only activated with high frequency in all

datasets (Table S2-S3, materials and methods). These activated miRNAs and genes were used to enrich the entire network (Figure 1) to get the activated network using methods as previously described [17]. The network activated by overexpressed miRNAs and down-regulated proteins (Table S2-S3) formed a subnetwork activated by up-regulated miRNAs in stem cells. We run the GO (gene ontology) functional analysis (www.geneontology.org/) of this subnetwork [21] by separately using references of both entire GO annotation database and a set of all up-regulated genes in stem cells to get the less biased enrichment results. Different references generated different enrichment p-value but produced the similar result that the entire subnetwork primarily functions for development, with corrected p-value < 6.517e-62 (Bonferroni correction using entire annotation, referred as corr, hereafter) (Figure S1A). This activated subnetwork was further enriched by the developmental GO term to obtain the developmental module (Figure 2A). This developmental module was decomposed into functional modules based on network topology [22] and it contained 6 sub-functional modules (Materials and Methods). All of these 6 modules primarily function for development (corr p-value < 8.2615E-83, Figure 2A), indicating that the primary function of over-expressed miRNAs at pluripotent state is to repress developmental modules.

The repressing role of up-regulated miRNAs was further evidenced by examining the targets of three primarily represented miRNA groups, the top overexpressed miRNA group, a miR-302/367 cluster, and a single miR-294. First, a total of 17 out of the 20 most important miRNAs, which were selected on the basis of the variance contribution

to the system (material and methods, Table S4), directly target a developmental sub-network (corr $p < 6.3512E-20$, Figure S2). Among the 17 miRNAs, the top 5 miRNAs also target a module that primarily functions for development (corr $p < 8.5158E-23$, Figure 2B-2C). Furthermore, the well-known miR-302 cluster and even a single miRNA, miR-294, also target modules functionally enriched in the developmental category with respectively corr p-value $< 1.4436E-21$ (Figure 3A-3B), and corr p-value $< 1.0157E-29$ (Figure 3C-3D). Therefore, overexpressed miRNAs at pluripotent state primarily repress development. Biologically, to maintain the self-renewal and pluripotency, stem cells have some ways to prevent development and differentiation. This repressing function of miRNAs can help miRNAs claim their contributions to the stem cell properties at the pluripotent state.

On the other hand, the down-regulated miRNAs in stem cells directly target genes that primarily function for metabolism and pluripotency (corr- $p < 3.5159E-42$, Figure S1B, Figure S3). These down-regulated miRNAs become up-regulated during differentiation and directly inhibit metabolism and pluripotency during this state. Together, miRNAs primarily and directly repress development during the pluripotent state while they repress metabolism and pluripotency during cell differentiation.

Paths from activated miRNAs to pluripotent core factors

MiRNAs like miR-302 cluster mediate pluripotency [7, 23], and it thus was assumed that top over-expressed miRNAs such as miR-302 and miRNA-290-295 cluster in stem cells

might directly or indirectly target pluripotent core factors [7, 23]. To investigate if these top miRNAs directly target the core factors, we systematically searched the shortest paths respectively from the top important miRNAs as described above (Materials and Methods, Table S4) to the three core factors (POU5F1, NANOG, and SOX2). Surprisingly, all these top miRNAs, including miR-302b, miR-367, miR-294, and miR-292, do not directly target any core factors (Figure 4A-4D). Actually, the direct basis of these miRNAs regulating the pluripotent core factors seemed blurred because all these miRNAs must go through at least 2 steps to reach any pluripotency core factor. These two steps include a miRNA and a protein, which are not consistently expressed with these miRNAs. This suggested that these top over-expressed miRNAs in stem cells do not directly mediate the pluripotency.

We then globally and unbiasedly searched for up-regulated miRNAs that target the pluripotent core factors (NANOG, POU5F1 and SOX2). Surprisingly, we only found one miRNA (miR-684) that barely up-regulated (~2 fold changed) in stem cells and directly binds to SOX2 (Figure 4E), which was also targeted by down-regulated miRNA-431. This indicated that activated miRNAs (>100miRNAs) do not primarily and directly target the pluripotent core factors. On the other hand, only limited miRNAs, regardless of expression, target POU5F1, while many miRNAs target SOX2 although they might not over-express in stem cells (Figure 1D), suggesting that the connection from miRNAs to the pluripotent core factors primarily go through SOX2, in contrast with the current thought that miRNAs should primarily target POU5F1 [10], a key factor for

reprogramming induced pluripotent stem cells.

Pluripotent network targeted by activated miRNAs

We next expanded the pluripotent gene list to all pluripotent genes uncovered by Hu et al [24]. We still focused on the direct miRNA target and searched the first neighbor of all overexpressed miRNAs (Figure S4A) and all down-regulated miRNAs (Figure S4B). Most of these targets are shared by up- and down-regulated miRNAs (Figure 5A-5B), indicating that the primary pluripotent genes in stem cells are carefully modulated by multiple up- and down-regulated miRNAs. The function of the entire shared network is primarily for extrinsic signal pathways associated with pluripotency (Figure 5B). For example, the highly connected nodes APC, RAD21 and EIF4G2 are involved in Wnt signaling and mitotic cell cycle pathways. Signaling pathways with similar functions were also found in the network targeted by over-expressed miRNAs only (Figure 5C) and in modules directly targeted by the represented miRNAs cluster in stem cells, such as miR-302/367 cluster (Figure 5D) and miR-294 (Figure 5E). This indicated that miRNAs in stem cells primarily function for modulating the balance of pluripotent signal pathways instead of directly targeting pluripotent core factors. This suggests that these regulations driven by miRNAs might go through multiple steps to the pluripotent core factors.

In contrast to the up-regulated miRNAs in stem cells, down-regulated miRNAs directly target the core pluripotency factors (Figure S5), suggesting that these miRNAs inhibit the core factors for pluripotency to facilitate differentiation when these down-regulated

miRNAs become up-regulated during differentiation. To summarize, miRNAs do not directly target pluripotent core factors during pluripotent state but miRNAs directly target and repress these core factors during differentiation.

MiRNAs abundantly target epigenetic system

The above results indicated that the number of miRNA binding (degree) to their targets (nodes) was very limited. The target with the highest degree, EIF4G2, was only attacked by ~20 up-regulated miRNAs (Figure 5C). It was expected that a certain group of nodes should be targeted by more than that. This drove us to further search the network hubs (the important nodes) in the entire network. We systematically ranked the miRNA targets by degree (miRNA directly binding only) and obtained the top hubs. The top hubs mostly function for RNA processing, but surprisingly, DNMT3A, a DNA-methyltransferase for de novo DNA methylation, was among the top hubs. DNMT3A actually holds more than 160 miRNA binding sites in 3'-UTR region based only on 8bp seed mapping and it was ranked within the top 1% of the up-regulated miRNA targets (Figure 6A). DNMT3A was even ranked higher than EIF4G2, the highest ranked node in the pluripotent genes (Figure 6A, Figure 5B-5C), indicating that DNMT3A should be a top important node in the network directly regulated by miRNAs in stem cells. This also indicated that miRNAs predominately target DNA methylation system, rather than the pluripotent genes. We extracted the network of DNMT3A directly targeted by miRNAs and found that the top miRNA clusters (Table S4), such as miR-302/367 and miR-290-295 cluster, were among the miRNAs that target DNMT3A (Figure 6B). Similarly, Many

well-known miRNA clusters (e.g. miR-290-295 and miR-302) in stem cells also target DNMT1 (Figure 6C), an enzyme predominately responsible for methylation in hemimethylated CpG islands. Many down-regulated miRNAs also target DNMT3A and DNMT1 (Figure S6), but their attacks would lead to differentiation instead of maintaining pluripotency in stem cells. These abundant overexpressed miRNAs that target the methylation system suggest that miRNAs predominately repress DNMTs in stem cells.

In addition, miRNAs directly and abundantly target a core histone modification complex (HDAC4-MEF2C-MEF2D, <http://www.ncbi.nlm.nih.gov/gene/9759>) (Figure 7), including MEF2C (myocyte enhancer factor 2C), which was targeted by the top over-expressed miRNA clusters including miR-290-295 and miR-302 cluster (Figure 7). Up-regulating MEF2C enhances stem cells differentiation [25], and down-regulated MEF2C should inhibit differentiation. The down-regulation of MEF2C targeted by the top over-expressed miRNA clusters suggests that miRNAs repress differentiation in stem cells. This is consistent with our discussion above on the miRNA repressing development and differentiation at the pluripotent state in stem cells (Figure 2-3). Together, miRNAs directly and abundantly target the epigenetic systems at the pluripotent state.

DNA methylation mediates the miRNA activation in stem cells

To search the mechanism controlling the miRNA activations, this study turned to the genome-wide sequencing of DNA methylation in stem cells and methylation-loss-stem

cells [26] (Table S1, materials and methods). A total of 2000bp in each upstream and downstream of start sites of all activated miRNAs were examined. While the DNA methylation in the downstream of up-regulated miRNAs is not different from that of down-regulated miRNAs ($p > 0.1899$), the down-regulated miRNAs hold significantly higher methylation upstream than up-regulated miRNAs ($p < 3.685e-05$, Figure 8A). Surprisingly, the biggest difference locates in ~1000bp up-stream instead of immediate up-stream ($p < 1.265e-06$, Figure 8B). Furthermore, these differences are overall negatively correlated to miRNA expressions with correlation coefficient of -0.35 and p -value < 0.05 (Figure 8C). This suggested that the difference in DNA methylation accounts for the miRNA activations. This parallels a recent observation showing that the loss of DNA methylation significantly increases miRNA expressions [29]. Therefore, miRNA activations and their network are mediated by DNA methylation in ~1000bp upstream regions.

Discussion

This study is the first, to our knowledge, to investigate the primary mechanism of miRNA functions in stem cells at systems level on the basis of a physical map constructed by direct interactions of miRNAs and proteins. All data employed in this study were collected from published biological experiments, and the core part of this study, the miRNA-target physical binding map, was inferred from the CLIP-seq (Figure 1). This gives our study several advantages. First, the interaction map should be more accurate than that predicted by pure computations, a motif prediction [27], which only produced

20% overlapped with the experimental data (data not shown). Secondly, the nature of map inferred here provides the direct physical binding linkages between miRNAs and their targets. This network(....) makes it possible for us to understand a precise mechanistic picture of miRNA interactions with their targets. In contrast, traditional functional studies like gene knockdown/knockout only provide the linking hint from a gene to phenotypes with unknown mechanisms. Finally, the map provides a path to appreciate the molecular mechanics at systems level, like the pattern and network module recognitions that are based on many components (genes in this case). Mistakes could be made during observations of individual components (genes) because these individual observations could vary with conditions, but the nature of modules and patterns are normally robust enough to buffer the individual variations and noises, and they would not be changed with conditions [28]. Therefore, the results and conclusions on the global roles of miRNAs on the basis of pattern and module recognition here should represent the real nature of miRNA functions in stem cells.

Current studies have demonstrated that miRNAs play critical roles in maintaining overall properties like pluripotency in stem cells [2-4]. However, the mechanisms still remain elusive. In this study, we systematically revealed that one of miRNA primary functions is to repress developmental modules during the pluripotent state while miRNAs directly target pluripotent core factors during differentiation state (Figure 2, Figure 3, and Figure S3). This suggests that miRNAs primarily repress development at pluripotent state to prevent stem cell differentiation and to keep stem cell pluripotency while another set of

miRNAs degrade pluripotent core factors to facilitate differentiation during differentiation state. This is consistent with the recent observation that overexpressions of miRNAs induce pluripotency [10] and miRNAs also facilitate stem cells differentiation [11].

The linkage between miRNAs and pluripotency has been widely investigated [7-9], but whether the linkage is direct or indirect still remains to be investigated. It has been consistently observed that gene expressions of the top over-expressed miRNAs are positive correlative to that of pluripotent core factors. Recent evidences also show that miRNA-302/367 cluster could induce pluripotency [10]. The observations lead to a speculation that miRNAs might directly target the pluripotent core factors. Although miRNAs could have many functions in certain conditions, the primary functions of miRNAs are for degrading and inhibiting their targets. If the overexpressed miRNAs directly target the core factors, these miRNAs would likely repress the core factors as previously evidenced [29], leading to down-regulations of these core factors. A negative correlation between them should show up, but the fact is that positive correlation has been consistently observed. This suggests that the top miRNAs might not directly target and degrade pluripotent core factors. Here, we utilized the power of our system network to exhaustively search the direct linkages between miRNAs and the pluripotent core factors. Our results revealed that the top miRNAs (Table S4) such as miR-290 and miRNA-302 cluster do not directly target any core pluripotent factors during the pluripotent state (Figure 4). Most of top miRNAs only directly target the extrinsic signal pathways associated with pluripotency (Figure 5). A signal from a signal pathway

normally travels through multiple steps to reach its targets. Therefore, the top miRNAs indirectly target the core factors and indirectly regulate the pluripotency.

In contrast to the indirect mechanism of miRNAs in targeting the genetic system, the top miRNAs, including miRNA-302/367 and miRNA-294 cluster, directly and abundantly target the core enzymes of DNA methylation system, DNMT3A and DNMT1 (Figure 6). Targeting these DNMTs by the top expressed miRNAs suggests inhibition of DNMTs, which highly express in somatic cells but lowly express in stem cells (Figure 6). This parallels the most recent observations showing that gene expressions of up-regulated miRNAs are negatively correlative with that of DNA methyltransferases (DNMTs) [30-31] and that microRNAs degrade DNMTs in stem cells [30].

Our result of miRNA directly repressing DNMTs can help to understand the positive relationship existing between up-regulated miRNAs and overexpressed core pluripotent factors in stem cells as discussed above. Over-expressed miRNAs directly target DNMTs, leading to DNA methylation reductions at genome-wide level, including the loci of the pluripotent core factors. This results in over-expression of these core factors responsible for pluripotency. This is consistent with the observation of less methylation in the loci of the core factors during induced pluripotent stem cell reprogramming, and it is also consistent with our recent finding that the demethylation level can be used as a variable for discriminating different stem cells [32]. Therefore, miRNAs primarily and directly target the epigenetic system that further activates pluripotent core factors in

stem cells. This parallels the most current report that miRNAs degrade DNMTs in stem cells [30].

DNA methylation might regulate expressions of a certain group of genes in stem cells [28]. Our data further showed that DNA methylation globally mediates the miRNA activations in stem cells (Figure 8). These miRNA activations by DNA methylation in turn repress the DNA methylation (Figure 6). Less methylation activates the miRNAs and pluripotent core factors again.

Together, we proposed a system-wide circuit to describe a part of miRNA primary roles in modulating pluripotency in pluripotent stem cells (Figure 9). In this circuit, miRNAs directly repress development and directly repress the DNA methylation system, while miRNAs indirectly regulate pluripotency genes. This repression of DNA methylation activates both pluripotent factors and miRNAs. The activations of pluripotent factors and repression of development contribute to pluripotency in stem cells, while the activations of miRNAs further inhibit both DNA methylation and development. This create an active system-wide circuit in stem cells to maintain the pluripotent state (Figure 9).

We here pay more attentions on the primary functions of miRNAs in pluripotent state than that in differentiation state and we do not exclude other functions of miRNAs in stem cells. The results made here are based on the current limited data and these miRNA functions drawn here only account for as a part of miRNA roles in stem cells.

With data accumulating, more functions of miRNAs will certainly be explored. However, understanding the fundamental systems roles of miRNAs studied here would broadly direct the future functional studies of miRNAs in stem cells and would guide the successful development of stem-cell–based therapies for regenerative medicine.

Materials and methods

Data resources

This study analyzed data generated by high-throughput methods, including CLIP-seq, ChIP-seq, RNA-seq, microarray, and bisulfite sequencing DNA (Table S1). The data was downloaded from GEO database (www.ncbi.nlm.nih.gov/geo/) and the details are shown in table S1.

Data bioinformatics analysis

To be consistent and comparable, all sequencing data were mapped to mouse genome (mm9). All fastq files (Table S1) from CLIP-seq, ChIP-seq and RNA-seq were aligned using BWA 0.6.2 with default parameters [33]. All PCR-duplicates were removed. Bisulfite sequencing DNA was aligned by Bismark 0.7.6 with tolerating one non-bisulfite mismatch per read (<http://www.bioinformatics.babraham.ac.uk/projects/bismark/>).

The clusters from CLIP-seq were generated by using GenomicRanges 1.12 under R 2.15 and were subjected to second noise quartile cutoff. The filtered clusters located in 3'UTRs and CDSs were used to search for miRNA bindings. The miRNA bindings were

searched against the perfect match of miRNA seed sequences (6-8nt) using home-made python scripts. The seed sequences were extracted from miRNA sequences downloaded from miRBase 19 (<http://microrna.sanger.ac.uk/>). Peaks from ChIP-seq were called using SISSRs 1.4 (<http://dir.nhlbi.nih.gov/papers/lmi/epigenomes/sissrs/>) and the peaks were annotated using ChIPpeakAnno 2.8 under R-2.15. The differential expressions from RNA-seq were performed using negative binomial model using edgeR 3.2 under R 2.15. Bisulfite DNA methylation was counted against the genome coordinates of miRNAs (Table S4). Microarray data were analyzed using limma 3.16 under R 2.15.

Activated genes and miRNAs selection

To minimize the biases from individual experiments and cell lines as well as the noises caused by high-throughput methods, we selected genes and miRNAs activated in different conditions by using different sets of data (Table S1). Expression data from various resources (Table S1) were employed. Genes coding for proteins with activating frequency >50% in all observations and miRNAs with activating frequency of more than 25% out of all observations were treated as activated genes and were selected (Table S2-S3). The up-regulation and down-regulation was based on comparison to somatic cells in each experiment (corrected $p < 0.01$ and fold change > 2).

The top important miRNAs (Table S4) were selected on the basis of their contributions

to network structure and variance by using the algorithm as we previously published [28]. Briefly, the top miRNAs were selected on the basis of their ranking scores calculated by the eigengene-based connectivity as defined below [28].

$$SCORE = \frac{d_i}{d_{max}} + 2 \times cor[X_i, E]$$

where d_i denotes the i^{th} node degree, and d_{max} denotes the maximum degree of a node in the entire network. $|Cor(x_i, E)|$ represents the absolute Pearson correlation coefficient, where x_i represents a vector of i^{th} node value, and E eigengene of the network.

Network construction and analysis

The network construction and analysis were performed by approaches as our previous reports [17, 34]. Briefly, we built the map (Figure 1,) by collecting the interactions of both miRNAs targets from the CLIP-seq and protein-binding promoters of miRNAs from ChIP-seq data. The interactions from CLIP-seq and ChIP-seq were signed as miRNA or proteins respectively (Figure 1). Only the direct interactions (first-neighbor) were selected and included, and thus this map is a physical binding network. The network was enriched by the activated genes and miRNAs selected above to get the activated network. Functional modules were further enriched by the functional genes based on gene ontology enrichment (e.g. Figure S1, <http://www.geneontology.org/>). Six sub-modules in functional developmental module (Figure 2A) were based on network topology to identify the densely connected modules. The target node ranking was based

on the degree of each miRNA node (Figure 6A-6B).

Acknowledgments

We specially thank Drs. Ying Du and Chunxiao Zhou for data collection.

Author contributions

A.W, conceived and designed the experiments, analyzed the data, contributed data/materials/analysis tools, wrote the manuscript. Y.Z., Q.H. contributed data/materials/analysis tools. All authors reviewed the manuscript.

Competing interests

No competing interests exist in this study

No financial conflict of interest in this study.

References

1. Kloosterman WP, Plasterk RH: **The diverse functions of microRNAs in animal development and disease.** *Dev Cell* 2006, **11**(4):441-450.
2. Kanellopoulou C, Muljo SA, Kung AL, Ganesan S, Drapkin R, Jenuwein T, Livingston DM, Rajewsky K: **Dicer-deficient mouse embryonic stem cells are defective in differentiation and centromeric silencing.** *Genes Dev* 2005, **19**(4):489-501.
3. Murchison EP, Partridge JF, Tam OH, Cheloufi S, Hannon GJ: **Characterization of Dicer-deficient murine embryonic stem cells.** *Proc Natl Acad Sci U S A* 2005,

- 102(34):12135-12140.**
4. Wang Y, Medvid R, Melton C, Jaenisch R, Blelloch R: **DGCR8 is essential for microRNA biogenesis and silencing of embryonic stem cell self-renewal.** *Nat Genet* 2007, **39(3):380-385.**
 5. Chen CZ, Li L, Lodish HF, Bartel DP: **MicroRNAs modulate hematopoietic lineage differentiation.** *Science* 2004, **303(5654):83-86.**
 6. Cordes KR, Sheehy NT, White MP, Berry EC, Morton SU, Muth AN, Lee TH, Miano JM, Ivey KN, Srivastava D: **miR-145 and miR-143 regulate smooth muscle cell fate and plasticity.** *Nature* 2009, **460(7256):705-710.**
 7. Hanina SA, Mifsud W, Down TA, Hayashi K, O'Carroll D, Lao K, Miska EA, Surani MA: **Genome-wide identification of targets and function of individual MicroRNAs in mouse embryonic stem cells.** *PLoS Genet* 2010, **6(10):e1001163.**
 8. Benetti R, Gonzalo S, Jaco I, Munoz P, Gonzalez S, Schoeftner S, Murchison E, Andl T, Chen T, Klatt P *et al*: **A mammalian microRNA cluster controls DNA methylation and telomere recombination via Rbl2-dependent regulation of DNA methyltransferases.** *Nat Struct Mol Biol* 2008, **15(3):268-279.**
 9. Sinkkonen L, Hugenschmidt T, Berninger P, Gaidatzis D, Mohn F, Artus-Revel CG, Zavolan M, Svoboda P, Filipowicz W: **MicroRNAs control de novo DNA methylation through regulation of transcriptional repressors in mouse embryonic stem cells.** *Nat Struct Mol Biol* 2008, **15(3):259-267.**
 10. Anokye-Danso F, Trivedi CM, Juhr D, Gupta M, Cui Z, Tian Y, Zhang Y, Yang W, Gruber PJ, Epstein JA *et al*: **Highly efficient miRNA-mediated reprogramming of mouse and human somatic cells to pluripotency.** *Cell Stem Cell* 2011, **8(4):376-388.**
 11. Melton C, Judson RL, Blelloch R: **Opposing microRNA families regulate self-renewal in mouse embryonic stem cells.** *Nature* 2010, **463(7281):621-626.**
 12. Stefani G, Slack FJ: **Small non-coding RNAs in animal development.** *Nat Rev Mol Cell Biol* 2008, **9(3):219-230.**
 13. Bartel DP: **MicroRNAs: target recognition and regulatory functions.** *Cell* 2009, **136(2):215-233.**
 14. Chen C, Ridzon D, Lee CT, Blake J, Sun Y, Strauss WM: **Defining embryonic stem cell identity using differentiation-related microRNAs and their potential targets.** *Mamm Genome* 2007, **18(5):316-327.**
 15. Rajewsky N: **microRNA target predictions in animals.** *Nat Genet* 2006, **38 Suppl:S8-13.**
 16. Xu N, Papagiannakopoulos T, Pan G, Thomson JA, Kosik KS: **MicroRNA-145 regulates OCT4, SOX2, and KLF4 and represses pluripotency in human embryonic stem cells.** *Cell* 2009, **137(4):647-658.**
 17. Wang A, Johnston SC, Chou J, Dean D: **A systemic network for Chlamydia pneumoniae entry into human cells.** *J Bacteriol* 2010, **192(11):2809-2815.**
 18. Leung AK, Young AG, Bhutkar A, Zheng GX, Bosson AD, Nielsen CB, Sharp PA: **Genome-wide identification of Ago2 binding sites from mouse embryonic stem cells with and without mature microRNAs.** *Nat Struct Mol Biol* 2011, **18(2):237-244.**
 19. Chi SW, Zang JB, Mele A, Darnell RB: **Argonaute HITS-CLIP decodes microRNA-mRNA interaction maps.** *Nature* 2009, **460(7254):479-486.**
 20. Marson A, Levine SS, Cole MF, Frampton GM, Brambrink T, Johnstone S, Guenther MG, Johnston WK, Wernig M, Newman J *et al*: **Connecting microRNA genes to the**

- core transcriptional regulatory circuitry of embryonic stem cells. *Cell* 2008, **134**(3):521-533.
21. Maere S, Heymans K, Kuiper M: **BiNGO: a Cytoscape plugin to assess overrepresentation of gene ontology categories in biological networks.** *Bioinformatics* 2005, **21**(16):3448-3449.
 22. Bader GD, Hogue CW: **An automated method for finding molecular complexes in large protein interaction networks.** *BMC Bioinformatics* 2003, **4**:2.
 23. Miyoshi N, Ishii H, Nagano H, Haraguchi N, Dewi DL, Kano Y, Nishikawa S, Tanemura M, Mimori K, Tanaka F *et al*: **Reprogramming of mouse and human cells to pluripotency using mature microRNAs.** *Cell Stem Cell* 2011, **8**(6):633-638.
 24. Hu G, Kim J, Xu Q, Leng Y, Orkin SH, Elledge SJ: **A genome-wide RNAi screen identifies a new transcriptional module required for self-renewal.** *Genes Dev* 2009, **23**(7):837-848.
 25. Cho EG, Zaremba JD, McKercher SR, Talantova M, Tu S, Masliah E, Chan SF, Nakanishi N, Terskikh A, Lipton SA: **MEF2C enhances dopaminergic neuron differentiation of human embryonic stem cells in a parkinsonian rat model.** *PLoS One* 2011, **6**(8):e24027.
 26. Meissner A, Mikkelsen TS, Gu H, Wernig M, Hanna J, Sivachenko A, Zhang X, Bernstein BE, Nusbaum C, Jaffe DB *et al*: **Genome-scale DNA methylation maps of pluripotent and differentiated cells.** *Nature* 2008, **454**(7205):766-770.
 27. Lewis BP, Burge CB, Bartel DP: **Conserved seed pairing, often flanked by adenosines, indicates that thousands of human genes are microRNA targets.** *Cell* 2005, **120**(1):15-20.
 28. Wang A, Huang K, Shen Y, Xue Z, Cai C, Horvath S, Fan G: **Functional modules distinguish human induced pluripotent stem cells from embryonic stem cells.** *Stem Cells Dev* 2011, **20**(11):1937-1950.
 29. Tay Y, Zhang J, Thomson AM, Lim B, Rigoutsos I: **MicroRNAs to Nanog, Oct4 and Sox2 coding regions modulate embryonic stem cell differentiation.** *Nature* 2008, **455**(7216):1124-1128.
 30. Guo X, Liu Q, Wang G, Zhu S, Gao L, Hong W, Chen Y, Wu M, Liu H, Jiang C *et al*: **microRNA-29b is a novel mediator of Sox2 function in the regulation of somatic cell reprogramming.** *Cell Res* 2013, **23**(1):142-156.
 31. Garzon R, Liu S, Fabbri M, Liu Z, Heaphy CE, Callegari E, Schwind S, Pang J, Yu J, Muthusamy N *et al*: **MicroRNA-29b induces global DNA hypomethylation and tumor suppressor gene reexpression in acute myeloid leukemia by targeting directly DNMT3A and 3B and indirectly DNMT1.** *Blood* 2009, **113**(25):6411-6418.
 32. Wang A, Du Y, He Q, Zhou C: **A quantitative system for discriminating induced pluripotent stem cells, embryonic stem cells and somatic cells.** *PLoS One* 2013, **8**(2):e56095.
 33. Li H, Durbin R: **Fast and accurate long-read alignment with Burrows-Wheeler transform.** *Bioinformatics* 2010, **26**(5):589-595.
 34. Wang A, Al-Kuhlani M, Johnston SC, Ojcius DM, Chou J, Dean D: **Transcription Factor Complex AP-1 Mediates Inflammation Initiated by Chlamydia pneumoniae Infection.** *Cell Microbiol* 2012.

Figure legends

Figure 1. Overall view. A, The concept of miRNA and protein interactions. B, Workflow of this study. C, Overall view of the entire network constructed by this study. D, A sample of entire network contents shows direct interactions between miRNAs and pluripotent core factors (NANOG, POU1F5, and SOX2). Nodes denote miRNAs or proteins as labeled; red node represents the gene up-regulation in pluripotent stem cells, green node as down-regulation, and white node as insignificant expression. Edges represent interactions; red edge represents miRNA targeting proteins and green edge represents the binding of proteins with the regulatory elements of miRNAs. This labeling strategy applies to all figures in this study.

Figure 2. MiRNAs primarily repress developmental processes in pluripotent stem cells. A, Up-regulated miRNAs primarily target a developmental module, which includes 6 primary sub-modules functioning for development as highlighted in cycles. B, The key modules targeted by the top 5 important miRNAs (Table S4). C, the key modules were significantly and abundantly enriched in the developmental category.

Figure 3. Modules targeted by miRNA-302/367 cluster and miRNA-294. A, modules targeted by the miRNA-302/367 cluster and their functional enrichment was shown in B. C, a module targeted by miRNA-294, and D, its functional enrichment.

Figure 4. Pathways from activated miRNAs to pluripotent core factors. A, B, C, D,

the shortest paths from miR-302b, miR-367, miR-294, and miR-292-5p respectively to the pluripotent core factors. None of these top overexpressed miRNAs directly binds to any of these core factors. E, Core factors directly targeted by activated miRNAs in stem cells.

Figure 5. Pluripotent genes targeted by over-expressed miRNAs in stem cells. A, Venn diagram of pluripotent genes targeted by both up- and down-regulated miRNAs. B, activated modules commonly targeted by both up- and down-regulated miRNAs. The enlarged nodes represent the highly connected targets that are targeted by both up- and down-regulated miRNAs in stem cells, and they work for pluripotent signal pathways. C, down-regulated modules targeted by over-expressed miRNAs in stem cells. D, activated modules targeted by the miRNA-302/367 cluster in stem cells. E, an activated network targeted by miR-294.

Figure 6. MiRNAs abundantly target DNA methylation systems. A, DNMT3A was targeted by 30 up-regulated miRNAs and was ranked in the top 1% of the up-regulated miRNA targets. The network nodes (miRNA targets) were plotted against the node degree (miRNA binding number only). B, the DNMT3A network targeted by miRNAs. The most important miRNAs, including the miRNA-302/367 and miR-290-295 cluster, are found in the DNMT3A network. C, miRNAs target another methylation enzyme DNMT1.

Figure 7. MiRNAs directly target a histone complex. The MiR-290-295 cluster heavily attacks the MEF2C.

Figure 8. DNA methylation mediates miRNA activations in stem cells. A, The methylation levels upstream and downstream from the miRNA start site. Down-regulated miRNAs have significantly higher methylation in upstream region when compared with up-regulated miRNAs. B, Detailed methylation profiling for regions that are 2000bp upstream from the activated miRNAs. The top 30 down-regulated miRNAs (Table S2) have higher methylation around -1000bp (green highlighted in middle panel) than that of the top 30 up-regulated miRNAs (Table S2, upper panel). The methylation profiling of a single down-regulated miRNA-133 as a representative example (bottom panel). C. Negative correlation between DNA methylation and miRNA expression. Red line represents regression line.

Figure 9. A system view of miRNAs primary mechanistic roles in maintaining pluripotency in pluripotent stem cells. MiRNAs directly repress both the DNA methylation system and the development, while indirectly regulate pluripotency genes. Reduced DNA methylation activates the miRNAs and pluripotent core factors for pluripotency. The activated circuit between miRNAs and DNA methylation, as well as the development inhibition, help stem cells to maintain the pluripotent state, see text for detail. The solid lines are directly derived from the present study and they represent direct interactions and the dash dark line represents indirect interactions, while dash green lines denote evidences from reference papers.

Systematically Dissecting the Global Mechanism of MiRNA functions in Stem Cells

Anyou Wang^{1*}, et al

Supplemental materials

Figure S1, Functional enrichments of all targets that are targeted by all up- and down-regulated miRNAs. A, Up-regulated miRNAs. B, down-regulated miRNAs.

Figure S2, total 17 out of top 20 important miRNAs target developmental genes.

Figure S3, down-regulated miRNAs mediate metabolism in stem cells.

Figure S4, all pluripotent genes targeted by up-regulated miRNAs (A) and down-regulated miRNAs (B) in stem cells.

Figure S5. An activated network targeted by down-regulated miRNAs.

Figure S6. Activated miRNAs targeting DNA methylation systems. A. Up-regulated miRNAs targeting DNMT3A. B, Down-regulated miRNAs targeting DNMT3A.

Table S1, Data sources


Table S2, a list of miRNAs that are frequently and significantly differential expressed between stem cells and somatic cells.

Table S3, a list of genes coding for proteins that are frequently and significantly differential expressed between stem cells and somatic cells.


Table S4, a list of top 20 important miRNAs

Figure 2


A




B




C




A




B



D



C









A**B****C****D****E**

Figure 6


A



B



C











Figure 8




B







A Pluripotent genes targeted by up-regulated miRNAs in stem cells



B Pluripotent genes targeted by down-regulated miRNAs in stem cells



A



B




Table S1 data resources

GEO number	Platform	measurement
GSE25310	CLIP-seq	ES miRNA targets
GSE11724	ChIP-seq	TF binding
GSE11431	ChIP-seq	TF binding
GSM278905	RRBS	ES DNA methylation
GSM278902	RRBS	TKO methylation
GSM539867	RNA-Seq	mouse embryonic fibroblast cells [09-002]
GSM539866	RNA-Seq	mouse embryonic stem cells [09-002]
GSE30012	Mouse430_2 array	ES miRNA expression
GSE25310	GPL11410	micrRNA and mRNA expression
GSM231739	Agilent-GPL2872	ES gene expression TKO/WT
GSE15267	AffymetrixGPL1261	Gene expression somatic cells vs stem cells
GSE14012	AffymetrixGPL1261	Gene expression somatic cells vs stem cells
GSE10871	AffymetrixGPL1261	Gene expression somatic cells vs stem cells
GSE14790	AffymetrixGPL1261	Gene expression somatic cells vs stem cells
GSE16062	IlluminaGPL6885	Gene expression somatic cells vs stem cells

Table S2 miRNAs differentially expressed between stem cell vs somatic cells with high frequency >25% of observations

geneID	foldChange.StemcellvsSomaticCells
mmu-miR-291a-3p	23.38772103
mmu-miR-292-5p	20.9773225
mmu-miR-295	18.61514609
mmu-miR-290	16.83638073
mmu-miR-291b-5p	16.74397919
mmu-miR-294	16.33979504
mmu-miR-293	15.71944751
mmu-miR-291a-5p	10.70302558
mmu-miR-291b-3p	8.775662583
mmu-miR-205	8.656636965
mmu-miR-292-3p	7.513040922
mmu-miR-200a	7.476828136
mmu-miR-20b	7.113430578
mmu-miR-429	6.970376253
mmu-miR-93	6.320621674
mmu-miR-302b	6.319533201
mmu-miR-124a	6.210366092
mmu-miR-141	6.112532011
mmu-miR-32	5.892597823
mmu-miR-25	5.664093286
mmu-miR-106a	5.660199796
mmu-miR-96	5.576097803
mmu-miR-690	5.566673137
mmu-miR-210	5.494180249
mmu-miR-182	5.380304319
mmu-miR-339	5.178218031
mmu-miR-33	5.150748639
mmu-miR-19b	5.130661066
mmu-miR-302d	4.786597786
mmu-miR-195	4.773097795
mmu-miR-297b	4.742119218
mmu-miR-499	4.699375408
mmu-miR-106b	4.67924418
mmu-miR-712*	4.616388282
mmu-miR-101b	4.615452785
mmu-miR-706	4.577798951
mmu-miR-672	4.425689091
mmu-miR-20a	4.355849211
mmu-miR-302	4.187271105
mmu-miR-142-3p	4.185529766
mmu-miR-19a	4.154314537
mmu-miR-450b	4.143615059
mmu-miR-200c	4.139625599
mmu-miR-17-5p	4.015734124

mmu-miR-669b	4.005643461
mmu-miR-363	4.004376577
mmu-miR-130b	3.921190227
mmu-miR-142-5p	3.771559152
mmu-miR-708	3.757978939
mmu-miR-301b	3.671137996
mmu-miR-297	3.649760196
mmu-miR-711	3.642487186
mmu-miR-135b	3.61287415
mmu-miR-145	-3.689012325
mmu-miR-125b	-3.750522829
mmu-miR-675-3p	-3.775797104
mmu-miR-31	-3.789955248
mmu-miR-431	-3.827836369
mmu-miR-133a	-3.931634792
mmu-miR-434-5p	-3.95970537
mmu-miR-29a	-4.128390595
mmu-miR-125a	-4.168574915
mmu-miR-27b	-4.181204463
mmu-miR-206	-4.1991836
mmu-miR-222	-4.206742949
mmu-miR-329	-4.226853284
mmu-let-7i	-4.253606445
mmu-miR-221	-4.277690509
mmu-miR-132	-4.327457901
mmu-miR-345	-4.362726416
mmu-miR-452	-4.493999108
mmu-let-7d	-4.733310501
mmu-miR-181a*	-4.890038681
mmu-miR-212	-5.03014549
mmu-miR-152	-5.091913825
mmu-miR-143	-5.251965264
mmu-miR-433-5p	-5.353386842
mmu-let-7e	-5.627257306
mmu-miR-322	-5.627467759
mmu-miR-380-5p	-5.840198038
mmu-miR-666	-5.963478682
mmu-miR-365	-6.053925857
mmu-miR-10a	-6.463178387
mmu-miR-24*	-6.501407687
mmu-miR-99b	-6.54057846
mmu-miR-98	-6.824098585
mmu-miR-28	-6.985496746
mmu-miR-23a	-7.258185943
mmu-miR-675-5p	-7.319121705
mmu-miR-224	-7.383165325
mmu-miR-133b	-8.331863729

mmu-miR-155

-12.36089174

Table S3. Genes differentially expressed between stem cells
and somatic cells with high frequency >50% of observations

GB_ACC	ENTREZ_GI	Gene.Symbol	logFoldChange.ESvsSC
BC006640	20315	Cxcl12	-7.932773867
BF225802	16011	Igfbp5	-7.7652649
AW550625	12825	Col3a1	-7.7405454
BF227507	12843	Col1a2	-7.50996995
BB542051	18295	Ogn	-7.13772715
U08020	12842	Col1a1	-7.025342217
NM_013655	20315	Cxcl12	-6.8464668
NM_011581	21826	Thbs2	-6.743730283
AV229424	12832	Col5a2	-6.6650371
BB051738	18933	Prrx1	-6.523803517
M65143	16948	Lox	-6.263793183
AF007248	14118	Fbn1	-6.234405867
NM_009933	12833	Col6a1	-6.142877467
NM_007993	14118	Fbn1	-6.087729233
BC019502	12111	Bgn	-6.042548083
AB015978	18414	Osmr	-6.018220117
BE197945	20324	Sdpr	-5.902171217
NM_007729	12814	Col11a1	-5.888984717
AW049660	18032	Nfix	-5.878165733
NM_010514	16002	Igf2	-5.805814817
BB197591	54216	Pcdh7	-5.793524283
AV226618	13837	Epha3	-5.703367667
AF378762	69538	Antxr1	-5.671588117
M68513	13837	Epha3	-5.646988233
BC002064	19242	Ptn	-5.631718783
NM_011340	20317	Serpinf1	-5.561155483
BB795075	18214	Ddr2	-5.5521091
BC023060	216616	Efemp1	-5.541377517
BF144658	21808	Tgfb2	-5.516599417
BB250384	22329	Vcam1	-5.511377133
AB029929	12389	Cav1	-5.478985417
D67076	11504	Adamts1	-5.469247617
BB315728	18032	Nfix	-5.460392167
BI794771	12842	Col1a1	-5.456734267
L06502	18933	Prrx1	-5.448142767
BG793483	21813	Tgfbr2	-5.434682533
BB532202	14264	Fmod	-5.394454433
BB361162	22771	Zic1	-5.383117233
AI931862	12111	Bgn	-5.378983617
BC014690	21809	Tgfb3	-5.302384367
NM_018865	22402	Wisp1	-5.29682725
AV064339	20324	Sdpr	-5.2953667
BG067986	54216	Pcdh7	-5.295023283
NM_030888	81799	C1qtnf3	-5.279631533

BB233297	21826 Thbs2	-5.26170495
BF168458	21858 Timp2	-5.239630733
BB259670	14362 Fzd1	-5.23704885
AA499047	18596 Pdgfrb	-5.226616983
BC026446	219151 Scara3	-5.1983228
NM_010714	16876 Lhx9	-5.18046395
NM_133859	99543 Olfml3	-5.167943083
BB067079	22418 Wnt5a	-5.160633383
AW555664	399558 Flrt2	-5.14692565
BC021484	16651 Sspn	-5.141520367
BC011063	15402 Hoxa5	-5.109899467
M93954	21858 Timp2	-5.0922978
AV310588	12444 Ccnd2	-5.076240083
NM_013589	16997 Ltbp2	-5.07117345
BB791906	14178 Fgf7	-5.0607749
BQ175880	12444 Ccnd2	-5.031714083
NM_009829	12444 Ccnd2	-5.03108855
BC022107	12558 Cdh2	-5.030543317
NM_011160	19091 Prkg1	-5.010924967
AK007904	12444 Ccnd2	-5.010597567
BB114398	12816 Col12a1	-5.001727783
BB377340	13612 Edil3	-4.984431667
BI455189	12834 Col6a2	-4.981937467
BC018425	22418 Wnt5a	-4.949481167
AF309564	80837 Rhoj	-4.942552217
AW744319	12831 Col5a1	-4.912884333
NM_010284	14600 Ghr	-4.907539267
AK003674	68588 Cthrc1	-4.9040982
NM_009610	11468 Actg2	-4.88703015
BM220945	239217 Kctd12	-4.885967567
BB658835	23794 Adamts5	-4.884421383
BC019836	16010 Igfbp4	-4.86110935
NM_018764	54216 Pcdh7	-4.826503217
AA500897	26362 Axl	-4.824049367
AV222756	259300 Ehd2	-4.811690783
NM_018884	55983 Pdzrn3	-4.770790783
BB759833	17300 Foxc1	-4.766068317
BQ177170	17112 Tm4sf1	-4.749287983
BM250666	12830 Col4a5	-4.738634083
BC025502	231532 Arhgap24	-4.7154329
NM_007392	11475 Acta2	-4.71167825
NM_008885	18858 Pmp22	-4.71162355
AV246911	12831 Col5a1	-4.70608125
NM_133918	100952 Emilin1	-4.685038
AV260198	83675 Bicc1	-4.676352617
BC021376	225207 Zfp521	-4.6747989
NM_011607	21923 Tnc	-4.6736706

NM_007833	13179 Dcn	-4.668888083
AF357006	16949 Loxl1	-4.661639267
BC011507	74480 Samd4	-4.6606821
BB493031	23888 Gpc6	-4.659793267
BI249259	18032 Nfix	-4.659746767
NM_054044	78560 Gpr124	-4.652361
BC001999	13732 Emp3	-4.648468717
BB041180	16876 Lhx9	-4.631340583
BI690209	57342 Parva	-4.603038383
AK018128	70717 6330406l15Ri	-4.599817333
AI595932	17260 Mef2c	-4.571089667
AI463873	11819 Nr2f2	-4.558937633
BG075699	399558 Flrt2	-4.550595967
AV149705	108927 Lhfp	-4.537199083
AK014221	73230 Bmper	-4.536276183
BB468025	54381 Pgcg	-4.535907333
BB363812	14051 Eya4	-4.52926855
AK020118	23888 Gpc6	-4.5247486
BM217996	83675 Bicc1	-4.521218667
NM_009636	11568 Aebp1	-4.509546617
AA987181	15405 Hoxa9	-4.496607467
NM_011766	22762 Zfpm2	-4.496262983
AV315205	11658 Alcam	-4.495887367
U95030	11658 Alcam	-4.4742135
BF147716	17390 Mmp2	-4.463715417
NM_008608	17387 Mmp14	-4.4584154
AV332957	20429 Shox2	-4.44107
AK008112	18933 Prrx1	-4.429082483
BB787243	16010 Igfbp4	-4.408829283
M27130	12505 Cd44	-4.408205433
NM_023118	13132 Dab2	-4.401669267
BB817332	399558 Flrt2	-4.401003567
D63423	11747 Anxa5	-4.3868292
BC014870	235587 Parp3	-4.368841483
U03425	13649 Egfr	-4.353990467
NM_011150	19039 Lgals3bp	-4.34716645
BB229377	13717 Eln	-4.3469066
BF681826	26360 Angptl2	-4.34243245
AV369812	13649 Egfr	-4.340586717
AI385532	21825 Thbs1	-4.3388455
AK012411	20317 Serpinf1	-4.334218667
NM_016900	12390 Cav2	-4.327267733
NM_007471	11820 App	-4.32309575
BC026153	13841 Epha7	-4.322886683
BE652876	60527 Fads3	-4.31762715
NM_022563	18214 Ddr2	-4.317137833
BC005490	11820 App	-4.311839833

AY083458	235505 Cd109	-4.310650267
AW146109	12505 Cd44	-4.309742383
AF022889	268977 Ltbp1	-4.293317933
BF780807	23972 Papss2	-4.2837158
NM_010222	14231 Fkbp7	-4.283483917
BB765827	74761 Mxra8	-4.28266945
BB369191	27528 DOH4S114	-4.258134683
NM_011658	22160 Twist1	-4.252172233
BC022679	52552 Parp8	-4.249244683
BC024375	14600 Ghr	-4.235411317
AW558570	13617 Ednra	-4.235029783
NM_010151	13865 Nr2f1	-4.23266835
BB053506	207181 Rbms3	-4.227119367
BB464523	12293 Cacna2d1	-4.2176231
AI325255	14230 Fkbp10	-4.2139772
NM_009365	21804 Tgfb1i1	-4.2037574
AW537708	18595 Pdgfra	-4.201673433
BG244279	26360 Angptl2	-4.19122825
AV021105	208647 Creb3l2	-4.189322833
NM_021474	58859 Efemp2	-4.172999333
AW558468	230103 Npr2	-4.170204383
BM230959	229731 Slc25a24	-4.15558595
NM_013519	14234 Foxc2	-4.150217817
BG248060	12153 Bmp1	-4.1456309
AK011935	67155 Smarca2	-4.1424534
BC009660	109042 Prkcdbp	-4.138494033
NM_010730	16952 Anxa1	-4.137948167
NM_017464	18003 Nedd9	-4.125438817
NM_020606	57342 Parva	-4.117250083
NM_054042	70445 Cd248	-4.1162995
AW107196	74480 Samd4	-4.089650683
NM_008520	16998 Ltbp3	-4.088398567
NM_007400	11489 Adam12	-4.087465183
BB138485	70676 Gulp1	-4.06900915
BB041237	17389 Mmp16	-4.0669223
NM_009152	20346 Sema3a	-4.06663545
AK007400	77889 Lbh	-4.065124233
BC027199	233328 Lrrk1	-4.0638856
BC003726	14789 Leprel2	-4.061919033
BM220576	53623 Gria3	-4.059143167
BM570006	20564 Slit3	-4.050010933
NM_007472	11826 Aqp1	-4.049342233
BB329489	12406 Serpinh1	-4.035949167
BC024358	22004 Tpm2	-4.027799067
BB371406	57265 Fzd2	-4.024201683
BI106777	12293 Cacna2d1	-4.016322467
BB484759	224024 Scarf2	-3.995337517

BC025145	19266 Ptprd	-3.96890555
AK004519	56726 Sh3bgrl	-3.915095233
NM_010221	14230 Fkbp10	-3.911402667
BB114067	21345 Tagln	-3.906768233
NM_010517	16010 Igfbp4	-3.895716167
AV164956	22418 Wnt5a	-3.889125017
U71189	13592 Ebf2	-3.883624733
AF339910	18605 Enpp1	-3.88293505
BQ173967	13640 Efna5	-3.882553467
AV246882	235505 Cd109	-3.87793775
AK003186	22004 Tpm2	-3.875057833
NM_009821	12394 Runx1	-3.8666507
NM_010743	17082 Il1rl1	-3.865813883
BB324823	16998 Ltbp3	-3.86059405
BC025514	102644 Oaf	-3.857056783
NM_007801	13036 Ctsh	-3.85513275
NM_010581	16423 Cd47	-3.854950117
AV238225	16905 Lmna	-3.8510058
AK018679	16423 Cd47	-3.832084267
BG073383	15410 Hoxb3	-3.820933467
NM_026405	67844 Rab32	-3.819808667
BF144687	70676 Gulp1	-3.81964405
NM_019391	16985 Lsp1	-3.81878645
NM_013586	16950 Loxl3	-3.813466317
AI415741	94352 Loxl2	-3.8120326
BI220012	12406 Serpinh1	-3.811311317
NM_007802	13038 Ctsk	-3.804638867
BB832504	18028 Nfib	-3.801083117
BC013463	15430 Hoxd10	-3.784799
M22479	22003 Tpm1	-3.783677167
BG963150	20563 Slit2	-3.783541217
BC013560	12827 Col4a2	-3.774358167
NM_008984	19274 Ptprm	-3.769132717
BB233088	21928 Tnfaip2	-3.7684148
BC008277	13617 Ednra	-3.767968717
BI111620	21859 Timp3	-3.76779495
BB475194	23794 Adamts5	-3.76137165
NM_021355	14264 Fmod	-3.736082733
NM_007585	12306 Anxa2	-3.731015333
BF578055	229731 Slc25a24	-3.725295933
BB547877	20563 Slit2	-3.716729817
NM_016753	17035 Lxn	-3.70538055
AV016275	20742 Spnb2	-3.701173833
NM_008495	16852 Lgals1	-3.6991015
AW551930	83675 Bicc1	-3.698414167
AK018466	207181 Rbms3	-3.68959625
NM_008987	19288 Ptx3	-3.687359367

AK003819	74777 Sepn1	-3.6822946
AW412729	12816 Col12a1	-3.663389833
BC020152	223272 Itgbl1	-3.657189217
Y07687	18028 Nfib	-3.657044833
NM_011985	26561 Mmp23	-3.644890267
NM_009378	21824 Thbd	-3.6371471
BF451748	107589 Mylk	-3.626035767
BM251152	13003 Vcan	-3.620597833
BB468082	18481 Pak3	-3.6195631
NM_007899	13601 Ecm1	-3.6077534
BB522674	20541 Slc8a1	-3.606644633
NM_019989	56726 Sh3bgrl	-3.60465905
BC023448	23888 Gpc6	-3.603901967
AK002516	11857 Arhgdib	-3.6024453
BF100813	13618 Ednrb	-3.594391017
NM_011777	22793 Zyx	-3.594158817
BB326929	58194 Sh3kbp1	-3.582396833
BB248904	56726 Sh3bgrl	-3.58152235
BB027759	231997 Fkbp14	-3.5728165
AU021035	15529 Sdc2	-3.570693117
AI481026	29817 Igfbp7	-3.56210855
NM_008393	16373 Irx3	-3.560717517
BB526042	320452 P4ha3	-3.5487859
BB100920	17118 Marcks	-3.53907495
BB097480	223254 Farp1	-3.534434933
BI687652	18028 Nfib	-3.524464033
BG868949	17122 Mxd4	-3.514140767
NM_020561	57319 Smpdl3a	-3.51306225
AF059567	12579 Cdkn2b	-3.5073857
NM_009472	22253 Unc5c	-3.505214767
NM_009794	12334 Capn2	-3.498667433
NM_009627	11535 Adm	-3.48734415
BQ175796	18481 Pak3	-3.4733871
BC005569	58809 Rnase4	-3.460646467
BB327018	14118 Fbn1	-3.456580617
NM_010212	14200 Fhl2	-3.453920533
BB313689	20541 Slc8a1	-3.452686383
AK019164	17475 Mpdz	-3.437966567
AW702161	11745 Anxa3	-3.4355384
BC027084	259300 Ehd2	-3.43459445
BC005686	13713 Elk3	-3.43373535
BB065799	240725 Sulf1	-3.429931133
NM_130449	140792 Colec12	-3.424621383
BB823350	19092 Prkg2	-3.412943967
AI642438	16852 Lgals1	-3.4127125
NM_007855	13345 Twist2	-3.412527233
BB787292	14302 Frk	-3.411315533

BB354684	217410 Trib2	-3.410151567
AF117951	94352 Loxl2	-3.400751667
BC025897	225791 Zadh2	-3.398803383
NM_019877	56358 Copz2	-3.386908433
AV162270	22240 Dpysl3	-3.3799255
D63383	20465 Sim2	-3.378190883
NM_008546	17150 Mfap2	-3.375675317
NM_008008	14178 Fgf7	-3.373739283
BQ174721	214791 Sertad4	-3.37195165
BB377873	121021 Cspg4	-3.368490867
BB542535	13036 Ctsh	-3.3666448
BC024876	223453 Dap	-3.364419667
L24755	12153 Bmp1	-3.360524783
X66083	12505 Cd44	-3.354735533
AW547821	17268 Meis1	-3.3441435
BC016447	26427 Creb3l1	-3.342256433
BM213516	20742 Spnb2	-3.33549655
BM220880	108655 Foxp1	-3.326646233
AK004853	50781 Dkk3	-3.314331467
NM_008809	18596 Pdgfrb	-3.31185265
AV245241	259300 Ehd2	-3.304100233
NM_020007	56758 Mbnl1	-3.289487817
BC028307	68792 SrpX2	-3.28413355
AA709993	20365 Serf1	-3.275465433
BC006737	12370 Casp8	-3.272708067
BB151715	23871 Ets1	-3.271100167
BI452727	14314 Fstl1	-3.26508275
BB003393	21923 Tnc	-3.2549266
BB535494	18003 Nedd9	-3.2516385
BB779859	209448 Hoxc10	-3.2440814
BI248947	109624 Cald1	-3.241609283
L26349	21937 Tnfrsf1a	-3.2395265
BC012674	19285 Ptrf	-3.228721367
U30244	13642 Efnb2	-3.225253617
AK004598	71703 Armcx3	-3.218577033
BB075247	19266 Ptprd	-3.217064717
BB766878	107581 Col16a1	-3.216479983
NM_019586	56228 Ube2j1	-3.216275217
NM_008761	18301 Fxyd5	-3.21540255
M24849	22352 Vim	-3.212612833
BB366804	18230 Nxn	-3.208821517
AV031691	22771 Zic1	-3.207325183
AK003303	19266 Ptprd	-3.1900006
NM_009252	20716 Serpina3n	-3.1897822
BM233698	12977 Csf1	-3.186553767
AK018666	50766 Crim1	-3.184091083
NM_013813	13823 Epb4.1l3	-3.17559455

BB097063	16826 Ldb2	-3.167842717
BG070361	18514 Pbx1	-3.16060585
NM_023564	70310 Plscr3	-3.156913317
BC016893	15414 Hoxb6	-3.149439067
BC025602	52377 Rcn3	-3.143586083
BE628275	68910 Zfp467	-3.129177833
BB751459	240725 Sulf1	-3.118822083
NM_023813	108058 Camk2d	-3.106603767
BE630020	19122 Prnp	-3.099244017
NM_015786	50708 Hist1h1c	-3.096520833
NM_010736	17000 Ltbr	-3.096244717
NM_008380	16323 Inhba	-3.090835667
AF080090	20350 Sema3f	-3.086323
AV220340	229534 Pbxip1	-3.08320105
NM_016846	19731 Rgl1	-3.083172067
AA510713	56468 Soccs5	-3.08271985
AV246759	21808 Tgfb2	-3.078846817
AV238718	233733 Galntl4	-3.07739925
BB311061	56876 Nelf	-3.07710745
BM231794	216188 Aldh1l2	-3.071789317
NM_053147	93893 Pcdhb22	-3.06913275
AV124445	17984 Ndn	-3.067003683
AI747133	12338 Capn6	-3.061495833
NM_009784	12293 Cacna2d1	-3.0553154
NM_008675	17965 Nbl1	-3.054173417
X58380	15364 Hmga2	-3.053552917
NM_009468	22240 Dpysl3	-3.0489959
NM_019390	16905 Lmna	-3.043480833
BB533903	50708 Hist1h1c	-3.028666667
BB250811	18542 Pcolce	-3.026433
BE951265	66180 1110036O03R	-3.02614895
AW546141	17118 Marcks	-3.021924833
BB825801	13713 Elk3	-3.021571583
NM_009255	20720 Serpine2	-3.02130325
BF158638	12826 Col4a1	-3.010500317
NM_016886	53623 Gria3	-3.003217433
NM_009925	12813 Col10a1	-3.001262167
AV278675	244448 Triml1	3.01398255
BG966751	104156 Etv5	3.0156244
X67668	97165 Hmgb2	3.020431667
BB089717	12449 Ccnf	3.022975367
BB292220	58180 Hic2	3.025155533
BG069191	50927 Nasp	3.027539567
BG066754	67204 Eif2s2	3.0295477
AK019115	68239 Krt42	3.03983065
AU015121	268697 Ccnb1	3.042294933
NM_008566	17218 Mcm5	3.058338333

NM_019812	93759 Sirt1	3.064211417
NM_020599	19771 Rlbp1	3.06822615
AV280841	219132 D14Ertd668e	3.078624667
BB493242	50927 Nasp	3.081964833
NM_022409	63872 Zfp296	3.082454667
NM_013873	29859 Sult4a1	3.0839579
AF302127	72388 Ripk4	3.086288367
NM_011597	21873 Tjp2	3.087218833
BF578266	227327 B3gnt7	3.0910465
A1195532	216456 Glc2	3.091450267
BE650359	13619 Phc1	3.103443667
BM247863	56380 Arid3b	3.10414725
BM935811	16403 Itga6	3.109543333
BG071670	75796 Cdyl2	3.110318667
BC006704	226419 Dyrk3	3.114882817
NM_008628	17685 Msh2	3.118807583
BM196098	14562 Gdf3	3.129241967
NM_020567	57441 Gmnn	3.129819017
AK011289	66953 Cdca7	3.133227817
NM_009426	22044 Trh	3.133924383
BB367422	239759 Liph	3.1340973
BC011474	16818 Lck	3.140572717
NM_011176	19143 St14	3.14224885
BB667216	22371 Vwf	3.143579333
BI739053	12725 Clcn3	3.159571817
NM_011487	20849 Stat4	3.169867333
BQ177140	218038 Amph	3.17188915
NM_008452	16598 Klf2	3.1728325
AF329833	66790 Grtp1	3.203848083
BB540053	217653 C79407	3.221807417
NM_007691	12649 Chek1	3.229280333
BC016095	228421 Kif18a	3.231334583
BC017621	223775 Pim3	3.23674225
U42190	17688 Msh6	3.242437417
NM_027288	110173 Manba	3.25011815
BE951628	217653 C79407	3.269844583
BB125424	242894 Actr3b	3.270038817
AK010648	70024 Mcm10	3.288071933
AK006509	75796 Cdyl2	3.304674333
NM_010174	14077 Fabp3	3.309141333
NM_053271	116838 Rims2	3.314384433
BG070553	110957 D1Pas1	3.318172517
NM_133664	16763 Lad1	3.323882083
BQ176661	54484 Mkrn1	3.3289175
AF237702	20425 Shmt1	3.331763533
NM_016851	54139 Irf6	3.333859083
AA543734	21386 Tbx3	3.333990817

NM_010346	14786 Grb7	3.34545575
BC016571	50799 Slc25a13	3.36053825
AK007201	14536 Nr6a1	3.360795183
NM_013514	13829 Epb4.9	3.361635267
U39203	16331 Inpp5d	3.380223367
AW552407	217837 Itpk1	3.380479
AK011462	72148 2610019F03R	3.381959433
BB397151	332221 Zscan10	3.400120667
U31625	12189 Brca1	3.4060535
BM225706	243574 Kbtbd8	3.415306283
NM_031386	83560 Tex14	3.416736917
NM_009171	20425 Shmt1	3.449195983
AK011986	112422 2610305D13R	3.475925817
AV305746	231327 Ppat	3.479371517
BI465857	12224 Klf5	3.483464667
NM_021308	57746 Piwil2	3.4897425
BB758095	21873 Tjp2	3.499096167
NM_016964	50878 Stag3	3.503349717
BG066919	17765 Mtf2	3.526635767
BE986504	71801 Plekhf2	3.556072867
AV277242	243963 Zfp473	3.576478483
NM_011799	23834 Cdc6	3.57881495
NM_026190	67486 Polr3g	3.6006572
NM_010357	14860 Gsta4	3.6286395
AU016127	67374 Jam2	3.629554333
BM230524	217944 Rapgef5	3.6361713
NM_022024	63986 Gmfg	3.640558983
AK014330	107435 Hat1	3.648040833
NM_008816	18613 Pecam1	3.660811183
BC003329	54484 Mkrn1	3.66147285
AK011596	22042 Tfrc	3.664959917
BB128741	319801 9630033F20R	3.701341
AF037370	12865 Cox7a1	3.709307883
BC006646	12224 Klf5	3.738650033
AV024881	52609 Cbx7	3.776539367
BB234337	101602 AI467606	3.805827767
AK013117	76974 1190003J15Ri	3.806894467
NM_010068	13436 Dnmt3b	3.83760305
NM_009886	12614 Celsr1	3.862550233
BE648070	58180 Hic2	3.868284683
NM_007469	11812 Apoc1	3.873187833
BC018383	226180 Ina	3.88687835
AB072269	13511 Dsg2	3.8942605
NM_013512	13824 Epb4.114a	3.902842033
BB473446	239157 Pnma2	3.946305617
AK010391	71988 Esco2	3.965438167
BC020144	13511 Dsg2	3.9713275

AK009960	69697 2310057J16Ri	3.9715563
AF285580	77485 Stk31	3.971668
NM_007397	11481 Acvr2b	3.974705033
BC003778	21420 Tcfap2c	4.010026117
NM_011934	26380 Esrrb	4.057861333
NM_016907	20732 Spint1	4.058743833
BC004654	19725 Rfx2	4.076663867
AV060866	18778 Pla2g1b	4.077706683
BE133749	54484 Mkrrn1	4.082165
AF425084	97848 Serpinb6c	4.1037625
AV213552	24074 Taf7	4.123073
BQ175337	72685 Dnajc6	4.13664
AF104416	11828 Aqp3	4.179715783
BM210256	14106 Foxh1	4.211211117
NM_023844	67374 Jam2	4.2200985
AK019319	11816 Apoe	4.313206417
NM_031383	83557 Lin28	4.338394817
BG084230	66824 Pycard	4.33866605
AK010826	67374 Jam2	4.358034267
AF330212	12310 Calca	4.358796767
BQ173923	12808 Cobl	4.384203133
M18775	17762 Mapt	4.42236835
BC027285	68713 Ifitm1	4.4224295
AK021186	77485 Stk31	4.436390117
NM_009903	12740 Cldn4	4.437383
NM_009749	12069 Bex2	4.45785005
NM_009434	22113 Phlda2	4.474681333
BC017609	18424 Otx2	4.492426433
BC023498	72027 Slc39a4	4.528567183
AA215046	67080 1700019D03R	4.535403217
BC004037	22256 Ung	4.580378817
NM_007515	11989 Slc7a3	4.595658
BB469300	243574 Kbtbd8	4.605572983
NM_016754	17907 Mylpf	4.613079
AV277440	18020 Nfatc2ip	4.6769755
BI556771	76974 1190003J15Ri	4.717042917
BB559861	60510 Syt9	4.853818083
BB218107	16792 Laptm5	4.885893
NM_030676	26424 Nr5a2	5.081557
NM_008108	14562 Gdf3	5.14758275
NM_133982	102614 Rpp25	5.190646483
BG245669	14275 Folr1	5.20975025
NM_008652	17865 Mybl2	5.210572917
AV095095	17865 Mybl2	5.318561367
NM_009482	22286 Utf1	5.421930317
AK006110	67080 1700019D03R	5.43684025
C79957	13511 Dsg2	5.656151683

BB739342	58198 Sall1	5.673027467
BB414515	20527 Slc2a3	5.699811883
BG092030	13511 Dsg2	6.010094817
BG064756	99377 Sall4	6.034042167
BC006054	103551 E130012A19R	6.06322625
AF220524	54427 Dnmt3l	6.105105033
NM_009477	22271 Upp1	6.128136833
BB067210	73693 Dppa4	6.187330683
M75135	20527 Slc2a3	6.2282474
X69698	20527 Slc2a3	6.23042505
AV029604	22773 Zic3	6.296894567
NM_021480	58865 Tdh	6.529936483
AF176530	50764 Fbxo15	6.66547415
AK005720	71827 Lrrc34	6.691872117
BB550860	21420 Tcfap2c	6.771666083
AK010400	406217 Bex4	6.816883167
AV212609	353283 Eras	6.834029183
AV298358	381269 Mreg	6.872264583
NM_009337	21432 Tcl1	6.879203683
BB732077	22773 Zic3	6.953696833
AV099404	245128 AU018091	7.054458833
U31967	20674 Sox2	7.296394917
BB709552	14175 Fgf4	7.45980965
AV333667	26380 Esrrb	7.469081817
NM_007430	11614 Nr0b1	7.59146485
BB041571	381591 L1td1	7.663746183
NM_009556	22702 Zfp42	7.9192687
NM_009864	12550 Cdh1	7.9907572
NM_013633	18999 Pou5f1	8.090839917
AK010332	71950 Nanog	8.503540967
AV294613	21667 Tdgf1	8.80162735

Table S4 Top 20 important miRNAs in stem cells

mmu-miR-290
mmu-miR-293
mmu-miR-291a-3p
mmu-miR-292-5p
mmu-miR-295
mmu-miR-294
mmu-miR-291a-5p
mmu-miR-291b-5p
mmu-miR-302b
mmu-miR-124a
mmu-miR-182
mmu-miR-200a
mmu-miR-20b
mmu-miR-302d
mmu-miR-106a
mmu-miR-183
mmu-miR-135b
mmu-miR-690
mmu-miR-142-3p
mmu-miR-302