

Systematically dissecting the mechanistic and functional roles of miRNAs in pluripotent stem cells

Anyou Wang^{1*} et al

¹Bioinformatics core and Lineberger Comprehensive Cancer Center, University of North Carolina, Chapel Hill, NC

²Public Health Sciences Division, Fred Hutchinson Cancer Research Center, Seattle, WA

³Department of Obstetrics and Gynecology, University of North Carolina at Chapel Hill, NC

*Corresponding author

Anyou Wang

5109 Bioinformatics Building

Bioinformatics core

University of North Carolina

Chapel Hill, NC 27599

anyou.wang@alumni.unc.edu

Running title: miRNA functions in stem cells

Key words: microRNAs; miRNA; stem cells; pluripotency; induced pluripotent stem cells; iPS, iPSCs; somatic cells; systems biology; systems network; module; development; metabolism; function; regulatory; network modules; protein complex; miRNAs

Abstract: MicroRNAs (miRNAs) critically modulate stem cell pluripotency and their differentiation, but the precise mechanistic mechanism remains largely unknown. This study systematically reveals the functional and mechanistic roles of miRNAs in mouse pluripotent stem cells by analyzing the genome-wide physical interactions between all activated miRNAs and their targets. Generally, miRNAs vary their physical targets and

functions with the switch of pluripotency to differentiation state. During pluripotency miRNAs primarily and mechanistically target and repress developmental processes, but surprisingly miRNAs do not directly target the pluripotent core factors but only mediate extrinsic signal pathways associated with pluripotency. During differentiation miRNAs mechanistically inhibit metabolism and directly repress the pluripotency. Interestingly, DNA methylation in enhancer regions mediates these miRNA activations. Together, under mediation by DNA methylation, miRNAs directly repress development and mechanistically modulate the pluripotent signal pathways to help stem cells maintain pluripotency; yet miRNAs directly repress both pluripotency and metabolism to facilitate cell differentiation.

Introduction

MicroRNAs (miRNAs), short (~22 nts) conserved endogenous non-coding RNAs, inhibit targeting messenger RNAs by repressing translation or reducing mRNA stability ¹. MiRNAs critically modulate many cellular events, including the balance between proliferation and differentiation during organ development ². In pluripotent stem cells (including induced pluripotent stem cells and embryonic stem cells, referred as stem cells thereafter), miRNAs critically modulate stem cell properties such as pluripotency ³⁻⁵ and their differentiation ⁶⁻⁹. These regulatory roles might be caused by miRNAs indirectly, possibly via downstream bioprocesses. In addition, these regulatory roles of

miRNAs have been derived from the individual miRNAs studies although high-throughput data might be employed. The results from these studies might be biased to individual miRNAs. The general mechanistic mechanism of overall miRNA functions in stem cells remains elusive.

MiRNAs normally execute their functions by targeting the transcription factors and key pathways ¹⁰. These miRNAs might directly interact with their targets together in a network style when they perform their functions. One miRNA may target more than 100 genes ¹¹⁻¹², and one gene can be repressed by multiple miRNA in a sequence-specific fashion ^{10-11,13}. In turn, proteins targeted by miRNA can physically bind to the promoters and enhancers of miRNAs to regulate miRNA activations ⁹. These physical interactions between miRNAs and proteins would form a mechanistic network in stem cells. Systematically revealing such as a physical network would help to understand the overall mechanistic basis of miRNA functions. However, in stem cells, the activation of all miRNAs (~130 miRNAs) ¹⁴⁻¹⁵, like miR-290 family ¹⁶, make this task challenging.

There are many ways to infer with network interactions such as those of physical binding and correlation relationship as we previously reported ¹⁷⁻²⁰. For the physical interactions, the motif binding and experimental binding data can be employed. While the motif-based approaches, like targetscan ²¹, have predicted miRNA targets for more than a decade, a new experiment method called CLIP-seq has recently been developed ²²⁻²³. The CLIP-seq measures RNA fragmentation sequences harvested from miRNA-mRNA

complexes precipitated by Argonaute, a key protein in miRNA binding system. Due to its direct experimental measurements, CLIP-seq should provide more reliable interaction data than computational predictions based on motif-binding. Most of these miRNA target interactions have been collected in the recent miRNA interaction database ^{21,24-25}, but interactions from the current database lack the concern of stem cell specific biases. Applications of these data to species-specific stem cells would produce a lot of noise. In addition, the miRNA target database largely lacks experimental protein binding interactions. Therefore, a mechanistic map containing physical bi-directional interactions of miRNAs and their targets in stem cells remains to be uncovered.

This study constructs a comprehensive physical map of miRNAs and their targets specific for stem cells. This network includes bi-directional interactions between miRNA and their targets. The interactions from miRNAs to their targets were generated from CLIP-seq data and the interactions from proteins to miRNAs were extracted from ChIP-seq data, which measure the protein-binding to miRNA regulatory elements. Analysis of the network enriched by expression data reveals that miRNAs physically repress development, metabolism, and pluripotency in different cellular states. Furthermore, DNA methylation in the upstream of miRNAs mediates these miRNA activations.

Results

Physical network of miRNA and protein interactions in stem cells

To systematically reveal the mechanistic and functional roles of miRNAs in stem cells,

this study first constructed a global physical map of mouse miRNA and protein interactions (Figure 1, materials and methods). This network includes interactions from miRNAs to their targets and interactions from proteins to miRNAs (Figure 1A). The interactions from miRNAs to their target were generated from CLIP-seq and those from proteins to miRNA promoters and enhancers were primarily from ChIP-seq (Figure 1B, Table S1 and materials and methods). The entire network contains ~7000 nodes and ~150,000 miRNA and protein interactions (Figure 1C). This network should be comprehensive for stem cells. For example, the comprehensive interactions between miRNAs and the pluripotent core factors (NANOG, POU5F1 and SOX2) could be extracted from this network (Figure 1D). Due to its natural interactions generated from experimental data, this network would provide true interactions between miRNA and their targets and results generated from this network would be reliable.

The general mechanistic and functional roles of miRNAs in stem cells

MiRNAs critically regulate stem cell properties³⁻⁴, but the general mechanistic mechanism remains largely unknown. To understand the overall mechanistic and functional roles of all activating miRNAs in stem cells, this study enriched the entire physical network with frequently activated (up or down-regulated) miRNAs and activated genes coding for proteins (Table S2-S3, materials and methods). The list of activated miRNAs and genes in stem cells were created when their gene expressions were compared to those of somatic cells. The up-regulated miRNAs and down-regulated

genes in stem cells (Table S2-S3) formed a mechanistic network that primarily functions for development (Figure 2A). The network contains 6 modules and all of them primarily function for development (Figure 2A).

To be simply, we turned to the most important miRNAs. A total of 17 out of the 20 top important miRNAs (Table S4, materials and methods), such as miR-290-295 cluster members, directly target a developmental subnetwork (Figure S1). Furthermore, even a single top miRNA, miR-294 for example here, targets genes functionally enriched in the developmental category (Figure 2B-2C). Therefore, the whole network targeted by over-expressed miRNA, from the large scale of the entire enriched network to the small scale of subnetworks, modules, and even individual miRNA targets, primarily functions for development, indicating that over-expressed miRNAs in stem cells primarily repress development in stem cells. Biologically, to maintain the self-renewal and pluripotency, stem cells should have some ways to prevent development and differentiation. This repressing function of miRNAs can help miRNAs claim their contributions to the stem cell properties at the pluripotent state.

On the other hand, the down-regulated miRNAs in stem cells directly target genes that primarily function for metabolism (Figure S2). These down-regulated miRNAs would become up-regulated during differentiation and they thus would directly inhibit metabolism during this state. Together, miRNAs mechanistically repress development and metabolism respectively during the pluripotent state and during cell differentiation.

MiRNA roles in mediating pluripotency

MiRNAs mediating pluripotency have been widely investigated ¹⁶. For example, miR-294 has been reported as a regulator for c-Myc ¹⁶, a transcription factor mediating pluripotency during cell state transition, but the mechanistic mechanisms still remain largely unclear. In addition, most of the current relative studies have focused on individual miRNAs. Results generated from these studies may not provide the general picture of how miRNAs regulate pluripotency. To systematically explore the regulatory roles of miRNAs in mediating the stem cell pluripotency, this study searched the miRNA direct targets associated with pluripotency. First, we investigated the shortest network path from an individual miRNA (miR-294) to the core pluripotency factors (POU5F1, NANOG, SOX2) and two other transcription factors (KLF4 and MYC) that have been used for producing induced pluripotent stem cells. MiR-294 was selected for this purpose because miR-294 is a top important miRNA in stem cells (Table S4) and it is highly activated in stem cells (Figure 2). Surprisingly, miR-294 does not directly target any above factors (Figure 3A). Actually, the mechanistic basis of miR-294 regulating the pluripotency core seemed dim because miR-294 must go through at least 2 steps to reach any pluripotency core factors. These two steps include a miRNA and a protein, which are not consistently expressed with miR-294. This suggested that miR-294 does not directly mediate the pluripotency.

The above result might be biased to the selected single miRNA. To be unbiased, this

study used all pluripotent genes ²⁶ to systematically search all direct targets that were targeted by all up-regulated miRNAs (Figure 3B). Surprisingly, the core pluripotency factors (e.g. NANGO and POU5F1) were not directly targeted by any up-regulated miRNAs. Instead, the over-expressed miRNAs directly bind to genes functioning for extrinsic signal pathways associated with pluripotency, such as the high connected nodes APC and RAD21 that involve in Wnt pathway and mitotic cell cycle pathway (Figure 3B). This indicated that up-regulated miRNAs might not directly regulate the core pluripotency, but only mediate signal pathways associated with pluripotency.

In contrast, down-regulated miRNAs directly target the core pluripotency factors and pluripotent signal pathways (Figure S3). Interestingly, most of these pluripotent signal pathways are also targeted by up-regulated miRNAs (Figure 3B), indicating that these pluripotent signal pathways are carefully modulated by multiple up- and down-regulated miRNAs. This suggested that the dynamic balance of these extrinsic pluripotent signals mediated by miRNAs might determine the cell states, such as staying in pluripotency or differentiation and lineage commitment.

To make the picture clear, these common signal pathway components were filtered out to reveal only the networks specifically targeted by up-regulated miRNAs (Figure 3C) or by down-regulated miRNAs (Figure 3D). The up-regulated specific network primarily consists of development and differentiation activators (ADK,ALDOA,EHMT1) and metabolism suppressors (CCRN4L) (Figure 3C), indicating that over-expressed miRNAs

still repress development activations by suppressing their activators and promote metabolism by repressing metabolism suppressors in the pluripotent state. This is consistent with the previous discussion (Figure 2). Yet, the down-regulated specific network exclusively includes the pluripotency core factors (Figure 3D), suggesting that when these down-regulated miRNAs become up-regulated during differentiation, these miRNAs would repress pluripotency and then facilitate differentiation.

In summary, miRNAs in the pluripotent state do not directly repress the activities of the core pluripotent factors, but only mediate signal pathways associated with pluripotency at the pluripotency state, while miRNAs physically repress pluripotency and facilitate differentiation during cell differentiation.

DNA methylation mediates the miRNA activation in stem cells

To search the mechanisms controlling the miRNA activations, this study turned to the genome-wide sequencing of DNA methylation in stem cells and methylation-loss-stem cells ²⁷ (Table S1, materials and methods). A total of 2000bp in each upstream and downstream of start sites of all activated miRNAs were examined. While the DNA methylation in the downstream of up-regulated miRNAs are not different from that of down-regulated miRNAs, the down-regulated miRNAs hold significantly higher methylation upstream than up-regulated miRNAs (Figure 4A). Surprisingly, the big difference is not in the promoter regions, but in the enhancer region, ~1000bp up-stream (Figure 4B). Furthermore, these differences are negatively correlated to miRNA

expressions (Figure 4C), suggesting that the difference in DNA methylation accounts for the miRNA activations. This parallels with a recent observation showing that the loss of DNA methylation significantly increases miRNA expressions²⁸. Therefore, miRNA activations and their network are mediated by DNA methylation in enhancer regions.

Discussion

This study first time, as our knowledge, investigates the general mechanistic and functional roles of miRNAs in stem cells. Previous studies have emphasized the regulatory roles of individual miRNAs. This leaves a big question to be answered: what are these ~130 activated miRNAs doing in stem cells in general? This study systematically revealed that miRNAs generally repress development, metabolism and pluripotency in a cellular state fashion. The results concluded from this present study should be reliable because they were generalized from the real experimental data, including physical binding data (CLIP-seq and CHIP-seq) and expression data (RNA-seq and microarray).

MiRNAs play critical roles in maintaining the overall properties in stem cells. Knocking out the key miRNA processing enzymes Dicer³⁻⁴ or DGCR8⁵ causes stem cells lose their pluripotency. However, the mechanistic mechanisms still remain elusive. In this study, we systematically revealed that miRNAs primarily repress developmental activators and developmental modules during the pluripotent state (Figure 2, Figure 3).

This suggested that the primary functions of miRNAs in stem cells are to repress development to prevent stem cell differentiations and to keep stem cells stay in the pluripotent state. On the other hand, miRNAs repress the metabolism suppressors (Figure 3) to maintain the high energy demands in the pluripotent state.

Many studies tried to find the link between miRNAs and pluripotency ^{1,16,29}. MiR-290 cluster has been proposed to regulate the core pluripotency factors like OCT4 ^{1,16,29}. MiRNAs have also been used to induce pluripotency ³⁰. These results are mostly derived from regulatory data and the link between miRNAs and pluripotency could be direct or indirect. This present study showed that mechanistically miRNAs do not directly target any core pluripotent factors during the pluripotent state (Figure 3). This suggested that miRNAs might not directly involve in the key pluripotent processes in stem cells. However, miRNAs directly target the extrinsic signal pathways associated with pluripotency (Figure 3). These signal pathways are actually regulated by both up- and down-regulated miRNAs (Figure 3 and Figure S3). The delicate regulation by these up- and down-regulated miRNAs could determine the cell state, either to pluripotency or to differentiation. Thus modulating pluripotent signal pathways would be the key mechanistic link between miRNA functions and pluripotency.

Individual miRNAs have been reported in regulating cell differentiation ^{9,31}. miR-143 regulates the differentiation of adipocytes ³¹; miR-181 is related to differentiation of B-lineage cells ⁶, while miR-155 is associated with development of immune system ⁷. miR-

miR-145 and miR-143 are involved in smooth muscle differentiation and proliferation⁸, and miR-145 functions as a critical switch in promoting cell differentiation⁹. However, the mechanistic mechanism remains unclear. This present study found that miRNAs directly target both the core pluripotent factors and metabolism genes in the differential state. This suggested that in the differential state miRNAs primarily repress both the pluripotency and metabolism to facilitate the cell differentiations. Repressing only either the pluripotency or metabolism alone might not sufficient to help stem cell differentiations.

DNA methylation might regulate expressions of a certain group of genes in stem cells³². Whether DNA methylation globally mediates the miRNA activations has not been examined. We analyzed the genome methylation sequencing data and revealed that the methylation levels of the upstream of miRNAs, especially the enhancer regions (around -1000bp), are anti-correlated to the miRNA activations (figure 4). This suggested that DNA methylation globally mediates the miRNA activations in stem cells.

Together, under mediated by DNA methylation, miRNAs contribute to maintain pluripotency by directly repressing development and mechanistically regulating the pluripotent signal pathways. On the other hand, to facilitate cell differentiation, miRNAs directly repress both pluripotency and metabolism during the differentiation stage. Understanding these general fundamental roles of miRNAs would broadly direct the future functional studies of miRNAs in stem cells and would guide the successful

development of stem-cell-based therapies for regenerative medicine.

Materials and methods

Data resources

This study analyzed data generated by high-throughput methods (Table S1), including CLIP-seq, CHIP-seq, RNA-seq, microarray, and bisulfite sequencing DNA Methylation. The data was downloaded from GEO database (www.ncbi.nlm.nih.gov/geo/).

Bioinformatics analysis

To be consistent and comparable, all sequencing data were mapped to mouse mm10 genome. All fastq files (Table S1) from CLIP-seq²²⁻²³, CHIP-seq and RNA-seq were aligned using BWA³³. Bisulfite sequencing DNA Methylation was aligned by Bismark (<http://www.bioinformatics.babraham.ac.uk/projects/bismark/>). Downstream analyses were performed by in-house programs using R, python, SciPy and Linux shell scripts. Microarray data were analyzed using R.

The network construction and analysis were performed by approaches as our previous reports^{17-18,32}. The original network included data from the miRNA and protein interaction database, but the overlap between the motif-based interactions and experimental ones were less than 20%. Thus, these motif-based miRNA target interactions were deleted from the network, and only miRNA to target interactions generated from experimental data were used for our analysis and to draw results and conclusions in this study. This practice ensures our conclusions are reliable.

The highly frequent genes were selected from all observations obtained by the study (Table S1). Genes with activating frequency >50% and miRNAs with frequency of more than 25% were selected (Table S2-S3). The top important genes (Table S4) were selected by using the algorithm that counts on network topology and eigengene score as we previously published ³².

Competing interests

No competing interests exist in this study
No financial conflict of interest in this study.

Acknowledgments

We specially thank Dr. Amanda Jackson for reading this manuscript.

References

- 1 Benetti, R. *et al.* A mammalian microRNA cluster controls DNA methylation and telomere recombination via Rbl2-dependent regulation of DNA methyltransferases. *Nat Struct Mol Biol* **15**, 998, doi:nsmb0908-998b [pii]
10.1038/nsmb0908-998b (2008).
- 2 Kloosterman, W. P. & Plasterk, R. H. The diverse functions of microRNAs in animal development and disease. *Dev Cell* **11**, 441-450, doi:S1534-5807(06)00402-3 [pii]
10.1016/j.devcel.2006.09.009 (2006).
- 3 Kanellopoulou, C. *et al.* Dicer-deficient mouse embryonic stem cells are defective in differentiation and centromeric silencing. *Genes Dev* **19**, 489-501, doi:19/4/489 [pii]
10.1101/gad.1248505 (2005).
- 4 Murchison, E. P., Partridge, J. F., Tam, O. H., Cheloufi, S. & Hannon, G. J. Characterization of Dicer-deficient murine embryonic stem cells. *Proc Natl Acad Sci U S A* **102**, 12135-12140, doi:0505479102 [pii]
10.1073/pnas.0505479102 (2005).
- 5 Wang, Y., Medvid, R., Melton, C., Jaenisch, R. & Blelloch, R. DGCR8 is essential for microRNA biogenesis and silencing of embryonic stem cell self-renewal. *Nat Genet* **39**, 380-385, doi:ng1969 [pii]

- 10.1038/ng1969 (2007).
- 6 Chen, C. Z., Li, L., Lodish, H. F. & Bartel, D. P. MicroRNAs modulate hematopoietic lineage differentiation. *Science* **303**, 83-86, doi:10.1126/science.1091903 1091903 [pii] (2004).
- 7 Turner, M. & Vigorito, E. Regulation of B- and T-cell differentiation by a single microRNA. *Biochem Soc Trans* **36**, 531-533, doi:BST0360531 [pii] 10.1042/BST0360531 (2008).
- 8 Cordes, K. R. *et al.* miR-145 and miR-143 regulate smooth muscle cell fate and plasticity. *Nature* **460**, 705-710, doi:nature08195 [pii] 10.1038/nature08195 (2009).
- 9 Xu, N., Papagiannakopoulos, T., Pan, G., Thomson, J. A. & Kosik, K. S. MicroRNA-145 regulates OCT4, SOX2, and KLF4 and represses pluripotency in human embryonic stem cells. *Cell* **137**, 647-658, doi:S0092-8674(09)00252-9 [pii] 10.1016/j.cell.2009.02.038 (2009).
- 10 Stefani, G. & Slack, F. J. Small non-coding RNAs in animal development. *Nat Rev Mol Cell Biol* **9**, 219-230, doi:nrm2347 [pii] 10.1038/nrm2347 (2008).
- 11 Bartel, D. P. MicroRNAs: target recognition and regulatory functions. *Cell* **136**, 215-233, doi:S0092-8674(09)00008-7 [pii] 10.1016/j.cell.2009.01.002 (2009).
- 12 Chen, C. *et al.* Defining embryonic stem cell identity using differentiation-related microRNAs and their potential targets. *Mamm Genome* **18**, 316-327, doi:10.1007/s00335-007-9032-6 (2007).
- 13 Rajewsky, N. microRNA target predictions in animals. *Nat Genet* **38 Suppl**, S8-13, doi:ng1798 [pii] 10.1038/ng1798 (2006).
- 14 Laurent, L. C. *et al.* Comprehensive microRNA profiling reveals a unique human embryonic stem cell signature dominated by a single seed sequence. *Stem Cells* **26**, 1506-1516, doi:2007-1081 [pii] 10.1634/stemcells.2007-1081 (2008).
- 15 Morin, R. D. *et al.* Application of massively parallel sequencing to microRNA profiling and discovery in human embryonic stem cells. *Genome Res* **18**, 610-621, doi:gr.7179508 [pii] 10.1101/gr.7179508 (2008).
- 16 Hanina, S. A. *et al.* Genome-wide identification of targets and function of individual MicroRNAs in mouse embryonic stem cells. *PLoS Genet* **6**, e1001163, doi:10.1371/journal.pgen.1001163 (2010).
- 17 Anyou Wang, L. R., and Hong Li. A Systemic Network Triggered by Human Cytomegalovirus Entry. *Advances in Virology* **2011**, 11 (2011).
- 18 Wang, A., Johnston, S. C., Chou, J. & Dean, D. A systemic network for Chlamydia pneumoniae entry into human cells. *J Bacteriol* **192**, 2809-2815, doi:JB.01462-09 [pii] 10.1128/JB.01462-09 (2010).

- 19 Wang, A. *et al.* Functional Modules Distinguish Human Induced Pluripotent Stem Cells from Embryonic Stem Cells. *Stem Cells Dev.* **in Press**, doi:10.1089/scd.2010.0574 (2011).
- 20 Wang, A. *et al.* Transcription Factor Complex AP-1 Mediates Inflammation Initiated by Chlamydia pneumoniae Infection. *Cell Microbiol*, doi:10.1111/cmi.12071 (2012).
- 21 Lewis, B. P., Burge, C. B. & Bartel, D. P. Conserved seed pairing, often flanked by adenosines, indicates that thousands of human genes are microRNA targets. *Cell* **120**, 15-20, doi:S0092867404012607 [pii] 10.1016/j.cell.2004.12.035 (2005).
- 22 Leung, A. K. *et al.* Genome-wide identification of Ago2 binding sites from mouse embryonic stem cells with and without mature microRNAs. *Nat Struct Mol Biol* **18**, 237-244, doi:nsmb.1991 [pii] 10.1038/nsmb.1991 (2011).
- 23 Chi, S. W., Zang, J. B., Mele, A. & Darnell, R. B. Argonaute HITS-CLIP decodes microRNA-mRNA interaction maps. *Nature* **460**, 479-486, doi:nature08170 [pii] 10.1038/nature08170 (2009).
- 24 Vergoulis, T. *et al.* TarBase 6.0: capturing the exponential growth of miRNA targets with experimental support. *Nucleic Acids Res* **40**, D222-229, doi:gkr1161 [pii] 10.1093/nar/gkr1161 (2012).
- 25 Yang, J. H. *et al.* starBase: a database for exploring microRNA-mRNA interaction maps from Argonaute CLIP-Seq and Degradome-Seq data. *Nucleic Acids Res* **39**, D202-209, doi:gkq1056 [pii] 10.1093/nar/gkq1056 (2011).
- 26 Hu, G. *et al.* A genome-wide RNAi screen identifies a new transcriptional module required for self-renewal. *Genes Dev* **23**, 837-848, doi:23/7/837 [pii] 10.1101/gad.1769609 (2009).
- 27 Meissner, A. *et al.* Genome-scale DNA methylation maps of pluripotent and differentiated cells. *Nature* **454**, 766-770, doi:nature07107 [pii] 10.1038/nature07107 (2008).
- 28 Yan, H., Choi, A. J., Lee, B. H. & Ting, A. H. Identification and functional analysis of epigenetically silenced microRNAs in colorectal cancer cells. *PLoS One* **6**, e20628, doi:10.1371/journal.pone.0020628 PONE-D-11-01853 [pii] (2011).
- 29 Sinkkonen, L. *et al.* MicroRNAs control de novo DNA methylation through regulation of transcriptional repressors in mouse embryonic stem cells. *Nat Struct Mol Biol* **15**, 259-267, doi:nsmb.1391 [pii] 10.1038/nsmb.1391 (2008).
- 30 Miyoshi, N. *et al.* Reprogramming of mouse and human cells to pluripotency using mature microRNAs. *Cell Stem Cell* **8**, 633-638, doi:S1934-5909(11)00221-9 [pii] 10.1016/j.stem.2011.05.001 (2011).
- 31 Esau, C. *et al.* MicroRNA-143 regulates adipocyte differentiation. *J Biol*

Chem **279**, 52361-52365, doi:C400438200 [pii]
10.1074/jbc.C400438200 (2004).

32 Wang, A. *et al.* Functional modules distinguish human induced pluripotent stem cells from embryonic stem cells. *Stem Cells Dev* **20**, 1937-1950, doi:10.1089/scd.2010.0574 (2011).

33 Li, H. & Durbin, R. Fast and accurate long-read alignment with Burrows-Wheeler transform. *Bioinformatics* **26**, 589-595, doi:btp698 [pii]
10.1093/bioinformatics/btp698 (2010).

Figure legends

Figure 1. Overall view. A, The concept of miRNA and protein interactions. B, Workflow of this study. C, Overall view of the entire network constructed by this study. D, A sample of entire network contents. Nodes denote miRNAs or proteins as labeled, red node represents the gene up-regulation in pluripotent stem cells, green node as down-regulation, white node as insignificant expression. Edges represent interactions, red edge represents miRNA targeting proteins and green edge represents protein binding the regulatory elements of miRNAs. This labeling strategy applies to all figures in this study.

Figure 2. MiRNAs primarily repress developmental processes in pluripotent stem cells. A, Up-regulated miRNAs primarily target genes associated with developmental processes, which include 6 primary modules (protein complexes) functioning for developments as highlighted in cycles. All these modules primarily function for developments. B, Targets of mmu-miR-294, a highly expressed miRNA in stem cell and a top miRNAs in this study (Table S4), function for development as shown in C. C, Developmental category is significantly and abundantly enriched by miRNA-294 targets.

Similar functional enrichments were observed for other up-regulated miRNA targets in stem cells.

Figure 3. MiRNAs indirectly target the key pluripotent factors in stem cells. A, The shortest path from miR-294 to the pluripotent core factors. MiR-294 does not directly bind to any these core factors. B, Up-regulated miRNAs directly target several pluripotent genes except the core factors, but these targets serve for only signal pathway associated with pluripotency and most of these up-regulated miRNA targets are also targeted by down-regulated miRNAs (Figure S3). The enlarged nodes represent the high connected targets that are targeted by both up- and down-regulated miRNAs in stem cells and they work for pluripotent signal pathways. C, Subnetworks specifically targeted by up-regulated miRNAs only. D. Subnetworks targeted by down-regulated miRNAs only.

Figure 4. DNA methylation mediates miRNA activations in stem cells. A, Methylation level of upstream and downstream of miRNA start site. Down-regulated miRNAs have a significantly high methylation in upstream when compared with up-regulated miRNAs. B, Detail methylation profiling of 2000bp upstream of activated miRNAs. The top 30 down-regulated miRNAs have specifically higher methylation around -1000bp (middle panel) than that of the top 30 up-regulated miRNAs (upper panel). The methylation profiling of a single down-regulated miRNA serves as a representative example (bottom panel). C. Negative correlation between DNA

methylation and miRNA expression.

Supplemental materials

Figure S1, total 17 out of top 20 important miRNAs target developmental genes.

Figure S2, down-regulated miRNAs mediate metabolism in stem cells.

Figure S3, pluripotent genes targeted by down-regulated miRNAs in stem cells.

Table S1-S4 were shown in sheet1 to sheet4 in one excel file

Table S1, Data sources


Table S2, a list of miRNAs that are frequently and significantly differential expressed between stem cells and somatic cells.

Table S3, a list of genes coding for proteins that are frequently and significantly differential expressed between stem cells and somatic cells.


Table S4, a list of top 20 important miRNAs

Figure 1


A. miRNA and protein interactions



B. Workflow



C. Entire network



D. A section of entire network






Figure 2

A



B



C






Figure 3 A




Shortest path from miRNA294

C




Specific for up-regulated miRNAs

D



Specific for down-regulated miRNAs


B




Pluripotent genes targeted by up-regulated miRNAs in pluripotent stem cells

Figure 4


A



C



B



Supplemental materials

Figure S1, total 17 out of top 20 important miRNAs target developmental genes.

Figure S2, down-regulated miRNAs mediate metabolism in stem cells.

Figure S3, pluripotent genes targeted by down-regulated miRNAs in stem cells.

Table S1, Data sources

Table S2, a list of miRNAs that are frequently and significantly differential expressed between stem cells and somatic cells.

Table S3, a list of genes coding for proteins that are frequently and significantly differential expressed between stem cells and somatic cells.

Table S4, a list of top 20 important miRNAs

Figure S1 Total 17 out of top 20 miRNAs target developmental genes




Figure S2. Down-regulated miRNAs mediate metabolism





Figure S3. Pluripotent genes targeted by down-regulated miRNAs



Sheet1

Table S1 data resources

GEO number	Platform
GSE25310	CLIP-seq
GSE11724	ChIP-seq
GSE11431	ChIP-seq
GSM278905	RRBS
GSM278902	RRBS
GSM539867	RNA-Seq
GSM539866	RNA-Seq
GSE30012	Mouse430_2 array
GSE25310	GPL11410
GSM231739	Agilent-GPL2872
GSE15267	AffymetrixGPL1261
GSE14012	AffymetrixGPL1261
GSE10871	AffymetrixGPL1261
GSE14790	AffymetrixGPL1261
GSE16062	IlluminaGPL6885

Sheet1

Table S2 miRNAs differentially expressed between stem cell vs somatic cells with high frequency >25% of observations

geneID

mmu-miR-291a-3p	mmu-miR-17-5p
mmu-miR-292-5p	mmu-miR-669b
mmu-miR-295	Mmu-miR-363
mmu-miR-290	mmu-miR-130b
mmu-miR-291b-5p	mmu-miR-142-5p
mmu-miR-294	mmu-miR-708
mmu-miR-293	mmu-miR-301b
mmu-miR-291a-5p	mmu-miR-297
mmu-miR-291b-3p	mmu-miR-711
mmu-miR-205	mmu-miR-135b
mmu-miR-292-3p	mmu-miR-145
mmu-miR-200a	mmu-miR-125b
mmu-miR-20b	mmu-miR-675-3p
mmu-miR-429	mmu-miR-31
mmu-miR-93	mmu-miR-431
mmu-miR-302b	mmu-miR-133a
mmu-miR-124a	mmu-miR-434-5p
mmu-miR-141	mmu-miR-29a
mmu-miR-32	mmu-miR-125a
mmu-miR-25	mmu-miR-27b
mmu-miR-106a	mmu-miR-206
mmu-miR-96	mmu-miR-222
mmu-miR-690	mmu-miR-329
mmu-miR-210	mmu-let-7i
mmu-miR-182	mmu-miR-221
mmu-miR-339	mmu-miR-132
mmu-miR-33	mmu-miR-345
mmu-miR-19b	mmu-miR-452
mmu-miR-302d	mmu-let-7d
mmu-miR-195	mmu-miR-181a*
mmu-miR-297b	mmu-miR-212
mmu-miR-499	mmu-miR-152
mmu-miR-106b	mmu-miR-143
mmu-miR-712*	mmu-miR-433-5p
mmu-miR-101b	mmu-let-7e
mmu-miR-706	mmu-miR-322
mmu-miR-672	mmu-miR-380-5p
mmu-miR-20a	mmu-miR-666
mmu-miR-302	mmu-miR-365
mmu-miR-142-3p	mmu-miR-10a
mmu-miR-19a	mmu-miR-24*
mmu-miR-450b	
mmu-miR-200c	

Sheet1

Table S3. Genes differentially expressed between stem cells
and somatic cells with high frequency >50% of observations

GB_ACC	ENTREZ_GENE	Gene.Symbol
BC006640	20315	Cxcl12
BF225802	16011	Igfbp5
AW550625	12825	Col3a1
BF227507	12843	Col1a2
BB542051	18295	Ogn
U08020	12842	Col1a1
NM_013655	20315	Cxcl12
NM_011581	21826	Thbs2
AV229424	12832	Col5a2
BB051738	18933	Prrx1
M65143	16948	Lox
AF007248	14118	Fbn1
NM_009933	12833	Col6a1
NM_007993	14118	Fbn1
BC019502	12111	Bgn
AB015978	18414	Osmr
BE197945	20324	Sdpr
NM_007729	12814	Col11a1
AW049660	18032	Nfix
NM_010514	16002	Igf2
BB197591	54216	Pcdh7
AV226618	13837	Epha3
AF378762	69538	Antxr1
M68513	13837	Epha3
BC002064	19242	Ptn
NM_011340	20317	Serpinf1
BB795075	18214	Ddr2
BC023060	216616	Efemp1
BF144658	21808	Tgfb2
BB250384	22329	Vcam1
AB029929	12389	Cav1
D67076	11504	Adamts1
BB315728	18032	Nfix
BI794771	12842	Col1a1
L06502	18933	Prrx1
BG793483	21813	Tgfbr2
BB532202	14264	Fmod
BB361162	22771	Zic1
AI931862	12111	Bgn
BC014690	21809	Tgfb3
NM_018865	22402	Wisp1
AV064339	20324	Sdpr
BG067986	54216	Pcdh7
NM_030888	81799	C1qtnf3
BB233297	21826	Thbs2
BF168458	21858	Timp2

Sheet1

BB259670	14362 Fzd1
AA499047	18596 Pdgfrb
BC026446	219151 Scara3
NM_010714	16876 Lhx9
NM_133859	99543 Olfm13
BB067079	22418 Wnt5a
AW555664	399558 Flrt2
BC021484	16651 Sspn
BC011063	15402 Hoxa5
M93954	21858 Timp2
AV310588	12444 Ccnd2
NM_013589	16997 Ltbp2
BB791906	14178 Fgf7
BQ175880	12444 Ccnd2
NM_009829	12444 Ccnd2
BC022107	12558 Cdh2
NM_011160	19091 Prkg1
AK007904	12444 Ccnd2
BB114398	12816 Col12a1
BB377340	13612 Edil3
BI455189	12834 Col6a2
BC018425	22418 Wnt5a
AF309564	80837 Rhoj
AW744319	12831 Col5a1
NM_010284	14600 Ghr
AK003674	68588 Cthrc1
NM_009610	11468 Actg2
BM220945	239217 Kctd12
BB658835	23794 Adamts5
BC019836	16010 Igfbp4
NM_018764	54216 Pcdh7
AA500897	26362 Axl
AV222756	259300 Ehd2
NM_018884	55983 Pdzn3
BB759833	17300 Foxc1
BQ177170	17112 Tm4sf1
BM250666	12830 Col4a5
BC025502	231532 Arhgap24
NM_007392	11475 Acta2
NM_008885	18858 Pmp22
AV246911	12831 Col5a1
NM_133918	100952 Emilin1
AV260198	83675 Bicc1
BC021376	225207 Zfp521
NM_011607	21923 Tnc
NM_007833	13179 Dcn
AF357006	16949 Loxl1
BC011507	74480 Samd4
BB493031	23888 Gpc6

Sheet1

BI249259	18032 Nfix
NM_054044	78560 Gpr124
BC001999	13732 Emp3
BB041180	16876 Lhx9
BI690209	57342 Parva
AK018128	70717 6330406I15Rik
AI595932	17260 Mef2c
AI463873	11819 Nr2f2
BG075699	399558 Flrt2
AV149705	108927 Lhfp
AK014221	73230 Bmper
BB468025	54381 Pgcg
BB363812	14051 Eya4
AK020118	23888 Gpc6
BM217996	83675 Bicc1
NM_009636	11568 Aebp1
AA987181	15405 Hoxa9
NM_011766	22762 Zfpm2
AV315205	11658 Alcam
U95030	11658 Alcam
BF147716	17390 Mmp2
NM_008608	17387 Mmp14
AV332957	20429 Shox2
AK008112	18933 Prrx1
BB787243	16010 Igfbp4
M27130	12505 Cd44
NM_023118	13132 Dab2
BB817332	399558 Flrt2
D63423	11747 Anxa5
BC014870	235587 Parp3
U03425	13649 Egfr
NM_011150	19039 Lgals3bp
BB229377	13717 Eln
BF681826	26360 Angptl2
AV369812	13649 Egfr
AI385532	21825 Thbs1
AK012411	20317 Serpinf1
NM_016900	12390 Cav2
NM_007471	11820 App
BC026153	13841 Epha7
BE652876	60527 Fads3
NM_022563	18214 Ddr2
BC005490	11820 App
AY083458	235505 Cd109
AW146109	12505 Cd44
AF022889	268977 Ltbp1
BF780807	23972 Papss2
NM_010222	14231 Fkbp7
BB765827	74761 Mxra8

Sheet1

BB369191	27528 D0H4S114
NM_011658	22160 Twist1
BC022679	52552 Parp8
BC024375	14600 Ghr
AW558570	13617 Ednra
NM_010151	13865 Nr2f1
BB053506	207181 Rbms3
BB464523	12293 Cacna2d1
AI325255	14230 Fkbp10
NM_009365	21804 Tgfb1i1
AW537708	18595 Pdgfra
BG244279	26360 Angptl2
AV021105	208647 Creb3l2
NM_021474	58859 Efemp2
AW558468	230103 Npr2
BM230959	229731 Slc25a24
NM_013519	14234 Foxc2
BG248060	12153 Bmp1
AK011935	67155 Smarca2
BC009660	109042 Prkcdp
NM_010730	16952 Anxa1
NM_017464	18003 Nedd9
NM_020606	57342 Parva
NM_054042	70445 Cd248
AW107196	74480 Samd4
NM_008520	16998 Ltbp3
NM_007400	11489 Adam12
BB138485	70676 Gulp1
BB041237	17389 Mmp16
NM_009152	20346 Sema3a
AK007400	77889 Lbh
BC027199	233328 Lrrk1
BC003726	14789 Leprel2
BM220576	53623 Gria3
BM570006	20564 Slit3
NM_007472	11826 Aqp1
BB329489	12406 Serpinh1
BC024358	22004 Tpm2
BB371406	57265 Fzd2
BI106777	12293 Cacna2d1
BB484759	224024 Scarf2
BC025145	19266 Ptprd
AK004519	56726 Sh3bgrl
NM_010221	14230 Fkbp10
BB114067	21345 Tagln
NM_010517	16010 Igfbp4
AV164956	22418 Wnt5a
U71189	13592 Ebf2
AF339910	18605 Enpp1

Sheet1

BQ173967	13640 Efna5
AV246882	235505 Cd109
AK003186	22004 Tpm2
NM_009821	12394 Runx1
NM_010743	17082 Il1rl1
BB324823	16998 Ltbp3
BC025514	102644 Oaf
NM_007801	13036 Ctsh
NM_010581	16423 Cd47
AV238225	16905 Lmna
AK018679	16423 Cd47
BG073383	15410 Hoxb3
NM_026405	67844 Rab32
BF144687	70676 Gulp1
NM_019391	16985 Lsp1
NM_013586	16950 Loxl3
AI415741	94352 Loxl2
BI220012	12406 Serpinh1
NM_007802	13038 Ctsk
BB832504	18028 Nfib
BC013463	15430 Hoxd10
M22479	22003 Tpm1
BG963150	20563 Slit2
BC013560	12827 Col4a2
NM_008984	19274 Ptprm
BB233088	21928 Tnfaip2
BC008277	13617 Ednra
BI111620	21859 Timp3
BB475194	23794 Adamts5
NM_021355	14264 Fmod
NM_007585	12306 Anxa2
BF578055	229731 Slc25a24
BB547877	20563 Slit2
NM_016753	17035 Lxn
AV016275	20742 Spnb2
NM_008495	16852 Lgals1
AW551930	83675 Bicc1
AK018466	207181 Rbms3
NM_008987	19288 Ptx3
AK003819	74777 Sepn1
AW412729	12816 Col12a1
BC020152	223272 Itgbl1
Y07687	18028 Nfib
NM_011985	26561 Mmp23
NM_009378	21824 Thbd
BF451748	107589 Mylk
BM251152	13003 Vcan
BB468082	18481 Pak3
NM_007899	13601 Ecm1

Sheet1

BB522674	20541 Slc8a1
NM_019989	56726 Sh3bgrl
BC023448	23888 Gpc6
AK002516	11857 Arhgdib
BF100813	13618 Ednrb
NM_011777	22793 Zyx
BB326929	58194 Sh3kbp1
BB248904	56726 Sh3bgrl
BB027759	231997 Fkbp14
AU021035	15529 Sdc2
AI481026	29817 Igfbp7
NM_008393	16373 Irx3
BB526042	320452 P4ha3
BB100920	17118 Marcks
BB097480	223254 Farp1
BI687652	18028 Nfib
BG868949	17122 Mxd4
NM_020561	57319 Smpd13a
AF059567	12579 Cdkn2b
NM_009472	22253 Unc5c
NM_009794	12334 Capn2
NM_009627	11535 Adm
BQ175796	18481 Pak3
BC005569	58809 Rnase4
BB327018	14118 Fbn1
NM_010212	14200 Fhl2
BB313689	20541 Slc8a1
AK019164	17475 Mpdz
AW702161	11745 Anxa3
BC027084	259300 Ehd2
BC005686	13713 Elk3
BB065799	240725 Sulf1
NM_130449	140792 Colec12
BB823350	19092 Prkg2
AI642438	16852 Lgals1
NM_007855	13345 Twist2
BB787292	14302 Frk
BB354684	217410 Trib2
AF117951	94352 Loxl2
BC025897	225791 Zadh2
NM_019877	56358 Copz2
AV162270	22240 Dpysl3
D63383	20465 Sim2
NM_008546	17150 Mfap2
NM_008008	14178 Fgf7
BQ174721	214791 Sertad4
BB377873	121021 Cspg4
BB542535	13036 Ctsh
BC024876	223453 Dap

Sheet1

L24755	12153 Bmp1
X66083	12505 Cd44
AW547821	17268 Meis1
BC016447	26427 Creb3l1
BM213516	20742 Spnb2
BM220880	108655 Foxp1
AK004853	50781 Dkk3
NM_008809	18596 Pdgfrb
AV245241	259300 Ehd2
NM_020007	56758 Mbnl1
BC028307	68792 Srpx2
AA709993	20365 Serf1
BC006737	12370 Casp8
BB151715	23871 Ets1
BI452727	14314 Fstl1
BB003393	21923 Tnc
BB535494	18003 Nedd9
BB779859	209448 Hoxc10
BI248947	109624 Cald1
L26349	21937 Tnfrsf1a
BC012674	19285 Ptrf
U30244	13642 Efnb2
AK004598	71703 Armcx3
BB075247	19266 Ptprd
BB766878	107581 Col16a1
NM_019586	56228 Ube2j1
NM_008761	18301 Fxyd5
M24849	22352 Vim
BB366804	18230 Nxn
AV031691	22771 Zic1
AK003303	19266 Ptprd
NM_009252	20716 Serpina3n
BM233698	12977 Csf1
AK018666	50766 Crim1
NM_013813	13823 Epb4.1l3
BB097063	16826 Ldb2
BG070361	18514 Pbx1
NM_023564	70310 Plscr3
BC016893	15414 Hoxb6
BC025602	52377 Rcn3
BE628275	68910 Zfp467
BB751459	240725 Sulf1
NM_023813	108058 Camk2d
BE630020	19122 Prnp
NM_015786	50708 Hist1h1c
NM_010736	17000 Ltbr
NM_008380	16323 Inhba
AF080090	20350 Sema3f
AV220340	229534 Pbxip1

Sheet1

NM_016846	19731 Rgl1
AA510713	56468 Socs5
AV246759	21808 Tgfb2
AV238718	233733 Galntl4
BB311061	56876 Nelf
BM231794	216188 Aldh1l2
NM_053147	93893 Pcdhb22
AV124445	17984 Ndn
AI747133	12338 Capn6
NM_009784	12293 Cacna2d1
NM_008675	17965 Nbl1
X58380	15364 Hmga2
NM_009468	22240 Dpysl3
NM_019390	16905 Lmna
BB533903	50708 Hist1h1c
BB250811	18542 Pcolce
BE951265	66180 1110036O03Rik
AW546141	17118 Marcks
BB825801	13713 Elk3
NM_009255	20720 Serpine2
BF158638	12826 Col4a1
NM_016886	53623 Gria3
NM_009925	12813 Col10a1
AV278675	244448 Triml1
BG966751	104156 Etv5
X67668	97165 Hmgb2
BB089717	12449 Ccnf
BB292220	58180 Hic2
BG069191	50927 Nasp
BG066754	67204 Eif2s2
AK019115	68239 Krt42
AU015121	268697 Ccnb1
NM_008566	17218 Mcm5
NM_019812	93759 Sirt1
NM_020599	19771 Rlbp1
AV280841	219132 D14Ertd668e
BB493242	50927 Nasp
NM_022409	63872 Zfp296
NM_013873	29859 Sult4a1
AF302127	72388 Ripk4
NM_011597	21873 Tjp2
BF578266	227327 B3gnt7
AI195532	216456 Glis2
BE650359	13619 Phc1
BM247863	56380 Arid3b
BM935811	16403 Itga6
BG071670	75796 Cdy12
BC006704	226419 Dyrk3
NM_008628	17685 Msh2

Sheet1

BM196098	14562 Gdf3
NM_020567	57441 Gmnn
AK011289	66953 Cdca7
NM_009426	22044 Trh
BB367422	239759 Liph
BC011474	16818 Lck
NM_011176	19143 St14
BB667216	22371 Vwf
BI739053	12725 Clcn3
NM_011487	20849 Stat4
BQ177140	218038 Amph
NM_008452	16598 Klf2
AF329833	66790 Grtp1
BB540053	217653 C79407
NM_007691	12649 Chek1
BC016095	228421 Kif18a
BC017621	223775 Pim3
U42190	17688 Msh6
NM_027288	110173 Manba
BE951628	217653 C79407
BB125424	242894 Actr3b
AK010648	70024 Mcm10
AK006509	75796 Cdy12
NM_010174	14077 Fabp3
NM_053271	116838 Rims2
BG070553	110957 D1Pas1
NM_133664	16763 Lad1
BQ176661	54484 Mkrm1
AF237702	20425 Shmt1
NM_016851	54139 Irf6
AA543734	21386 Tbx3
NM_010346	14786 Grb7
BC016571	50799 Slc25a13
AK007201	14536 Nr6a1
NM_013514	13829 Epb4.9
U39203	16331 Inpp5d
AW552407	217837 Itpk1
AK011462	72148 2610019F03Rik
BB397151	332221 Zscan10
U31625	12189 Brca1
BM225706	243574 Kbtbd8
NM_031386	83560 Tex14
NM_009171	20425 Shmt1
AK011986	112422 2610305D13Rik
AV305746	231327 Ppat
BI465857	12224 Klf5
NM_021308	57746 Piwil2
BB758095	21873 Tjp2
NM_016964	50878 Stag3

Sheet1

BG066919	17765 Mtf2
BE986504	71801 Plekhhf2
AV277242	243963 Zfp473
NM_011799	23834 Cdc6
NM_026190	67486 Polr3g
NM_010357	14860 Gsta4
AU016127	67374 Jam2
BM230524	217944 Rapgef5
NM_022024	63986 Gmfg
AK014330	107435 Hat1
NM_008816	18613 Pecam1
BC003329	54484 Mkrn1
AK011596	22042 Tfrc
BB128741	319801 9630033F20Rik
AF037370	12865 Cox7a1
BC006646	12224 Klf5
AV024881	52609 Cbx7
BB234337	101602 Al467606
AK013117	76974 1190003J15Rik
NM_010068	13436 Dnmt3b
NM_009886	12614 Celsr1
BE648070	58180 Hic2
NM_007469	11812 Apoc1
BC018383	226180 Ina
AB072269	13511 Dsg2
NM_013512	13824 Epb4.1l4a
BB473446	239157 Pnma2
AK010391	71988 Esco2
BC020144	13511 Dsg2
AK009960	69697 2310057J16Rik
AF285580	77485 Stk31
NM_007397	11481 Acvr2b
BC003778	21420 Tcfap2c
NM_011934	26380 Esrrb
NM_016907	20732 Spint1
BC004654	19725 Rfx2
AV060866	18778 Pla2g1b
BE133749	54484 Mkrn1
AF425084	97848 Serpinb6c
AV213552	24074 Taf7
BQ175337	72685 Dnajc6
AF104416	11828 Aqp3
BM210256	14106 Foxh1
NM_023844	67374 Jam2
AK019319	11816 Apoe
NM_031383	83557 Lin28
BG084230	66824 Pycard
AK010826	67374 Jam2
AF330212	12310 Calca

Sheet1

BQ173923	12808 Cobl
M18775	17762 Mapt
BC027285	68713 Ifitm1
AK021186	77485 Stk31
NM_009903	12740 Cldn4
NM_009749	12069 Bex2
NM_009434	22113 Phlda2
BC017609	18424 Otx2
BC023498	72027 Slc39a4
AA215046	67080 1700019D03Rik
BC004037	22256 Ung
NM_007515	11989 Slc7a3
BB469300	243574 Kbtbd8
NM_016754	17907 Mylpf
AV277440	18020 Nfatc2ip
BI556771	76974 1190003J15Rik
BB559861	60510 Syt9
BB218107	16792 Laptm5
NM_030676	26424 Nr5a2
NM_008108	14562 Gdf3
NM_133982	102614 Rpp25
BG245669	14275 Folr1
NM_008652	17865 Mybl2
AV095095	17865 Mybl2
NM_009482	22286 Utf1
AK006110	67080 1700019D03Rik
C79957	13511 Dsg2
BB739342	58198 Sall1
BB414515	20527 Slc2a3
BG092030	13511 Dsg2
BG064756	99377 Sall4
BC006054	103551 E130012A19Rik
AF220524	54427 Dnmt3l
NM_009477	22271 Upp1
BB067210	73693 Dppa4
M75135	20527 Slc2a3
X69698	20527 Slc2a3
AV029604	22773 Zic3
NM_021480	58865 Tdh
AF176530	50764 Fbxo15
AK005720	71827 Lrrc34
BB550860	21420 Tcfap2c
AK010400	406217 Bex4
AV212609	353283 Eras
AV298358	381269 Mreg
NM_009337	21432 Tcl1
BB732077	22773 Zic3
AV099404	245128 AU018091
U31967	20674 Sox2

Sheet1

BB709552	14175 Fgf4
AV333667	26380 Esrrb
NM_007430	11614 Nr0b1
BB041571	381591 L1td1
NM_009556	22702 Zfp42
NM_009864	12550 Cdh1
NM_013633	18999 Pou5f1
AK010332	71950 Nanog
AV294613	21667 Tdgf1

Table S4 Top 20 important miRNAs in stem cells

mmu-miR-290
mmu-miR-293
mmu-miR-291a-3p
mmu-miR-292-5p
mmu-miR-295
mmu-miR-294
mmu-miR-291a-5p
mmu-miR-291b-5p
mmu-miR-302b
mmu-miR-124a
mmu-miR-182
mmu-miR-200a
mmu-miR-20b
mmu-miR-302d
mmu-miR-106a
mmu-miR-183
mmu-miR-135b
mmu-miR-690
mmu-miR-142-3p
mmu-miR-302