

---

# THE LIPID-RNA WORLD: A MEMBRANE BOUND ORIGIN OF RIBOSOME

Saurav Mallik<sup>1\*</sup> and Sudip Kundu<sup>1\*</sup>

<sup>1</sup>Department of Biophysics, Molecular Biology and Bioinformatics, University of Calcutta

Address: 92, APC Road, Kolkata; India; Zip: 700009; Email: Saurav Mallik: saurav.bmb@gmail.com; Sudip Kundu: skbmbg@caluniv.ac.in

---

“The only way of discovering the limits of the possible is to venture a little way past them into the impossible.”

- Arthur C. Clarke (*Profiles of the Future*, 1962).

## ABSTRACT

The simplest possible beginning of abiogenesis has been a riddle from the last century, which is most successfully solved by the Lipid World hypothesis. However, origin of the next stages of evolution starting from lipids is still in dark. We propose a ‘Lipid-RNA World Scenario’ based on the assumption that modern stable lipid-RNA interactions are molecular fossils of an ancient stage of evolution when RNA World originated from Lipid World. In accordance to the faint young sun conditions, we present an ‘ice-covered hydrothermal vent’ model of Hadean Ocean. Our hypothetical model suggests that faint young sun condition probably provided susceptible physical conditions for an evolutionary route from Lipid-World to Protein-RNA World, through an intermediate Lipid-RNA World. Ancient ribozymes were ‘protected’ by lipids assuring their survival in prebiotic ocean. The origin of natural selection ensures transition of Lipid-RNA World to Protein-RNA World after the origin of ribosome. Assuming the modern peptidyl-transferase as the proto-ribosome structure, we have presented a hypothetical translation mechanism: proto-ribosome randomly polymerized amino acids being attached to the inner layer of a lipid-vesicle, using only physical energies available from our Hadean Ocean model. In accordance to the strategy of chemical evolution, we also have described the possible evolutionary behavior of this proto-ribosome, which explains the contemporary three-dimensional structure of 50S subunit and supports the predictions regarding the ancient regions of it. It also explains the origin of membrane-free ‘minimal ribosome’ in the time of LUCA.

## 1 INTRODUCTION

### 1.1 THE RNA WORLD HYPOTHESIS

The possible mechanism of early evolution and origin of life on earth has been a central problem for evolutionists for centuries. In modern form of life, it is observed that DNA functions as genetic material (for some virus, genetic material is RNA) and proteins are employed for structural and enzymatic purposes. RNA links between two separate biomolecules; messenger RNAs are transcribed from DNA, which are then translated to functional and structural proteins. Lipids and sugars are the fourth and fifth kind of biomolecules, which have a very important role in compartmentalization of cells/organelles and metabolism.

The modern form of life operates by means of sequence specificity, which is responsible for structural diversity of biomolecules. Various structures are assigned with various enzymatic functions and heredity. Thus, the sequence specificity seems to be the fundamental property of life. Due to the detection of enzymatic function (1-6) of RNAs and presence of them as genetic materials in some virus, and its self-replicating property (1, 2, 7), RNA was hypothesized to initiate the earliest form of life. This led to an increased interest in the hypothesis that an RNA World, a term introduced by Gilbert (8), preceded the DNA/RNA/Protein world (9). A detailed discussion of this hypothesis can be found in Supplementary Materials section 1.1.

In spite of its popularity, the RNA World hypothesis had to face strong arguments against its possible existence. Firstly, RNA oligomers itself seems unlikely to have emerged spontaneously in a primordial environment (9, 10). Secondly, according to geological researches (11-13), in a hot, acidic prebiotic environment RNA breaks upon hydrolysis (14). Fourthly, primitive earth was completely exposed to UV and gamma ray radiations (no ozone layer). The aromatic bases of RNA absorb strongly in the ultraviolet region, which makes them more susceptible to damage by background radiation (15). Fifthly, catalytic activities of RNAs are very limited (16) and are properties of relatively long RNA chains (17). We have presented a detailed critique on the RNA World in Supplementary Materials section 1.2.

### 1.2 ALTERNATIVE HYPOTHESIS: THE LIPID WORLD

On this context, some alternative hypotheses were proposed: for example ‘proteins first’ (18-23), virus world (24), lipid world (25) and prion-RNA world (15). However, except the lipid World hypothesis, all the others are inadequate in the sense that they claim the origin of functional proteins prior to the origin of a functional ribosome. This seems unlikely. We shall show later that the environmental conditions of Hadean Ocean (~4.4 Ga) only permit production of small non-ribosomal oligomers of amino acids. Only the Lipid World hypothesis probably provides the most intuitive description of the possibilities in Hadean Eon. Scenarios in which replication is a collective property of a loose molecular assembly could be as likely to reflect the earliest stages of biogenesis (26-29). The notion that lipids and other amphiphiles could serve as intermediates in prebiotic evolution has also been specifically elaborated (30-32). It has been proposed that lipid membranes may have a hereditary potential, as most membranes are generated from other membranes but not created de novo (33, 34). Lipid World hypothesis proposes the existence of self-aggregating lipid vesicles that have the heredity potential in terms of generating ‘daughter’ vesicles through fission and fusion interactions. A capacity for

---


\*To whom correspondence should be addressed.

high fidelity transfer of compositional information could be acquired gradually, after many growth and division cycles, leading to a process akin to self-reproduction (35). Catalytic activities of lipids have also been proposed (36). The possible mechanism of polynucleotide or polypeptide synthesis leading from lipid aggregation was also proposed, which will be further developed later.

## 2 THE BEGINNING OF ABIOGENESIS

### 2.1 THE HADEAN OCEAN ENVIRONMENT

The Study of zircons has found that liquid hydrosphere (and so the atmosphere) must have existed as long ago as 4.4 Gya, very soon after the formation of the Earth (37). Several hypothesized models show that the early evolution of life was linked to ancient hydrothermal activities in the Hadean and the early Archean (38-40). On the other hand, following the faint young sun paradox (41) a number of models were proposed proposing about a cool early earth (37, 42). A diplomatic solution for the two contradictory hypothesis was proposed by Bada *et al.* (43) in 1994 and Cleaves *et al.* in 1998 (44) that Hadean ocean were frozen only at the top, having approximately 300-1000 meter thick ice layer, which protected the early biomolecules from UV and gamma ray radiation. This model also provides a cold environmental situation where an RNA World can emerge and evolve.



**Figure-1:** The 'ice-covered hydrothermal vent' model of Hadean Ocean is schematically represented here. For a detailed discussion, see Supplementary Materials section 2.1.

Merging together the proposals of Bada *et al.* (43) and Cleaves *et al.* (44) that Hadean Ocean should have a frozen top due to faint young sun and should also have a Hydrothermal vent core, we propose an 'ice-covered hydrothermal vent' model of Hadean ocean. This is a simple modification of a common hydrothermal vent system with a frozen top. Submarine volcanoes establish a biologically rich alkaline (pH 9-10) high temperature (~350°C) sea floor, where the ocean top is frozen due to the faint young sun. Distant regions from volcanoes are acidic (pH 4-5). The hot water of sea floor will continuously experience an upstream motion towards the top, where they will lose temperature due to the ice layer and again come back to the sea floor. This will give rise to a continuous circular current of water from top to bottom. This continuous current will protect the organic materials from UV by taking

them deep into water repeatedly. In addition, the ice layer at the top surface of ocean provides an extra protection against UV. In fact, Jupiter's moon Europa is believed to have an oceanic environment similar to our model. A detailed discussion about this model is included in supplementary materials section 2.1.

### 2.2 ORIGIN OF NATURAL SELECTION

A general tendency of the biological evolution is a gradual complexation of the living systems, by "fine tuning" the enzymatic functions and heredity mechanisms through continuous cycles of self-reproduction at a large timescale. Assuming the abiogenesis started with a prebiotic Lipid World, the diverse biochemical reaction network (property of cellular life) was absent at that time, which is contemporary to the modern life. Nevertheless, when there was no cellular life, there should not be any evolutionary pressure on maintaining any biochemical function. Thus, in Lipid World or RNA World scenario, the natural selection of mutations could not be processed in the contemporary way. The strategy of natural selection should be dependent and consistent with the physical conditions of the habitat of the biomolecules (consistency with habitat is the basic feature of natural selection).

An ice covered hydrothermal vent system with water temperature ranging from ~350°C at the bottom and ~0°C at the top and a strong pH gradient requires structural consistency of the earliest biomolecules. Otherwise, they would have denatured soon in this hostile condition. In fact, the evolutionary strategy of biomolecules should be solely dependent on their structural stability. If a new monomer joins a polymer structure at a certain stage, its persistency will be determined by the structural stability it is providing to the polymer. If the monomer destabilizes the polymer, it should be dissociated soon. On the other hand, if it provides structural stability, its persistency will be high. Same kind of natural selection strategy should also be observed in case of isoenzymes: between two isoenzymes, the one with higher structural stability and higher catalytic activity should undergo natural selection. Thus, during abiogenesis, only those chemical reactions will undergo natural selection, whose components are structurally consistent and product yields are high. Eventually such reactions will organize and biochemical pathways will evolve. Therefore, we can identify a Darwinian picture in chemical evolution. We shall show later that this hypothetical strategy of natural selection fits perfectly with the evolutionary picture of ribosome, a remnant of RNA World. A detailed discussion on this topic is included in supplementary materials section 2.2.

## 3 CONSTRUCTING THE IDEA OF A LIPID-RNA WORLD

### 3.1 LIPIDS IN PREBIOTIC WORLD

Bangham *et al.* (45) first reported that phospholipids have the capacity to self-assemble. It was found that a variety of single-chain amphiphiles (medium- and long-chain monocarboxylic acids, alcohols, amines, alkyl phosphates, and alkyl sulfates) could self-assemble into bilayers under certain conditions. The minimal chain length for the microscopic appearance of closed vesicles at physiological temperatures was observed to be 10 carbons. Researches by

Ourisson and Nakatani (31) and others (46, 47) have shown that isopentenol and dipolyprenol phosphates, either individually or as mixtures, are able to produce stable bilayer vesicles and membranes. Some other interesting studies have shown that when fatty acid micelles are added to a buffered solution containing a dispersion of particles (clay mineral), vesicles are produced at an accelerated rate, the particles become coated with amphiphiles, and particles are often encapsulated in the vesicles (45, 48). A more recent study (49) experimentally demonstrates that a microcapillary thermal diffusion column can concentrate dilute solutions of nucleotides, oligonucleotides, and fatty acids. Upon concentration, the self-assembly of large vesicles containing encapsulated DNA occurs in regions where the critical aggregate concentration of the fatty acid is exceeded.

Molecular fossils have become a matter of serious scientific attention since last decade for Astrobiologists. The common presence of terpenoid and hopanoid derivatives as molecular fossils (31) show direct evidence that they were present in prebiotic world and were utilized by microorganisms. It is well known that such compounds, either individually or in mixture, are able to produce bilayers (46, 47). Deamer and his coworkers (50) described experiments in which they extracted samples of the Murchison meteorite with organic solvents, separated the mixture of extracted organic components, and then examined the structures formed mixing the separated components with water. Many different structures were observed, most interestingly the membranous vesicles obtained with some but not all of the separated components. In a few cases, vesicles were observed that seemed to consist of an interior compartment surrounded by a double membrane; fluorescent molecules could be trapped in the interior of the structure. Dworkin and coworkers have demonstrated very similar results with amphiphilic material obtained in simulations of organic synthesis in interstellar ices subjected to UV irradiation (51). Self-assembling lipids can be synthesized in simulated interstellar/precometary ices, as described by Deamer *et al.* (50). These studies suggest that the prebiotic formation of vesicles (and even vesicle within a vesicle) was possible.

### 3.2 VESICLE ENCAPSULATION AND ITS BIOLOGICAL ADVANTAGES IN PREBIOTIC WORLD

The ice covered hydrothermal vent model enables us to observe the Hadean Ocean as a large thermal column with circulating water. The biologically rich environment and pH gradient is a susceptible condition for formation of a large number of lipid vesicles. In fact, our ocean model represents a situation analogous to the experimental conditions described by Hargreaves *et al.* (48) and Budin *et al.* (49). Presence of montmorillonite in prebiotic ocean (10) would enhance the lipid aggregation rate, generating an extensive amount of vesicles encapsulating montmorillonite and other organic materials inside. Segré *et al.* (25) in their Lipid-World hypothesis described such an encapsulation event. The biological advantages of the prebiotic biomolecules of being vesicle encapsulated are pointed out in the following.

**3.2.1 Local phases of biologically rich environment:** The vesicle encapsulation of biological materials generates local phases of biologically rich environment (which means high concentration of organic materials inside). Earlier works (52, 53) calculated the

concentration of organic compounds in Archean seawater and ended up with a conclusion that they were not significantly different from modern seawater concentrations. Therefore, in the acidic primitive ocean, the concentrations of organic materials were potentially too low for conducting covalent chemical reactions (25). On this aspect, Segré *et al.* (25) described one important advantage of vesicle encapsulation that it generates local phases of biologically rich environment (45, 48, 49). Such vesicle-encapsulated phases are more capable for biochemical reactions forming covalent bonds.

**3.2.2 Polymerization of nucleic acid and amino acids inside vesicles:** The formation of vesicles in a hydrothermal column causes encapsulation of montmorillonite, amino- and nucleic acids (see Supplementary Materials section 1.2.2-1.2.3). The montmorillonite is known to catalyze nucleic acid polymerization (54, 55). Tsukahara *et al.* in 2002 (56) experimentally showed that amino acids are polymerized on or inside lipid vesicles if they are subjected to the circulating water current in a hydrothermal vent with top and bottom temperatures being 0° and 180°C respectively. Thus, the hydrothermal vent model enables the Lipid World to synthesize nucleic acid and peptide oligomers. We shall show later that this lipid-mediated oligopeptide formation had a very important role in proto-ribosome function.

**3.2.3 Catalytic activity of Lipids:** Lipid assemblies are known to have some catalytic activity. They can enhance the rates of some chemical reactions (33, 57). Lipid vesicles potentially enhance rates of many biochemical reactions (58, 59). This rate enhancement can even lead to autocatalytic growth of such vesicles/assemblies (60). These catalytic functions were later taken over by ribozymes when an RNA World established.

**3.2.4 Vesicle encapsulation protects from UV:** Vesicle encapsulation can protect the fragile prebiotic RNA segments from UV (a strong argument against RNA World hypothesis) and other radioactive radiations. Once RNA enters inside a vesicle (mechanism will be described later), it is somewhat protected from the cosmic radiations of early earth (UV and Gamma), since lipids absorb such radiations (61, 62). Lipid peroxidation due to UV radiation might be delayed in our ‘ice-covered hydrothermal vent’ model of Hadean Ocean. The thin ice covering at the top of ocean provides initial protection. The continuous circular flow of water (along with organic materials) from top to bottom can protect the lipid vesicle from peroxidation and provide a long lifetime.

**3.2.5 Lipids as stabilizing agents:** Lipids can even work as stabilizing agents; proteins and RNAs are commonly expressed on membranes and liposomes. Membrane interactions with DNA/RNA and stability analysis of such complexes have been a matter of good research interest because of the use of liposomes as vectors of gene delivery (63, 64). The history of stability analysis of DNA-lipid complexes is quite old (65-71). Examples of lipid-RNA binding are not that diverse as DNA (72-76). However, Tajmir-Riahi *et al.* (77) have reported that in case of lipid-tRNA binding, RNA have a higher binding constant for membrane (i.e. more stable association) than those of the lipid-DNA complexes (78). This indefinitely states that even in a prebiotic world, there was a high possibility that the structural stability to the ancient ribozymes were provided by vesicle membranes they were encapsulated.

### 3.3 MODERN LIPID-RNA INTERACTIONS

RNA can bind to both interior (73, 79) and exterior (78) of liposomes. Exterior RNA binding with liposome membrane is linear over 1000-fold range of RNA concentrations (73). RNAs that can bind, change the permeability, and disrupt phospholipid bilayers have been isolated using selection–amplification (73, 74). The oligomerizing heterotrimer RNA(9:9:10)<sub>n</sub> (Fig. 1, ref. 73) was one particular example, found to bind stably to the exposed face of a phosphatidylcholine vesicle. Many RNA complexes with similar activities were isolated, with no sequences in common. It is observed that many RNA structures exist that bind phospholipid membranes (80). Membrane affinity is, in this sense, a simple molecular property, so it is likely that mutation would continuously produce such RNAs (80). This argument could be a rationale of the origin of ancient membrane-bound RNAs in prebiotic earth. Moreover, it has also been shown that RNAs can bind to lipid vesicles and alter the membrane permeability (both positive and negative regulation, ref. 74). Such membrane permeability alteration can take place in primitive vesicles also, allowing inflow of small charged biological molecules and metal ions into the vesicle from water body, thus creating an ion gradient across the membrane.

RNAs normally bind to the polar side of the membrane, probably through Hydrogen bonds and van der Waals contacts. RNA 9, RNA 10, RNA 67-2, RNA 10<sub>TRP</sub>, RNA 80N, RNA 10Arg(D)5, and RNA Trp70-93 are established membrane-interacting ribonucleotide polymers of ~100-residue length (72). Janas *et al.* (72) have shown that a greater fraction of RNA structures bind bilayers as phospholipid order increases: fluid<raft<ripple<gel. We shall show that the proto ribosome once polymerized amino acids based on this molecular property. RNA binding to lipid ripple-gel phase (at the phase transition temperature) unexpectedly shows no significant RNA-structure specificity. On the other hand, a state of intermediate order (the liquid-ordered or rafted state, believed to occur in biological membranes) shows RNA-structure dependent affinity. To account for this, Khvorova *et al.* in 1999 (74) and Janas *et al.* in 2006 (72) predicted secondary structures of the membrane-interacting RNA sequences. All the predicted secondary structures contained one or more helical segments and, in a few cases, a centroid region (Table-1, ref. 68 and Figure-4, ref. 74). In addition, it is obvious from the predicted secondary structures that the three-dimensional structure of these RNA segments must be planar, which can “fit” on the membrane surface and cause binding. On this context, we argue that any ancient ribozyme having a planar three-dimensional structure and having a centroid and/or at least one linear helical secondary structural segment could possibly have a membrane-bound origin.

In the next sections, we shall show that predicted proto-ribosome structure fits excellently with these structural requirements. In fact, this discussion, in accordance with our hypothesis, can lead to an intuitive argument that modern RNA-lipid complexes are molecular fossils of an ancient stage of evolution when lipids provided RNAs structural stability essential to survive in prebiotic environment. In the hot primitive ocean, it is quite intuitive to imagine the mutual stability of ribozymes interacting with lipid vesicles. Such a stabilized assembly should be more susceptible for a directional evolution compared to independent presence of RNAs and lipids,

in a Darwinian picture. In addition, such assemblies can result in the ribozyme take over of lipid-catalytic activities and can initiate the transition of Lipid World to a Lipid-RNA World.

### 3.4 EVOLUTIONARY ROUTE TOWARDS A LIPID-RNA WORLD

The transition of Lipid World to the next stages of evolution should have occurred by means of Lipid-mediated polymerization of biomolecules. Segré *et al.* (25) provided a hypothesis about the possible mechanism of polypeptide and polynucleotide synthesis from lipid vesicles/membranes. According to this hypothesis, headgroups of ancient lipids likely had a large chemical diversity like today. Even amino acids and nucleotides might appear as headgroups. As these lipids aggregate and form vesicle and membranes, the headgroups can oligomerize on the planes of membrane. Such oligomerized headgroups are more akin to the contemporary biopolymers. They may have enhanced the catalytic activity within the vesicles (leading to vesicle growth) and templating capacities within the assemblies. In future, some of them could detach from membrane and become soluble. However, there are possible alternative mechanisms of polymerization of nucleotides. Hanczyc *et al.* in 2003 (81) reported that montmorillonite catalyze the formation of closed vesicles from micelles composed of simple aliphatic carboxylic acids and that particles of the clay become encapsulated within the vesicles. Since montmorillonite are excellent catalysts for the oligomerization of a number of activated nucleotides, this might point to a route to a nucleic acid synthesizing system enclosed within a vesicle (10). We have already discussed that the thermal diffusion column-like behavior of Hadean ocean might provide a susceptible environment for encapsulation of both montmorillonite and organic materials.

The importance of montmorillonite in RNA structures was discussed in detail by Pyle in 2002 (82). Steitz and coworkers in 2004 (83) reported the abundance and structural significance of cations in stabilizing 23S rRNA structure. They identified 116 Mg<sup>2+</sup> ions and 88 monovalent cations (K<sup>+</sup>, Na<sup>+</sup>) bound principally to 23S rRNA. Both divalent and monovalent cations stabilize the tertiary structure of 23S rRNA by mediating interactions between its structural domains. Bound metal ions are particularly abundant in the region surrounding the peptidyl transferase center (PTC), where stabilizing ribosomal proteins are notably absent. They proposed that this might point to the importance of metal ions for the stabilization of specific RNA structures in the evolutionary period prior to the appearance of proteins. In fact, Fox presented a number of logical arguments (84) to establish that PTC originated in a protein-free world. After vesicular intake of RNAs attached with montmorillonite occurred in prebiotic environment, ribonucleotides were further polymerized within vesicles, with polymerization rate and stabilization enhanced by both metallic ions and lipids. In a high temperature environment, stabilization provided by lipid was necessary, but might not be sufficient for ribozymes to hold their three-dimensional structure. The metal ions not only aided the polymerization of ribonucleotides, but also provided structural stability to maintain their three-dimensional structures in a high temperature environment.

## 4 THE ORIGIN OF RIBOSOME

So far, we have discussed in detail that prebiotic polymerization and stabilization of RNAs by lipids and metallic cations are possible. As we attempt to draw the hypothetical picture of events in a Lipid-RNA World, we have to keep accordance of it with two essential requirements: one, catalytic activity initially mediated by lipids must be taken over by ribozymes with course of time; and two, ribozyme actions should lead to amino acid polymerization, so that with course of time, a protein-RNA World originates and proteins now take over most of the ribozyme functions. Ribozyme take over of enzymatic functions from lipids is obvious from the picture we have drawn so far. As the RNAs are encapsulated, polymerized and stabilized within vesicles, they can show catalytic activity, which are far stronger than that of lipid-catalysis. However, a high rate of amino acid polymerization seems to be impossible without ribosome. Thus, we need to understand the origin and evolution of ribosome in a Lipid-RNA World. A detailed discussion about the origin of ribosome in accordance to several prior hypothetical pictures is presented in the Supplementary Materials section 3. In addition, we have presented there a critique on the previous predictions about the nature of translation machinery in a prebiotic environment. A conclusion of that discussion is demonstrated in the following.


### 4.1 MODELLING A PROTO-RIBOSOME IN LIPID-RNA WORLD

In Supplementary Materials section 3, we have presented a detailed discussion on functional importance of ribosomal proteins and the predictions of proto-ribosome structure. The concluding remark is PTC is the oldest part of 50S subunit. This RNA segment is self-folding and its stabilization strategy clearly indicates to its probable origin in a protein-free world. In accordance with these arguments, we suggest that the PTC RNA-segment evolved in the Lipid-RNA World. Random polymerization of nucleotides within a large vesicle pool in Hadean Ocean can bear such an ancient structure. In addition, the secondary and tertiary structure of the PTC segment indicates a possible membrane bound-origin of it. It contains two helical and one centroid region in secondary structure and two parallel helical segments in tertiary structure. We can recall that these are the structural requirements necessary for RNAs to interact with membranes. The two parallel helices of PTC define a convex bottom surface, which, likewise the hypothesis of Smith *et al.* (85) seems to be attached to a concave lipid membrane (we do not support their prediction of membrane interacting surface: Supplementary Materials section 3.5). In fact, we propose a hypothesis that the convex bottom surface of PTC is the molecular fossil of an ancient stage of evolution when this surface was attached to the inner surface of a lipid-vesicle (Figure-3, D). Available contemporary PTC structure of various species ensures that this structure is universally conserved. Furthermore, *Thermus thermophilus* PTC structures shows that the stabilization strategy of PTC (just by metallic cations) can ensure its structural constancy even at temperatures close to  $\sim 100^\circ\text{C}$ . If this structure is further stabilized by interacting with the polar headgroups of lipid vesicle, we can expect that it would survive at even higher temperatures. On the con-


text of a Hadean Ocean environment, without such a structural consistency a ribozyme could not undergo natural selection under a Darwinian picture.

### 4.2 MINIMAL REQUIREMENTS FOR RIBOSOMAL PROTEIN SYNTHESIS

Since this proto-ribosome alone cannot hold tRNAs, it is expected that early proteins were synthesized randomly, without the sequence specificity defined by mRNA in contemporary translation mechanism. However, it is difficult to imagine how PTC can proceed to polymerization of amino acids being attached to vesicle. In the next section, we shall suggest a possible mechanism of protein synthesis by vesicle-bound proto-ribosome. We have presented a discussion about the contemporary protein synthesis mechanism and predictions about the ancient translation mechanism in Supplementary Materials (section 3.6, 3.7). Remark of our discussion entails the minimal requirements for amino acid polymerization by ribosome. The mechanism of modern protein synthesis enables us to speculate how the proto-ribosomes synthesized polypeptides in their lipid-bound state. Figure-2 (A), schematically represents the chemical reactions occurring in PTC when a peptide bond is formed between already synthesized amino acid chain and a new amino acid brought by aminoacylated tRNA (82).



**Figure-2:** Comparison between chemistry of contemporary ribosomal polypeptide synthesis and the hypothetical polypeptide synthesis mechanism in Lipid-RNA World. (A) During contemporary peptide bond formation, the alpha-amino group of aminoacyl-tRNA (shown as gray cartoon view, taken from 50S subunit structure PDB-code: 3KNI; ref. 89) in the A-site attacks the carbonyl carbon of P-site bound peptidyl tRNA. The tetrahedral transition state now decomposes into the reaction products. One deacylated tRNA at the P-site and peptidyl-tRNA carrying an additional amino acid (+1 aa) at the A-site are produced. (B) The possible peptide bond formation by proto-ribosome in Lipid-RNA World: lipids are presented as a combination of gray head and black tails. Lipids emerging out from membrane plane can deliver the amino acids to the PTC and can hold the upcoming amino acid by an ester bond. Thus, membrane lipids functionally mimic aminoacylated tRNAs. This image is generated adopting the concept of Figure-1, ref.-86.



**Figure-3:** The 23S rRNA structures describing the possibilities of membrane-bound origin. (A) The position of self-folding PTC site (defined by Smith *et al.* in 2008, ref. 85) in 50S subunit structure of *Haloarcula marismortui* is represented here. The PTC RNA segment is shown as orange cartoon view and rest of the 23S rRNA is presented as gray stick view. Proteins are represented as space-fill views (see ref. 85 for a detailed picture). The RNA segment in green cartoon representation is another self-folding RNA segment from 2670 to 2830. The arrow shows the direction of relative motion of amino acids with respect to PTC site (see section 4.2), as the polypeptide chain from one tRNA is transferred to the next aminoacylated tRNA when a new peptide bond forms. The direction of relative motion of tRNAs with respect to PTC site is just antiparallel to this. This image is produced by using 50S structure of PDB-id 1S72 (87). (B) The three-dimensional position of Domain-V (cyan, cartoon view) and Domain-IV (orange, cartoon view) in 23S rRNA structure of *Escherichia coli* is shown. Rest of the 23S rRNA is shown in gray stick view. The two helices of PTC and the hypothetical membrane-interacting surface are highlighted. Here we can see that both implicate a beehive-like growth on a vesicle internal surface. In fact, the position of Domain-IV exactly over Domain-V provides us a clear indication that this domain is newer compared to Domain-V. This image is produced using the 50S subunit structure having PDB-id 2AWB (88). (C) This image describes figure-B from its 30S interface side. The picture depicts that domain-IV and Domain-V jointly fill the membrane interacting plane and provides the “base” for membrane-free growth of ribosome in post- Lipid-RNA World. In other words, the lateral size of the 50S subunit was determined before LUCA. (D) The *Thermus thermophilus* PTC RNA segment is presented as orange cartoon view. The hypothetical membrane interacting surface is highlighted in the two images (right hand side image is 90° anticlockwise-rotated form of the left one, about z-axis). This image is produced using the 50S subunit structure having PDB-id 3KNI (89). Above images are produced using The PyMOL Molecular Graphics System (90) and POV-Ray program (91).

The formation of a peptide bond in PTC does not require additional energy, since energy is stored in an ester bond linking the peptidyl residue to the tRNA 3'-hydroxyl group. This ester bond is a simple single bond between the 5'-oxygen atom of tRNA to the carbonyl carbon of the amino acid. Creation of this bond is 'energized' by ATP hydrolysis that drives aminoacylation of a tRNA by the respective aminoacyl-tRNA synthetase (86). This molecular

mechanism clearly demonstrates the minimal requirements for protein synthesis. The chemical requirement is that, the carbonyl carbon of the upcoming amino acid is attached with an oxygen atom via an ester bond. In addition, there are physical requirements. The amino acids are to be placed properly in PTC so that the chemical reaction can occur. As the ribosome propagates in 5'→3' direction of mRNA, aminoacylated tRNAs bring the amino

acids to the PTC. There, the already synthesized polypeptide chain is transferred to the next tRNA and a peptide bond forms between the amino acid of the new tRNA and already synthesized polypeptide. This relative displacement of already synthesized polypeptide chain with respect to PTC is parallel to the direction of ribosome motion on mRNA. This relative motion is essential for formation of a peptide bond and is depicted in Figure-3 (A). Here we can identify another physical requirement for peptide bond formation by PTC: peptide bond between two amino acids will form only if the second amino acid is brought to the first following this relative motion. Thus, we can identify that one chemical and two physical requirements are the basis of ribosomal peptide bond formation. The other constituents of modern translation system (30S subunit, mRNA, tRNAs and ribosomal proteins) are employed to assure accuracy and sequence specificity of the resulting polypeptide chain. We can argue that the earliest translation mechanism was deprived of such complications; the proto-ribosome must have evolved accomplishing the three minimal requirements only.


The idea that protein synthesis by a proto-ribosome eventually made to evolve a Protein-RNA World from the preceding RNA World, leaves a problem for us. That is, according to this hypothesis, the protein synthesis mechanism has to be at least as old as 4.2 Gyr. No cellular life was present, which means there was no continuous source of chemical energy (ATP/GTP). Thus, the ancient protein synthesis mechanism has to be driven by physical energy sources. In modern translation mechanism, ATPs are used for aminoacylation of tRNA and GTPs are required for movement of tRNA through the A, P and E sites. Without any chemical energy source, and even without tRNAs, the PTC (attached to the inner surface of lipid vesicle) can polymerize amino acids if some other carrier molecule (less complicated than tRNAs) delivers the amino acids to the PTC fulfilling the physical and chemical requirements described above.

### 4.3 HYPOTHETICAL PROTEIN SYNTHESIS MECHANISM BY PROTO-RIBOSOME

We suggest this hypothesis that in Lipid-RNA World, these carrier molecules were vesicle lipids. Akin to modern liposomes, ancient vesicles are expected to contain a high concentration of amino acids, many of which can interact with polar head groups of lipids. We mentioned in section 3.2.2 that amino acids are oligomerized on or inside lipid vesicles subjected to circular water current within a hydrothermal vent (56). These small mono- and oligopeptides can bind to the headgroups of vesicle lipids. The chemistry of peptide bond formation requires that the lipid headgroup would hold the amino acid by an oxygen atom having an ester bond with the carbonyl carbon atom of the new amino acid (thus, functionally mimicking an aminoacylated tRNA). This requirement is very elementary with respect to the expected prebiotic diversity of lipid headgroups (25). This chemical mechanism is depicted in Figure-2 (B), which suggests chemically such a reaction is possible. Therefore, now we can investigate whether the lipids can satisfy the physical requirements of amino acid polymerization discussed before.

If such mono- or oligopeptides, attached to lipid headgroups emerge out of the vesicle membrane surface along with the at-

tached lipid (this is possible during rapid phase changes of membrane upon temperature fluctuations), then their continuous relative motion through the PTC site in the proper direction can result in polymerization of those amino acids. Nevertheless, this hypothesis leads to another problem for consideration. How could such relative motions be possible in Lipid-RNA World?



**Figure-4:** The hypothetical protein synthesis mechanism in Lipid-RNA World by proto-ribosome is presented. Proto-ribosome is represented as contemporary PTC structure of *Thermus thermophilus* (green cartoon view). The tertiary structure of PTC RNA resembles an ‘RNA fork,’ which continuously collects amino acids emerging out from the membrane plane during rapid phase changes of lipids upon temperature fluctuations and polymerizes them (image is not according to molecular scale). See text for a detailed description.

It has been reported that lipid vesicles undergo structural changes upon fluctuation of temperature, both experimentally (92, 93) and theoretically by Coarse-Grained MD simulation studies (94). Such studies clearly show that upon high temperature treatment, the vesicle shape is stretched, leading to budding initiation, and in low temperature, its shape shrinks and restores in spherical form. Therefore, in regions of rapid temperature fluctuation (common event in an ice-covered hydrothermal vent system), it is possible that ancient vesicles would undergo rapid structural changes (stretch and shrink). PTC being a large RNA segment (178 residues for *H. marismortui*), it should experience a continuous relative displacement (slide) with respect to the vesicle membrane surface during such rapid structural changes. The physical reasons of sliding are the fluctuations in membrane surface tension and pressure differences in the vesicle interior due to budding initiation. The binding efficiency of PTC with membrane decreases as lipids experience fluid state at high temperatures, and it can slide fast on the membrane surface. The reason follows from the work of Janas *et al.* (72), that a greater fraction of RNA structures binds bilayers as phospholipid order increases: fluid <raft <ripple <gel. Sliding will slow down when vesicles are back in low temperature and lipids gain gel state again. Thus, such a relative motion can cause a continuous flow of amino acids through the PTC site as temperature varies rapidly. PTC RNA tertiary structure resembles an ‘RNA fork,’ (Figure-4) which, during such relative displacements, continuously collects the amino acids from lipid molecules emerging out of the membrane plane and polymerizes them. The emergence of lipid molecule along with the attached amino acid is expected in such a rapid phase-changing lipid membrane. There is one distinguishable advantage of polymerizing amino acids in this mechanism: if at a certain stage of polymerization process, amino acid-holding lipid is unavailable at the neighborhood of PTC, then

PTC can drag the latest lipid (already synthesized polypeptide attached to its headgroup) through the fluid membrane until next lipid with an amino acid is available. Thus, this mechanism can synthesize long enough polypeptides.

This model (depicted in Figure-4) is acceptable in prebiotic scenario when mRNA, tRNAs and 30S subunit not likely evolved or assisted amino acid polymerization. This hypothetical mechanism satisfies all the physical and chemical ‘minimal requirements’ of ribosomal protein synthesis. Most importantly, this hypothetical model enables membrane-bound proto-ribosomes to polymerize amino acids without any chemical energy source. PTC polymerizes amino acids just using physical energies obtained from a hydrothermal vent system. Though this was not a true ‘translation process’; rather PTC was a random amino acid polymerizer. This picture is in agreement with the arguments of Noller (95) that the ancient translation mechanism did not evolve initially to make functional proteins, instead it produced random polypeptides. There are two experimental supports entailing the fact that our model of PTC displacement through vesicle membrane surface is possible. The lateral and vertical displacement of integral membrane proteins during temperature-mediated lipid phase changes has been confirmed by Armond and Staehelin (96). Another study has reported that lipid fluidity changes can cause vertical displacement of membrane proteins (97). Though these are results for membrane proteins, but such physical phenomena can cause to membrane bound RNAs also. In fact, the displacement of membrane-attached PTC segment is likely to be faster compared to integral proteins during lipid phase changes.

#### 4.4 BEEHIVE-LIKE GROWTH: ORIGIN OF MINIMAL RIBOSOME IN LUCA

Proto-ribosome probably originated by random polymerization of RNA segments inside lipid vesicles. Though it is very hard to understand how such specific sequences originated at that time since probability of formation of such a ~150-residue chain by means of random oligomerization of nucleotides is very small. However, once synthesized, it adopted a specific tertiary structure stabilized by metallic cations (since this is self-folding RNA) and rested upon the internal vesicle surface. There it could polymerize proteins in the previously discussed mechanism.

Ribozyme mediated RNA transcription probably was the basis of ribozyme replication in RNA World. The high sequence conservation of PTC (98) indicates the fact that modern ribosome of every species is the result of a divergent evolution starting from a unique ancestor. This means if we accept the idea that PTC was the true ancestor of ribosome, then we have to accept that the probability of synthesizing such a unique sequence (~150 residues long) just by random polymerization of nucleotides is extremely small. This entails the fact that if PTC RNA was the actual proto-ribosome, then it had to undergo repeated replication cycles. Production of a large amount of proto-ribosome (which is structurally consistent) by repeated cycles of replication causes a large amount of protein yield within vesicles. This can prepare the evolutionary route to a protein-RNA World. As we discussed in section 2.2, high structural consistency and high protein yield of the proto-ribosome enables it to undergo natural selection. Replication of proto-ribosome (by ribozymes, which were RNA dependent RNA polymerases) in

every cycle should be associated with joining or deletion of ribonucleotides with its ancient core. In a Darwinian picture, the residues providing high stability to the proto-ribosome upon addition will stay in the structure (be selected by nature to persist). On the other hand, residues providing less stability or disability to the polymer will soon be dissociated in the upcoming replication cycles. This can lead to a gradual growth of the proto-ribosome on the membrane, similar to the growth of a beehive on a wooden wall. This ‘beehive-like growth’ of proto-ribosome results in the origin of oldest domains of 50S subunit.

Such a growth is likely to be continued with the course of time; specific proteins will evolve that can bind to the ribosome properly and provide necessary stability. It is obvious that RNA binding affinity of proteins is higher than lipid binding affinity of RNA. In addition, with the growing size of ribosome, it would be difficult for a lipid bilayer to hold such a giant structure. For these two reasons, eventually, one time did come, when ribosome detached from the membrane and became a free-living ribonucleoprotein complex. Protein binding and metallic ions offered this free complex all the structural stability it required. Such an event definitely occurred before LUCA (99). Possible timing of LUCA is speculated to be about 3.8 Gyr. Interestingly, geological records confirm an Archean ocean of approximately 39°C surface temperature (100). That means ‘ice-covered hydrothermal vent’ model of oceans was no longer applicable and ribosome must have evolved contemporary translation mechanism. Our hypothetical translation model is no longer applicable in Archean ocean with almost uniform temperature. Thus, though ribosome originated as a membrane-bound organelle, the free cytosolic ribonucleoprotein complex structure is selected in evolution.

We also conducted a study on the possible ribosome structure, which could be detached from lipid vesicle. If ribosome had grown on a hypothetical vesicle surface, then it is expected that the oldest regions of ribosome will maintain a planar, or, rather, a convex shape on its membrane bound side. In addition, it is expected that when ribosome experienced a continuous growth bound to membrane, all the nucleotides added to the PTC, must have positioned to the opposite direction of the membrane bound side (beehive-like growth). This means, the oldest domain of 50S subunit (Domain-V) will be positioned exactly on the hypothetical membrane interacting surface, and the domain originated next to it (Domain-IV) will be positioned on Domain-V. In fact, Domain-V and Domain-IV, the two oldest domains of 23S ribosomal RNA, reported by Hury *et al.* (101), fulfill these two requirements. This is demonstrated in Figure-3, B (see figure legend). The positioning of Domain-IV just over Domain-V clearly indicates the fact that Domain-IV originated after Domain-V, by virtue of a beehive like growth of ribosome, on the internal surface of a lipid vesicle, starting from PTC. We can see another important behavior from three-dimensional placing of Domain-IV and Domain-V. These two domains jointly fill the membrane interacting plane and provide the ‘base’ for membrane-free growth of ribosome after detachment. In other words, the lateral size of the 50S subunit was determined before LUCA, i.e. before membrane detachment. Interestingly, this ancient 50S subunit was almost that of the minimal ribosome defined by Mears *et al.* (102).



Our picture of ribosome origin speculates that residues being closer to the modern PTC site of a ribosome are likely to have an older origin and persisted due to their structural importance. If this is true, then, we must observe a slower evolutionary rate in the core regions of 50S subunit (i.e. regions close to PTC) compared to distant residues or surface residues. Since structurally important residues evolve slowly. Such an evolutionary trend has been confirmed earlier (103, 104). The starting of ribosome evolution from such an ancestor core was also proposed by Bokov and Steinberg in 2009 (105).

## 5 FROM LIPID-WORLD TO PROTEIN-RNA WORLD

The “Primordial soup” theory (106-108) speculates about an ancient stage of evolution when organic materials were concentrated in prebiotic oceans (4.4 Gyr). Bernal coined the term *biopoiesis* for chemical evolution starting from such a soup, and suggested that there were a number of clearly defined “stages” that could be recognized in explaining the origin of life (108): first stage, when biological monomers originated, second stage, when monomers were polymerized and the third stage, when polymers evolved into proto-cells. If compositional complexity of a system is visualized as the diversity of polymer structures and possible biochemical reactions, then the whole process can be visualized as a gradual increment of complexity.

From such a ‘Primordial soup,’ self-aggregation of lipids can form early membranes and vesicles. Experimental works of Hill and coworkers (109) shows that sufficiently long negatively charged oligomers (e.g., RNA) adsorb almost irreversibly on anion-exchanging minerals. Since template-directed synthesis occurs without interference on many mineral surfaces (110), mineral particles might support replication until they became saturated with the descendents of a single ancestral RNA molecule. We have already discussed that montmorillonite within minerals can catalyze ribonucleotide polymerization. Moreover, when lipid vesicles are formed in a thermal gradient solution containing mineral clay, those clay particles are often encapsulated by vesicles (49). We propose this as a possible molecular process by which oligonucleotides were encapsulated inside lipid vesicles: being attached to clay particles in prebiotic oceans. Eventually, they might be incorporated on inner layer of vesicles, gaining extra stability by interacting with polar headgroups of lipid. Further polymerization of the RNA segments attached to metallic cations and lipids can originate a Lipid-RNA World. If the vesicles are exposed to an environment of fluctuating temperature, this will lead to rapid fission-fusion interaction (111, 112) between vesicles, which will eventually produce a vesicle pool where daughter vesicles are produced with high fidelity of compositional identity. This can be viewed as a somewhat complicated alternative of the ‘evolution like behavior’ described by Segré *et al.* (25) in their Lipid World hypothesis. Probably by this time, ribozymes that were capable of RNA-dependent RNA transcription evolved. This is essential for high fidelity heredity in a Lipid-RNA World. Nelson *et al.* (113) and Wochner *et al.* (114) described such ribozymes.

Thus, we have described the Lipid-RNA World as a large vesicle pool containing diverse types of ribozymes and the proto-

ribosome, where vesicles are undergoing rapid budding and fission-fusion interactions upon temperature fluctuation. However, vesicles subjected to hydrothermal vent are also capable of generating small oligopeptides on and inside them (56). Synthesis of random polypeptides in a large vesicle pool maintains the potential of synthesizing many functional proteins, which could have played important roles in early evolution. Protein-RNA complexes originated by the random interactions between protein and RNA segments. As we described before, such complexes being structurally more stable than RNA-only structures, were subjected to higher advantage in natural selection in a Darwinian picture. This was the beginning of ‘protein take over’ of the ribozyme functions.

Pool *et al.* in 1998 (115) presented a model for the protein take over of ribozyme functions. This was far developed by Cech in 2009 (116). According to this model, Primordial ribozymes were capable of purely RNA catalysis, but given random peptides in their environment, selection for fitness may have occurred at the level of the ribonucleoprotein (RNP). Formation of protein-RNA complexes was a more advanced stage in evolution. Peptides and polypeptides were synthesized that enhanced RNP stability and function. RNP enzymes have continued to evolve by adding (and subtracting) RNA elements that either contribute directly at the RNA level or bind to additional protein subunits. In most cases, catalysis has gradually been taken over by protein enzymes.

Some RNA viruses (poliovirus) are known to mediate vesicle budding of eukaryotic host (117). Though this process is associated with action of many proteins, but the RNA binding with membrane finally results in a replication complex which results in both vesicle synthesis and viral RNA replication (118). One theoretical study by Atilgan and Sun (119) shows a critical concentration of conical membrane proteins or proteins with nonzero spontaneous curvature can drive the formation of small vesicles. The driving force of vesicle budding stems from the preference of proteins to gather in regions of high curvature. A sufficiently high concentration of proteins therefore can influence the topology of the membrane. In the limit of mechanical equilibrium with no thermal fluctuations, it was found that depending on their shapes, pairs of proteins can repel each other with long range forces (120, 121). With membrane thermal fluctuations, a slight attraction is introduced, although the mechanical repulsion still dominates (122). Moreover, protein interactions depend on the curvature of the membrane (122). All these lead to the hypothesis that proteins can induce membrane shape changes, budding and even fission-fusions. We suggest that on the context of Lipid-RNA World to Protein-RNA World transition, these results entail that proteins could have played an important role in vesicle replication and establishment of a ‘self-replicating organism’ pool. Probably there was a transition phase in chemical evolution when vesicle fission-fusion interactions were turning to be more and more protein action-dependent and less temperature dependent.

As the vesicles within a vesicle pool undergo self-replication processes, compositional informations are transferred to the next progeny. If the compositional complexity is slowly increasing, then after a large number of cycles of self-replication, a vesicle pool with high fidelity heredity will be established. However, as proteins were synthesized by the lipids and proto-ribosome, rate of

complexity increment with time will be enhanced. Diversity of protein structures and functions might be the reason of it. Thus, as the Lipid-RNA World to Protein-RNA World transition occurs, high fidelity heredity is no longer possible. However, we can expect a balance between increasing complexity and high fidelity heredity in such transition phases. Proteins mediating vesicle budding and fission-fusion interactions might evolve as the cell-division related proteins in future and such vesicles can evolve as proto-cells. A schematic representation of lipid-RNA World scenario is schematically shown in Supplementary Materials Figure-2 (A-F).

## 6 SUMMARY

It is almost certain that sequence specificity of life cannot appear in prebiotic earth from inorganic materials (rejecting the idea about extra-terrestrial origin of life) just by sudden. The idea of RNA World strongly requires this sequence specificity. Hence, it is rational to imagine that chemical evolution had an even simpler beginning, which can eventually lead to sequence specificity. The Lipid World hypothesis deals with such a possibility. Assembly formation, catalytic activities and evolution-like behavior of lipids, along with the presence of lipid vesicles as molecular fossils strongly support this idea. RNA World seems to be the second stage of life when sequence specificity first ever coped with enzymatic function and heredity. It is an intuitive argument that proteins, constructed by twenty different types of components (amino acids) and having large structural variations are too complex to be the first enzymes during abiogenesis. On this context, ‘protein-first’ hypothesis does not seem to be an accurate description of prebiotic origin of life. On the other hand, RNA, in comparison to proteins has simpler monomers with less structural diversity that could originate well before proteins. Nevertheless, RNA is structurally fragile. Therefore, without the help of stabilizing agents it must face rapid degradation in hot acidic prebiotic oceans. Lipids and metallic ions could indefinitely serve this job as stabilizing agents. Hence, this logical picture enables us to redevelop the concepts of early evolution. In addition, we have tried to provide a molecular mechanism of protein synthesis in Lipid-RNA World, which would eventually take up the Lipid-RNA World to the next stage of chemical evolution: the Protein-RNA World. This model suggests that proto-ribosome could have polymerized amino acids without the need of any chemical energy source, just by utilizing physical forces generated by thermal fluctuations. It is possible that self-replicating vesicles of the prebiotic vesicle pool would grow with course of time and gain more complexity. Synthesis of random polypeptides in a large vesicle pool maintains the potential of synthesizing many functional proteins, which could have played important roles in early evolution. It can be speculated that with course of time such vesicles turned into proto-cells.

## ACKNOWLEDGEMENTS

## REFERENCES

1. Woese, C. (1967) *The Genetic Code, the Molecular Basis for Genetic Expression*. New York: Harper and Row.
2. Orgel, L.E. (1968). Evolution of the genetic apparatus. *J. Mol. Biol.*, 38:381–393.
3. Orgel, L. and Sulston, J. 1971. Polynucleotide replication and the origin of life. In *Prebiotic and Biochemical Evolution*, pp. 89–94. Kimball, A., and Oro, J., Eds., Amsterdam: North-Holland Publishing Company.
4. White, H.B., 3rd. (1976) Coenzymes as fossils of an earlier metabolic state. *J Mol Evol* 7:101–104.
5. Kruger, K., Grabowski, P.J., Zaug, A.J., Sands, J., Gottschling, D.E., and Cech, T.R. (1982) Self-splicing RNA: autoexcision and autocyclization of the ribosomal RNA intervening sequence of *Tetrahymena*. *Cell* 31:147–157.
6. Guerrier-Takada, C., Gardiner, K., Marsh, T., Pace, N., and Altman, S. (1983) The RNA moiety of ribonuclease P is the catalytic subunit of the enzyme. *Cell* 35:849–857.
7. Crick, F.H.C. (1968) The origin of the genetic code. *J. Mol. Biol.*, 38:367–379.
8. Gilbert, W. (1986) The RNA World. *Nature* 319:618.
9. Shapiro, R. (1984), *The Improbability of Prebiotic Nucleic Acid Synthesis*, *Origins Life Evol. Biosphere* 14, 565–570.
10. Orgel, L. E., (2004). *Prebiotic Chemistry and the Origin of the RNA World*, *Critical Reviews in Biochemistry and Molecular Biology*, 39:99–123, 2004.
11. Gesteland, R.F., Cech, T.R., and Atkins, J.F. (1999) *The RNA World*, 2nd ed. Cold Springs Harbor, NY: Cold Springs Harbor Press.
12. Knauth, L.P., Lowe, D.R. (1978). Oxygen isotope geochemistry of cherts from the Onverwacht Group (3.4 billion years), Transvaal Group, South Africa, with implications for secular variations in the isotopic composition of cherts. *J. Geol.*, 41, 209–222.
13. Tajika, E., Matsui, T. (1993). Degassing history and carbon cycle of the Earth: From an impact-induced steamatmosphere to the present atmosphere. *Lithos*, 30, 267–280.
14. de Ronde, C.E.J., Channer, D.M.D.R., Faure, K., Bray, C. J., Spooner, T. C. (1997). Fluid chemistry of Archean seafloor hydrothermal vents: Implications for the composition of circa 3.2 Ga seawater. *Geochim. Cosmochim. Acta*, 61, 4025–4042.
15. Singh, P. P., Banerji, A., (2011). Case for an RNA-prion world: a hypothesis based on conformational diversity, *J Biol Phys* (2011) 37:185–188.
16. Lindahl, T. (1993). Instability and decay of the primary structure of DNA. *Nature* 362(6422), 709–715.
17. Benner, S. A., Ellington, A. D., Tauer, A. (1989). Modern metabolism as a palimpsest of the RNA world. *Proc Natl Acad Sci USA* 1989, 86:7054–7058.
18. Bernhardt, H. S. (2012). The RNA world hypothesis: the worst theory of the early evolution of life (except for all the others), *Biol Direct*, 7(1):23.
19. Egel, R. (2009). Peptide-dominated membranes preceding the genetic takeover by RNA: latest thinking on a classic controversy. *Bioessays* 2009, 31:1100–1109.
20. Kurland, C. G. (2010). The RNA dreamtime: modern cells feature proteins that might have supported a prebiotic polypeptide world but nothing indicates that RNA world ever was. *Bioessays* 2010, 32:866–871.
21. Caetano-Anollés, D., Kim, K. M., Mittenthal, J. E., Caetano-Anollés, G. (2011). Proteome evolution and the metabolic origins of translation and cellular life. *J Mol Evol* 2011, 72:14–33.
22. Caetano-Anollés, G., Kim, K. M., Caetano-Anollés, D. (2012). The phylogenomic roots of modern biochemistry: origins of proteins, cofactors and protein biosynthesis. *J Mol Evol* 2012, 74:1–34.
23. Harish, A., Caetano-Anollés, G. (2012). Ribosomal history reveals origins of modern protein synthesis. *PLoS One* 2012, 7:e32776.
24. Koonin, E. V., Senkevich, T. G., and Dolja, V. V. (2006). The ancient Virus World and evolution of cells, *Biology Direct* 2006, 1:29.
25. Segré, D., Ben-Eli, D., Deamer, D. W., and Lancet, D. (2001). *The Lipid World, Origins of Life and Evolution of the Biosphere* 31: 119–145.
26. Dyson, F. (1985). *Origins of Life*, Cambridge, Cambridge University Press.
27. Farmer, J. D., Kauffman, S. A. and Packard, N. H. (1986). Autocatalytic Replication of Polymers, *Physica* 22D, 50–67.
28. Morowitz, H. J. (1992). *Beginnings of Cellular Life*, London, Yale University Press.

29. Kauffman, S. A. (1993). *The Origins of Order – Self-Organization and Selection in Evolution*, Oxford, Oxford University Press.
30. Segré, D., Lancet, D., Kedem, O. and Pilpel, Y. (1998a). Graded Autocatalysis Replication Domain(GARD): Kinetic Analysis of Self-Replication in Mutually Catalytic Sets, *Origins Life Evol. Biosphere* 28, 501–514.
31. Ourisson, G. and Nakatani, Y. (1994). The Terpenoid Theory of the Origin of Cellular Life: The Evolution of Terpenoids to Cholesterol, *Chemistry & Biology* 1, 11–23.
32. Norris, V. and Raine, D. J. (1998). A Fission-Fusion Origin for Life, *Origins Life Evol. Biosphere* 28,523–537.
33. Luisi, P. L., Walde, P. and Oberholzer, T. (1999). Lipid Vesicles as Possible Intermediates in the Origin of Life, *Current Opinions in Colloid & Interface Science* 4, 33–39.
34. Cavalier-Smith, T. (1995). Biodiversity and Evolution, M. Kato and Y. Doi. Tokyo, National Science Museum Foundation.
35. Szathmáry, E.: 1999, *Chemes, Genes, Memes: A Revised Classification of Replicators*, *Lectures on Mathematics in the Life Sciences*, 26, 1–10.
36. Segré, D. and Lancet, D. (1998b). Mutually Catalytic Amphiphiles: Simulated Chemical Evolution and Implications to Exobiology, in J. Chela-Flores and F. Raulin, *Exobiology: Matter, Energy and Information in the Origin and Evolution of Life in the Universe*, Trieste, Italy, Kluwer, pp. 123–131.
37. Valley, J. W., Peck, W. H., King, E. M., Wilde, S. A. (2002). A cool early Earth, *Geology*, v. 30, no. 4, p. 351–354
38. Russell, M.J., Arndt, N.T., (2005). Geodynamic and metabolic cycles in the Hadean. *Biogeosciences* 2, 97–111.
39. Takai, K., Nakamura, K., Suzuki, K., Inagaki, F., Neelson, K.H., Kumagai, H., (2006). Ultramafics–Hydrothermalism–Hydrogenesis–HyperSLiME (UltraH3) linkage: a key insight into early microbial ecosystem in the Archean deep-sea hydrothermal systems. *Paleontological Research* 10, 269–282.
40. Russell, M.J., (2007). The alkaline solution to the emergence of life: energy, entropy and early evolution. *Acta Biotheoretica* 55, 133–179.
41. Sagan, C., Mullen, G. (1972). Earth and Mars: Evolution of Atmospheres and Surface Temperatures, *Science, New Series*, Vol. 177, No. 4043, pp. 52–56.
42. Valley, J. W. (2005). A cool early Earth? *Sci. American*,293(4):58-65.
43. Bada, J. L., Bigham, C., Miller, S. L. (1994). Impact melting of frozen oceans on the early Earth: implications for the origin of life. *Proc Natl Acad Sci U S A*, 91:1248-50.
44. Cleaves, H. J., Miller, S. L. (1998). Oceanic protection of prebiotic organic compounds from UV radiation, *Proc Natl Acad Sci U S A*. 1998 Jun 23;95(13):7260-3.
45. Bangham, A. D., Standish, M. M. and Watkins, J. C. (1965). Diffusion of Univalent Ions Across the Lamellae of Swollen Phospholipids, *J. Mol. Biol.* 13, 238.
46. Hargreaves, W. R. and Deamer, D. W. (1978). Liposomes from Ionic, Single-Chain Amphiphiles, *Biochemistry* 17, 3759–3768.
47. Walde, P., Wick, R., Fresta, M., Mangone, A. and Luisi, P. L. (1994). Autopoietic Self-Reproduction of Fatty Acid Vesicles, *J. Am. Chem. Soc.* 116, 11649–11654.
48. Hargreaves, W. R., Mulvihill, S. and Deamer, D. W. (1977). Synthesis of Phospholipids and Membranes in Prebiotic Conditions, *Nature* 266, 78–80.
49. Budin, I., Bruckner, R. J., and Szostak, J.W. (2009). Formation of Protocell-like Vesicles in a Thermal Diffusion Column, *J. Am. Chem. Soc.*, 2009, 131 (28), pp 9628–9629
50. Deamer, D.W. and Pashley, R.M. (1989). Amphiphilic components of the Murchison carbonaceous chondrite: surface properties and membrane formation. *Orig Life Evol Biosph* 19:21–38.
51. Dworkin, J., Deamer, D., Sandford, S., and Allamandola, L. (2001). Selfassembling amphiphilic molecules: synthesis in simulated interstellar/precometary ices. *Proc Natl Acad Sci USA* 98:815–819.
52. Lasaga, A. C., Holland, H. D. and Dwyer, M. J.: 1971, *Primerial Oil Slick*, *Science* 174, 53–55.
53. Channer, D.M.D.R., de Ronde, C.E.J., Spooner, E.T.C. (1997). The Cl-Br-I composition of ~3.23 Ga modified seawater: implications for the geological evolution of ocean halide chemistry. *Earth Planet. Sci. Lett.*, 150, 325–335.
54. Lohrmann, R. and Orgel, L.E. (1980). Efficient catalysis of polycytidylic acid-directed oligoguanylate formation by Pb<sup>2+</sup>. *J Mol Biol* 142:555–567.
55. Bridson, P.K. and Orgel, L.E. (1980). Catalysis of accurate poly(C)- directed synthesis of 3'-5'-linked oligoguanylates by Zn<sup>2+</sup>. *J Mol Biol* 144:567–577.
56. Tsukahara, H., Imai, E., Honda, H., Hatori, K., Matsuno, K. (2002). Prebiotic oligomerization on or inside lipid vesicles in hydrothermal environments, *Orig Life Evol Biosph.* 2002 Feb;32(1):13-21.
57. Blocher, M., Liu, D.,Walde, P. and Luisi, P. L. (1999). Liposome-Assisted Selective Polycondensation of A-Amino Acids and Peptides, ISSOL'99, San Diego, CA, USA, Abstract c1.7.
58. Cuccovia, I. M., Quina, F. H. and Chaimovich, H. (1982) A Remarkable Enhancement of the Rate of Ester Thiolysis by Synthetic Amphiphile Vesicles, *Tetrahedron* 38(7), 917–920.
59. Kust, P. R. and Rathman, J. F. (1995). Synthesis of Surfactants by Micellar Autocatalysis: N,Ndimethyldodecylamine N-oxide, *Langmuir* 11, 3007–3012.
60. Bachmann, P. A., Luisi, P. L. and Lang, J. (1992). Autocatalytic Self-Replicating Micelles as Models for Prebiotic Structures, *Nature* 357, 57–59.
61. De, A. K., Ghosh, J. J., Mandal, T. K. (2006). Ultraviolet radiation-induced lipid peroxidation in liposomal membrane: Modification by capsaicin, *Phytotherapy Research* Volume 7, Issue 1, pages 87–89.
62. Bose, B., Agarwal, S., Chatterjee, S. N. (1990). Membrane lipid peroxidation by UV-A: mechanism and implications, *Biotechnol Appl Biochem.* 1990 Oct;12(5):557-61.
63. Felgner, P. L., Gadek, T. R., Holm, M., Roman, R., Chan, H. W., Wenz, M., Northrop, J. P., Ringold, G. M., and Danielsen, M. (1987). Lipofectin: a highly efficient, lipid-mediated DNA transfection procedure. *Proc. Natl. Acad. Sci. USA.* 84:7413–7417.
64. Felgner, P. L., and Ringold, G. M. (1989). Cationic liposome mediated transfection. *Nature.* 331:461– 462.
65. Gershon, H., Ghirlando, R., Guttman, S. B., and Minsky, A. (1993). Mode of formation and structural features of DNA-cationic liposome complexes used for transfection. *Biochemistry.* 32:7143–7151.
66. Gustafsson, J., Arvidson, G., Karlsson, G., and Almgren, M. (1995). Complexes between cationic liposomes and DNA visualized by cryo-TEM. *Biophys. Biochim. Acta.* 1235:305–312.
67. Lasic, D. D., Strey, H., Stuart, M. C. A., Podgornik, R., and Frederik, P. M. (1997). The structure of DNA-liposome complexes. *J. Am. Chem. Soc.* 119:832– 833.
68. Zuidam, N. J., and Barenholz, Y. (1997). Electrostatic parameters of cationic liposomes commonly used for gene delivery as determined by 4-heptadecyl- 7-hydroxycoumarin. *Biophys. Biochim. Acta.* 1329:211–222.
69. Hui, S. W., Langner, M., Zhao, Y-L., Patrick, R., Hurley, E., and Chan, K. (1996). The role of helper lipids in cationic liposome-mediated gene transfer. *Biophys. J.* 71:590 –599.
70. Mok, K. W. C., and Cullis, P. R. (1997). Structural and fusogenic properties of cationic liposomes in the presence of plasmid DNA. *Biophys. J.* 73:2534 –2545.
71. Harries, D., May, S., Gelbart, W. M., and Ben-Shaul, A. (1998). Structure, Stability, and Thermodynamics of Lamellar DNA-Lipid Complexes, *Biophysical Journal*, Vol.75, pp. 159–173
72. Janas, T., Janas, T. and Yarus, M. (2006) Specific RNA binding to ordered phospholipids bilayers. *Nucleic Acids Res.*, 34, 2128–2136.
73. Vlassov, A., Khvorova, A. and Yarus, M. (2001) Binding and disruption of phospholipid bilayers by supramolecular RNA complexes. *Proc. Natl Acad. Sci. USA*, 98, 7706–7711.
74. Khvorova, A., Kwak, Y.G., Tamkun, M., Majerfeld, I. and Yarus, M. (1999) RNAs that bind and change the permeability of phospholipid membranes. *Proc. Natl Acad. Sci. USA*, 96, 10649–10654.
75. Weisman, S., Hirsch-Lerner, D., Barenholz, Y. and Talmon, Y. (2004) Nanostructure of cationic lipid-oligonucleotide complexes. *Biophys. J.*, 87, 909–914.
76. Zhdanov, R. I., Kuvichkin, V. V., Shmyrina, A. S., Jdanov, A. R. and Tverdislov, V. A. (2002) Role of lipid membrane-nucleic acid interactions, DNA-membrane contact and metal(II) cations in origination of initial cells and in evolution of prokaryotes to eukaryotes. *Bioelectrochemistry*, 58, 41–46.
77. Marty, R., N'soukpoe'-Kossi, C. N., Charbonneau, D. M., Kreplak, L., and Tajmir-Riahi, H. A. (2009). Structural characterization of cationic lipid-tRNA complexes, *Nucleic Acids Research*, 2009, 1–11.
78. Marty, R., N'soukpoe'-Kossi, C.N., Charbonneau, D., Weinert, C.M., Kreplak, L. and Tajmir-Riahi, H.A. (2009) Structural analysis of DNA complexation with cationic lipids. *Nucleic Acids Res.*, 37, 749–757.

79. Suga, K., Umakoshi, H., Tomita, H., Tanabe, T., Shimanouchi, T. and Kuboi, R. (2010) Liposomes destabilize tRNA during heat stress. *Biotechnol. J.*, 5, 526–529.
80. Janas, T., and Yarus, M. (2003). Visualization of membrane RNAs, *RNA* (2003), 9:1353–1361.
81. Hanczyc, M.M., Fujikawa, S.M., and Szostak, J.W. (2003). Experimental models of primitive cellular compartments: encapsulation, growth, and division. *Science* 302:618–622.
82. Pyle, A.M. (2002). Metal ions in the structure and function of RNA. *J. Biol. Inorg. Chem.* 7: 679–690.
83. Klein, D. J., Moore, P.B., and Steitz, T. A. (2004). The contribution of metal ions to the structural stability of the large ribosomal subunit, *RNA*, 10:1366–1379.
84. Fox, G. E. (2010). Origin and Evolution of the Ribosome, *Cold Spring Harb Perspect Biol.*:2(9):a003483.
85. Smith, T. F., Lee, J. C., Gutell, R. R., Hartman, H. (2008). The origin and evolution of the ribosome, *Biol Direct.* 2008 Apr 22;3:16.
86. Polacek, N., Mankin, A. (2005). The ribosomal peptidyl transferase center: Structure, function, evolution, inhibition. *Crit Rev Biochem Mol Biol*, 40(5):285-311.
87. Klein, D.J., Moore, P.B., Steitz, T.A. (2004). The roles of ribosomal proteins in the structure assembly, and evolution of the large ribosomal subunit, *J.Mol.Biol.* 340: 141-177.
88. Schuwirth, B.S., Borovinskaya, M.A., Hau, C.W., Zhang, W., Vila-Sanjurjo, A., Holton, J.M., Cate, J.H. (2005). Structures of the bacterial ribosome at 3.5 Å resolution, *Science* 310: 827-834.
89. Stanley, R.E., Blaha, G., Grodzicki, R.L., Strickler, M.D., Steitz, T.A. (2010). The structures of the anti-tuberculosis antibiotics viomycin and capreomycin bound to the 70S ribosome, *Nat.Struct.Mol.Biol.* 17: 289-293.
90. The PyMOL Molecular Graphics System, Version 1.5.0.4 Schrödinger, LLC.
91. POV-Ray, Persistence of Vision Pty. Ltd. (2004). Persistence of Vision Raytracer (Version 3.6) [Computer software]. Retrieved from <http://www.povray.org/download/>.
92. Sanson, C., Le Meins, J.-F., Schatz, C., Soumab, A. and Lecommandoux, S. (2010). Temperature responsive poly(trimethylene carbonate)-block-poly(L-glutamic acid) copolymer: polymersomes fusion and fission, *Soft Matter*, 6, 1722–1730
93. Dobereiner, H.-G., Kas, J., Noppl, D., Sprenger, I., and Sackmann, E. (1993). Budding and Fission of Vesicles, *Biophysical Journal*, Vol.65, pp. 1396-1403.
94. Zheng, C., Liu, P., Li, J., and Zhang, Y. W. (2010). Phase Diagrams for Multi-Component Membrane Vesicles: A Coarse-Grained Modeling Study, *Langmuir*, 26(15), 12659–12666.
95. Noller, H. F. (2004). The driving force for molecular evolution of translation, *RNA*:10(12):1833-7.
96. Armond, P. A., Staehelin, L. A. (1979). Lateral and vertical displacement of integral membrane proteins during lipid phase transition in *Anacystis nidulans*, *Proc Natl Acad Sci U S A*:76(4):1901-5.
97. Borochof, H., Shinitzky, M. (1976). Vertical displacement of membrane proteins mediated by changes in microviscosity, *Proc Natl Acad Sci U S A*:73(12):4526-30.
98. Bashan, A., Yonath, A. (2008). Correlating ribosome function with high-resolution structures. *Trends Microbiol* 9:326–335.
99. Woese, C. (1998). The universal ancestor, *Proc. Natl. Acad. Sci. USA*. 1998 June 9; 95(12): 6854–6859.
100. de Ronde, C.E.J., Channer, D.M.D.R., Faure, K., Bray, C. J., Spooner, T. C. (1997). Fluid chemistry of Archean seafloor hydrothermal vents: Implications for the composition of circa 3.2 Ga seawater. *Geochim. Cosmochim. Acta*, 61, 4025–4042.
101. Hury, J., Nagaswamy, U., Larios-Sanz, M., Fox, G. E. (2006). Ribosome origins: The relative age of 23S rRNA domains. *Orig Life Evol Biosphere* 36: 421–429.
102. Mears, J. A., Cannone, J. J., Stagg, S. M., Gutell, R. R., Agrawal, R. K., Harvey, S. C. (2002). Modeling a minimal ribosome based on comparative sequence analysis. *J Mol Biol* 321: 215–234.
103. Smit, S., Widmann, J., Knight, R. (2007). Evolutionary rates vary among rRNA structural elements, *Nucleic Acids Res*:35(10):3339-54.
104. Wuyts, J., Van de Peer, Y., De Wachter, R. (2001). Distribution of substitution rates and locations of insertion sites in the tertiary structure of ribosomal RNA. *Nucleic Acids Res* 29: 5017–5028.
105. Bokov, K., and Steinberg, S. V. (2009). A hierarchical model for evolution of 23S ribosomal RNA, *Nature*:457(7232):977-80.
106. Bernal, J.D. (1969). *Origins of Life*. London: Wiedenfeld and Nicholson.
107. Bryson, B. (2004). *A short history of nearly everything*. London: Black Swan. pp. 300–302.
108. Bernal J.D. (1951) *The physical basis of life*, (Routledge and Keganb Paul).
109. Hill, A.R., Jr., Bohler, C., and Orgel, L.E. (1998). Polymerization on the rocks: negatively-charged alpha-amino acids. *Orig Life Evol Biosph* 28:235–243.
110. Schwartz, A. W. and Orgel, L. E. (1984). Template-directed polynucleotide synthesis on mineral surfaces. *J Mol Evol* 21:299–300.
111. Sanson, C., Le Meins, J.-F., Schatz, C., Soumab, A. and Lecommandoux, S. (2010). Temperature responsive poly(trimethylene carbonate)-block-poly(L-glutamic acid) copolymer: polymersomes fusion and fission, *Soft Matter*, 6, 1722–1730
112. Ellens, H., Bentz, J., Szoka, F. C. (1986). Fusion of phosphatidylethanolamine-containing liposomes and mechanism of the L alpha-III phase transition, *Biochemistry*, 25(14):4141-7.
113. Nelson, K.E., Robertson, M.P., Levy, M., and Miller, S.L. (2001). Concentration by evaporation and the prebiotic synthesis of cytosine. *Orig Life Evol Biosph* 31:221–229.
114. Wochner, A., Attwater, J., Coulson, A., Holliger, P. (2011). Ribozyme-catalyzed transcription of an active ribozyme. *Science*, 332:209–212.
115. Poole, A.M., Jeffares, D.C., and Penny, D. (1998). The path from the RNA world. *J. Mol. Evol.* 46: 1–17.
116. Cech, T. R. (2009). *Crawling Out of the RNA World*, *Cell*, 136, DOI 10.1016
117. Cuconati, A., Molla, A., Wimmer, E. (1998). Brefeldin A inhibits cell-free, de novo synthesis of poliovirus, *J Virol*:72(8):6456-64.
118. Atilgan, E., Sun, S. X. (2007). Shape transitions in lipid membranes and protein mediated vesicle fusion and fission, *J Chem Phys*:126(9):095102.
119. Weikl, T. R., Kozlov, M. M., Helfrich, W. (1998). Interaction of conical membrane inclusions: Effect of lateral tension, *Physical Review E*, Volume 57, Issue 6, pp.6988-6995.
120. Goulian, M., Bruinsma, R., and Pincus, P. (1993). Long-Range Forces in Heterogeneous Fluid Membranes, *Europhys. Lett.* vol. 22, pp. 145.
121. Park, J. and Lubensky, T. C. (1996). Interactions between membrane inclusions on Fluctuating Membranes, *J. Phys. I France*, vol. 6, pp.1217-1235.
122. Atilgan, E., Wirtz D. and S. X. Sun. (2005). Mechanics and Dynamics of Actin-Driven Thin Membrane Protrusions, *Biophys J*: 90(1): 65–76.

# SUPPLEMENTARY MATERIALS

## 1 INTRODUCTION

### 1.1 THE RNA WORLD HYPOTHESIS

The possible mechanism of early evolution and origin of life on earth has been a central problem for evolutionists for centuries. Knowing that cell operates through a series of organized chemical reactions, investigations started to find the origins of such systems. Oparin's experiment with coacervate droplets (1, 2) was the first attempt to find the possible origins of biochemical compounds in primitive earth. In modern form of life, it is observed that DNA functions as genetic material (for some virus, genetic material is RNA) and proteins are employed for structural and enzymatic purposes. RNA links between two separate biomolecules, providing translations of functional and structural proteins from informations contained in DNA. Lipids and sugars are the fourth kind of biomolecules, which have a very important role in compartmentalization of cells/organelles and metabolism.

The search of origin of life requires identification of the basic feature of life. Knowing that cell operates through a network of chemical reactions, we can see that for performing the biochemical reaction corresponding to a function, we need a specific structure of the reactant or of the enzyme. In addition, the structure comes from the respective sequence. However, there are arguments against this. In case of natively unfolded proteins, we can see isomeric behavior between similar sequences (3), leading to the conclusion that a protein adopting several different conformations might be employed to many functional roles (4). However, keeping in mind the essential simplicity of earliest form of life, we can argue that the origin of molecular alphabets (which means polymerization) is a fundamental requirement for origin of life. However, such molecular alphabets must also couple with heredity, in order to transfer compositional information to the next progeny. In other words, the molecular alphabet has to function as a self-replicating system. It seems that nucleic acids are more plausible as self-replicating biomolecules, compared to proteins. Watson and Crick model on DNA (5) provided us the idea about such a self-replication mechanism, where one strand of a polynucleotide can work as a template for synthesis of its complement strand from respective monomers (5-7). The authors of the three early papers clearly recognized that an autonomous 'RNA organism' would be possible only if RNA could take on several of the functions presently performed by proteins. For example, the functions of RNA polymerases and nucleases. They speculated that coenzymes incorporating nucleotides in their structure were fossils from a time when RNA functioned without the help of proteins (5, 7, 8). This idea was subsequently developed in some detail by White (9). The unanticipated discovery of catalytic RNA molecules, ribozymes, which perform enzyme-like reactions (10, 11) marked the beginning of the present interest in a protein-less biological world. Following Cech and Altman's discoveries, ribozymes were shown to be able to catalyze a significant number of diverse chemical reactions. This led to an increased interest in the hypothesis that an RNA World, a term introduced by Gilbert (12), preceded the DNA/RNA/Protein world (13). After which, it has been shown that under defined conditions RNA can be made to evolve new catalytic activities through molecular Darwinian selection (14-16). The RNA World hypothesis was much enriched by the discovery of ribosome structure at molecular level, revealing the fact that it is a truly RNA machine or a ribozyme (17).

In biological systems, it is known that DNA can be synthesized from RNA components (18). It is also known that ribosome, the truly RNA machine (19) synthesizes almost all the functional proteins. However, there are some examples of non-ribosomal protein synthesis (20, 21), but those mechanisms require proteins which are previously synthesized by ribosome. By this way, it seems that RNA is the 'true ancestor' of all the biomolecules, from which DNA and proteins originated. The basis of RNA World hypothesis lays in the fact that enzymatic and self-replication properties were coupled with RNA at the earliest stages of abiogenesis, which were later taken over by proteins. This does not mean that RNA was the first biomolecule to appear and polymerize. In fact, RNAs have less diverse structural variations compared to proteins, which were likely to be coupled with enzymatic functions at the earliest stages of evolution. This picture is in accordance with the common evolutionary scenario that biological complexity generally increases with stages of evolution. We shall show in our work that the conditions of Hadean Ocean were susceptible for oligomerization of both amino acids and nucleotides.

### 1.2 A CRITIQUE ON RNA WORLD HYPOTHESIS

Mills and Kenyon in 2011 (22) and Bernhardt in 2012 (23) presented two reviews on the problems with RNA World hypothesis. The present discussion elaborates these two review articles with some new ideas. We have presented arguments against RNA World hypothesis along with their acceptability in the light of some previously neglected fields. We also have discussed how we can puzzle out a majority of these problems by introducing a Lipid-RNA world hypothesis.

#### 1.2.1 Origin of a Beta-D-ribonucleotide pool

The first problem regarding RNA World hypothesis is the debate upon the prebiotic origin of its monomers. Prebiotic origin of RNA World requires a pool of beta-D-ribonucleotides existing in the Hadean Eon. Origin of such ribonucleotides again requires purine or pyrimidine bases, ribose sugar and phosphate groups. However, origin of ribose sugars seems difficult in prebiotic conditions (24, 25). The short half life of decomposition of ribose and other sugars at neutral pH entails it was probably unavailable in prebiotic conditions (26). The hot, acidic, radiation-exposed prebiotic conditions (27-29) seem unfavourable both for sugar (26) and for some nitrogen bases (30). In spite of that, some mechanisms of prebiotic ribonucleotide synthesis have been suggested (31-33). However, almost all of these processes are very slow and occur at low temperatures (33).

The basis of this argument lies in the commonly accepted hot, acidic, radiation-exposed scenario of Hadean Eon, in which nucleotides are unlikely to emerge. The unavailability of enough geological record in Hadean Eon makes it difficult to identify the exact environmental conditions. Supporters of faint young sun conditions (34) claim the Hadean earth to be extremely cool (35, 36); though, geologists say otherwise. Using oxygen isotope analysis and other methods, ocean temperature at 4.0 Ga has been reported to be about 100°C (27). Study of zircons has found that liquid water must have existed as long ago as 4.4 Gyr, very soon after the formation of the Earth (35). This indicates formation of atmosphere by that time. In fact, there are some geological evidences about a CO<sub>2</sub> dominated Hadean atmosphere (28, 37), which was picked up by Astrobiologists as a solution for faint young sun paradox (38, 39). However, this leads to another problem. CO<sub>2</sub>-rich atmospheres have been shown to be inefficient in the prebiotic synthesis of organic compounds (40, 41). In addition, the faint young sun paradox has not been solved completely by this CO<sub>2</sub>-rich atmosphere (42) and possibility of an extremely cold early earth remains. Thus, it is difficult to ‘choose a side:’ whether to accept a hot, acidic prebiotic environment or a cool early earth surface with frozen oceans (43).

In fact, the frozen ocean hypothesis might cope with faint young sun paradox and might provide hypothetical environment for ribonucleotide formation (33), but the melting of such frozen oceans in Archean Eon is confusing. From geological evidences, we are sure that the environmental temperature was warmer in Archean Eon compared to the present day (34). To explain this, one hypothesis was suggested that continuous meteorite impact in the beginning of Archean Eon caused melting of the frozen oceans (35, 43), which, in our personal view, is not an adequate hypothesis. Because this hypothesis is unable to explain how frozen oceans can establish under much higher rate of meteorite bombardment in Hadean Eon (44). In summary, the possibility of establishment of a beta-D-ribonucleotide pool in Hadean Eon remains in darkness.

### **1.2.2 Spontaneous polymerization of Beta-D ribonucleotides**

The second problem regarding RNA World hypothesis is the argument that Beta-D ribonucleotides spontaneously form polymers linked together by 3', 5'-phosphodiester linkages. It was shown that Evaporation of acidic solutions of nucleotides and subsequent heating leads to the formation of complex mixtures of very short oligonucleotides, in which 2'-5'-, or 3'-5'-phosphodiester linkages occur almost randomly (45). Preactivated nucleotides show a good yield of polymerization and was claimed to be prebiotic (46). However, in modern cells, nucleotides are activated before polymerization by ATP. In prebiotic environment, it is unclear whether such a high amount of ATP was available or not (18). Thus, the source of free energy to drive the uphill polymerization of nucleotides remains in darkness (33).

There are some ‘realistic’ arguments favoring the polymerization reaction, that the metallic ions can catalyze the nucleoside 5'-phosphorimidazolides oligomerize in aqueous solution (47-49). This kind of polymerization can even lead to the generation of long homo- and hetero-oligomers (50). Predominantly 2'-5'-linked (51, 52). [UO<sub>2</sub>]<sup>+</sup> is also an efficient catalysts in aqueous solution for ribonucleotide polymerization, producing predominantly 2'-5'-linked oligomers (51, 52). In fact, James Ferris and his coworkers (53) have investigated in considerable detail a remarkable series of reactions in which an abundant clay mineral, montmorillonite, catalyzes the synthesis of long oligonucleotides from relatively dilute solutions of nucleoside phosphoramidates including the 5'-phosphorimidazolides. It was reported that montmorillonite (clay with a layered aluminosilicate structure) expands to permit large molecules to enter its interlayers, and it is probable that new phosphodiester bonds are formed when the substrates are absorbed in the interlayers (54). Existence of catalytic sites within montmorillonite was also confirmed (55) from the work of Wang and Ferris. In summary, this discussion leads to the possible conclusion that once a prebiotic pool of nucleotides was available, the montmorillonite clay and metallic cations could catalyze the oligomerization interactions. From the hypothesized possibilities in prebiotic earth, it is unclear whether chemical sources of energy (ATP/GTP) were abundant; most believe that they were not. Therefore, we should concentrate on the possibilities excluding the need of any chemical sources of energy (or at least utilizing minimum possible chemical energy) to explain the events of abiogenesis. Here, montmorillonite and metallic ion catalysis provides a ‘realistic’ possibility in prebiotic earth.

### **1.2.3 RNAs are self-replicating biomolecules**

The third argument against a possible RNA World is RNAs are incapable of functioning as a self-replicating biomolecule. Certainly, this template-directed RNA synthesis in Hadean Eon was devoid of proteins, which was described by Orgel (56) as “seems probable, but by no means certain.” The problem of energy source for polymerization is again unclear.

However, there are experimental evidences that if a polynucleotide is incubated with an appropriate mixture of complementary mononucleotides or short oligonucleotides at a sufficiently low temperature, in many cases a double- or triple-helical complex is formed (57). These are structurally interrupted, but still are structurally similar to normal double- or triple-stranded nucleic acid helices. However, most of the molecular mechanisms (58, 59) seem not to be appropriate in a RNA World scenario (33). In other studies, Mg<sup>++</sup>, Pb<sup>++</sup>, Zn<sup>++</sup> have been observed to catalyze the template based polymerization of nucleotides (60, 61). A detailed discussion can be found in reference 33. From the discussion so far, we have picked up two important points: both the random polymerization and template-directed polymerization of nucleotides can be mediated by metallic cations/ montmorillonite. Since these cations are positively charged, it is possible that they will interact with the negatively charged backbone of RNA-oligomers and provide structural stabilization. In fact, such events have been confirmed by many observations. We will show the importance of this event on the context of a Lipid-RNA World in the next sections.

### **1.2.4 Inherent unstable nature of RNAs**

The inherent unstable nature of RNAs provides another strong argument against the RNA World hypothesis. The conspicuous presence of hydroxyl group at the 2'-position of the ribose sugar in RNA makes it a bit unstable (62). This argument stands against the RNA World hypothesis, which requires prebiotic accumulation of RNA oligomers. Since in a hot, acidic prebiotic ocean RNA can easily hydrolyze, some believe that the RNA world may have evolved on ice, possibly in the eutectic phase (43, 63-66). The activity of the ribozymes maximizes at  $-7$  to  $-8$  °C temperature. Geological studies report (37) that approximates pH of Hadean Ocean was about 4-5. In general, phosphodiester bonds are stable in this pH, which led to a proposal that RNA World might evolve in acidic pH (67). In addition, it is commonly argued that the ozone layer did not exist 4.0 billion years ago, which means primitive earth was completely exposed to UV and gamma ray radiations. The aromatic bases of RNA absorb strongly in the ultraviolet region, which makes them more susceptible to damage by background radiation (68). Thus, this radiation exposed prebiotic environment again seems to be incompatible for establishment of an RNA World. However, a theoretical model (43, 69) utilizes faint young sun hypothesis and predicts an oceanic protection strategy for RNA World. This model assumes a frozen ocean, with approximately 300-1000 meter thick ice layer (43), which would have offered complete shielding of dissolved organic compounds from damaging UV radiation by scattering of the UV light. The ice layer on Europa (satellite of Jupiter) offers similar protection.

The importance of low temperature environment in the establishment of RNA World is almost certain from this discussion. In spite of the unstable nature of RNAs, a cold oceanic environment resulted by faint young sun condition could have saved them from degradation and ensured susceptible environment for their functional activity. In summary, an ancient RNA World has to be established in a cold environment.

### 1.2.5 Catalytic activity is seen in long RNA chains only

Another argument against the RNA World hypothesis is the limited catalytic activity of ribozymes. Many of the ribozymes are long RNA chains and probability of emergence of such long chains by random polymerization of nucleotides is very small. A ribozyme catalyzed RNA transcription was reported by Wochner *et al.* in 2011; where the replicase ribozyme was  $\sim 190$  nucleotides long (70). Although, now we have a number of known small ribozymes (64) that can enhance rates of some chemical reactions. Yarus and coworkers have also reported about ribozymes with few nucleotides length having catalytic activity (71). In fact they observed that 3-5 nucleotide long RNA segments can aminoacylate a substrate variant with the sequence GCCA (similar to the universal aminoacylated 3' terminus of tRNA). Thus, it is adequate to imagine an early stage of evolution when catalytic activity of small RNA segments initiated an RNA World. At the earliest stages of abiogenesis, we should not expect biomolecules with high structural and functional diversity.

## 1.3 CONCLUSION

Modern life follows genetic code to ensure the sequence specific synthesis of functional biomolecules. DNA is the genetic material from which messenger RNA and ribosomal RNA are transcribed. Ribosomal RNA translates messenger RNA into functional proteins. On the other hand, RNA can be reversely transcribed into DNA. Ribozymes alone can also catalyze the transcription of RNA (70). All these indicate to the conclusions that RNA could be the ancestor of all the biomolecules. In fact, the non-ribosomal peptides are small amino acid oligomers with very limited structural and functional diversity. This leads to a crucial conclusion that although small oligopeptides existed in the time of RNA World, but large functional proteins could only be synthesized after the emergence of ribosome. On this basis, we strongly deny 'Proteins First' (72-77) hypothesis.

The emergence of RNA World faces some strong opposing arguments from which it seems that RNA World might not be the earliest evolutionary stage. Origin of a beta-D ribonucleotide pool and their polymerization should be assisted by some preceding biomolecules. The Lipid World hypothesis (78) deals with such a possibility. To understand the emergence of RNA World starting from a preceding Lipid World, firstly we need to understand the exact environmental conditions of the system where all these taken place. In summary, first we need to understand environmental conditions of Hadean Ocean and the strategy of Natural Selection during abiogenesis. The RNA World cannot exist in a hot, acidic radiation exposed environment. At first we have to investigate whether a cold and radiation free environment was available in Hadean Ocean or not.

## 2 THE BEGINNING OF ABIOGENESIS IN HADEAN OCEAN

### 2.1 ENVIRONMENTAL CONDITIONS OF HADEAN OCEAN

#### 2.1.1 Formation of Earth

The modern planet earth was formed approximately 4.5 Gya by the collision between earth and a planet having size of Mars (79). This giant impact resulted in the formation of the earth and moon system and is responsible for earth's spin as well (80). This hypothesis is known as Giant Impact hypothesis and is commonly accepted because the predicted earth spin and moon orbit from this hypothesis matches well with observation.

#### 2.1.2 Origin of Atmosphere

The Earth's atmosphere is thought to be derived from mantle degassing (81). Excesses of  $^{129}\text{Xe}$  in mantle-derived samples relative to the atmosphere have been interpreted to indicate the presence of live parent  $^{129}\text{I}$  in the deep Earth following early degassing of the atmosphere (82, 83) since  $^{129}\text{I}$  decays to  $^{129}\text{Xe}$  with a half life of only 16 million years. If true, then it is argued that the present atmosphere and hydrosphere must have formed by  $\sim 4.4$  Gyr (84). An elegant discussion about about the origin of earth and atmosphere can be found in NASA Astrobiology Institute research page (URL: <http://astrobiology.ucla.edu/pages/research.html>).

### 2.1.3 Geological Approach to the Hadean Ocean Environment

Using oxygen isotope analysis and other methods, ocean temperature at 4.0 billion years ago has been reported to be about 100°C (27). Study of zircons has found that liquid water must have existed as long ago as 4.4 Gya, very soon after the formation of the Earth (35). This indicates formation of atmosphere by that time. In fact, there are some geological evidences about a CO<sub>2</sub> dominated Hadean atmosphere (28, 37). Based on such geological studies, hydrothermal vent models were proposed to be the earliest systems where abiogenesis began. The basic idea is, in 4.4 Gyr, liquid water was available on earth surface at an enormous amount, and early earth crust was very thin. This means the earth surface had to be volcano prone. Submarine volcanoes ejected molten rock, lava and extremely hot gases which could have created hydrothermal vents in Hadean Ocean like today.

Several hypothesized models show that the early evolution of life was linked to ancient hydrothermal activities in the Hadean and the early Archean (85-87). Some models further propose that the hot (<200°C), alkaline hydrothermal discharges might prepare the earliest stage of the origin of life in the Hadean ocean (88, 89) because the alkaline hydrothermal vent environments would be feasible as a cradle of the most ancient life forms sustained by the primitive ATP-generating machinery that was based on the mineral compartments (primordial cells) intervening the chemiosmotic (pH, proton, and other materials) gradient in the hydrothermal mixing zones (89). We shall show in this paper that such hydrothermal vent systems actually provided the necessary physical conditions for emergence of a Lipid-RNA World from Lipid World and later a Protein-RNA World from Lipid-RNA World. The discovery of warm (<100 °C), alkaline H<sub>2</sub>-rich hydrothermal fluids in the Lost City hydrothermal field off the Mid-Atlantic Ridge has also greatly inspired the idea that the ancient ultramafic rock-hosted hydrothermal activities could sustain H<sub>2</sub>-driven chemolithotrophic microbial ecosystems (90, 91). Shibuya *et al.* in 2010 (92) presented a Hydrothermal vent model of Archean ocean by a combination of experiment and thermodynamic calculations. In their model, the volcano-prone marine base is the source of a very high temperature at the oceanic base (~350°C), whereas the water temperature is quite low at top (~0°C).

### 2.1.4 Biochemical advantages of a hydrothermal vent system on the context of abiogenesis

Martin *et al.* in 2008 (89) and Shibuya *et al.* in 2010 (92) presented elegant discussions on the biochemical advantages of hydrothermal vent systems on the context of abiogenesis. We are pointing out some of them on the context of our hypothesis.

In the present hydrothermal systems, in situ pH and redox conditions of high-temperature hydrothermal fluids are strongly buffered by the mineral assemblage of plagioclase, epidote and anhydrite at the hydrothermal reaction zone. Anhydrite influences the redox condition of the hydrothermal fluid together with plagioclase and epidote (93). The anorthite component in plagioclase and the clinozoisite component in epidote control the  $a\text{Ca}^{2+}/a^2\text{H}^+$  ratio of the hydrothermal fluid (94), while Ca<sup>2+</sup> concentration is also controlled by metastable, igneous calcic plagioclase in fresh rocks (95). This process maintains the value of in situ pH around 5.0 that is close to the neutral pH at high-temperature condition (94), which was confirmed by direct in situ pH measurements of deep-sea high temperature hydrothermal fluid (96). Acidification of the hydrothermal fluid is caused by precipitation of metal sulfides, dissociation of H<sup>+</sup>-bearing complexes due to decreasing temperature during emission on the seafloor (96).

Archean seafloor basalts show extensive alteration and strong carbonatization attributed to the high-CO<sub>2</sub> concentration in seawater/hydrothermal fluid (97-100), which led to almost complete albitization of igneous calcic plagioclase and calcite-bearing secondary mineral assemblages even at high-temperature conditions of ~350–400°C (97, 98, 100). We have accepted this model.

There are striking parallels between the chemistry of the H<sub>2</sub>-CO<sub>2</sub> redox couple that is present in hydrothermal systems and the core energy metabolic reactions of some modern prokaryotic autotrophs (89). Probably this is the reason that submarine hydrothermal vents are geochemically reactive habitats that harbor rich microbial communities, the energy source of which stems mainly from mid-ocean-ridge volcanism (101, 102). The ~350°C sulphide chimneys of the vent systems are primordial environments that are reminiscent of early Hadean Oceanic conditions. The volcanic eruptions produce black smokers and associated diffuse flow systems, that host dense and diverse biological communities (103). Black smoker chimneys emit hot (up to 405°C) and chemically modified seawater (104). Beneath the fissured sea floor, during its circulation from the ocean floor downwelling seawater comes into close contact with the magma chamber. Effluent at black smokers is typically acidic (pH 2–3) and rich in dissolved transition metals (105), such as Fe(II) and Mn(II). The black smoker systems are fuelled by volcanoes, and so these fluids commonly contain high concentrations of magmatic CO<sub>2</sub> (4–215 mmol per kg), H<sub>2</sub>S (3–110 mmol per kg) and dissolved H<sub>2</sub> (0.1–50 mmol per kg), with varying amounts of CH<sub>4</sub> (0.05–4.5 mmol per kg) that is formed both through biogenic and abiogenic processes (106). A range of temperatures exists, from the hot interior of black smokers to the interface with cold (2°C), oxygenated seawater.

There are examples (Lost City, ref. 89) of submarine hydrothermal vent systems where volcanic effluent is alkaline (pH 9-11). This produces a pH gradient in the vent-ocean interface. In case of a Hadean Ocean scenario, naturally chemiosmotic nature of alkaline hydrothermal vents with pH 9–10 and the outer walls of chimneys, bathed in ocean fluids with a pH of 5–6, were essential for the origin of free-living cells (107-109).

Submarine volcanoes in Hadean Ocean were much active compared to those of the modern, since early earth had a thin crust and sea floor was very close to the mantle. This could have established a hydrothermal vent system in the entire Hadean ocean, which an approximated ~350°C sea floor temperature. Volcanic eruptions ejected a large amount of biochemical elements, which established biologically rich



oceanic water. In addition, close to the volcanic regions, there were alkaline phases of seawater. Such biochemically rich environments could have been very important in the beginning of abiogenesis.

### 2.1.5 Astrophysical approach to Hadean Ocean environment: The Faint Young Sun paradox

In 1972, Carl Sagan and George Mullen (34) showed that according to the standard solar model, the sun was about 30% less luminous in Hadean Eon compared to its present luminosity. This solar output is insufficient to maintain a liquid hydrosphere. Therefore, according to the standard solar model, Hadean Ocean should have been completely frozen. This is contradictory to the geological evidences, which demonstrate the presence of liquid water on earth surface as early as 4.4 Gyr. This contradiction is known as the faint young sun paradox. Following this paradox, a number of models were proposed proposing about a cool early earth (35, 36). Scientists tried to solve this faint young sun paradox by approaching from diverse angles, like CO<sub>2</sub> dominated Hadean atmosphere (28, 37), Green House gas clouds (39), elevated OCS-rich environment (110) etc. However, none of them solves this paradox completely.

A diplomatic solution was proposed by Bada *et al.* (43) in 1994 and Cleaves *et al.* in 1998 (69) that Hadean ocean were frozen only at the top, having approximately 300-1000 meter thick ice layer, which protected the early biomolecules from UV and gamma ray radiation. This model also provides a cold environmental situation where an RNA World can emerge and evolve. However, the melting of ice is confusing. From geological evidences, we are sure that the environmental temperature was warmer in Archean Eon compared to the present day (34). To explain this, one hypothesis was suggested that continuous meteorite impact in the beginning of Archean Eon caused melting of the frozen oceans (35, 43), which, in our personal view, is not an adequate hypothesis. Because this hypothesis is unable to explain how frozen oceans can establish under much higher rate of meteorite bombardment in Hadean Eon (44).

### 2.1.6 The Ice-Covered Hydrothermal Vent Model

Merging together the proposals of Bada *et al.* (43) in 1994 and Cleaves *et al.* in 1998 (69) that Hadean Ocean should have a frozen top due to faint young sun and should also have a Hydrothermal vent core (85-87), we propose a 'ice-covered hydrothermal vent' model of Hadean ocean. This is a simple modification of a common hydrothermal vent system with a frozen top. The earth surface temperature should be very low due to faint young sun; however, the submarine volcanoes will maintain an extremely hot sea floor. The hot water of sea floor will continuously experience an upstream motion towards the top, where they will lose temperature due to the ice layer and again come back to the sea floor. This will give rise to a continuous circular current of water from top to bottom. This continuous current will protect the organic materials from UV by taking them deep into water repeatedly. In addition, the ice layer at the top surface of ocean provides an extra protection against UV. Thickness of this ice layer should not be as much as predicted by Bada *et al.* (43). Because the upstream extremely hot water will keep melting ice continuously and the faint young sun will keep freezing the seawater. There will be a balance between these two events. The resulting ice layer thickness should be small enough to be melted during the high meteorite impact rate at the beginning of Archean Eon. We shall show that such a system provides the necessary physical conditions for establishment of an RNA World and can prepare an evolutionary route towards a Protein-RNA World. This picture is demonstrated in Figure-1 in the manuscript.

## 2.2 NATURAL SELECTION IN THE BEGINNING OF ABIOGENESIS

A general tendency of the biological evolution is a gradual complexation of the living systems, by "fine tuning" the enzymatic functions and heredity mechanisms through continuous cycles of self-reproduction at a large timescale. Biological complexation leads to high energy, far-from-equilibrium systems, rather than the lower energy, equilibrium systems that are the target of non-biological complexation (111). On the other hand, non-biological complexation, in the context of a closed system, drives the system toward minimum free energy, i.e. the equilibrium state. The process can be chemical in nature, as manifest in the bonding between atoms to create molecules, or physical in nature, as exemplified by aggregation processes such as micelle or vesicle formation and crystallization (111). In attempting to bridge between biology and chemistry, Eigen in 1971 (112) proposed that the very same process of Darwinian evolution that operated at the biological level must have initially also been operational at the prebiotic molecular level. Based on this idea, the chemical evolution can be visualized as a process when the random chemical reactions were organized within a specific boundary and sequential chemical pathways originated.

The era of chemical evolution was not assigned with cellular life and thereby versatile structural and functional complexity of biomolecules. Assuming the abiogenesis started with a prebiotic Lipid World, the diverse biochemical reaction network (property of cellular life) was absent at that time, which is contemporary to the modern life. Mutations result increment of disorder in this network system. Only those mutations are naturally selected which keep core reactions intact and modify the side reactions; and thus causes 'fine tuning' of existing biochemical reactions. The more a chemical reaction being a core reaction, the higher is the evolutionary pressure to keep that one unaffected.

Nevertheless, when there was no organized biochemical network (i.e. no cellular life), there should not be any evolutionary pressure on maintaining any biochemical function. In the Lipid World or RNA World scenario, the chemical reactions were completely random, which were organized later to give rise to specific pathways and initiated cellular life. Thus, in Lipid World or RNA World scenario, the natural selection of mutations could not be processed in the contemporary way. The strategy of natural selection should be dependent and consistent on the physical conditions of the habitat of the biomolecules (consistency with habitat is the basic feature of natural selection). An ice covered hydrothermal vent system with water temperature ranging from ~350°C at the bottom and ~0°C at the top and a strong pH gradient requires structural consistency of the earliest biomolecules. Otherwise, they would have denatured soon in this hostile condition. One ar-

gument seems possible: the earliest biomolecules must have originated by means of random polymerization of monomers and have grown in size with course of time. Such an event should be consistent with the physical conditions and must have continued ensuring high structural stability. In fact, the evolutionary strategy of biomolecules should be solely dependent on structural stability. Let us consider an amino acid or nucleic acid polymer growing in size with time; more monomers are joining the structure. If a new monomer joins the polymer at a certain stage, its persistency will be determined by the structural stability it is providing to the polymer. If the new monomer destabilizes the whole polymer structure, it should be dissociated soon. On the other hand, if it provides a structural stability, its persistency will be high. We shall show later that this strategy of natural selection fits perfectly with the evolutionary picture of ribosome, a remnant of RNA World. Same kind of natural selection strategy should also be observed in case of isoenzymes: between two isoenzymes, the one with higher structural stability and higher catalytic activity should undergo natural selection. Therefore, we can identify a Darwinian picture in chemical evolution. We shall show that the ribosome bears the signature of such an evolutionary strategy.

### **3 THE ANCESTOR RIBOSOME AND ANCIENT TRANSLATION MECHANISM**

#### **3.1 ANCIENT AND MODERN RIBOSOMAL PROTEINS**

Arcady Mushegian in 2005 (113) identified the distribution of ribosomal proteins across the three phylogenetic domains and presented a set of proteins that could be the ancestor proteins, possibly present in Last Universal Common Ancestor, LUCA (114). Such an analysis was conducted again in 2008 by Roberts *et al.* (115), where they correlated the molecular signatures regarding coevolution of ribosomal proteins with ribosomal RNAs. Vishwanath *et al.* 2004 (116) argued that all parts of the r-proteins are probably not equally old. Experimental studies have shown that ribosomal components are assembled in a reproducible manner, which might recapitulate to a significant extent the history of the ribosome and thereby provide timing information. It is hypothesized that the oldest proteins would assemble first and be at the core of the process whereas newer proteins would be incorporated into later stages of assembly and the newest proteins would be last (117). Based on this context, Fox in 2010 (118) predicted the relative ages of the ribosomal proteins. He argued that the assembly order of the universal r-proteins speaks to their relative age. He grouped the universal ribosomal proteins of the 50S subunit into four clusters ranging from oldest to most recent. These four groups are (1): L2, L3, L4; (2): L22, L23, L24; (3): L5, L6, L10, L11, L13, L18, L29; and (4): L9, L31, L32-L34. Smith *et al.* in 2008 (119) identified that all three predicted oldest proteins (L2, L3 and L4) interact with the PTC region of 50S. Apart from them, other universal PTC-interacting proteins are L6, L14, L10e/L16 and L15e/L31.

#### **3.2 RIBOSOMAL PROTEINS ARE NOT ESSENTIAL FOR TRANSLATION**

Brodersen and Nissen in 2005 (120) presented an elegant minireview on the ribosomal proteins and their functions. A number of small and large subunit proteins were shown to be involved in the translation mechanism directly (Table-1, ref. 120). In addition, there are translation factor proteins. However, there are many experimental evidences showing that none of them is essential for translation mechanism, though absence of some proteins might result in greatly reduced rate of amino acid polymerization.

Polypeptide synthesis was shown to initiate in the absence of initiation factors, by manipulating the ionic conditions (121). Aminoacyl-tRNA can bind to the ribosome in the absence of elongation factor EF-Tu, though, the binding rate is greatly reduced (122). Peptide bond formation is catalyzed by the large ribosomal subunit (17). Even translocation of tRNA was shown to occur without EF-G (123, 124). Nomura *et al.*, in 1969 (125) showed that many small subunit ribosomal proteins can be singly omitted without hampering the function. Hoang *et al.* 2004 (126) experimentally demonstrated that S9 and S13- the two small subunit proteins interacting with P-site tRNA are not essential for ribosome function; however, deletion of their interacting site results viable, but 40% reduced growth rate in *Escherichia coli* phenotypes. In fact, ribosomal proteins, though not essential for translation, but they are responsible for accuracy and speed of translation. (127). For example, S4 and S5 improve the accuracy of translation and their mutations increase error frequencies (128, 129). In fact, the inessentiality of ribosomal proteins in translation mechanism leads to the conclusion that protein synthesis started in a protein-free world. Which means, whatever the proto-ribosome structure was, at was an RNA only structure, with no protein present. However, probably translation mechanism was not a 'true translation', but was random polymerization of amino acids. In the next sections, we shall try to explore this concept.

#### **3.3 ANCIENT REGIONS OF 23S rRNA AND THE MINIMAL RIBOSOME**

The substantial structural and sequence conservation seen in comparisons of the rRNAs from all three Domains of life suggest that they reached their modern size early in the development of the ribosome (119). However, it is extremely unlikely that all parts of the rRNAs are of the same age. Instead, researchers have argued that like some of the ribosomal proteins, they have increased in size over time, perhaps beginning as an amalgamation of smaller fragments (129, 130). Cogent arguments have been presented that certain portions of the rRNAs are older than rest of the parts (131-135). Hury *et al.* in 2006 (135) argued that older regions of 23S rRNA would have more time to be integrated into the structure and hence would show greater connectivity to other regions than newer additions. On this basis, they found Domain V and Domain IV to be the oldest parts of ribosome, while Domain V containing the functional PTC region. Interestingly, these ancient regions are observed to be in a good alignment with the structure of the proposed Minimal Ribosome by Mears *et al.* in 2002 (133). Mears *et al.* (133) have defined minimal ribosomal RNA as the set of rRNA regions conserved in all living forms and two types of organelles. This minimal rRNA is approximated by two extremely deleted rRNAs found in mitochondria of nematodes. The small-subunit rRNA in nematode mitochondria has lost ~55% of its nucleotides compared to an average bacterial homolog, the large-subunit rRNA has lost ~74%, and 5S RNA is missing. The remaining, universally conserved rRNA domains include most of the intersubunit interface, as well

as the peptidyltransferase center and the sites of interaction with tRNAs. The similarity between predicted ancient ribosomal regions with the minimal ribosome indicates that this minimal ribosome could have possibly existed in the time of LUCA (118). However, it is extremely unlikely that this was the ribosome of Lipid-RNA World because of its structural complexity and large size. Such a large size can easily denature in high temperature environment (bottom of Hadean ocean), unless proteins are there to provide it necessary stabilization.

### 3.4 PEPTIDYL TRANSFERASE: RIBOSOME IS A RIBOZYME

Structures of the 30S and 50S subunits at 3.0Å and 2.4Å resolution, respectively, provided detailed descriptions of the folding of the RNA and protein components of the ribosome for the first time (136, 137). Most of the proteins are distributed on the backside of the two subunits, whereas the interface side is almost devoid of proteins, with a few exceptions. From the distribution pattern of ribosomal proteins, Noller argued that an ancient RNA structure has gradually incorporated a number of proteins over evolutionary time, but has not allowed them to impinge on its crucial functional centers. In fact, no protein is found within 17Å distance of the catalytic site of 50S subunit. This observation led to the conclusion that ribosome is a truly RNA machine (17), or, in other words, it is a ribozyme. The long, positively charged extensions of ribosomal proteins may serve to hold together rRNA components of the PTC (136), which explains the lack of catalytic activity of protein-free 23S rRNA (138). A detailed description on the PTC structure can be found in reference 137.

### 3.5 PEPTIDYL TRANSFERASE: STRUCTURE AND PREDICTION OF ORIGIN

According to the definition of PTC segment (139), it is composed of two main helical segments (A-loop and P-loop) and a large centroid region. In three-dimensional structure (119), there are two long parallel helices (A-loop and P-loop) and an unstructured region (centroid segment). This structure is conserved in all the crystal structures of ribosome available yet. Polacek and Mankin (139) argued that the main function of PTC is the proper alignment of the reaction components and hence, peptide bond formation. Therefore, PTC cannot be a mere rigid mold for the positioning of peptidyl- and aminoacyl-tRNAs. Rather, it presents a flexible and sensitive environment, which might be achieved through some kind of induced fit in response to the binding of the reaction substrates or auxiliary factors. Mutations in PTC, binding of inhibitors, which affect accuracy of translation, probably prevent the conformational changes ongoing during translation (140-143). They have further argued that the pH-dependence of translation is another support for the general flexibility of PTC structure.


Since proteins are absent from PTC region, Mg<sup>++</sup> ions are the main stabilizing agents of this region. In many cases, the phosphate oxygen atoms act as inner sphere Mg<sup>++</sup> ligand (144, 145). It can be speculated that such a stabilization strategy was an essential condition for survival of ribozymes in a protein free world. Fox argued that this stabilization strategy indicates that the PTC structure evolved before the availability of proteins (118). Smith *et al.* in 2008 (119) published an elegant structural analysis of the ribosome structure and found that there is no self-folding RNA segment in 16S rRNA. However, interestingly, it was found that PTC region of 23S rRNA (2472 through 2650, *H. marismortui* numbering) is a self-folding RNA segment (having lowest free energy when the folded secondary structure is in its native fold). They have identified one additional 23S RNA segment of which predicted minimum free energy secondary structure is also in its native structure. This is the segment from 2670 to 2830 (*H. marismortui* numbering) forming a cruciform-like structure in the extant LSU. This cruciform structure effectively forms part of the Smith and coworkers observed that this self-folding RNA along with PTC-interacting ribosomal protein segments define an almost planar surface. This surface was defined by PTC RNA (2472-2650) and L3 (1-21, 205-262), L4 (42-100), L6 (139-172), L15 (1-57) and L22 (115-143) proteins (Figure-3, ref. 119). All residue numbers described here are for *H. marismortui*. They presented a hypothesis that the planar structure of the PTC along with these protein segments probably indicates a membrane bound origin of 50S subunit.

The hypothesis of Smith *et al.* (119) describes the origin of 50S subunit as the PTC RNA and some small protein segments attached to a membrane surface (may be a vesicle), from which the growth of RNA and proteins (coevolution) resulted the contemporary structure. In spite of this courageous speculation (from which our idea of Lipid-RNA World originated), we strongly argue that the membrane interacting plane speculated by Smith *et al.* is not possible. Their hypothesis is unable to explain the following arguments.

- i> It is speculated that the PTC RNA segment and the earliest translation mechanism initiated in a protein-free world. This means the earliest ribosome structure could not have any protein segment attached to it.
- ii> It is unclear how such a PTC would polymerize amino acids. Authors did not suggest any mechanism.
- iii> The protein segments speculated by Smith *et al.* have minor interactions with PTC (each with only few hydrogen bonds and van der Waals contacts). Without a strong attachment, such a structure could not be subjected to natural selection in a Darwinian picture.
- iv> If proteins assisted the earliest 50S subunit structure, they should have evolved with higher functional diversity. Yet, only L2 and L3 are required for PTC in modern 50S structure, rest all are employed in peptide exit channel and tRNA interaction (which means not essential in PTC structure or function). If after four billion years of evolution not a single protein is essential in PTC structure of function, then how could they be essential at the earliest stages of ribosome?
- v> RNAs are capable of membrane-interaction. PTC RNA alone fulfills all the structural requirements necessary for membrane-interaction. Then why does it require proteins where it alone can bind to membrane?

- vi> We have described the hypothetical scenario of growth of 50S subunit in its membrane-bound conformation in manuscript. It appears if ribosome experiences a growth in size, bound to the membrane as speculated by Smith *et al.*, then the possible growth cannot explain the contemporary arrangement of the different domains of 50S subunit. In Figure-1, we have shown the oldest two domains (Domain-V and IV) of 50S subunit. The three-dimensional arrangement of these two domains can only be explained if the membrane interacting surface of PTC were as per our prediction (see manuscript).

**Figure-1:** The hypothetical membrane interacting surface predicted by Smith *et al.* (119) is presented as a red curve, whereas our prediction of



membrane interacting surface is presented as a yellow curve. Domain-V and Domain-IV, the two oldest domains of 23S rRNA (135) are shown as cyan and orange cartoon view respectively. The Smith *et al.* prediction cannot explain the origin of Domain-IV over the membrane interacting surface. Whereas, our prediction shows a beehive-like growth of ribosome starting from PTC can result the formation of Domain-IV over the oldest Domain of 23S rRNA (Domain-V). Some small parts of the two oldest domains are above our predicted surface, which are, probably structural modification after membrane detachment. This image is the reproduction of Figure-3 (B) in the manuscript. This image is produced using the 50S subunit structure having PDB-id 2AWB (146).

However, the prediction of a possible membrane origin seems to be correct. We have shown in the manuscript (Figure-3) that a different membrane-interacting surface of PTC can resolve all the ambiguities and can explain the contemporary structure of ribosome as well. This surface is also presented in Figure-1 here. Smith *et al.* prediction of the membrane-interacting surface demands protein help for membrane attachment of PTC. On the other hand, our membrane-interaction model does not require any protein; PTC binds to membrane on its own. This model is more 'compatible' in an RNA World scenario.

### 3.6 CONTEMPORARY MOLECULAR MECHISM OF TRANSLATION


Polacek and Mankin (139) presented a detailed discussion on the contemporary mechanism of peptide bond formation in ribosome. During contemporary peptide bond formation, the alpha-amino group of aminoacyl-tRNA in the A-site attacks the carbonyl carbon of P-site bound peptidyl tRNA. The tetrahedral transition state decomposes into the reaction products, deacylated tRNA at the P-site and peptidyl-tRNA carrying an additional amino acid (+1 aa) at the A-site. The formation of a peptide bond does not require additional energy, since energy is stored in an ester bond linking the peptidyl residue to the tRNA 3'-hydroxyl group. Creation of this bond is "energized" by ATP hydrolysis

that drives aminoacylation of a tRNA by the respective aminoacyl-tRNA synthetase. Hydrolysis of one Mole of ATP releases about 8.0 kcal energy, which is almost completely transferred to the aminoacyl-tRNA. However, only ~0.5 kcal/mol is required for amide bond formation (147). In the absence of ribosome, the uncatalyzed reaction occurs very slowly in solution, with about one peptide bond formation per 30 hours (148). The ribosome accelerates the rate of peptide bond formation approximately  $10^5$  to  $10^7$ -fold compared with the uncatalyzed reaction (147, 148). Detailed description of translation process can be accessed from any standard textbook.

### **3.7 ANCIENT TRANSLATION MECHISM**

Gesteland *et al.* in 1999 (149) proposed the hypothesis that translation mechanism evolved in a protein-free world. Woese in 1967 (5) argued that the ancient translation mechanism did not evolve initially to make functional proteins, because of the vanishingly small probability that the first attempts at polypeptide synthesis by a primitive translational apparatus could yield a protein with any useful enzymatic activity. Noller in 2005 (150) argued that, until the first active proteins emerged, no selective advantage would exist for evolution of translational machinery, if its only purpose were to synthesize functional proteins. Noller pointed out a possible alternative that early primitive peptides could have enhanced the functions of existing RNAs by providing additional functional groups such as positively charged or hydrophobic side chains, acting essentially as cofactors for ribozymes (151). A more subtle possibility is that primitive peptides synthesized in the RNA world could have played a “chaperone-like” role in binding to and stabilizing RNA structures (152).

## **4 THE LIPID-RNA WORLD SCENARIO**



**Figure-2:** The Lipid-RNA World scenario (A-F). Lipids are shown in gray, amino acids and nucleotides are represented as pink and green balls respectively. Description of the picture is as follows: (A) The primordial soup containing various types of biomolecules (amino acids, nucleotides and lipids), existing in Hadean ocean. (B) Lipids aggregate and forms membranes, vesicles etc.; vesicles are formed with local phases of biologically rich environment within their interior, which is susceptible for biochemical reactions and polymerizations. (C) The biologically rich environment leads to metallic ion-mediated polymerization of ribonucleic acids, thus forming functional ribozymes. A Lipid-RNA World originates. The proto-ribosome (contemporary PTC RNA) evolves (shown in green cartoon view). (D) The hypothetical protein synthesis mechanism in Lipid-RNA World by proto-ribosome is presented. Proto-ribosome is represented as contemporary PTC structure of *Thermus thermophilus* (green cartoon view). The tertiary structure of PTC RNA resembles an ‘RNA fork,’ which continuously collects amino acids emerging out from the membrane plane during rapid phase changes of lipids upon temperature fluctuations and polymerizes them (image is not according to molecular scale). See text for a detailed description. (E) The random synthesis of proteins and polymerization of nucleic acids increases compositional complexity within prebiotic vesicles. They experience exponential growth. Protein-RNA complex forms which being structurally more stable compared to RNAs are selected for evolution in a Darwinian picture. At each replicating cycle, proto-ribosome experiences a beehive-like growth being attached to the internal surface of vesicle. (F) Gradually, a self-replicating vesicle pool originates with biological complexity increasing in their interior with each repeated replicating cycle. With course of time, proteins gradually took over enzymatic functions from RNA, leading to a Protein-RNA World. Eventually these vesicles were turned into protocells (see text). We used *Thermus thermophilus* 50S ribosome structure for showing PTC RNA tertiary structure in this figure having PDB-id: 3KNI (153).

## REFERENCES

1. Oparin, A. I. (1957), The Origin of Life on the Earth, London, Oliver and Boyd.
2. Oparin, A. I., Orlovskii, A. F., Bukhlaeva, V. Y. and Gladilin, K. L. (1976), Influence of the Enzymatic Synthesis of Polyadenylic Acid on a Coacervate System, Dokl. Akad. Nauk SSSR 226, 972–974.
3. Uversky, V.N.(2002) Natively unfolded proteins: a point where biology waits for physics. Prot. Sci. 11, 739–756.
4. Ma, B., Shatsky, M., Wolfson, H.J., Nussinov, R.(2002) Multiple diverse ligands binding at a single proteinsite: a matter of pre-existing populations. Prot. Sci. 11, 184–197.
5. Woese, C. (1967) The Genetic Code, the Molecular Basis for Genetic Expression. New York: Harper and Row.

6. Crick, F.H.C. (1968) The origin of the genetic code. *J. Mol. Biol.*, 38:367–379.
7. Orgel, L.E. (1968). Evolution of the genetic apparatus. *J. Mol. Biol.*, 38:381–393.
8. Orgel, L. and Sulston, J. (1971). Polynucleotide replication and the origin of life. In *Prebiotic and Biochemical Evolution*, pp. 89–94. Kimball, A., and Oro, J., Eds., Amsterdam: North-Holland Publishing Company.
9. White, H.B., 3rd. (1976) Coenzymes as fossils of an earlier metabolic state. *J Mol Evol* 7:101–104.
10. Kruger, K., Grabowski, P.J., Zaug, A.J., Sands, J., Gottschling, D.E., and Cech, T.R. (1982). Self-splicing RNA: autoexcision and autocyclization of the ribosomal RNA intervening sequence of *Tetrahymena*. *Cell* 31:147–157.
11. Guerrier-Takada, C., Gardiner, K., Marsh, T., Pace, N., and Altman, S. (1983) The RNA moiety of ribonuclease P is the catalytic subunit of the enzyme. *Cell* 35:849–857.
12. Gilbert, W. (1986) The RNA World. *Nature* 319:618.
13. Gesteland, R.F., Cech, T.R., and Atkins, J.F. (1999) *The RNA World*, 2nd ed. Cold Springs Harbor, NY: Cold Springs Harbor Press.
14. Eigen, M. and Schuster, P. (1982), *Stages of Emerging Life-Five Principles of Early Organization*, *J. Mol. Evol.* 19(1), 47–61.
15. Beaudry, A. A. and Joyce, G. F. (1992), Directed Evolution of an RNA Enzyme, *Science* 342, 255–257.
16. Wilson, C. and Szostak, J. W. (1994), In vitro Evolution of a Self-Akylating Ribozyme, *Nature* 374, 777–782.
17. Steitz, T.A. and Moore, P.B. (2003) RNA, the first macromolecular catalyst: the ribosome is a ribozyme. *Trends Biochem Sci* 28:411–418.
18. Forterre, P., Grosjean, H. (2009). The interplay between RNA and DNA modifications: back to the RNA world. In *DNA and RNA Modification enzymes: Structure, Mechanism, Function and Evolution*. Edited by Grosjean H. Austin: Landes Bioscience, 259–274.
19. Steitz, T.A. and Moore, P.B. (2003) RNA, the first macromolecular catalyst: the ribosome is a ribozyme. *Trends Biochem Sci* 28:411–418.
20. Koglin, A., Walsh, C. T. (2009) Structural insights into nonribosomal peptide enzymatic assembly lines. *Nat Prod Rep* 2009, 26:987–1000.
21. Strieker, M., Tanovic, A., Marahiel, M. A. (2010). Nonribosomal peptide synthetases: structures and dynamics. *Curr Opin Struct Biol* 2010, 20:234–240.
22. Mills, G. C., Kenyon, D. (Retrieved 10 Sept 2011). *The RNA World: A Critique*, *Origins & Design* 17:1.
23. Bernhardt, H. S. (2012). The RNA world hypothesis: the worst theory of the early evolution of life (except for all the others), *Biol Direct*;7(1):23.
24. Shapiro, R. (1984). The improbability of prebiotic nucleic acid synthesis, *Origins of Life*, 14: 565-570.
25. Shapiro, R. (1988). Prebiotic ribose synthesis: a critical analysis, *Origins of Life*, 18: 71-85.
26. Larralde, R., Robertson, M. P., and Miller, S. L. (1995). Rates of decomposition of ribose and other sugars: Implications for chemical evolution, *Proc. Natl. Acad. Sci. USA* 92: 8158-8160.
27. Knauth, L.P., Lowe, D.R. (1978). Oxygen isotope geochemistry of cherts from the Onverwacht Group (3.4 billion years), Transvaal Group, South Africa, with implications for secular variations in the isotopic composition of cherts. *J. Geol.*, 41, 209–222.
28. Tajika, E., Matsui, T. (1993). Degassing history and carbon cycle of the Earth: From an impact-induced steamatmosphere to the present atmosphere. *Lithos*, 30, 267–280.
29. de Ronde, C.E.J., Channer, D.M.D.R., Faure, K., Bray, C. J., Spooner, T. C. (1997). Fluid chemistry of Archaean seafloor hydrothermal vents: Implications for the composition of circa 3.2 Ga seawater. *Geochim. Cosmochim. Acta*, 61, 4025–4042.
30. Shapiro, R. (1995). The prebiotic role of adenine: a critical analysis, *Origins of Life and the Evolution of the Biosphere* 25: 83-98
31. Sanchez, R. A., Orgel, L. E. (1970). Studies in prebiotic synthesis: V. Synthesis and photoanomerization of pyrimidine nucleosides. *J Mol Biol*, 47:531–543.
32. Powner, M. W., Gerland, B., Sutherland, J. D. (2009). Synthesis of activated pyrimidine ribonucleotides in prebiotically plausible conditions. *Nature* 2009, 459:239–242.
33. Orgel, L. E., (2004). Prebiotic Chemistry and the Origin of the RNA World, *Critical Reviews in Biochemistry and Molecular Biology*, 39:99–123, 2004.
34. Sagan, C., Mullen, G. (1972). Earth and Mars: Evolution of Atmospheres and Surface Temperatures, *Science, New Series*, Vol. 177, No. 4043, pp. 52-56.
35. Valley, J. W., Peck, W. H., King, E. M., Wilde, S. A. (2002). A cool early Earth, *Geology*, v. 30, no. 4, p. 351–354
36. Valley, J. W. (2005). A cool early Earth? *Sci. American*, 293(4):58-65.
37. Sleep, N.H., Zahnle, K. (2001). Carbon dioxide cycling and implications for climate on ancient Earth. *J. Geophys. Res.*, 106, 1373–1399.

38. Goldblatt, C., Zahnle, K. J. (2011). Clouds and the Faint Young Sun Paradox, *Clim. Past*, 7, 203–220
39. Rosing, M. T., Bird, D. K., Sleep, N.H. and Bjerrum, C. J. No climate paradox under the faint early Sun. *Nature* 464, 744–747 (2010).
40. Schlesinger, G., Miller, S. L. (1983). Prebiotic synthesis in atmospheres containing CH<sub>4</sub>, CO, and CO<sub>2</sub>. I. Amino acids. *J Mol Evol.*;19(5):376-82.
41. Stribling, R. and S. L. Miller. (1987). Energy yields for hydrogen cyanide and formaldehyde syntheses: the HCN and amino acid concentrations in the primitive ocean. *Origins of Life and Evolution of the Biosphere* 17: 261-273.
42. Goldblatt, C., Zahnle, K. J. (2011). Faint young Sun paradox remains. *Nature*,474(7349):E3-4, discussion E4-5.
43. Bada, J. L., Bigham, C., Miller, S. L. (1994). Impact melting of frozen oceans on the early Earth: implications for the origin of life. *Proc Natl Acad Sci U S A*, 91:1248-50.
44. Hartmann, W.K., Ryder, G., Dones, L., and Grinspoon, D., (2000). The time-dependent intense bombardment of the primordial Earth/Moon system, in Canup, R.M., and Righter, K., eds., *Origin of the Earth and Moon*: Tucson, University of Arizona Press, p. 493–512.
45. Moravek, J. (1967). Formation of oligonucleotides during heating of a mixture of uridine 2'(3')-phosphate and uridine. *Tetrahedron Lett* 18:1707–1710.
46. Lohrmann, R. (1977). Formation of nucleoside 5'-phosphoramidates under potentially prebiological conditions. *J Mol Evol* 10:137–154.
47. Sawai, H. and Orgel, L. E.(1975). Oligonucleotide synthesis catalyzed by the Zn<sup>2+</sup> ion. *J Am Chem Soc* 97:3532-3533.
48. Sawai, H. (1976). Catalysis of internucleotide bond formation by divalent metal ions. *J Am Chem Soc* 98:7037–7039.
49. Sleeper, H. L. and Orgel, L. E. (1979). The catalysis of nucleotide polymerization by compounds of divalent lead. *J Mol Evol* 12:357–364.
50. Kanavarioti, A., Monnard, P. A., and Deamer, D. W. (2001). Eutectic phases in ice facilitate nonenzymatic nucleic acid synthesis. *Astrobiology* 1:271–281.
51. Sawai, H., Karoda, K., and Hojo, T. (1989). Uranyl ion as a highly effective catalyst for internucleotide bond formation. *Bull Chem Soc Jpn* 62:2018–2023.
52. Sawai, H., Higa, K., and Kuroda, K. (1992). Synthesis of cyclic and acyclic oligocytidylates by uranyl catalysis in aqueous solution. *Chem Soc Perkin Trans* 1:505–508.
53. Ferris, J.P., Joshi, P.C., Wang, K.J., Miyakawa, S., and Huang, W. (2004). Catalysis in prebiotic chemistry: application to the synthesis of RNA oligomers. *Advances in Space Research*, 33, 100–105.
54. Ertem, G. and Ferris, J.P. (1998). Formation of RNA oligomers on montmorillonite: site of catalysis. *Orig Life Evol Biosph* 28:485–499.
55. Wang, K.J. and Ferris, J.P. (2001). Effect of inhibitors on the montmorillonite clay-catalyzed formation of RNA: studies on the reaction pathway. *Orig Life Evol Biosph* 31:381–402.
56. Orgel, L.E. (2003). Some consequences of the RNA world hypothesis. *Orig Life Evol Biosph* 33:211–218.
57. Howard, F.B., Frazier, J., Singer, M.F., and Miles, H.T. (1966). Helix formation between polyribonucleotides and purines, purine nucleosides and nucleotides. II. *J Mol Biol* 16:415–439.
58. Naylor, R. and Gilham, P.T. (1966). Studies on some interactions and reactions of oligonucleotides in aqueous solution. *Biochemistry* 5:2722–2728.
59. Shabarova, Z.A. (1988). Chemical development in the design of oligonucleotide probes for binding to DNA and RNA. *Biochimie* 70:1323–1334.
60. Lohrmann, R. and Orgel, L.E. (1980). Efficient catalysis of polycytidylic acid-directed oligoguanylate formation by Pb<sup>2+</sup>. *J Mol Biol* 142:555–567.
61. Bridson, P.K. and Orgel, L.E. (1980). Catalysis of accurate poly(C)- directed synthesis of 3'-5'-linked oligoguanylates by Zn<sup>2+</sup>. *J Mol Biol* 144:567–577.
62. Singh, P. P., Banerji, A., (2011). Case for an RNA–prion world: a hypothesis based on conformational diversity, *J Biol Phys* (2011) 37:185–188.
63. Kanavarioti, A., Monnard, P. A., Deamer, D. W. (2001). Eutectic phases in ice facilitate nonenzymatic nucleic acid synthesis. *Astrobiology* 2001, 1:271–281.
64. Vlassov, A. V., Kazakov, S. A., Johnston, B. H., Landweber, L. F. (2005). The RNA world on ice: a new scenario for the emergence of RNA information. *J Mol Evol* 2005, 61:264–273.



65. Kazakov, S. A., Balatskaya, S. V., Johnston, B. H. (2006). Ligation of the hairpin ribozyme in cis induced by freezing and dehydration. *RNA* 2006, 12:446–456.
66. Attwater, J., Wochner, A., Pinheiro, V. B., Coulson, A., Holliger, P. (2010). Ice as a protocellular medium for RNA replication. *Nat Commun* 2010, 1:76.
67. Bernhardt, H. S., Tate, W. P. (2012). Primordial soup or vinaigrette: did the RNA world evolve at acidic pH? *Biol Direct*, 7:4.
68. Lindahl, T. (1993). Instability and decay of the primary structure of DNA. *Nature* 362(6422), 709–715.
69. Cleaves, H. J., Miller, S. L. (1998). Oceanic protection of prebiotic organic compounds from UV radiation, *Proc Natl Acad Sci U S A*. 1998 Jun 23,95(13):7260-3.
70. Wochner, A., Attwater, J., Coulson, A., Holliger, P. (2011). Ribozyme-catalyzed transcription of an active ribozyme. *Science*, 332:209–212.
71. Turk, R. M., Chumachenko, N. V., Yarus, M. (2010). Multiple translational products from a fivenucleotide ribozyme. *Proc Natl Acad Sci USA*, 107:4585–4589.
72. Bernhardt, H. S. (2012). The RNA world hypothesis: the worst theory of the early evolution of life (except for all the others), *Biol Direct*;7(1):23.
73. Egel, R. (2009). Peptide-dominated membranes preceding the genetic takeover by RNA: latest thinking on a classic controversy. *Bioessays* 2009, 31:1100–1109.
74. Kurland, C. G. (2010). The RNA dreamtime: modern cells feature proteins that might have supported a prebiotic polypeptide world but nothing indicates that RNA world ever was. *Bioessays* 2010, 32:866–871.
75. Caetano-Anollés, D., Kim, K. M., Mittenthal, J. E., Caetano-Anollés, G. (2011). Proteome evolution and the metabolic origins of translation and cellular life. *J Mol Evol* 2011, 72:14–33.
76. Caetano-Anollés, G., Kim, K. M., Caetano-Anollés, D. (2012). The phylogenomic roots of modern biochemistry: origins of proteins, cofactors and protein biosynthesis. *J Mol Evol* 2012, 74:1–34.
77. Harish, A., Caetano-Anollés, G. (2012). Ribosomal history reveals origins of modern protein synthesis. *PLoS One* 2012, 7:e32776.
78. Segré, D., Ben-Eli, D., Deamer, D. W., and Lancet, D. (2001). The Lipid World, *Origins of Life and Evolution of the Biosphere* 31: 119–145.
79. Halliday, A. N. (2000). Terrestrial accretion rates and the origin of the Moon, *Earth and Planetary Science Letters*, Volume 176, Issue 1, 28 February 2000, Pages 17–30
80. Mackenzie, D. (2003). *The Big Splat, or How The Moon Came To Be*. John Wiley and Sons. ISBN 978-0-471-15057-2.
81. Brown, H. (1952). Rare gases and the formation of the Earth's atmosphere, *The Atmospheres of the Earth and Planets*. University of Chicago Press, Chicago, IL, 2nd edn., pp. 258-266.
82. Staudacher T. and Allegre C.J. (1982). Terrestrial xenology. *Earth Planet. Sci. Lett.* 60, 389-406.
83. Allegre C.J., Staudacher T., Sarda P. and Kurz M. (1983): Constraints on evolution of Earth's mantle from rare gas systematics. *Nature* 303, 762-766.
84. Podesk, F. A. (1970). Dating of meteorites by the high-temperature release of iodine-correlated Xe 129. *Geochim. Cosmochim. Acta* 34, 341-365.
85. Russell, M.J., Arndt, N.T., (2005). Geodynamic and metabolic cycles in the Hadean. *Biogeosciences* 2, 97–111.
86. Takai, K., Nakamura, K., Suzuki, K., Inagaki, F., Nealson, K.H., Kumagai, H., (2006). Ultramafics–Hydrothermalism–Hydrogenesis–HyperSLiME (UltraH3) linkage: a key insight into early microbial ecosystem in the Archean deep-sea hydrothermal systems. *Paleontological Research* 10, 269–282.
87. Russell, M.J., (2007). The alkaline solution to the emergence of life: energy, entropy and early evolution. *Acta Biotheoretica* 55, 133–179.
88. Russell, M.J., Daniel, R.M., Hall, A.J., (1993). On the emergence of life via catalytic iron-sulphide membranes. *Terra Nova* 5, 343–347.
89. Martin, W., Baross, J., Kelley, D., Russell, M.J., (2008). Hydrothermal vents and the origin of life. *Nature Reviews Microbiology* 6, 805–814.
90. Kelley, D.S., Karson, J.A., Blackman, D.K., Früh-Green, G.L., Butterfield, D.A., Lilley, M.D., Olson, E.J., Schrenk, M.O., Roe, K.K., Lebon, G.T., Rivizzigno, P., Party, t.A.-S., (2001). An off-axis hydrothermal vent field near the Mid-Atlantic Ridge at 30°N. *Nature* 412, 145–149.
91. Kelley, D.S., Karson, J.A., Früh-Green, G. L., Yoerger, D.R., Shank, T.M., Butterfield, D.A., Hayes, J.M., Schrenk, M.O., Olson, E.J., Proskurowski G., Jakuba, M., Bradley, A., Larson, B., Ludwig, K., Glickson, D., Buckman, K., Bradley, A.S., Brazelton, W.J., Roe, K.,

- Elend, M.J., Delacour, A., Bernasconi, S.M., Lilley, M.D., Baross, J.A., Summons, R.E., Sylva, S.P., (2005). A serpentinite-hosted ecosystem: the Lost City hydrothermal field. *Science* 307, 1428–1434.
92. Shibuya, T., Komiya, T., Nakamura, K., Takai, K., Maruyama, S. (2010). Highly alkaline, high temperature hydrothermal fluids in the early Archean ocean, *Precambrian Research*, 182: 230-238.
93. Seyfried Jr., W.E., Ding, K., (1995). Phase equilibria in seafloor hydrothermal systems: a review of the role of redox, temperature, pH and dissolved Cl on the chemistry of hot spring fluids at mid-ocean ridges. *Seafloor Hydrothermal Systems: Physical, Chemical, Biological, and Geological Interactions. Geophysical Monograph. American Geophysical Union, Washington, DC*, pp. 248–272.
94. Seyfried Jr., W.E., Ding, K., Berndt, M.E., (1991). Phase equilibria constraints on the chemistry of hot spring fluids at mid-ocean ridge. *Geochimica et Cosmochimica Acta* 55, 3559–3580.
95. Berndt, M.E., Seyfried Jr., W.E., (1993). Calcium and sodium exchange during hydrothermal alteration of calcic plagioclase at 400 °C and 400 bars. *Geochimica et Cosmochimica Acta* 57, 4445–4451.
96. Ding, K., Seyfried Jr., W.E., Zhang, Z., Tivey, M.K., Von Damm, K.L., Bradley, A.M., (2005). The in situ pH of hydrothermal fluids at mid-ocean ridge. *Earth and Planetary Science Letters* 237, 167–174.
97. Kitajima, K., Maruyama, S., Utsunomiya, S., Liou, J.G., (2001). Seafloor hydrothermal alteration at an Archean mid-ocean ridge. *Journal of Metamorphic Geology* 19, 583–599.
98. Terabayashi, M., Masuda, Y., Ozawa, H., (2003). Archean ocean-floor metamorphism in the North Pole area, Pilbara Craton, Western Australia. *Precambrian Research* 123, 167–180.
99. Nakamura, K., Kato, Y., (2004). Carbonatization of oceanic crust by the seafloor hydrothermal activity and its significance as a CO<sub>2</sub> sink in the Early Archean. *Geochimica et Cosmochimica Acta* 68, 4595–4618.
100. Shibuya, T., Kitajima, K., Komiya, T., Terabayashi, M., Maruyama, S., (2007). Middle Archean ocean ridge hydrothermal metamorphism and alteration recorded in the Cleaverville area, Pilbara Craton, Western Australia. *Journal of Metamorphic Geology* 25, 751–767.
101. Corliss, J.B., Dymond, J., Gordon, L.I., Edmond, J.M., Von Herzen, R.P., Ballard, R.D., Green, K., Williams, D., Bainbridge, A., Crane, K., van Andel, T.H. et al. (1979). Submarine thermal springs on the Galapagos rift. *Science* 203, 1073–1083.
102. Spiess, F.N., Macdonald, K.C., Atwater, T., Ballard, R., Carranza, A., Cordoba, D., Cox, C., Garcia, V.M., Francheteau, J., Guerrero, J. et al. (1980). East Pacific rise: hot springs and geophysical experiments. *Science* 207, 1421–1433.
103. Delaney, J.R., Kelley, D.S., Lilley, M.D., Butterfield, D.A., Baross, J.A., Wilcock, W.S.D., Embley, R.W., Summit, M. (1998). The Quantum Event of Oceanic Crustal Accretion: Impacts of Diking at Mid-Ocean Ridges, *Science*. 1998 Jul 10;281(5374):222-30.
104. von Damm, K.L., Lilley, M.D., Shanks, W.C., Brockington, M., Bray, A.M., O'Grady, K.M., Olson, E., Graham, A., Proskurowski, G. (2003). Extraordinary phase separation and segregation in vent fluids from the southern East Pacific Rise, *Earth and Planetary Science Letters*, Volume 206, Issue 3-4, p. 365-378.
105. Von Damm, K. L. in *Physical, Chemical, Biological, and Geological Interactions within Seafloor Hydrothermal Systems* (eds Humphris, S., Zierenberg, R., Mullineau, L. & Thomson R.) 222–247 (American Geophysical Union, Washington DC, 1995).
106. Kelley, D.S., Baross, J.A. and Delaney, J.R. (2002). Volcanoes, fluids, and life at mid-ocean ridge spreading centers. *Annu. Rev. Earth Planet Sci.* 30, 385–491 .
107. Martin, W. and Russell, M. J. (2007). On the origin of biochemistry at an alkaline hydrothermal vent. *Philos. Trans. R. Soc. Lond. B* 367, 1187–1925.
108. Russell, M. J. and Hall, A. J. (1997). The emergence of life from iron monosulphide bubbles at a submarine hydrothermal redox and pH front. *J. Geol. Soc. London* 154, 377–402.
109. Russell, M. J., Daniel, R. M., Hall, A. J. and Sherringham, J. (1994). A hydrothermally precipitated catalytic iron sulphide membrane as a first step toward life. *J. Mol. Evol.* 39, 231–243.
110. Ueno, Y., Johnson, M. S., Danielache, S. O., Eskebjerg, C., Pandey, A., Yoshida, N. (2009). Geological sulfur isotopes indicate elevated OCS in the Archean atmosphere, solving faint young sun paradox, *Proc Natl Acad Sci U S A.*;106(35):14784-9.
111. Pross, A. (2005). On the emergence of biological complexity: life as a kinetic state of matter, *Origins of Life and Evolution of Biospheres* (2005) 35: 151–166
112. Eigen, M. (1971). Self-Organization of Matter and the Evolution of Biological Macromolecules, *Naturwissenschaften* 58, 465–523.
113. Mushegian, A. (2005). Protein content of minimal and ancestral ribosome, *RNA*, 11:1400–1406.
114. Woese, C. (1998). The universal ancestor, *Proc. Natl. Acad. Sci. USA*. 1998 June 9; 95(12): 6854–6859.

115. Roberts, E., Sethi, A., Montoya, J., Woese, C. R., Luthey-Schulten, Z. (2008). Molecular signatures of ribosomal evolution, *Proc Natl Acad Sci U S A*;105(37):13953-8.
116. Vishwanath, P., Favaretto, P., Hartman, H., Mohr, S. C., Smith, T. F. (2004). Ribosomal protein-sequence block structure suggests complex prokaryotic evolution with implications for the origin of eukaryotes. *Mol Phylo Genet Evol* 33:615–625.
117. Fox, G. E., Naik, A. K. (2004). The evolutionary history of the ribosome, In *The genetic code and the origin of life* (Ribas de Pouplana L. ed), Landes Bioscience Chapter 6, pp 92–105.
118. Fox, G. E. (2010). Origin and Evolution of the Ribosome, *Cold Spring Harb Perspect Biol.*;2(9):a003483.
119. Smith, T. F., Lee, J. C., Gutell, R. R., Hartman, H. (2008). The origin and evolution of the ribosome, *Biol Direct*. 2008 Apr 22;3:16.
120. Brodersen, D. E., Nissen, P. (2005). The social life of ribosomal proteins, *FEBS J.*;272(9):2098-108.
121. Nirenberg, M., Leder, P. (1964). RNA code words and protein synthesis. *Science* 145: 1399–1407.
122. Lill, R., Robertson, J. M., Wintermeyer, W. (1986). Affinities of tRNA binding sites of ribosomes from *Escherichia coli*. *Biochemistry* 25: 3245–3255.
123. Pestka, S. (1968). Studies on the formation of transfer ribonucleic acid-ribosome complexes. 3. The formation of peptide bonds by ribosomes in the absence of supernatant enzymes. *J Biol Chem* 243: 2810–2820.
124. Gavrillova, L. P., Kostiaschkina, O. E., Koteliansky, V. E., Rutkevitch, N. M., Spirin, A. S. (1976). Factor-free (“non-enzymic”) and factor-dependent systems of translation of polyuridylic acid by *Escherichia coli* ribosomes. *J Mol Biol* 101: 537–552.
125. Nomura, M., Mizushima, S, Ozaki M., Traub, P., Lowry, C. V. (1969). Structure and function of ribosomes and their molecular components. *Cold Spring Harbor Symp Quant Biol* 34: 49–61.
126. Hoang, L., Fredrick, K., Noller, H. F. (2004). Creating ribosomes with an all-RNA 30S subunit P site. *Proc Natl Acad Sci* 101: 12439–12443.
127. Noller, H.F. (2012). Evolution of protein synthesis from an RNA world, *Cold Spring Harb Perspect Biol.*;4(4).
128. Davies, J., Nomura, M. (1972). The genetics of bacterial ribosomes. *Ann Rev Genet* 6: 203–234.
129. Kurland, C.G., Jorgensen, F., Richter, A. A., Ehrenberg, M., Bilgin, N., Rojas, A. M. (1990). Through the Accuracy Window. in *The Ribosome: Structure, function, and evolution*, pp. 513–526. American Society of Microbiology, Washington, D.C.
130. Clark, C.G. (1987). On the evolution of ribosomal RNA. *J. Mol. Evol.* 25: 343–350.
131. Gray, M. W., Schnare, M. N. (1996). Evolution of rRNA gene organization, in *Ribosomal RNA Structure, Evolution, Processing, and Function in Protein Biosynthesis* (eds R.A. Zimmerman, and A.E. Dahlberg), CRC Press, Boca Raton FL. pp49–69.
132. Wuyts, J., Van de Peer, Y., De Wachter, R. (2001). Distribution of substitution rates and locations of insertion sites in the tertiary structure of ribosomal RNA. *Nucleic Acids Res* 29: 5017–5028.
133. Mears, J. A., Cannone, J. J., Stagg, S. M., Gutell, R. R., Agrawal, R. K., Harvey, S. C. (2002). Modeling a minimal ribosome based on comparative sequence analysis. *J Mol Biol* 321: 215–234.
134. Caetano-Anolles G. (2002). Tracing the evolution of RNA structure in ribosomes. *Nucleic Acids Res* 30: 2575–2587.
135. Hury, J., Nagaswamy, U., Larios-Sanz, M., Fox, G. E. (2006). Ribosome origins: The relative age of 23S rRNA domains. *Orig Life Evol Biosphere* 36: 421–429.
136. Ban, N., Nissen, P., Hansen, J., Moore, P. B., Steitz, T. A. (2000). The complete atomic structure of the large ribosomal subunit at 2.4 Å resolution, *Science*;289(5481):905-20.
137. Wimberly, B. T., Brodersen, D. E., Clemons, W. M. Jr., Morgan-Warren, R. J., Carter, A. P., Vonrhein, C., Hartsch, T., Ramakrishnan, V. (2000). Structure of the 30S ribosomal subunit, *Nature*;407(6802):327-39.
138. Khaitovich, P. and Mankin, A.S. (1999). Effect of antibiotics on large ribosomal subunit assembly reveals possible function of 5 S rRNA. *J Mol Biol* 291:1025–1034.
139. Polacek, N., Mankin, A. (2005). The ribosomal peptidyl transferase center: Structure, function, evolution, inhibition. *Crit Rev Biochem Mol Biol*, 40(5):285-311.
140. Weiss-Brummer, B., Zollner, A., Haid, A., and Thompson, S. (1995). Mutation of a highly conserved base in the yeast mitochondrial 21S rRNA restricts ribosomal frameshifting. *Mol Gen Genet* 248:207–216.
141. O’Connor, M., Brunelli, C.A., Firpo, M.A., Gregory, S.T., Lieberman, K.R., Lodmell, J.S., Moine, H., Van Ryk, D.I., and Dahlberg, A.E. (1995). Genetic probes of ribosomal RNA function. *Biochem Cell Biol* 73:859–868.
142. Thompson, J., Kim, D.F., O’Connor, M., Lieberman, K.R., Bayfield, M.A., Gregory, S.T., Green, R., Noller, H.F., and Dahlberg, A.E. (2001). Analysis of mutations at residues A2451 and G2447 of 23S rRNA in the peptidyltransferase active site of the 50S ribosomal subunit. *Proc Natl Acad Sci USA* 98:9002–9007.
143. Thompson, J., O’Connor, M., Mills, J.A., and Dahlberg, A.E. (2002). The protein synthesis inhibitors, oxazolidinones and chloramphenicol, cause extensive translational inaccuracy in vivo. *J Mol Biol* 322:273–279.

144. Hsiao C, Williams LD. 2009. A recurrent magnesium-binding motif provides a framework for the ribosomal peptidyl transferase center. *Nucl Acids Res* 37: 3134–3142.
145. Hsiao C, Mohan S, Kalahar BK, Williams LD. 2009. Peeling the onion: Ribosomes are ancient molecular fossils. *Mol Biol Evol* 26: 2415–2425.
146. Schuwirth, B.S., Borovinskaya, M.A., Hau, C.W., Zhang, W., Vila-Sanjurjo, A., Holton, J.M., Cate, J.H. (2005). Structures of the bacterial ribosome at 3.5 Å resolution, *Science* 310: 827–834.
147. Krayevsky, A.A. and Kukhanova, M.K. 1979. The peptidyltransferase center of ribosomes. *Prog Nucleic Acid Res Mol Biol* 23:1–51.
148. Nierhaus, K.H., Schulze, H., and Cooperman, B.S. (1980). Molecular mechanisms of the ribosomal peptidyltransferase center. *Biochemistry International* 1:185–182.
149. Sievers, A., Beringer, M., Rodnina, M.V., and Wolfenden, R. (2004). The ribosome as an entropy trap. *Proc Natl Acad Sci USA* 101:7897–7901.
150. Noller, H. F. (2004). The driving force for molecular evolution of translation, *RNA*;10(12):1833–7.
151. Szathmáry, E. and Maynard Smith, J. (1997). From replicators to reproducers: The first major transitions leading to life. *J. Theor. Biol.* 187: 555–571.
152. Poole, A.M., Jeffares, D.C., and Penny, D. (1998). The path from the RNA world. *J. Mol. Evol.* 46: 1–17.
153. Stanley, R.E., Blaha, G., Grodzicki, R.L., Strickler, M.D., Steitz, T.A. (2010). The structures of the anti-tuberculosis antibiotics viomycin and capreomycin bound to the 70S ribosome, *Nat. Struct. Mol. Biol.* 17: 289–293.