Continuous intracavity monitoring of the dynamical Casimir effect

A V Dodonov

Instituto de Física and International Center for Condensed Matter Physics, Universidade de Brasília, PO Box 04455, 70910-900, Brasília, DF, Brazil ITAMP, Harvard–Smithsonian Center for Astrophysics, Cambridge, MA 02138, USA

Dynamical Casimir effect (DCE) is the name assigned to the process of generating quanta from vacuum due to an accelerated motion of macroscopic neutral bodies (mirrors) or time-modulation of cavity material properties, as well as the simulation of such processes. Here I review the theoretical results on the detection of DCE using intracavity quantum detectors, such as multi-level atoms, atomic networks and harmonic oscillators. I also stress the mathematical equivalence of this problem to the physics of optical parametric oscillators interacting with atoms or quantum wells, studied in Quantum Nonlinear Optics.

PACS numbers: 42.50.Pq, 32.80.-t, 42.50.Ct, 42.50.Hz

I. INTRODUCTION

Dynamical Casimir effect (DCE) is the term used nowadays for a rather wide group of phenomena whose common feature is the creation of quanta from the initial vacuum state of some field due to time-modulation of material properties or boundary conditions of some macroscopic system. In the majority of cases considered so far this corresponds to the creation of photon pairs from the electromagnetic vacuum due to the motion of a mirror in empty space, oscillation of a cavity wall or modulation of the dielectric properties of the medium inside it. Non-electromagnetic analogs of the DCE are also possible, such as the recent report on generating correlated pairs of elementary excitations from the initial thermal state in a trapped Bose-Einstein condensate (BEC) [1], where phonons were created instead of photons. For short reviews on the history and recent achievements in the DCE research see [2, 3] and references therein.

The detection, characterization and, ultimately, manipulation of excitations created due to the DCE are as important as the very generation from vacuum. For photons generated in empty space or in cavities one can distinguish at least four different measurement schemes. In the *extracavity detection* the photons propagate away from the source (mirror or leaky cavity) and are ultimately detected using standard optical or microwave techniques. This approach was discussed for photon emission from a semiconductor microcavity [4] and superconducting resonator [5] due to time-modulation of the Rabi frequency. For superconducting coplanar waveguides terminated by a quantum interference device (SQUID) the measurement setups were discussed in [3, 6– 8], and the experimental detection of radiation produced due to an analog of a moving mirror was described in [9]. Preliminary results on the detection of DCE photons produced in a lossy resonator composed of Josephson metamaterial were described in [10].

For DCE implemented in high-Q cavities a natural al-



FIG. 1: Artistic view of the continuous intracavity monitoring of DCE (illustration by *Porfírio*). A slow unexcited atom (quantum detector) crosses the cavity whose frequency is time-modulated by an experimentalist who hits its wall at a constant rate. The photon generation from vacuum is monitored by continuously measuring the atomic state using some external measurement device (in the picture the atomic spontaneous emissions are registered by an antenna). Reproduced from http://iopscience.iop.org/0953-4075/labtalk-article/48481.

ternative is to detect the photons inside the resonator prior to their escape. In what I call the *posterior intracavity detection* the detector is inserted inside the cavity (or turned on) after the time-modulation of the cavity properties ceased. One of advantages of this approach is that the photon statistics can be determined at chosen times, but the detector timing must be accurate to avoid disturbing the generation process and loosing photons due to damping. There are detailed analyzes of detection of DCE photons in microwave cavities employing antennas [11, 12], ensembles of population-inverted alkali-metal atoms [13, 14], single Rydberg atoms [15] and superconducting qubits [16]. A proposal based on measuring the change in kinetic energy of electrons passing through the cavity was given in [17].

In the continuous intracavity monitoring the intracavity detector is active from the very beginning of the timemodulation, as illustrated by the artistic view in figure 1. The upside is that no complex timing techniques are required, however the detector may disturb significantly the photon generation process or even completely inhibit it if not properly accounted for. This line started with papers [18, 19] for harmonic time-modulation of the cavity frequency, where the detector was modeled either as a 2-level atom or a harmonic oscillator. In [20-22] nonharmonic perturbations in the presence of a 2-level atom were analyzed. A simultaneous implementation and detection of a DCE analog in superconducting circuits using an artificial 2-level atom with time-varying transition frequency or coupling parameter was discussed in [23, 24]. Quite recently new results were found for the harmonic modulation of cavity frequency in the presence of lossy multi-level atoms [25–30] and harmonic oscillators [31– 33].

Finally, the method of *back-action detection* aims monitoring the back-action of the DCE on the external agent responsible for the time-modulation of the system parameters. For example, the energy of the photons created due to a moving boundary comes from the mechanical energy, so the wall experiences a friction force by the quantum vacuum. Reference [34] analyzed this approach in the case of DCE induced by the Rabi oscillations of a 3-level emitter illuminated by a strong laser while coupled to the cavity field mode, and it was shown that a suppression of the absorption by the emitter can pinpoint the DCE.

In this paper I review the recent results on the continuous intracavity monitoring of DCE, when the cavity field interacts coherently with a multi-level quantum detector that can be coupled to an external (classical) measurement device in order to read out its state. It will be shown that the presence of quantum detector leads to novel regimes of photon generation, so besides detecting the Casimir photons one could produce novel cavity field states by parametrically tuning the periodicity of modulation and applying post-selection techniques. I shall also point out the mathematical equivalence of DCE in the presence of detectors to the research in the area of Quantum Nonlinear Optics (QNO), where nonlinear pumped crystals interact with multi-level atoms and quantum wells.

This paper is organized as follows. The formalism of continuous intracavity monitoring is outlined in section II, and the modeling of the detector as N-level ladder atom is described in section III. The relationship of this topic to the area of QNO is outlined in section IV. The main results and discussion for different values of N and for networks of 2-level atoms are given in section V. Finally, the sections VI and VII contain some perspectives of future developments and the summary of this brief review.

II. CONTINUOUS INTRACAVITY MONITORING

I shall consider the simplest case of a single resonant cavity mode whose angular frequency ω_t is rapidly modulated in time around its bare frequency ω_0 according to the harmonical law $\omega_t = \omega_0 + \varepsilon \sin(\eta t)$ with a small modulation depth $|\varepsilon| \ll \omega_0$. The Hamiltonian describing this cavity DCE implementation is [35] (I set $\hbar = 1$)

$$\hat{H}_c = \omega_t \hat{n} + i\chi_t (\hat{a}^{\dagger 2} - \hat{a}^2), \tag{1}$$

where $\chi_t = (4\omega_t)^{-1} d\omega_t / dt$ is a so-called squeezing coefficient, \hat{a} and \hat{a}^{\dagger} are the cavity annihilation and creation operators satisfying the bosonic commutation relation $[\hat{a}, \hat{a}^{\dagger}] = 1$, and $\hat{n} = \hat{a}^{\dagger}\hat{a}$ is the photon number operator. In the presence of intracavity quantum detector the total Hamiltonian is $\hat{H}' = \hat{H}_c + \hat{H}_d$, where \hat{H}_d denotes the detector free Hamiltonian plus the detectorfield interaction. To find the dynamics one has to solve the von Neumann equation $i\partial \hat{\rho}'/\partial t = [\hat{H}', \hat{\rho}']$ for the total density operator $\hat{\rho}'$ of the detector-field system. Furthermore, due to the detector-field entanglement the detector may serve to produce novel cavity field states via postselection procedures. If the information about the detector state is completely discarded, then the field state at time t is simply $\hat{\rho}'_{f}(t) = \operatorname{Tr}_{d}[\hat{\rho}'(t)]$, where Tr_{d} denotes the partial trace operation over the detector's degrees of freedom. If the detector state is instead read in a single-shot projective measurement at the time t, with the outcome described by the projector \hat{P}_d on the detector's Hilbert space, then the cavity field state collapses to the state $\hat{\rho}_f'(t_+) = \text{Tr}_d[\hat{P}_d\hat{\rho}'(t)]/\text{Tr}[\hat{P}_d\hat{\rho}'(t)], \text{ and } \text{Tr}[\hat{P}_d\hat{\rho}'(t)] \text{ gives}$ the probability for this outcome.

When some external measurement device (EMD) continuously reads out the detector state, the time evolution of the field-detector system becomes non-unitary as the system is now open. The dynamics can then be described by the Continuous Photodetection Model [36] or closely related Quantum Trajectories approach [37].When the EMD emits a "click" as a result of absorbing one excitation from the detector, it disturbs the detector-field system by promoting the detector to a lower energy state. Formally this effect is described by the action of the "Quantum Jump superoperator" (QJS) \hat{J} on the density operator $\hat{\rho}'$, and the probability of a click during the infinitesimal time interval $[t, t + \Delta t]$ is $\mathrm{Tr}[J\hat{\rho}'(t)]\Delta t$. The system state immediately after the click becomes $\hat{\rho}'(t_{+}) = \hat{J}\hat{\rho}'(t)/\mathrm{Tr}[\hat{J}\hat{\rho}'(t)]$. Between the clicks the time evolution is also modified due to continuous leak of information from the system to the EMD, and the dynamics during the time interval $[t_0, t)$ must be described by the non-unitary "No-count" superoperator $\hat{S}_{t-t_0}\hat{\rho}'(t_0) \equiv \hat{s}_{t-t_0}\rho'(t_0)\hat{s}_{t-t_0}^{\dagger}$. Here $\hat{\rho}'(t_0)$ is the system density operator at the time t_0 and the operator \hat{s}_t obeys the differential equation $i\partial \hat{s}_t/\partial t = (\hat{H}' - i\hat{R}/2)\hat{s}_t$, where \hat{R} is a positive operator defined by the relation $\operatorname{Tr}[\hat{J}\hat{\rho}'] = \operatorname{Tr}[\hat{\rho}'\hat{R}]$ (for example, if $\hat{J}\hat{\rho}' = \sum_i \hat{L}_i \hat{\rho}' \hat{L}_i^{\dagger}$ then

 $\hat{R} = \sum_i \hat{L}_i^{\dagger} \hat{L}_i$ [36]. Combining the superoperators \hat{J} and \hat{S}_t one may evaluate the system state after an arbitrary sequence of detection events and the probability of such "quantum trajectory". For deeper discussion regarding the DCE under continuous monitoring by a 2-level detector see [25, 26].

III. N-LEVEL LADDER ATOM

In this paper the quantum detector is modeled as a dissipationless N-level atom in the ladder configuration (hence the terms "atom" and "quantum detector" will be used interchangeably). Assuming that the atom couples to the field via the dipole interaction, the Hamiltonian \hat{H}_d is

$$\hat{H}_d = \sum_{j=1}^N E_j \hat{\sigma}_j + \sum_{j=1}^{N-1} g_j (\hat{a} + \hat{a}^{\dagger}) (\hat{\sigma}_{j+1,j} + \hat{\sigma}_{j,j+1}), \quad (2)$$

where E_j is the energy of the *j*-th atomic state $|\mathbf{j}\rangle$, $\hat{\sigma}_j \equiv |\mathbf{j}\rangle\langle \mathbf{j}|$ and $\hat{\sigma}_{k,j} \equiv |\mathbf{k}\rangle\langle \mathbf{j}|$ are the generalized Pauli operators and g_j (assumed real) is the coupling parameter between the atomic states $\{|\mathbf{j}\rangle, |\mathbf{j}+\mathbf{1}\rangle\}$ via the cavity field. For $|\varepsilon| \ll \omega_0$ the modulation is only relevant for the squeezing coefficient, so one may write $\omega_t \simeq \omega_0$ and $\chi_t \simeq (\varepsilon \eta/4\omega_0) \cos(\eta t)$ in the equation (1). For the "empty cavity" (without the detector) the maximal photon generation occurs for the modulation frequency $\eta = 2\omega_0$, so in the following I shall write $\eta = 2\omega_0 + 2r$, where 2r is the resonance shift from the standard DCE resonance. In the interaction picture defined by the timedependent unitary transformation $\hat{\rho}'(t) = \hat{V}(t)\hat{\rho}(t)\hat{V}^{\dagger}(t)$ with $\hat{V}(t) = \exp[-\frac{1}{2}it(\eta\hat{n} + \sum_{k=0}^{N-1}(2E_1 + k\eta)\hat{\sigma}_{k+1})]$ the rotated statistical operator $\hat{\rho}$ evolves according to the Hamiltonian $\hat{H} = \hat{V}^{\dagger}(t)\hat{H}'\hat{V}(t) - i\hat{V}^{\dagger}(t)\partial\hat{V}(t)/\partial t$

$$\hat{H} = i\beta_r \left(\hat{a}^{\dagger 2} - \hat{a}^2 \right) - r\hat{n} - \sum_{l=1}^{N-1} \sum_{j=1}^{l} (\Delta_j + r)\hat{\sigma}_{l+1} + \sum_{l=1}^{N-1} g_l (\hat{a}\hat{\sigma}_{l+1,l} + e^{-i\eta t} \hat{a}\hat{\sigma}_{l,l+1} + H.c.), \quad (3)$$

where $\beta_r = (1 + r/\omega_0)\varepsilon/4$, $\Delta_j = \omega_0 - (E_{j+1} - E_j)$ is the detuning between the cavity bare frequency and a given atomic transition frequency $(j = 1, \ldots, N -$ 1), and *H.c.* stands for the Hermitian conjugate. For $|g_j|, |\Delta_j| \ll \omega_0$ one can neglect the rapidly oscillating term $(e^{-i\eta t} \hat{a} \hat{\sigma}_{i,i+1} + H.c.)$ in the so called Rotating Wave Approximation (RWA), which we adopt from now on.

In the case of empty cavity, null resonance shift (r = 0)and initial vacuum state $|0\rangle_{field}$ [64] the average number of photons increases with time as $\langle \hat{n}(t) \rangle = \sinh^2(2\beta_0 t)$, so for large times $\langle \hat{n}(t) \rangle \simeq \exp(4\beta_0 t)/4$. The field goes to the squeezed vacuum state, for which only even photon numbers are present and the field quadrature operators $\hat{x}_{\pm} = (\hat{a} \pm \hat{a}^{\dagger})/\sqrt{\pm 2}$ have zero averages and variances $\langle (\Delta \hat{x}_{\pm})^2 \rangle = \exp(\pm 4\beta_0 t)/2$. The statistics of this state is called sometimes "super-chaotic", because the Mandel factor, $Q \equiv [\langle (\Delta \hat{n})^2 \rangle - \langle \hat{n} \rangle]/\langle \hat{n} \rangle = 1 + 2\langle \hat{n}(t) \rangle$, is roughly twice bigger (for $\langle \hat{n} \rangle \gg 1$) than its value $Q_{therm} = \langle \hat{n} \rangle$ in the "usual chaotic" thermal state. It will be shown in the next section that even more "chaotic" states, with $Q > 1 + 2\langle \hat{n} \rangle$ (also called "hyper-Poissonian" [26]), can be produced when DCE is implemented in the presence of the quantum detector.

IV. RELATIONSHIP WITH QUANTUM NONLINEAR OPTICS

The interaction of 2-level atoms with squeezed states of light has attracted enormous interest in the Quantum Optics community since the prediction by Gardiner [38] in 1986 that the two polarization components of the atom would be damped at different rates when it interacts with a broadband squeezed vacuum, leading to a longer relaxation time for one of these components when compared to normal vacuum radiative decay. Thereafter a large number of papers addressed different aspects of the problem of atom-field interaction in the presence of squeezing and external driving. In the so called "passive" scheme [39] the atom is driven by a squeezed light produced externally [e.g., by the subthreshold degenerate optical parametric oscillator (DOPO)] or the atom-field system is damped by a broadband squeezed reservoir [40– 47]; this setup parallels the posterior intracavity detection of DCE. Contrary, in the "active" scheme the atom is located inside the squeezing generator (such as DOPO) during the entire operation in order to enhance its coupling to the squeezed field modes and eliminate the propagation losses [39, 48–58]; this design formally resembles the continuous intracavity monitoring of DCE.

Different aspects of this subject were investigated over the last 25 years, including: analytical description of the dynamics and quantum statistical properties of the system [51–57, 59], narrowing and hole burning in the spectrum of the fluorescent and transmitted light [40, 42, 44, 45, 47, 50], suppression or enhancement of the atomic population decay [40, 41], steady state behavior of atomic population inversion and cavity field correlation functions [43, 48, 55, 56], collapse of the atom into a pure state [45], bistability in the steady-state intracavity intensity versus pumping intensity [49], enhancement of the intracavity squeezing in lossy DOPO containing twolevel atoms [39], etc. It is impossible to account for all the pertinent literature in this short article, so I refer the interested reader to the reviews [60, 61] and references in [45, 46, 50].

Apart from eventual terms describing the external driving and dissipation, the chief Hamiltonian describing the Quantum Nonlinear Optics setups with multilevel atoms or harmonic oscillators (such as other cavity field modes [62] or quantum wells [58]) is similar to the Hamiltonian (3) describing the cavity DCE in the presence of quantum detectors. However the scope of investigations and the adopted regimes of parameters have been rather different in these areas. In the cavity DCE research the focus has been the dynamics of photon generation from vacuum (and other low-excitation initial states) and the detector-field entanglement under the assumption of small dissipative losses. On the other hand, in the QNO schemes the majority of studies addressed the steady-state behavior, assumed strong damping or external driving of the atom or the cavity, and the dynamics was usually studied for nonvacuum initial states. Hence, these two physically distinct phenomena are complementary from the mathematical standpoint, and future investigations on the continuous intracavity monitoring of the DCE can take advantage of the vast literature concerning the interaction of atoms with light inside optical parametric oscillators.

V. ANALYTICAL AND NUMERICAL RESULTS

A. Resonant *N*-level atom

Let us first analyze the detector's influence in the resonant case, when all atomic transitions are resonant with the field mode: $\Delta_j = 0$ for $j = 1, \ldots, N-1$. The strong modulation regime occurs when $|\varepsilon| \gg |g_j|$; it can be easily achieved in the circuit QED setup by reducing the atomfield couplings g_j , which currently are $|g_j|/\omega_0 \sim 10^{-2}$ but can be made small at will [63]. Performing the timeindependent unitary transformation $\hat{\rho} = \hat{U}\hat{\rho}_1\hat{U}^{\dagger}$, where $\hat{U} = \exp[-i\sum_{l=1}^{N-1} \xi_l(\hat{a}\hat{\sigma}_{l,l+1} + H.c.)]$ and $\xi_l = g_l/2\beta_r$, the rotated density operator $\hat{\rho}_1$ evolves according to the Hamiltonian $\hat{U}^{\dagger}\hat{H}\hat{U}$. To the second order in the small parameter of the order $\mathcal{O}(\xi)$ (assuming $\mathcal{O}(\xi_j) \sim \mathcal{O}(\xi)$ for all j) for the resonance shift r = 0 one obtains the effective Hamiltonian

$$H_{eff} \simeq i\beta_0 [\hat{\theta}\hat{a}^{\dagger 2} + \sum_{l=1}^{N-2} \xi_l \xi_{l+1} \hat{\sigma}_{l,l+2} - H.c.], \qquad (4)$$

where $\hat{\theta} = 1 + \sum_{l=1}^{N-1} \xi_l^2 (\hat{\sigma}_{l+1} - \hat{\sigma}_l)$. It shows that many photons are generated, but the photon generation rate is slightly affected by the atom depending on its initial state; moreover, the atomic levels become coupled via the cavity field. This Hamiltonian was thoroughly investigated for two-level [27] and three-level [29] detectors, and exact expressions for the photon number distribution and other relevant observable quantities were obtained.

The weak modulation regime, $|\varepsilon| \ll |g_j|$, is achieved by increasing the atom-field couplings beyond the modulation depth; in this case the detector can substantially modify the photon generation process. In the absence of damping and for initial state $|\mathbf{1}, 0\rangle \equiv |\mathbf{1}\rangle_{atom}|0\rangle_{field}$ the system state remains pure throughout the evolution, so instead of the von Neumann equation in the following I shall consider the Schrödinger equation $i\partial |\psi\rangle/\partial t = \hat{H}|\psi\rangle$ for the wavefunction $|\psi\rangle$, where the Hamiltonian H is given by the expression (3) under the RWA. For the null resonance shift, if $\varepsilon = 0$ and the number of levels Nis odd, then for m > 0 excitations the Hamiltonian \hat{H} has one null eigenvalue $\lambda_{m,0} = 0$ corresponding to the eigenstate $|\phi_{m,0}\rangle = \mathcal{N}_m^{-1} \sum_{k=0}^{(N-1)/2} \alpha_k |\mathbf{2k} + \mathbf{1}, m - 2k\rangle$, where \mathcal{N}_m is the normalization constant and $\alpha_{k+1} =$ $-\alpha_k g_{2k+1} \sqrt{m-2k} / (g_{2k+2} \sqrt{m-2k-1})$ [65]. The other eigenstates with m excitations are denoted by $|\phi_{m,k\neq 0}\rangle$ and the corresponding eigenvalues $\lambda_{m,k}$ are functions of g_j $(j = 1, \ldots, N - 1)$, so they are large compared to β_0 . Making the ansatz $|\psi\rangle = \sum_{m,k} e^{-it\lambda_{m,k}} A_{m,k}(t) |\phi_{m,k}\rangle$ one arrives at the set of differential equations

$$\frac{\partial A_{m,j}}{\partial t} = \beta_0 \sum_{n,k} e^{-it\lambda_{n,k}} \langle \phi_{m,j} | (\hat{a}^{\dagger 2} - \hat{a}^2) | \phi_{n,k} \rangle A_{n,k}$$
(5)

with the initial condition $A_{0,0}(0) = 1$. Neglecting the rapidly oscillating terms, for which $|\lambda_{n,k}| \gg$ $|\beta_0 \langle \phi_{m,j}| (\hat{a}^{\dagger 2} - \hat{a}^2) | \phi_{n,k} \rangle|$, one finds that all the coefficients $\{A_{m,0}, A_{m+2,0}\}$ for $m = 0, \ldots, \infty$ become resonantly coupled, while the coefficients $A_{m,k\neq 0}$ remain close to zero throughout the evolution [30]. Thus the number of created photons is unlimited and only the states $|\phi_{m,0}\rangle$ become significantly populated. For an even number N there is a null eigenvalue for the m-excitation eigenstate $|\phi_{m,0}\rangle$ if $m \leq N-2$, therefore at most N-2photons can be created in this case [30]. Such behavior is shown in figure 2 for different numbers N.

B. Harmonic oscillator

In the special case when $N \to \infty$ and $g_l = g\sqrt{l}$ the detector becomes a simple harmonic oscillator (H.O.) if one associates the atom level $|1\rangle$ with the oscillator ground (zero energy) state and makes the replacement $\sum_{l=1}^{\infty} \sqrt{l}\hat{\sigma}_{l,l+1} = \hat{b}$, where \hat{b} is the annihilation operator associated with the detector obeying bosonic commutation relation $[\hat{b}, \hat{b}^{\dagger}] = 1$. Then the Heisenberg equations of motion for operators $\hat{a}(t)$ and $\hat{b}(t)$ can be solved exactly [66], and the quadrature variances become [30, 32]

$$\langle (\Delta \hat{x}_{\pm})^2 \rangle = e^{\pm 2\beta_0 t} \left(\frac{1}{2} \pm \frac{\beta_0}{2\gamma} \sin(2\gamma t) + \frac{\beta_0^2}{\gamma^2} \sin^2(\gamma t) \right) ,$$
(6)

where $\gamma = \sqrt{g^2 - \beta_0^2}$. For $|g| \gg |\beta_0|$ the rate of photon generation becomes roughly twice smaller than for the empty cavity and the ratio $\langle \hat{b}^{\dagger} \hat{b} \rangle / \langle \hat{a}^{\dagger} \hat{a} \rangle$ is close to unity [30]. For $\beta_0 t \gg 1$ the average photon number $\langle \hat{n}(t) \rangle \approx \langle (\Delta \hat{x}_+)^2 \rangle / 2$ increases exponentially at the rate $2\beta_0$, modulated by some oscillations with the frequency 2γ (leading to appearance of almost horizontal "shelves" seen in figure 2), and the Mandel factor is $Q(t) \approx 2\langle \hat{n}(t) \rangle$. Nonetheless, the state of the field mode is not exactly the vacuum squeezed one since the uncertainty product $\langle (\Delta \hat{p})^2 \rangle \langle (\Delta \hat{x})^2 \rangle = 1/4 + (g\beta_0/\gamma^2)^2 \sin^4(\gamma t)$ is larger than the minimal possible value 1/4.



FIG. 2: Average photon number as function of dimensionless time εt for different number of levels N and "harmonic coupling" $g_l = g\sqrt{l}$. H.O. stands for harmonic oscillator. Parameters: $\Delta_l = 0, g/\omega_0 = 10^{-2}$ and $\varepsilon/\omega_0 = 10^{-3}$.

C. Resonances for $r \neq 0$

Besides the resonance r = 0, at least two photons can be created from vacuum for the resonance shift $2r = \pm \sqrt{2g_1^2 + g_2^2}$, though the concrete details depend on the concrete form of coupling coefficients $\{g_l\}$. For harmonic oscillator one recovers the resonances $r = \pm g$, first discovered in [18] and analyzed in details in [32, 33], for which the average photon number grows as $\langle \hat{n}(t) \rangle =$ $\sinh^2(\beta_0 t)/2$. For a 3-level atom [67] the nonzero eigenvalues for $\varepsilon = 0$ are $\pm \lambda_n$ with eigenvectors $|\phi_{n,\pm}\rangle =$ $(\sqrt{ng_1}|\mathbf{1},n\rangle \pm \lambda_n|\mathbf{2},n-1\rangle + g_2\sqrt{n-1}|\mathbf{3},n-2\rangle)/\sqrt{2\lambda_n}$, where $\lambda_n = \sqrt{ng_1^2 + (n-1)g_2^2}$. Thus for $2r = \pm \lambda_2$ there is an oscillation between the states $|\mathbf{1},0\rangle$ and $|\phi_{2,\pm}\rangle$ with the angular frequency $\beta_r[1 + (g_2/2g_1)^2]^{-1/2}$ [29].

D. Atomic network

One can easily extend the above results to a network of (N-1) identical two-level atoms with transition frequencies Ω coupled to the field mode with the same coupling constant g [28, 30]. The Hamiltonian reads

$$\hat{H}_{a} = \sum_{j=1}^{N-1} [\Omega \hat{\sigma}_{2}^{(j)} + g \left(\hat{a} + \hat{a}^{\dagger} \right) \left(\hat{\sigma}_{1,2}^{(j)} + \hat{\sigma}_{2,1}^{(j)} \right)], \quad (7)$$

where the upper index labels the *j*-th atom. Defining the collective operators $\hat{S}_z = \sum_{j=1}^{N-1} \hat{\sigma}_2^{(j)}$, $\hat{S}_+ = \sum_{j=1}^{N-1} \hat{\sigma}_{2,1}^{(j)}$ and $\hat{S}_- = \sum_{j=1}^{N-1} \hat{\sigma}_{1,2}^{(j)}$ the Hamiltonian (7) becomes $\hat{H}_a = \Omega \hat{S}_z + g(\hat{a} + \hat{a}^{\dagger})(\hat{S}_+ + \hat{S}_-)$. For identical atoms one can introduce the normalized Dicke state $|\mathbf{j}\rangle = \sqrt{[(N-1)!]^{-1}(j-1)!(N-j)!} \sum_p |\mathbf{2}^{(1)}\rangle |\mathbf{2}^{(2)}\rangle \cdots |\mathbf{2}^{(j-1)}\rangle |\mathbf{1}^{(j)}\rangle \cdots |\mathbf{1}^{(N-1)}\rangle$ with (j-1) excitations, where the sum runs over all the allowed permutations of excited and non-excited atoms and $|\mathbf{k}^{(j)}\rangle$ denotes

the state (k = 1, 2) of the *j*-th atom. Using the known properties $\hat{S}_{-}|\mathbf{j}\rangle = \sqrt{(j-1)(N-j+1)}|\mathbf{j}-\mathbf{1}\rangle$, $\hat{S}_{+}|\mathbf{j}\rangle = \sqrt{j(N-j)}|\mathbf{j}+\mathbf{1}\rangle$ and $\hat{S}_{z}|\mathbf{j}\rangle = (j-1)|\mathbf{j}\rangle$, in the Dicke basis the Hamiltonian (7) becomes identical to the Hamiltonian (2) if one identifies $E_{j} = \Omega(j-1)$ and $g_{j} = g\sqrt{j(N-j)}$. Thus the network of (N-1) identical two-level atoms is equivalent to an equidistant N-level ladder atom.



FIG. 3: Dynamical behavior for 2-level detector and parameters $\Delta_1 = 8g_1$, $g_1/\omega_0 = 10^{-2}$, $\varepsilon/\omega_0 = 3 \times 10^{-4}$. Correctional shifts are: a) $y = -\varepsilon/2$ and b) $y = -2\varepsilon$. In the bottom are shown the photon statistics for $\varepsilon t = 5$.

E. Two-level atom

Finally I outline the case of a 2-level quantum detector. Performing the unitary transformation $\hat{V}_2(t) = \exp[irt(\hat{n} + \hat{\sigma}_2)]$ on the equation (3) the new Hamiltonian reads (under the RWA)

$$\hat{H}_2 = -\Delta_1 \hat{\sigma}_2 + (i\beta_r \hat{a}^{\dagger 2} e^{-2irt} + g_1 \hat{a} \hat{\sigma}_{2,1} + H.c.) . \quad (8)$$

Without the modulation, $\varepsilon = 0$, this is the celebrated Jaynes-Cummings model, whose ground state is $|\phi_0\rangle = |\mathbf{1}, 0\rangle$ with eigenenergy $\lambda_0 = 0$, and the excited eigenstates with n > 0 excitations are $|\phi_{n,+}\rangle = \sin \theta_n |\mathbf{1}, n\rangle + \cos \theta_n |\mathbf{2}, n - 1\rangle$, $|\phi_{n,-}\rangle = \cos \theta_n |\mathbf{1}, n\rangle - \sin \theta_n |\mathbf{2}, n - 1\rangle$ with eigenenergies $\lambda_{n,\pm} = -\Delta_1/2 \pm z_n$, where $z_n = \sqrt{(\Delta_1/2)^2 + g_1^2 n}$ and $\theta_n = \arctan \sqrt{(z_n + \Delta_1/2)/(z_n - \Delta_1/2)}$.

Writing the ansatz

$$|\psi_2(t)\rangle = A_0(t)|\phi_0\rangle + \sum_{n>0} \sum_{\mathcal{S}=\pm} e^{-it\lambda_{n,\mathcal{S}}} A_{n,\mathcal{S}}(t)|\phi_{n,\mathcal{S}}\rangle$$
(9)

one can check that for $|g_1| \gg |\varepsilon|$ the coefficient A_0 couples resonantly to $\{A_{2n,S}\}$ for the resonance shift $2r = Sz_2 - \Delta_1/2 + y$ (where $S = \pm$ and y is a small correctional shift of the order of ε), where n = 1 if $\Delta/|\Delta| = -S$ and $n \ge 1$ if $\Delta/|\Delta| = S$. In the resonant regime, $|\Delta_1| \ll |g_1|$, at most two photons can be generated [18, 25]. On the other hand, in the dispersive regime, when $(\Delta_1/2)^2 \gg g_1^2 n$ for all relevant n, one has $z_{n>0} \simeq |\Delta_1|/2 + |\delta| n$ and $\lambda_{n+2,\pm} - \lambda_{n,\pm} = \pm 2 |\delta|$, where $\delta = g_1^2/\Delta_1$ is the cavity dispersive shift due to the detector; hence many photons can be created from the initial state $|\mathbf{1}, 0\rangle$ for the resonance shift $2r = 2\delta$ [28].

The most interesting case occurs when $|\Delta_1| \sim |g_1|$, as in this case the detector acts as saturable limiter and exotic field states can be produced [23–26]. In figure 3 are plotted $\langle \hat{n} \rangle$, Q, the atomic excitation probability P_2 and the photon number distribution at some time instant for two different values of y. One can see that the system dynamics is very sensitive to small changes in the modulation frequency, and the "hyper-Poissonian" states appear quite naturally (figure 3b). Detailed studies of this regime for different initial states and in the presence of EMD can be found in [25, 26].

VI. DISCUSSION AND FUTURE DIRECTIONS

An important actual task is to study how the detector's temperature and damping influence the photon generation regimes under the continuous intracavity monitoring (the cavity damping was considered in numerous papers, see [2] for a brief review). Preliminary numerical results for the case when the detector can be modeled as a 2-level atom were given in [25], while the paper [29] analyzed numerically the situation for 3-level atoms in the ladder or V configurations. Owing to dissipation the dynamics of photon generation from vacuum is qualitatively different from the lossless case, as system states that could not be populated via the unitary evolution now can become populated via dissipative channels. Moreover, for weak dissipation, the photon number probabilities exhibit oscillatory behavior as function of time before attaining the steady-state values [29], so both the dynamics and the stationary properties deserve further investigations. In this connection the rich literature on the steady-state behavior of atoms inside squeezing generators within the QNO area may be useful to foresee the stationary features in the realistic continuous intracavity monitoring of DCE.

The modeling of detector as harmonic oscillator also seems to be rather realistic in the so called "Motion Induced Radiation" (MIR) experiment [11, 12], where the microwave quanta created via DCE are supposed to be detected by means of a small antenna put inside the cavity. Since the inductive antenna (a wire loop) used in that experiment is a part of a LC-contour, it can be reasonably approximated as a harmonic oscillator. To our knowledge the effects of dissipation on the photon production via DCE have not been addressed for this type of detector, although a thorough mathematical analysis of a similar problem was carried out within the scope of QNO [58] and the results could be easily transposed to the DCE scenario.

Another interesting perspective is to exploit the rich eigenvalue spectrum of the atom-field interaction Hamiltonian to implement multi-modulation regimes, when the modulation consists of multiple sinusoids whose frequencies may adiabatically vary with time [24]. In such a way one could selectively couple different sets of the dressed states of the atom-field system and control the amount of photons generated from vacuum. Finally, the implementation of analogs of DCE in highly controllable solid state systems may lead to novel schemes of generating nonclassical states of light and manipulating the lightmatter interaction.

VII. SUMMARY

I summarized the main results on the continuous intracavity monitoring of the dynamical Casimir effect using multi-level quantum detectors that may be coupled to classical external measurement devices. It was shown that the photon generation from vacuum can be severely affected by the detector, therefore the periodicity of external modulation must be parametrically tuned to induce different dynamical regimes. In the absence of dissipation and for specific modulation frequencies the unitary evolution from vacuum can lead to field-detector entangled states with: (i) at most two photons, (ii) a small amount of photons (> 2) or (iii) unlimited photon number. Thus at the expense of reducing the degree of squeezing and the number of created photons (or the rate of exponential photon growth), the photon generation from vacuum can be continuously monitored, and qualitatively new quantum states could be generated using the post-selection procedures.

Acknowledgments

The present work was supported by CNPq, Conselho Nacional de Desenvolvimento Científico e Tecnolgico – Brazil. Partial support by Decanato de Pesquisa e Pós-Graduação (DPP–UnB, Brazil) is acknowledged.

- Jaskula J -C, Partridge G B, Bonneau M, Lopes R, Ruaudel J, Boiron D and Westbrook C I 2012 Acoustic Analog to the Dynamical Casimir Effect in a Bose-Einstein Condensate *Phys. Rev. Lett.* **109** 220401
- [2] Dodonov V V 2010 Current status of the dynamical Casimir effect Phys. Scr. 82 038105
- [3] Nation P D, Johansson J R, Blencowe M P and Nori F 2012 Colloquium: Stimulating uncertainty: Amplifying

the quantum vacuum with superconducting circuits *Rev.* Mod. Phys. **84** 1–24

- [4] De Liberato S, Ciuti C and Carusotto I 2007 Quantum vacuum radiation spectra from a semiconductor microcavity with a time-modulated vacuum Rabi frequency *Phys. Rev. Lett.* **98** 103602
- [5] De Liberato S, Gerace D, Carusotto I and Ciuti C 2009 Extracavity quantum vacuum radiation from a single qubit *Phys. Rev.* A 80 053810
- [6] Johansson J R, Johansson G, Wilson C M and Nori F 2009 Dynamical Casimir effect in a superconducting coplanar waveguide *Phys. Rev. Lett.* **103** 147003
- [7] Johansson J R, Johansson G, Wilson C M and Nori F 2010 Dynamical Casimir effect in superconducting microwave circuits *Phys. Rev.* A 82 052509
- [8] Johansson J R, Johansson G, Wilson C M, Delsing P and Nori F 2012 Nonclassical microwave radiation from the dynamical Casimir effect *preprint* arXiv:1207.1988
- [9] Wilson C M, Johansson G, Pourkabirian A, Simoen M, Johansson J R, Duty T, Nori F and Delsing P 2011 Observation of the dynamical Casimir effect in a superconducting circuit Nature 479 376–9
- [10] Lähteenmäki P, Paraoanu G S, Hassel J and Hakonen P J 2011 Dynamical Casimir effect in a Josephson metamaterial *preprint* arXiv:1111.5608v2
- [11] Braggio C, Bressi G, Carugno G, Del Noce C, Galeazzi G, Lombardi A, Palmieri A, Ruoso G and Zanello D 2005 A novel experimental approach for the detection of the dynamic Casimir effect *Europhys. Lett.* **70** 754
- [12] Braggio C, Bressi G, Carugno G, Della Valle F, Galeazzi G and Ruoso G 2009 Characterization of a low noise microwave receiver for the detection of vacuum photons *Nucl. Instr. and Meth.* A 603 451–5
- [13] Kim W J, Brownell J H and Onofrio R 2006 Detectability of dissipative motion in quantum vacuum via superradiance Phys. Rev. Lett. 96 200402
- [14] Brownell J H, Kim W J and Onofrio R 2008 Modelling superradiant amplification of Casimir photons in very low dissipation cavities J. Phys. A: Math. Theor. 41 164026
- [15] Kawakubo T and Yamamoto K 2011 Photon creation in a resonant cavity with a nonstationary plasma mirror and its detection with Rydberg atoms *Phys. Rev.* A 83 013819
- [16] Fujii T, Matsuo S, Hatakenaka N, Kurihara S and Zeilinger A 2011 Quantum circuit analog of the dynamical Casimir effect *Phys. Rev. B* 84 174521
- [17] Sarkisyan E M, Petrosyan K G, Oganesyan K B, Hakobyan A A, Saakyan V A, Gevorkian S G, Izmailyan N Sh, and Hu C K 2008 Detection of Casimir photons with electrons Las. Phys. 18 621–4
- [18] Dodonov V V 1995 Photon creation and excitation of a detector in a cavity with a resonantly vibrating wall *Phys. Lett.* A **207** 126–32
- [19] Dodonov V V and Klimov A B 1996 Generation and detection of photons in a cavity with a resonantly oscillating boundary *Phys. Rev.* A 53 2664–82
- [20] Lozovik Yu E, Narozhny N B and Fedotov A M 2000 Excitation of an atom in a nonstationary cavity JETP Lett. 72 234–7
- [21] Fedotov A M, Narozhny N B, Lozovik Yu E 2000 'Shaking' of an atom in a non-stationary cavity *Phys. Lett.* A 274 213–22
- [22] Narozhny N B, Fedotov A M, Lozovik Yu E 2001 Dynamical Lamb effect versus dynamical Casimir effect Phys.

Rev. A 64 053807

- [23] Dodonov A V, Céleri L C, Pascoal F, Lukin M D and Yelin S F 2008 Photon generation from vacuum in nonstationary circuit QED preprint arXiv:0806.4035v3
- [24] Dodonov A V 2009 Photon creation from vacuum and interactions engineering in nonstationary circuit QED J. Phys.: Conf. Ser. 161 012029
- [25] Dodonov A V, Lo Nardo R, Migliore R, Messina A and Dodonov V V 2011 Analytical and numerical analysis of the atom-field dynamics in non-stationary cavity QED J. Phys. B: At. Mol. Opt. Phys. 44 225502
- [26] Dodonov A V and Dodonov V V 2011 Strong modifications of the field statistics in the cavity dynamical Casimir effect due to the interaction with two-level atoms and detectors *Phys. Lett.* A **375** 4261–7
- [27] Dodonov A V and Dodonov V V 2012 Approximate analytical results on the cavity dynamical Casimir effect in the presence of a two-level atom *Phys. Rev.* A 85 015805
- [28] Dodonov A V and Dodonov V V 2012 Dynamical Casimir effect in two-atom cavity QED Phys. Rev. A 85 055805
- [29] Dodonov A V and Dodonov V V 2012 Dynamical Casimir effect in a cavity in the presence of a three-level atom *Phys. Rev.* A 85 063804
- [30] Dodonov A V and Dodonov V V 2012 Dynamical Casimir effect in a cavity with an N-level detector or N-1 two-level atoms Phys. Rev. A **86** 015801
- [31] Zhang X, Zheng T -Y, Tian T and Pan S -M 2011 The dynamical Casimir effect versus collective excitations in atom ensemble *Chin. Phys. Lett.* 28 064202
- [32] Dodonov A V and Dodonov V V 2013 Photon statistics in the dynamical Casimir effect modified by a harmonic oscillator detector *Phys. Scr.* T (in press)
- [33] De Castro A S M, Cacheffo A and Dodonov V V 2012 Influence of the field–detector coupling strength on the dynamical Casimir effect *preprint* arXiv:1212.2156v1
- [34] Carusotto I, De Liberato S, Gerace D and Ciuti C 2012 Back-reaction effects of quantum vacuum in cavity quantum electrodynamics *Phys. Rev.* A 85 023805
- [35] Law C K 1994 Effective Hamiltonian for the radiation in a cavity with a moving mirror and a time-varying dielectric medium *Phys. Rev.* A 49 433–7
- [36] Srinivas M D and Davies E B 1981 Photon counting probabilities in Quantum Optics Opt. Acta 28 981–96
- [37] Carmichael H 1993 An Open System Approach to Quantum Optics (Springer, Berlin)
- [38] Gardiner C W 1986 Inhibition of atomic phase decays by squeezed light: a direct effect of squeezing *Phys. Rev. Lett.* 56 1917–20
- [39] Jin S and Xiao M 1994 Extra intracavity squeezing of a degenerate optical parametric oscillator coupling with N two-level atoms *Phys. Rev.* A **49** 499–505
- [40] Parkins A S 1990 Rabi sideband narrowing via strongly driven resonance fluorescence in a narrow-bandwidth squeezed vacuum *Phys. Rev.* 42 4352–65
- [41] Cirac J I and Sanchez-Soto L L 1991 Suppression of spontaneous emission by squeezed light in a cavity *Phys. Rev.* A 44 1948–55
- [42] Vyas R and Singh S 1992 Resonance Suorescence with squeezed-light excitation Phys. Rev. A 45 8095–109
- [43] Cirac J I 1992 Interaction of a two-level atom with a cavity mode in the bad-cavity limit Phys. Rev. A 46 4354–62
- [44] Rice P R and Pedrotti L M 1992 Fluorescent spectrum of a single atom in a cavity with injected squeezed vacuum J. Opt. Soc. Am. B 9 2008–14

- [45] Smyth W S and Swain S 1996 Anomalous resonance fluorescence from an atom in a cavity with injected squeezed vacuum *Phys. Rev.* A 53 2846–54
- [46] Turchette Q A, Georgiades N Ph, Hood C J, Kimble H J and Parkins A S 1998 Squeezed excitation in cavity QED: experiment and theory *Phys. Rev.* A 58 4056-77
- [47] Smyth W S and Swain S 1999 Complete quenching of fluorescence from a two-level atom driven by a weak, narrow-band, nonclassical light field *Phys. Rev.* A 59 R2579–82
- [48] Agarwal G S 1989 Effects of optical gain and cavity-mode squeezing on the Mollow spectrum Phys. Rev. A 40 4138– 41
- [49] Jin S and Xiao M 1993 Optical spectra from a degenerate optical parametric oscillator coupling with N two-level atoms Phys. Rev. A 48 1473–82
- [50] Clemens J P, Rice P R, Rungta P K and Brecha R J 2000 Two-level atom in an optical parametric oscillator: Spectra of transmitted and fluorescent fields in the weakdriving-field limit *Phys. Rev.* A 62, 033802
- [51] Obada A S -F, Abdalla M S and Khalil E M 2004 Statistical properties of two-mode parametric ampliler interacting with a single atom *Phys.* A 336 433–53
- [52] Abdalla M S, Obada A -S F and Abdel-Aty M 2005 Von Neumann entropy and phase distribution of two mode parametric amplifier interacting with a single atom Ann. Phys. **318** 266–85
- [53] Abdalla M S, Khalil E M and Obada A -S F 2005 Statistical properties of a two-photon cavity mode in the presence of degenerate parametric amplifier Ann. Phys. **322** 2554–68
- [54] Abdalla M S, Hassan S S and Abdel-Aty M 2005 Entropic uncertainty in the Jaynes–Cummings model in presence of a second harmonic generation Opt. Commun. 244 431– 43
- [55] Strimbu C E, Leach J and Rice P R 2005 Conditioned

homodyne detection at the single-photon level: Intensityfield correlations for a two-level atom in an optical parametric oscillator *Phys. Rev.* A **71** 013807

- [56] Alebachew E and Fesseha K 2007 Interaction of a twolevel atom with squeezed light Opt. Commun. 271 154– 61
- [57] Batarfi H A, Al-Khayat I A, Abdalla M S and Hassan S. S. 2009 Spectra of the Jaynes-Cummings model in presence of a second harmonic generation *Int. J. Mod. Phys.* B 23 2215–32
- [58] Sete E A and Eleuch H 2010 Interaction of a quantum well with squeezed light: Quantum-statistical properties *Phys. Rev.* A 82 043810
- [59] Rice P R 2005 Entanglement, teleportation, and singlephoton storage with two-level atoms inside an optical parametric oscillator J. Opt. Soc. Am. B 22 1561–5
- [60] Parkins A S 2007 The interaction of squeezed light with atoms Advances in Chemical Physics: Modern Nonlinear Optics, Part 2, Vol 85 ed M Evans and S Kielich (New York: Wiley) p 607–66
- [61] Dalton B J, Ficek Z and Swain S 1999 Atoms in squeezed light fields J. Mod. Opt. 46 379–474
- [62] Abdalla M S 1996 Quantum mechanics of two modes two photons coupled oscillators *Phys.* A 234 554–70
- [63] Schoelkopf R J and Girvin S M 2008 Wiring up quantum systems Nature 451 664–9
- [64] The field states are expressed in the Fock basis throughout the paper.
- [65] For m = 0 the eigenvalue is 0 and the eigenstate is $|\mathbf{1}, 0\rangle$.
- [66] These equations were solved in the context of QNO in [58], while the solution for $\langle \hat{n} \rangle$ was obtained independently in [31].
- [67] 3-level atom also in the Lambda configuration was analyzed in [29].