Holographic Symmetries and Generalized Order Parameters for Topological Matter

Emilio Cobanera,¹ Gerardo Ortiz,² and Zohar Nussinov³

¹Instituut-Lorentz, Universiteit Leiden, P.O. Box 9506, 2300 RA Leiden, The Netherlands^{*}

²Department of Physics, Indiana University, Bloomington, IN 47405, USA

³Department of Physics, Washington University, St. Louis, MO 63160, USA

(Dated: September 28, 2018)

We introduce a universally applicable method, based on the bond-algebraic theory of dualities, to search for generalized order parameters in disparate systems including non-Landau systems with topological order. A key notion that we advance is that of *holographic symmetry*. It reflects situations wherein global symmetries become, under a duality mapping, symmetries that act solely on the system's boundary. Holographic symmetries are naturally related to edge modes and localization. The utility of our approach is illustrated by systematically deriving generalized order parameters for pure and matter-coupled Abelian gauge theories, and for some models of topological matter.

PACS numbers: 05.30.Rt,75.10.Kt,11.15.Ha

Introduction.— Landau's concept of an order parameter (OP) and spontaneous symmetry breaking are central in physics [1]. In systems with long-range Landau orders, two-point correlation functions of an OP field $\mathcal{O}(\mathbf{r})$, in their large distance limit, tend to a finite (i.e., non-zero) value, $\lim_{|\mathbf{r}-\mathbf{r'}|\to\infty} \lim_{N^d\to\infty} \langle \mathcal{O}(\mathbf{r})\mathcal{O}^{\dagger}(\mathbf{r'}) \rangle \neq 0$, where Nis the linear size of the d-dimensional system, and $\mathcal{O}(\mathbf{r})$ is local in the (spatial) variable \mathbf{r} . It is in Landau's spirit to use the OP as a macroscopic variable characterizing the ordered phase and as an indicator of a possible phase transition (classical or quantum) to a disordered state where the OP becomes zero.

There is much experience, including systematic methods [2, 3], for deriving Landau OPs and their effective field theories [1]. Landau's ideas of a (*local*) OP cannot be extended to topological states of matter as, by definition [4, 5], these lie beyond Landau's paradigm. However, the notion of long-range order, or the design of a *witness correlator* (i.e., a correlator discerning the existence of various phases and related transitions), can be extended to topological phases - phases that can only be meaningfully examined by *non-local probes* [5]. Topological orders appear in gauge theories, quantum Hall and spin liquid states (when defined as deconfined phases of emergent gauge theories [6]), including well-studied exactly solvable models [7, 8].

In this paper we demonstrate that generalized nonlocal OPs may diagnose topological phases of matter. Most importantly, we outline a method based on bondalgebraic duality mappings to search systematically for generalized OPs. Dualities have the striking capability of mapping Landau to topological orders and viceversa for essentially two reasons. First, dualities in general represent non-local transformations of elementary degrees of freedom [9] and may even perform transmutation of statistics [10]. Second, bond-algebra techniques [10–12] allow for the generation of dualities in finite and infinite size systems. As we will show, in systems with a boundary, dualities realize a form of holography [13] capable of transforming a global symmetry that may drive spontaneous symmetry breakdown into a boundary symmetry. We term these distinguished boundary symmetries holographic. They are under suitable further conditions connected to edge (boundary) states. To illustrate the method, we derive explicitly a (non-local) witness correlator and a generalized OP, suited to diagnose the transition between deconfined and confined phases of mattercoupled gauge theories, undetectable by standard OPs or Wilson loops. Other examples are reported in Ref [18].

The search for generalized order parameters.— A natural mathematical *language* to describe a physical system is that for which the system's degrees of freedom couple *locally.* This simple observation is key to understanding that topological order is a property of a state(s) relative to the algebra of observables (defining the language) used to probe the system experimentally [5]. In the language in which the system is topologically ordered, it is also robust (at zero temperature [16]) against perturbations local in that language. Spectral properties are invariant under unitary transformations of the local Hamiltonian H governing the system, $H \mapsto UHU^{\dagger}$. If UHU^{\dagger} corresponds to a sensible local theory then the unitary transformation U establishes a *duality* [10]. A duality may map a system that displays topological order to one that does not [5]. Dualities for several of Kitaev's models [7, 8, 17] epitomize this idea [5, 12, 16].

Since dualities are unitary transformations (or, more generally, partial isometries) [10] they cannot in general change a phase diagram, only its interpretation. This leads to a central point of our work: A duality mapping a Landau to a topologically ordered system must map the Landau OP to a generalized OP characterizing the topological order. Our method for searching for generalized OPs, combines this observation with the advantages of the bond-algebraic theory of dualities [10]. In this framework, dualities in arbitrary size (finite or infinite) systems can be systematically searched for as alternative local representations of bond algebras of interactions associated to a Hamiltonian H. Hence it is possible for *any* system possessing topological order to systematically search for a duality mapping it to a Landau order. When a dual Landau theory is found, the dual system's OP can be mapped back to obtain a generalized OP for the topologically ordered system. In what follows and in Ref. [18], we study various quantum gauge and topologically ordered theories, and their duals, to illustrate our ideas.

Holographic symmetries and edge states: the gauged Kitaev wire.— We next illustrate the concept of holographic symmetry and its relation to generalized OPs and edge modes. Consider the Kitaev wire Hamiltonian [17] with open boundary conditions, here generalized to include a \mathbb{Z}_2 gauge field (termed the gauged Kitaev wire),

$$H_{\mathsf{GK}} = -ih \sum_{m=1}^{N} b_m a_m - \sum_{m=1}^{N-1} [iJb_m \sigma^z_{(m;1)} a_{m+1} + \kappa \sigma^x_{(m;1)}],$$
(1)

where $a_m = a_m^{\dagger}$, $b_m = b_m^{\dagger}$ denote two Majorana fermions $(\{a_m, a_n\} = 2\delta_{mn} = \{b_m, b_n\}, \{a_m, b_n\} = 0)$ placed on each site of an open chain with N sites. The Pauli matrices $\sigma_{(m;1)}^{\alpha}$, $\alpha = x, z$, placed on the links (m; 1) connecting sites m and m + 1 represent a \mathbb{Z}_2 gauge field. For the gauged Kitaev wire, fermionic parity is obtained as the product of the local (gauge) \mathbb{Z}_2 symmetries $ib_1a_1\sigma_{(1;1)}^x$, $\sigma_{(N-1;1)}^x ib_N a_N$, and $\sigma_{(m;1)}^x ib_{m+1}a_{m+1}\sigma_{(m+1;1)}^x$ $(m = 1, \dots, N-2)$. Just like the standard Kitaev wire, $H_{\mathsf{GK}}[h=0]$ has two free edge modes a_1 and b_N .

The gauged Kitaev wire holds two important dualities. It is dual to the one-dimensional \mathbb{Z}_2 Higgs model [19]

$$H_{\mathsf{H}} = -h \sum_{i=1}^{N} \sigma_i^x - \sum_{i=1}^{N-1} [J \sigma_i^z \sigma_{(i;1)}^z \sigma_{i+1}^z + \kappa \sigma_{(i;1)}^x], \quad (2)$$

with Pauli matrices σ_i^{α} placed on sites *i*. Moreover, the gauge-reducing [10] duality mapping Φ_d

$$ib_m a_m \xrightarrow{\Phi_d} \sigma_m^z \sigma_{m+1}^z, \quad m = 1, \cdots, N, \quad (3)$$

$$ib_m \sigma^z_{(m;1)} a_{m+1} \xrightarrow{\Phi_d} \sigma^x_{m+1}, \quad m = 1, \cdots, N-1,$$

 $\sigma^x_{(m;1)} \xrightarrow{\Phi_d} \sigma^z_{m+1}, \quad m = 1, \cdots, N-1,$

transforms H_{GK} into a spin-1/2 system

$$H_{\rm GK}^D = -h \sum_{m=1}^N \sigma_m^z \sigma_{m+1}^z - \sum_{m=2}^N [J \sigma_m^x + \kappa \sigma_m^z].$$
(4)

defined on (N + 1) sites. The fermionic parity P maps to a holographic symmetry under this duality, since $P = \prod_{m=1}^{N} i b_m a_m \xrightarrow{\Phi_d} \sigma_1^z \sigma_{N+1}^z$, i.e., the product of two (commuting) boundary symmetries. Holography is a relational phenomenon (see [18]). A duality that uncovers a holographic symmetry links a global (higherdimensional) symmetry of a system to a boundary (lowerdimensional) symmetry of its dual. Boundary symmetries need not in general be duals of global symmetries.

What is the physical consequence of having an holographic symmetry? Consider the not uncommon situation in which the holographic symmetry is supplemented by an additional (non-commuting) boundary symmetry in some region of the phase diagram. By definition, holographic symmetries are boundary symmetries which are dual to global symmetries. Thus, global symmetries linking degenerate states (and properties in the broken symmetry phase) in the dual system have imprints in their holographic counterparts. Then, the many-body level degeneracy of the ground state may be ascribed to boundary effects. If the couplings are now changed, the ground state degeneracy may get removed, together with some boundary symmetries. However, so long as the system remains in a topological phase dual to the (broken symmetry) ordered phase, the low energy state splitting will be exponentially small in the system size, so that in the thermodynamic limit ground state degeneracy is restored.

The language providing the most local operator description of the ground-state manifold is the one realizing the edge modes, which are expected to be exponentially localized to the boundary. Thus, as long as the thermodynamic-limit degeneracy remains, a suitable local probe will detect localization on the boundary for those states. Conversely, non-commuting edge mode operators in a gapped phase reflect the existence of lowenergy many-body states with energy splittings vanishing exponentially in the system size. Many-body (zero*energy*) edge states are thus simply a natural consequence of a degenerate ground state manifold in a gapped system. They are witnesses of an ordered (degenerate) phase described in a most local language. Note that boundary operators that commute with the Hamiltonian at special values of the coupling(s) are a necessary but not sufficient condition to realize *exact* (zero-energy) edge modes.

The duality $H_{\rm H} \rightarrow H_{\rm GK}$ maps a global symmetry of $H_{\mathsf{H}}[\kappa, h = 0]$ to a boundary symmetry of $H_{\mathsf{GK}}[\kappa, h = 0]$, i.e., $\sigma_1^x \cdots \sigma_{N-1}^x \sigma_N^y \to b_N$, and one boundary symmetry to another, $\sigma_1^z \to a_1$. If we now turn on h < J, keeping $\kappa = 0, \text{ the edge mode operators } a_1, b_N, \text{ evolve respec tively into } \Gamma_1 = \sum_{m=1}^N (-h/J)^{m-1} a_m (\prod_{s=1}^{m-1} \sigma_{(s;1)}^z) \text{ and}$ $\Gamma_2 = \sum_{m=1}^N (h/J)^{N-m} b_m (\prod_{s=m}^{N-1} \sigma_{(s;1)}^z).$ These modes are exponentially localized as long as the system is in the ordered gapped phase within a gauge sector [23]. The Majorana language affords a *local* boundary description of these (partly non-local in the Higgs language) zero-energy modes. For h > J, and/or $\kappa > 0$, the ground state is unique, even in the thermodynamic limit, as we learn from the phase diagram of the onedimensional Higgs model [19]. Hence the zero-energy modes disappear together with the ground-state degeneracy. For $\kappa > 0$, they disappear despite the fact that fermionic parity remains an exact symmetry and cannot be spontaneously broken [21]. Consider now H_{CK}^D of Eq. (4). At h = 0, it has zero-energy edge mode operators $\sigma_1^z, \sigma_1^x, \sigma_{N+1}^z, \sigma_{N+1}^x$. For h > 0, and $\kappa = 0$, two of these remain unchanged, and the other two evolve into $\Sigma_1 = \sigma_1^x + \sum_{m=1}^{N-1} (h/J)^m \sigma_1^y (\prod_{s=2}^m \sigma_s^x) \sigma_{m+1}^y$ and $\Sigma_2 = \sigma_{N+1}^x + \sum_{m=2}^N (h/J)^m \sigma_m^y (\prod_{s=m+1}^N \sigma_s^x) \sigma_{N+1}^y$. These behave just as their Majorana relatives, yet they are recognized as non-local. The Majorana language distinguishes itself as the most local one for zero modes.

To obtain a generalized OP for the gauged Kitaev wire, notice that $H^D_{\mathsf{GK}}[\kappa = 0]$ reduces to the transverse-field Ising (TI) chain. Hence it exhibits a second-order phase transition at $J = h, \kappa = 0$. For $H^D_{\mathsf{GK}}[\kappa = 0]$, this transition is witnessed by the Landau OP correlator $\lim_{|i-j|\to\infty} \langle \mathsf{TI}|\sigma_i^z \sigma_j^z|\mathsf{TI}\rangle$. (From now on |label> represents the ground state of H_{label}). Our duality maps this correlator back to a generalized OP for the gauged Kitaev wire, the string correlator $\lim_{|i-j|\to\infty} \langle \mathsf{GK}|ib_ia_iib_{i+1}a_{i+1}\cdots ib_ja_j|\mathsf{GK}\rangle$.

Generalized OPs in higher-dimensional theories— We next show how to systematically *derive* generalized OPs in higher space dimensions. Our main goal is to illustrate the methodology in the challenging case of the Abelian (U(1)) matter-coupled gauge (Higgs) theory. Previous works [14, 15] *conjectured* generalized OPs for matter-coupled gauge theories and were numerically implemented, for instance, in Ref. [6]. Unfortunately, a systematic mathematical *derivation* was missing and this is what our work is about. Our (non-local) witness correlator for the Higgs model turns out to be the one conjectured in Ref. [14]. In Ref. [18], we study several other examples (displaying also holographic symmetries), including Ising and \mathbb{Z}_p gauge and Higgs theories, the \mathbb{Z}_p extended toric code [20] as an interesting example of topological order, and the XY model on the frustrated Kagome lattice. Non-Abelian extensions of our ideas based on Ref. [11] are currently under investigation.



FIG. 1. The Z gauge theory exactly dual to the quantum XY model must satisfy special boundary conditions and possesses a boundary symmetry. The lattice corresponding to the XY model is shown in thick lines, for N = 4.

To derive the generalized OP for the Abelian Higgs theory, our starting point is the XY model defined in terms of continuous U(1) degrees of freedom $s_r \equiv e^{-i\theta_r}, \ \theta_r \in [0, 2\pi)$, placed at sites $r = ie_1 + je_2 = (i, j)$ of a square lattice. The model's Hamiltonian reads

$$H_{XY} = h \sum_{i,j=1}^{N} L_{(i,j)}^{2}$$

$$+ \frac{J}{2} \Big[\sum_{i=1}^{N-1} \sum_{j=1}^{N} S_{(i,j;1)} + \sum_{i=1}^{N} \sum_{j=1}^{N-1} S_{(i,j;2)} + h.c. \Big],$$
(5)

with $L_{\mathbf{r}} \equiv -i\partial/\partial\theta_{\mathbf{r}}$, and $S_{(\mathbf{r};\mu)} \equiv s_{\mathbf{r}}s_{\mathbf{r}+e_{\mu}}^{\dagger}$. The XY model is dual to a \mathbb{Z} (solid-on-solid like) gauge theory also defined on a square lattice, but with degrees of freedom X and R associated to links $(\mathbf{r};\mu=1,2)$. (In mattercoupled gauge theories we will also have operators acting on sites \mathbf{r} .) These operators satisfy $X|m\rangle = m|m\rangle$, $R|m\rangle = |m-1\rangle$, $R^{\dagger}|m\rangle = |m+1\rangle$, with $m \in \mathbb{Z}$, and commute on different links (and/or sites). Then, the *exact* dual of H_{XY} for open boundary conditions reads

$$H_{\mathsf{ZG}} = h \sum_{i,j=1}^{N} b_{(i,j)}^{2}$$

$$+ \frac{J}{2} \Big[\sum_{i=2}^{N} \sum_{j=1}^{N} R_{(i,j;2)} + \sum_{i=1}^{N} \sum_{j=2}^{N} R_{(i,j;1)} + h.c. \Big].$$
(6)

We will call system indices the link indices $(i, j; \mu = 1, 2)$ labeling R operators that explicitly appear in H_{ZG} , and extra indices the remaining link indices. In the bulk, the plaquette operator $b_{(i,j)}$ reads

$$b_{(i,j)} \equiv X_{(i,j;1)} + X_{(i+1,j;2)} - X_{(i,j+1;1)} - X_{(i,j;2)}.$$
 (7)

On the lattice boundary, the plaquette operators are set by two rules: (i) $b_{(1,N)} = X_{(1,N;1)} - X_{(2,N;2)} - X_{(1,N+1;1)}$. Thus, $b_{(1,N)}$ involves one degree of freedom $X_{(1,N+1;1)}$ labelled by an extra link index. (ii) The remaining boundary plaquettes are determined by Eq. (7) provided operators labelled by extra link indices are omitted. With these definitions in tow, the mapping of bonds

$$b_{(i,j)} \xrightarrow{\Phi_{d}} L_{(i,j)}, \qquad 1 \le i, j \le N,$$

$$R_{(i,j;1)} \xrightarrow{\Phi_{d}} S_{(i,j-1;2)}^{\dagger}, \ 1 \le i \le N, \ 2 \le j \le N,$$

$$R_{(i,j;2)} \xrightarrow{\Phi_{d}} S_{(i-1,j;1)}, \ 2 \le i \le N, \ 1 \le j \le N,$$
(8)

implements the duality transformation $H_{\mathsf{ZG}} \xrightarrow{\Phi_{\mathsf{d}}} H_{\mathsf{XY}}$. As the operators $R_{(1,N+1;1)}, R_{(1,N+1;1)}^{\dagger}$ do not appear in H_{ZG} , the operator $X_{(1,N+1;1)}$ constitutes a boundary symmetry of H_{ZG} . Similar to the duality between the one-dimensional theories of Eqs. (1) and (4), this is a gauge-reducing duality. The gauge symmetries of H_{ZG} , given by $A_{(i,j)} = R_{(i,j;1)}R_{(i,j;2)}R_{(i-1,j;1)}^{\dagger}R_{(i,j-1;2)}^{\dagger}, 2 \leq i, j \leq N$, are removed by the mapping since $A_{(i,j)} \xrightarrow{\Phi_{\mathsf{d}}} \mathbb{1}$.

In the thermodynamic $(N \to \infty)$ limit, the stronglycoupled $(J \gg h)$ phase of the XY model displays spontaneous symmetry breakdown of its global U(1) symmetry with generator $L_{XY} = \sum_{i,j=1}^{N} L_{(i,j)}$, as evinced by a nonvanishing $\langle XY | s_r s_{r'}^{\dagger} | XY \rangle$ in the limit $|r - r'| \to \infty$. By virtue of being dual to the XY system, the gauge theory displays a non-analyticity in its ground state energy as his varied and its symmetry is broken. However, the phase transition in the gauge theory cannot be characterized by a local OP. So, how can the duality connecting the two models bridge the drastic gap separating the physical interpretation of their common phase diagram? The answer lies in our notion of holography, since

$$-X_{(1,N+1;2)} = \sum_{i,j=1}^{N} b_{(i,j)} \xrightarrow{\Phi_{\mathsf{d}}} L_{\mathsf{XY}}.$$
 (9)

Thus, the global symmetry of the XY model is holographically dual to the (local) boundary symmetry $X_{(1,N+1;2)}$ of its dual gauge theory and cannot be spontaneously broken in this dual theory [21]. This is how holographic symmetries explain the non-Landau nature of critical transitions in the \mathbb{Z} gauge theory. There are no edge modes nor localization associated with this holographic symmetry as the ordered phase of the XY model is gapless.

We now derive a generalized OP for the \mathbb{Z} gauge theory. Let Γ be an *oriented* path from r to r' made of directed links $l \in \Gamma$, and we adopt the convention that $S_l \equiv S_{(r;\mu)}$ if l points from r to $r + e_{\mu}$, or $S_l \equiv S_{(r;\mu)}^{\dagger}$ if l points oppositely from $r + e_{\mu}$ to r. Then $s_r s_{r'}^{\dagger} = \prod_{l \in \Gamma} S_l$. Also let Γ^* denote the set of links l^* such that $\Phi_d(R_{l^*}) = S_l$ (Γ^* need not be continuous, see Fig. 2). Then

$$\langle \mathsf{ZG} | \prod_{l^* \in \Gamma^*} R_{l^*} | \mathsf{ZG} \rangle \xrightarrow{\Phi_{\mathsf{d}}} \langle \mathsf{XY} | s_r s_{r'}^{\dagger} | \mathsf{XY} \rangle,$$
 (10)

and so the string correlator on the left-hand side is a generalized OP for the \mathbb{Z} gauge theory, displaying long-range order in the ordered phase. On a closed path, $\prod_{l^* \in \Gamma^*} R_{l^*}$ reduces to a product of gauge symmetries.



FIG. 2. Dual sets of links Γ^* and Γ .

Finally, we couple the \mathbb{Z} gauge theory to a \mathbb{Z} matter field (defined on sites r), $H_{\mathsf{ZH}} = H_{\mathsf{ZG}} + H_{\mathsf{M}}$, with

$$H_{\mathsf{M}} = \sum_{\boldsymbol{r}} \left[\lambda(R_{\boldsymbol{r}} + R_{\boldsymbol{r}}^{\dagger}) + \kappa \sum_{\mu=1,2} l_{(\boldsymbol{r};\mu)}^2 \right], \qquad (11)$$

and $l_{(\boldsymbol{r};\mu)} \equiv X_{\boldsymbol{r}+\boldsymbol{e}_{\mu}} - qX_{(\boldsymbol{r};\mu)} - X_{\boldsymbol{r}}$. The resulting mattercoupled theory H_{ZH} is dual to the Abelian Higgs model [19] with Hamiltonian

$$H_{\mathsf{AH}} = \sum_{\boldsymbol{r}} \left[\lambda (B_{\boldsymbol{r}} + B_{\boldsymbol{r}}^{\dagger}) + hL_{\boldsymbol{r}}^{2} + \sum_{\mu=1,2} \left(\kappa L_{(\boldsymbol{r};\mu)}^{2} + \frac{J}{2} (S_{(\boldsymbol{r},\mu)}^{(q)} + S_{(\boldsymbol{r},\mu)}^{(q)\dagger}) \right) \right].$$
(12)

Here $S_{(\mathbf{r},\mu)}^{(q)} \equiv s_{\mathbf{r}} s_{(\mathbf{r};\mu)}^{q} s_{\mathbf{r}+\mathbf{e}_{\mu}}^{\dagger}$ includes a coupling with integer charge q to the U(1) gauge field $s_{(\mathbf{r};\mu)} \equiv e^{-i\theta_{(\mathbf{r};\mu)}}$, $s_{(\mathbf{r};\mu)}^{q} \equiv e^{-iq\theta_{(\mathbf{r};\mu)}}$, and $B_{\mathbf{r}} \equiv s_{(\mathbf{r};1)} s_{(\mathbf{r}+\mathbf{e}_{1};2)} s_{(\mathbf{r}+\mathbf{e}_{2};1)}^{\dagger} s_{(\mathbf{r};2)}^{\dagger}$. The correspondence between the two models, established by the mapping of bonds

which holds only on physical gauge-invariant states. The reason is that Φ_d preserves all commutation relations while "trivializing" all gauge symmetries. More precisely, H_{ZH} 's gauge symmetries $G_r = R_r A_r$ map to $\Phi_d(G_r) = 1$, while H_{AH} 's gauge generators $g_r = L_{(r;1)} + L_{(r;2)} - L_{(r-e_1;1)} - L_{(r-e_2;2)} - qL_r$ map to $\Phi_d^{-1}(g_r) = 0$ as follows from Eqs. (13) (Φ_d^{-1} is the mapping obtained from Eqs. (13) by reversing all the arrows).

If the \mathbb{Z} matter field is weakly coupled to the \mathbb{Z} gauge field, the string correlator of Eq. (10) will still change analytic behavior across transitions. Then from Eqs. (13)

$$\langle \mathsf{Z}\mathsf{H}| \prod_{\boldsymbol{l}^* \in \Gamma^*} R_{\boldsymbol{l}^*} | \mathsf{Z}\mathsf{H} \rangle \xrightarrow{\Phi_\mathsf{d}} \langle \mathsf{A}\mathsf{H}| s_{\boldsymbol{r}} s_{\boldsymbol{r}'}^\dagger \prod_{\boldsymbol{l} \in \Gamma} s_{\boldsymbol{l}}^q | \mathsf{A}\mathsf{H} \rangle, \quad (14)$$

we obtain a witness correlator for the Abelian Higgs model that reduces to a Wilson loop on closed contours $(\boldsymbol{r} = \boldsymbol{r}')$ (here $s_l^q = s_{(\boldsymbol{r};\mu)}^q$ if a link l points from \boldsymbol{r} to $\boldsymbol{r} + \boldsymbol{e}_{\mu}$ and $s_l^q = s_{(\boldsymbol{r};\mu)}^{q\dagger}$ otherwise). This non-local correlator is directly related to intuitively motivated generalized OPs like $\langle \mathsf{AH}|s_{\boldsymbol{r}}s_{\boldsymbol{r}'}^{\dagger}\prod_{l\in\Gamma}s_l^q|\mathsf{AH}\rangle/\langle\mathsf{AH}|\prod_{l\in\Gamma_C}s_l^q|\mathsf{AH}\rangle$ conjectured in earlier work [6, 14, 15] (Γ_C denotes a closed loop roughly twice as long as Γ and containing it).

Outlook.— As demonstrated, holographic symmetries and generalized OPs appear in numerous systems once boundary conditions are properly accounted for in the framework of bond-algebraic dualities. By providing a systematic methodology and many examples, our results might bring the theory of generalized OPs and topological orders to a new level of development closer to that of Landau's theory. More key problems need to be tackled. First, the sufficient conditions under which a given topological order may be mapped to a Landau order and viceversa should be understood. Second, the problem of associating effective field theories to generalized OPs should be studied systematically.

Acknowledgements. This work was supported by the Dutch Science Foundation NWO/FOM and an ERC Ad-

vanced Investigator grant, and, in part, under grants No. NSF PHY11-25915 and CMMT 1106293.

- * Electronic address: ecobaner@indiana.edu
- H. Nishimori and G. Ortiz, *Elements of Phase Transitions* and *Critical Phenomena* (Oxford University Press, Oxford, 2011).
- [2] C. D. Batista, G. Ortiz, and J. E. Gubernatis, Phys. Rev. B 65, 180402(R) (2002).
- [3] S. Furukawa, G. Misguich, and M. Oshikawa, Phys. Rev. Lett. 96, 047211 (2006).
- [4] X-G. Wen, Int. J. Mod. Phys. B 4, 239 (1990).
- [5] Z. Nussinov and G. Ortiz, Annals of Physics 324, 977 (2009); Proceedings of the National Academy of Sciences 106, 16944 (2009).
- [6] K. Gregor, D. A. Huse, R. Moessner, and S. L. Sondhi, New J.Phys. 13, 025009 (2011).
- [7] A. Yu Kitaev, Ann. Phys. **303**, 2 (2003).
- [8] A. Yu Kitaev, Ann. Phys. **321**, 2 (2006).
- [9] F. J. Wegner, J. Math. Phys. 12, 2259 (1971).
- [10] E. Cobanera, G. Ortiz, and Z. Nussinov, Phys. Rev. Lett.

104, 020402 (2010); Adv. Phys. 60, 679 (2011).

- [11] E. Cobanera, G. Ortiz, and E. Knill, arXiv:1206.1367v1 [cond-mat.stat-mech] (2012).
- [12] Z. Nussinov and G. Ortiz, Phys. Rev. B 79, 214440 (2009).
- [13] Z. Nussinov, G. Ortiz, and E. Cobanera, Ann. Phys. 327, 2491 (2012).
- [14] J. Bricmont and J. Frölich, Phys. Lett. B 122, 73 (1983).
- [15] K. Fredenhagen and M. Marcu, Phys. Rev. Lett. 56, 223 (1986); Comm. Math. Phys. 92, 81 (1983).
- [16] Z. Nussinov and G. Ortiz, Phys. Rev. B 77, 064302 (2008).
- [17] A. Yu Kitaev, Phys.-Usp. 44, 131 (2001).
- [18] Supplemental material.
- [19] E. Fradkin and S. H. Shenker, Phys. Rev. D 19, 3682 (1979).
- [20] M. D. Schulz, S. Dusuel, R. Orus, J. Vidal, and K. P. Schmidt, New J. Phys. 14, 025005 (2012).
- [21] S. Elitzur, Phys. Rev. D 12, 3978 (1975).
- [22] I. Herbut, A Modern Introduction to Critical Phenomena (Cambridge University Press, Cambridge, 2010).
- [23] A precise estimate follows from $[H_{\mathsf{GK}}[\kappa = 0], \Gamma_1] = -2iJ(-h/J)^N b_N \sigma^z_{(N-1;1)} \cdots \sigma^z_{(1;1)}$, and $[H_{\mathsf{GK}}[\kappa = 0], \Gamma_2] = -2iJ(h/J)^N a_1 \sigma^z_{(1;1)} \cdots \sigma^z_{(N-1;1)}$.

SUPPLEMENTAL MATERIAL

Introduction.— In the main text, we illustrated key ideas by examining several examples: (i) the quantum one-dimensional Ising matter coupled gauge theory (the \mathbb{Z}_2 Higgs model) and its dual quantum spin chain in an external field, (ii) a two-dimensional quantum XY model and its solid-on-solid (\mathbb{Z}) type gauge theory dual, and (iii) the quantum U(1) matter coupled (or Abelian Higgs) gauge theory and its \mathbb{Z} matter field coupled gauge dual.

In what follows, we illustrate how our concepts can be similarly worked out in other examples: (iv) the twodimensional transverse field Ising model and its quantum Ising gauge theory dual, (v) the self-dual two-dimensional quantum \mathbb{Z}_2 Higgs model, (vi) the two-dimensional quantum *p*-clock model and its \mathbb{Z}_p gauge theory dual, (vii) the two-dimensional \mathbb{Z}_p Higgs theory which we earlier illustrated to be self-dual for all *p*, (viii) the two-dimensional \mathbb{Z}_p Higgs theory and its Extended Toric Code dual, and (ix) the quantum XY model on the kagome lattice and its \mathbb{Z} gauge theory dual. Taken together, these examples demonstrate how holographic symmetries and generalized OPs and associated correlators can be systematically derived in disparate theories.

The two-dimensional transverse field Ising model and its quantum Ising gauge theory dual.— As in the main text, we will perform dualities for finite size systems and explicitly label the Cartesian coordinates $1 \le i, j \le N$ of sites $\mathbf{r} = (i, j)$ on an $N \times N$ square lattice. The planar transverse-field Ising model for such any $N \times N$ square lattice system is given by

$$H_{I} = -h \sum_{i,j=1}^{N} \sigma_{(i,j)}^{x}$$

$$-J \sum_{i=1}^{N-1} \sum_{j=1}^{N} \sigma_{(i,j)}^{z} \sigma_{(i+1,j)}^{z} - J \sum_{i=1}^{N} \sum_{j=1}^{N-1} \sigma_{(i,j)}^{z} \sigma_{(i,j+1)}^{z}.$$
(15)

The first term denotes the transverse field along the x direction while last two terms correspond to Ising interactions (along the internal σ^z direction) associated with horizontal and vertical links of the lattice respectively. This transverse field Ising model is dual to (the gaugeinvariant sector of) the \mathbb{Z}_2 Ising gauge theory in the infinite size limit [1]. For open boundary conditions, we obtain via a bond-algebraic duality the *exact* dual of H_1 which reads

$$H_{IG} = -h \sum_{i,j=1}^{N} B_{(i,j)}$$
(16)
$$-J \sum_{i=2}^{N} \sum_{j=1}^{N} \sigma_{(i,j;2)}^{x} - J \sum_{i=1}^{N} \sum_{j=2}^{N} \sigma_{(i,j;1)}^{x}.$$

Following our earlier conventions, the link indices $(i, j; \mu) = (\mathbf{r}; \mu), \ \mu = 1, 2$, that appear *explicitly* in H_{IG}

labeling spins σ^x will be termed*allowed indices*, with all other link indices being *forbidden*. This convention will be invoked to define the plaquette product $B_{(i,j)}$ which will omit forbidden links. The plaquette operators will be of a similar nature to those of Fig. 1 of the main text. In the "bulk", the plaquette operator $B_{(i,j)}$ reads

$$B_{(i,j)} \equiv \sigma_{(i,j;1)}^{z} \sigma_{(i+1,j;2)}^{z} \sigma_{(i,j+1;1)}^{z} \sigma_{(i,j;2)}^{z}.$$
 (17)

At the topmost left site (see Fig. 1), $B_{(1,N)} = \sigma_{(1,N;1)}^z \sigma_{(2,N;2)}^z \sigma_{(1,N+1;1)}^z$. Hence $B_{(1,N)}$ involves one spin $\sigma_{(1,N+1;1)}^z$ labelled by a forbidden link index (see Fig. 1). Elsewhere on the boundary of the square lattice, the remaining boundary plaquettes are determined by Eq. (17) provided spins labelled by forbidden link indices are omitted.

We now examine to the bond algebra- the set of algebraic relations between the interaction terms in the Hamiltonian of Eqs. (15, 16). This will allow us to establish a duality mapping between the transverse field Ising model and the Ising gauge theory. A natural partition of the transverse field Ising Hamiltonian $H_{\rm I}$ of Eq. (15) is into three type of bonds: (a) the transverse fields $\{\sigma_{(i,j)}^x\}$, (b) horizontal Ising interactions $\{\sigma_{(i,j)}^z\sigma_{(i,j+1)}^z\}$. The bond algebra of this system is exhausted by the following relations:

(1) All bonds of type (b) and (c) commute amongst themselves and each other.

(2) Single site bonds of type (a) anticommute with the two-site bonds of type (b) [or (c)] whenever any two such bonds share a common site on the lattice; this is a trivial consequence of the anticommutation relation $\{\sigma_r^x, \sigma_r^z\} = 0.$

(3) Whenever bonds of type (a) share no common lattice sites with bonds of type (b) or (c) then any such bonds commute with one another. That this is so is readily seen as $[\sigma_{\boldsymbol{r}}^{z}, \sigma_{\boldsymbol{r}'}^{z}] = 0$ whenever $\boldsymbol{r} \neq \boldsymbol{r'}$.

(4) The square of any bond is the identity operator $((\sigma_{\boldsymbol{r}}^{x})^{2} = (\sigma_{\boldsymbol{r}'}^{z})\sigma_{\boldsymbol{r}'+\mu}^{z})^{2} = 1)).$

(5) The product of any horizontal/vertical bonds of types (b) and type (c) around any closed plaquette is equal to the identity operator. That is, as $(\sigma^{\boldsymbol{r}})^2 = 1$ for any site \boldsymbol{r} , for all $2 \leq i, j \leq N$,

$$\mathbb{1} = [\sigma_{(i,j)}^{z} \sigma_{(i-1,j)}^{z}] [\sigma_{(i-1,j)}^{z} \sigma_{(i-1,j-1)}^{z}] \\
\times [\sigma_{(i-1,j-1)}^{z} \sigma_{(i,j-1)}^{z}] [\sigma_{(i,j-1)}^{z} \sigma_{(i,j)}^{z}]$$
(18)

Upon removing gauge redundancies, an identical set of relations appears for the bonds of the Ising gauge theory of Eq. (16). To underscore this, we partition Eq. (16) into three types of bonds as follows: (a') the plaquette operators $\{B_{(i,j)}\}$, (b') gauge fields $\sigma_{i,j;2}^x$ on vertical and (c') gauge fields $\sigma_{i,j;1}^x$ on longitudinal links. That similarity of algebraic relations underlies the duality mapping between the theories. That the two theories of Eqs. (15, 16) are indeed dual to each other on an $N \times N$ square lattice (with of an size arbitrary N) is readily seen by the mapping of the bonds $(a,b,c) \rightarrow (a',b',c')$, or

$$B_{(i,j)} \xrightarrow{\Phi_{\mathsf{d}}} \sigma_{(i,j)}^x, \tag{19}$$

$$\sigma_{(i,j;1)}^x \xrightarrow{\Phi_{\mathbf{d}}} \sigma_{(i,j-1)}^z \sigma_{(i,j)}^z, \quad j = 2, \cdots, N, \quad (20)$$

$$\sigma_{(i,j;2)}^x \xrightarrow{\Phi_d} \sigma_{(i-1,j)}^z \sigma_{(i,j)}^z, \quad i = 2, \cdots, N, \qquad (21)$$

once the local gauge symmetries of the Ising gauge theories are removed and set to unity,

$$A_{(i,j)} = \sigma_{(i,j;1)}^x \sigma_{(i,j;2)}^x \sigma_{(i-1,j;1)}^x \sigma_{(i,j-1;2)}^x \xrightarrow{\Phi_{d}} \mathbb{1}.$$
(22)

With the constraint of Eq. (22) implemented, redundant gauge degrees of freedom are eliminated- applying a gauge transformation of the form of Eq. (22) does not lead to a new state. As the mapping of Eqs. (19, 22) preserves all of the algebraic relations amongst the bonds in both theories of Eqs. (15, 16), this mapping implements an duality $H_{\mathsf{IG}} \xrightarrow{\Phi_{\mathsf{d}}} H_{\mathsf{I}}$ for any $N \times N$ square lattice. Similar to the main text, we briefly elaborate on the change of number of degrees of freedom once Eq. (22) is implemented and how this leads to an identical number of degrees of freedom in both the transverse field Ising and Ising gauge theory. The counting of the number of degrees of freedom is identical to that of the U(1) system examined in the main text. We repeat it anew here for the Ising systems at hand. The Hamiltonian H_{IG} describes 2(N-1)N+1 spins and possesses $(N-1)^2$ gauge symmetries which are removed by the constraint of Eq. (22). This leaves a total of $2(N-1)N+1-(N-1)^2 = N^2$ independent spins. This is precisely the number of spins in $H_{\rm I}$.

Now, we turn to a quintessential feature seen by this duality- that of the holographic symmetry. As, by virtue of our definitions above of forbidden indices, $\sigma_{(1,N+1;1)}^z$ makes an appearance only in $B_{(1,N)}$ (see Fig. 1) and $\sigma_{1,N+1;1}^x$ appears nowhere in the Hamiltonian of Eq. (16), the operator $\sigma_{(1,N+1;1)}^z$ constitutes a *boundary symmetry* of H_{IG} . What is this boundary symmetry's origin? The answer is afforded by a simple calculation,

$$\sigma_{(1,N+1;2)}^{z} = \prod_{i,j=1}^{N} B_{(i,j)} \xrightarrow{\Phi_{\mathsf{d}}} Q_{\mathsf{l}}.$$
 (23)

Thus, the boundary symmetry $\sigma_{(1,N+1;2)}^{z}$ is holographic. As seen by Eq. (23), this symmetry is a "hidden" dual of the global \mathbb{Z}_2 symmetry of the Ising model. Its recognition became possible with the specific choice of operators and use of bond algebras which suited to treat exactly for finite size systems and their boundaries. We conclude our examination of this duality and its consequences by deriving the OP of the Ising gauge theory. As we will establish now, this OP will be given by string products- products of operators along connected links. To write this OP, we need to set a few preliminaries. Towards this end, as in the main text, we let Γ denote a path (set of concatenated links) in the Ising model's lattice, starting at site \mathbf{r}_1 and ending at \mathbf{r}_2 so that $\prod_{(\mathbf{r},\mu)\in\Gamma} \sigma_{\mathbf{r}}^z \sigma_{\mathbf{r}+\mathbf{e}_{\mu}}^z = \sigma_{\mathbf{r}_1}^z \sigma_{\mathbf{r}_2}^z$. We furthermore let $(\mathbf{r}^*, \mathbf{e}^*_{\mu})$ denote that link in the lattice of the Ising gauge theory such that $\Phi_{\mathsf{d}}(\sigma_{(\mathbf{r}^*,\mathbf{e}_{\mu}^*)}^x) = \sigma_{\mathbf{r}}^z \sigma_{\mathbf{r}+\mathbf{e}_{\mu}}^z$. From these definitions, it follows that

$$\mathcal{G}_{\Gamma^*} \equiv \langle \mathsf{IG} | \prod_{(\boldsymbol{r}^*, \boldsymbol{e}_{\boldsymbol{\mu}}*) \in \Gamma^*} \sigma^x_{(\boldsymbol{r}^*, \boldsymbol{e}_{\boldsymbol{\mu}}*)} | \mathsf{IG} \rangle \xrightarrow{\Phi_{\mathsf{d}}} \langle \mathsf{I} | \sigma^z_{\boldsymbol{r}_1} \sigma^z_{\boldsymbol{r}_2} | \mathsf{I} \rangle.$$

$$(24)$$

In Eq. (24), $|\mathsf{IG}\rangle$ and $|\mathsf{I}\rangle$ denote, respectively, the ground states of the Ising gauge (Eq. (16)) and Transverse field Ising model (Eq. (15)). Thus, putting all of the pieces together, the (infinite separation limit of the) string correlator \mathcal{G}_{Γ^*} is the generalized OP of the Ising gauge theory. This OP is the dual of the Landau order as it appears in Eq. (??).

The self-dual \mathbb{Z}_2 Higgs model in two dimensions.— We next examine the self-dual quantum Ising matter coupled gauge theory on the square lattice. Similar to its onedimensional variant of Eq. (1), the Hamiltonian of this system is given by

$$H_{\mathsf{H}} = -\sum_{\boldsymbol{r}} (hB_{\boldsymbol{r}} + \lambda \sigma_{\boldsymbol{r}}^x) \tag{25}$$

$$-\sum_{\boldsymbol{r}}\sum_{\mu=1,2}(J\tau^{\boldsymbol{x}}_{(\boldsymbol{r};\mu)}+\kappa\sigma^{\boldsymbol{z}}_{\boldsymbol{r}}\tau^{\boldsymbol{z}}_{(\boldsymbol{r};\mu)}.\sigma^{\boldsymbol{z}}_{\boldsymbol{r}+\boldsymbol{e}_{\mu}}).$$
 (26)

Similar to the main text, In the \mathbb{Z}_2 Higgs Hamiltonian of Eq. (25), $\sigma_{\boldsymbol{r}}^{x,z}$ are Pauli operators that reside on the lattice site \boldsymbol{r} (which play the role of quantum matter fields) while the quantum gauge fields $\tau_{(\boldsymbol{r};\mu)}^{x,z}$ are Pauli operators that lie on a square lattice link that connects the site \boldsymbol{r} with a neighboring site site $\boldsymbol{r} + \mu$ along one of the two ($\mu = 1, 2$) square lattice directions. Following all that we have stated thus far, the reader can verify that the correspondence of bonds

$$B_r \xrightarrow{\Phi_{\mathsf{sd}}} \sigma_r^x,$$
 (27)

$$\sigma_{(\boldsymbol{r};1)}^x \xrightarrow{\Phi_{\rm sd}} \sigma_{\boldsymbol{r}-\boldsymbol{e_2}}^z \tau_{(\boldsymbol{r}-\boldsymbol{e_2};2)}^z \sigma_{\boldsymbol{r}}^z, \qquad (28)$$

$$\sigma_{(\boldsymbol{r};2)}^{x} \xrightarrow{\Phi_{\mathsf{sd}}} \sigma_{\boldsymbol{r}-\boldsymbol{e_1}}^{z} \tau_{(\boldsymbol{r}-\boldsymbol{e_1};1)}^{z} \sigma_{\boldsymbol{r}}^{z}, \tag{29}$$

along with analogous relations for the matter fields [6] proves that H_{Higgs} is self-dual in its gauge-invariant sector. It is straightforward to check that Γ^* maps to the path Γ just as before under the self-duality mapping of Eqs. (28,29). Hence,

$$\langle H| \prod_{(r;\mu)\in\Gamma^*} \sigma^x_{\boldsymbol{r};\mu} |H\rangle \xrightarrow{\Phi_{\mathsf{sd}}} \langle \mathsf{H}| \sigma^z_{\boldsymbol{r}_1} \sigma^z_{\boldsymbol{r}_2} \prod_{(\boldsymbol{r};\mu)\in\Gamma} \sigma^z_{(\boldsymbol{r};\mu)} |\mathsf{H}\rangle,$$
(30)

where $|H\rangle$ denotes the ground state of the Higgs Hamiltonian of Eq. (25). The gauge-invariant correlator on the right-hand side of Eq. (30) is related to known string OPs for Higgs theories [2, 3]. The reader is referred to these works for details concerning asymptotic behaviors of string operators in the confined and deconfined phases. For completeness, we briefly reiterate that in these OPs [2, 3], the expectation value of Eq. (30) is divided by string [2] (or Wilson loop [3] products which are derived from Γ . This division enables the extraction of the nontrivial asymptotic large Γ behavior of the string correlator of Eq. (30); This factor is, equivalently, related to the normalization of the state generated, in an imaginary time representation, by the application of the string product on the ground state [3, 4]. Other relations to effective line tension resulting from combined bare line (borne by matter coupling) and surface (generated by pure gauge fields) tensions are discussed in [4].

The \mathbb{Z}_p Gauge Model.— We may place the \mathbb{Z}_2 theories of the previous sections and the U(1) theories in the main part of the paper at the ends of an infinite sequence of models with \mathbb{Z}_p symmetry, $p = 2, 3, \dots, \infty$.

A "clock model" spin is associated to the group \mathbb{Z}_p is described by two unitary $p \times p$ matrices U, V satisfying $U^p = \mathbb{1} = V^p$ and $VU = \omega UV$, with $\omega = e^{i2\pi/p}$ a *p*th root of unity [5]. When p = 2, these correspond to the Pauli operators $\sigma^{x,z}$. Thus the Hamiltonian for the quantum *p*-clock model can be written as

$$H_{pC} = -\left[\frac{J}{2}\sum_{i=1}^{N-1}\sum_{j=1}^{N}U_{(i,j)}U_{(i+1,j)}^{\dagger}\right]$$
(31)
$$+\frac{J}{2}\sum_{i=1}^{N}\sum_{j=1}^{N-1}U_{(i,j)}U_{(i,j+1)}^{\dagger} + \frac{h}{2}\sum_{i,j=1}^{N}V_{(i,j)}\right] + h.c. .$$

Similar to the \mathbb{Z}_2 theories, the *p*-clock is dual to (the gauge-invariant sector of) the \mathbb{Z}_p gauge theory

$$H_{pG} = -\left[\frac{J}{2}\sum_{i=2}^{N}\sum_{j=1}^{N}V_{(i,j;2)}\right]$$
(32)
$$+\frac{J}{2}\sum_{i=1}^{N}\sum_{j=2}^{N}V_{(i,j;1)} + \frac{h}{2}\sum_{i,j=1}^{N}B_{(i,j)}\right] + h.c. .$$

Following, verbatim, the same convention as earlier (as also marked in Fig. 1)) yet now applied to these operators, the plaquette operator $B_{(i,j)}$ for plaquettes in the "bulk" is given by

$$B_{(i,j)} \equiv U_{(i,j;1)} U_{(i+1,j;2)} U_{(i,j+1;1)}^{\dagger} U_{(i,j;2)}^{\dagger}.$$
 (33)

On the boundary, $B_{(1,N)} = U_{(1,N;1)}U^{\dagger}_{(2,N;2)}U^{\dagger}_{(1,N+1;1)}$. The remaining boundary plaquettes are determined by Eq. (33) provided operators labelled by forbidden link indices (those extend beyond the system boundaries) are omitted, see Fig. 1)). The correspondence of bonds

$$B_{(i,j)} \xrightarrow{\Phi_{\mathbf{d}}} V_{(i,j)}^{\dagger}, \tag{34}$$

$$V_{(i,j;1)} \xrightarrow{\Phi_{\mathsf{d}}} U_{(i,j-1)}^{\dagger} U_{(i,j)}, \quad j = 2, \cdots, N, \quad (35)$$

$$V_{(i,j;2)} \xrightarrow{\Phi_{\mathsf{d}}} U_{(i-1,j)} U_{(i,j)}^{\dagger}, \quad i = 2, \cdots, N,$$
 (36)

establishes the (gauge-reducing) duality $H_{pG} \xrightarrow{\Phi_d} H_{pC}$. The operator $U^{\dagger}_{(1,N+1;1)}$ is an *holographic symmetry* of H_{pG} , since

$$U_{(1,N+1;2)}^{\dagger} = \prod_{i,j=1}^{N} B_{(i,j)} \xrightarrow{\Phi_{\mathsf{d}}} \prod_{i,j=1}^{N} V_{(i,j)}^{\dagger}.$$
 (37)

The operator on the right-hand side of Eq. (37) is the global \mathbb{Z}_p symmetry of the quantum *p*-clock model of Eq. (31). As we see from the left-hand side of Eq. (37), this operator is dual to the *boundary symmetry* of the gauge theory, $U_{(1,N+1;2)}^{\dagger}$. Thus, this symmetry, $U_{(1,N+1;2)}^{\dagger}$, is holographic.

As we have in earlier examples, we now systematically devise an order parameter for the gauge theory. This Abelian gauge theory exhibits, as most others, both confined and deconfined phases. The program for methodically deriving a generalized OP diagnostic will replicate our calculations in earlier examples. First, we write down a Landau correlator which directly captures the long range order in the Landau type system (in this case, the clock model of Eq. (31)). Relying on the duality between the quantum clock model and the quantum \mathbb{Z}_p gauge theory, we then determine the corresponding correlator which serves as a generalized OP for the gauge theory of Eq. (32).

The Landau theory of the quantum p-clock exhibits long range order as seen by the finite value of the twopoint correlation function, in the limit of infinite separation,

$$\langle \mathsf{pC}|U_{(i,j)}U_{(i',j')}^{\dagger}|\mathsf{pC}\rangle, \quad \sqrt{(i-i')^2 + (j-j')^2} \to \infty,$$
(38)

where $|\mathbf{p}\mathbf{C}\rangle$ denotes the ground state of the p-clock model of Eq. (31). We now map this two-point correlator onto a generalized OP for the gauge theory by invoking the duality that we found. Towards this end, our steps will replicate, yet once again, those that we invoked in earlier examples. Let Γ be an *oriented* path from \mathbf{r} to \mathbf{r}' , and $\mathbf{l} \in \Gamma$ its links oriented accordingly. Furthermore, we employ the convention that $U_{\mathbf{l}} = U_{\mathbf{r}}U_{\mathbf{r}+\mathbf{e}_{\mu}}^{\dagger}$ if \mathbf{l} points from \mathbf{r} to $\mathbf{r} + \mathbf{e}_{\mu}$, and $U_{\mathbf{l}} = U_{\mathbf{r}}^{\dagger}U_{\mathbf{r}+\mathbf{e}_{\mu}}$ if \mathbf{l} points oppositely from $\mathbf{r} + \mathbf{e}_{\mu}$ to \mathbf{r} . With this conventions,

$$U_{\boldsymbol{r}}U_{\boldsymbol{r}'}^{\dagger} = \prod_{\boldsymbol{l}\in\Gamma} U_{\boldsymbol{l}}.$$
(39)

If we define Γ^* as the set of links mapping to the path Γ

under duality and $\Phi_{\mathsf{d}}(V_{l^*}) = U_l$, then we will find that

$$\mathcal{G} = \langle \mathsf{pG} | \prod_{\boldsymbol{l}^* \in \Gamma^*} V_{\boldsymbol{l}^*} | \mathsf{pG} \rangle \xrightarrow{\Phi_{\mathsf{d}}} \langle \mathsf{pC} | U_{\boldsymbol{r}} U_{\boldsymbol{r}'}^{\dagger} | \mathsf{pC} \rangle, \qquad (40)$$

where $|\mathbf{pG}\rangle$ is the ground state of the gauge theory $H_{\mathbf{pG}}$. In the confined phase of the gauge theory, the generalized OP \mathcal{G} assumes a non-zero value in the limit of infinite Γ^* . By contrast, in its deconfined phase, the gauge theory of Eq. (32) has a vanishing value of \mathcal{G} as Γ^* is made asymptotically large. This result constitutes a generalization of our earlier Ising (i.e., p = 2) results. Thus, similar to our earlier examples, putting all of the pieces together, the (infinite separation limit of the) string correlator \mathcal{G} is a generalized OP for the \mathbb{Z}_p gauge theory distinguishing confinement from deconfinement, and derived from simple considerations of spontaneous symmetry breakdown, duality, and holographic symmetries.

The self-dual \mathbb{Z}_p Higgs model.— We now extend the earlier Ising results to the \mathbb{Z}_p system, review the selfduality of the \mathbb{Z}_p gauge theory when it is coupled to \mathbb{Z}_p matter fields, and derive new string operators for this theory. This matter coupled gauge theory is the quantum \mathbb{Z}_p Higgs theory. That is, we will augment the (pure) gauge theory H_{pG} of Eq. (32) by terms that minimally couple the gauge fields on the lattice links $U_{r,\mu}$ to matter fields U_r and $U_{r+e_{\mu}}$ at sites that form the endpoints of those links as well as by additional matter only terms V_r ,

$$H_{\text{matter}} = -\sum_{r} \left[\frac{1}{2} V_{r} + \frac{\kappa}{2} \sum_{\mu=1,2} U_{r} U_{(r,\mu)}^{q} U_{r+e_{\mu}}^{\dagger} \right] + h.c..$$
(41)

From the minimal coupling term, where q is the power of the gauge field on the link that is coupled to the matter field, similar to the U(1) example in the main text, it is evident that q plays the role of a charge. As $U^q = U^{(p-q)\dagger}$, this charge can assume the values $q = 1, \dots, Int[p/2]$ (where $Int[\cdot]$ denotes the integer par of its argument). On augmenting the Hamiltonian of Eq. (32) by the terms in Eq. (41), the resulting model Hamiltonian H_{pH} - the (charge q) \mathbb{Z}_p Higgs model- is selfdual in its gauge invariant sector [6]. Specifically, this self-duality is given by the mapping

$$B_{\boldsymbol{r}} \stackrel{\Phi_{\mathrm{sd}}}{\longrightarrow} V_{\boldsymbol{r}}^{\dagger} \tag{42}$$

$$V_{(\boldsymbol{r},1)} \xrightarrow{\Psi_{sd}} U_{\boldsymbol{r}-\boldsymbol{e_2}}^{\dagger} U_{(\boldsymbol{r}-\boldsymbol{e_2},2)}^{q\dagger} U_{\boldsymbol{r}}, \qquad (43)$$

$$V_{(\boldsymbol{r},2)} \xrightarrow{\Phi_{\rm sd}} U_{\boldsymbol{r}-\boldsymbol{e_1}} U^q_{(\boldsymbol{r}-\boldsymbol{e_1},1)} U^\dagger_{\boldsymbol{r}}, \qquad (44)$$

for the gauge fields, and the transformation

$$V_{\boldsymbol{r}}^{\dagger} \xrightarrow{\Phi_{\mathsf{sd}}} B_{\boldsymbol{r}-\boldsymbol{e_1}-\boldsymbol{e_2}}^{\dagger}$$
 (45)

$$U_{\boldsymbol{r}}U^{q}_{(\boldsymbol{r},1)}U^{\dagger}_{\boldsymbol{r}+\boldsymbol{e_{1}}} \xrightarrow{\Phi_{\mathsf{sd}}} V^{\dagger}_{(\boldsymbol{r}-\boldsymbol{e_{2}},2)}, \qquad (46)$$

$$U_{\boldsymbol{r}}^{\dagger}U_{(\boldsymbol{r},2)}^{q\dagger}U_{\boldsymbol{r}+\boldsymbol{e_2}} \xrightarrow{\Phi_{\mathrm{sd}}} V_{(\boldsymbol{r}-\boldsymbol{e_1},1)}^{\dagger}, \qquad (47)$$

for the matter fields.

The string operator in Eq. (40), becomes, on invoking Eqs. (43, 44),

$$\langle \mathsf{pH} | \prod_{l^* \in \Gamma^*} V_{l^*} | \mathsf{pH} \rangle \xrightarrow{\Phi_{\mathsf{sd}}} \langle \mathsf{pH} | U_{\boldsymbol{r}} U_{\boldsymbol{r}'}^{\dagger} \prod_{l \in \Gamma} U_l^q | \mathsf{pH} \rangle.$$
(48)

In Eq. (48), $|\mathbf{pH}\rangle$ denotes the ground state of the Higgs Hamiltonian (given by $(H_{\mathbf{pG}} + H_{\mathbf{matter}})$ - the sum of the Hamiltonians in Eqs. (32, 41)). Similar to our earlier conventions, in Eq. (48) we set $U_{l} = U_{(\boldsymbol{r},\mu)}^{q}$ if l points from \boldsymbol{r} to $\boldsymbol{r} + \boldsymbol{e}_{\mu}$, and $U_{l} = U_{(\boldsymbol{r},\mu)}^{q\dagger}$ otherwise.

As in our discussions for the other examples of Higgs theories concerning string OPs and their normalization, in the infinite separation (i.e., Γ) limit, the gaugeinvariant correlator on the right-hand side is related to a normalized generalized OP proposed in Ref. [2, 3]. Taken on a *closed* path, the right-hand side of Eq. (48) reduces to a Wilson loop.

The \mathbb{Z}_p Extended Toric Code model.— The Toric code model [7] and its extensions, e.g., [6, 8, 9] have gained much interest in recent years. Their original motivation was to serve as a caricature of toy models in which some of the basic concepts of topological quantum computing can be explained [7]. Since the model and its variants have been investigated heavily large on their own right. The original Toric code model had an Ising symmetry. In the current section, we will further employ its \mathbb{Z}_p variant.

In the current context, our interest in this system derives from the fact that the \mathbb{Z}_p Higgs model of the previous section is dual to a generalization of the extended \mathbb{Z}_p Toric Code model (pETC). This model is given by

$$H_{\mathsf{pETC}} = -\sum_{\boldsymbol{r}} \left[\frac{1}{2} A_{\boldsymbol{r}}^{1/q} + \frac{h}{2} B_{\boldsymbol{r}} \right]$$
(49)

+
$$\sum_{\mu=1,2} \left(\frac{J}{2} V_{(\boldsymbol{r},\mu)} + \frac{\kappa}{2} U^{q}_{(\boldsymbol{r},\mu)} \right) \right] + h.c. ,$$
 (50)

on the square lattice, with $A_{\mathbf{r}} = V_{(\mathbf{r},1)}V_{(\mathbf{r},2)}V_{(\mathbf{r}-\mathbf{e}_1,1)}^{\dagger}V_{(\mathbf{r}-\mathbf{e}_2,2)}^{\dagger}$. The standard extended Toric Code is recovered for p = 2, in which case q = 1 is the only possible value. The \mathbb{Z}_p toric code discussed in Ref. [8] is recovered for q = 1.

A gauge-reducing duality connecting H_{pH} to H_{pETC} is given by

while leaving the gauge fields invariant , $B_r \xrightarrow{\Phi_d} B_r$ and $V_{(r,\mu)} \xrightarrow{\Phi_d} V_{(r,\mu)}$. This duality removes degrees of freedom that lie at lattice sites. While the Hamiltonian H_{pETC} has no global symmetries, the \mathbb{Z}_p Higgs model has one global symmetry. Similar to our earlier discussions, the duality $H_{\mathsf{pH}} \xrightarrow{\Phi_d} H_{\mathsf{pETC}}$ maps a global symmetry of H_{pH} to a holographic symmetry of H_{pETC} .

The generalized OPs introduced in the previous section for the \mathbb{Z}_p Higgs model can be mapped to (dual) generalized OPs for the (self-dual) extended Toric Code model. Employing Eqs. (51) on the correlator of Eq. (48),

$$\langle \mathsf{pH} | \prod_{l^* \in \Gamma^*} V_{l^*} | \mathsf{pH} \rangle \xrightarrow{\Phi_{\mathsf{d}}} \langle \mathsf{pETC} | \prod_{l^* \in \Gamma^*} V_{l^*} | \mathsf{pETC} \rangle, \quad (52)$$

and

$$\langle \mathsf{pH}|U_{\boldsymbol{r}}U_{\boldsymbol{r}'}^{\dagger}\prod_{\boldsymbol{l}\in\Gamma}U_{\boldsymbol{l}}^{q}|\mathsf{pH}\rangle \xrightarrow{\Phi_{\mathsf{d}}} \langle \mathsf{pETC}|\prod_{\boldsymbol{l}\in\Gamma}U_{\boldsymbol{l}}^{q}|\mathsf{pETC}\rangle,$$
(53)

where $|\mathsf{pETC}\rangle$ is the ground state of the extended toric code model of Eq. (49) and, as in Eq. (48), $|\mathsf{pH}\rangle$ denotes the ground state of the Higgs model. As H_{pH} is self-dual in its gauge-invariant sector, H_{pETC} is indeed also selfdual. The two string operators of Eqs. (52,53) transform into each other under this self-duality transformation.

The quantum XY model on the kagome lattice.— We now examine a frustrated spin system with a continuous symmetry: the quantum XY model on the open kagome lattice [see Fig. 3,we will denote this lattice by Λ_K],

$$H_{\mathsf{XYK}} = \sum_{\boldsymbol{r} \in \Lambda_K} \frac{1}{2I} L_{\boldsymbol{r}}^2 + \sum_{\langle \boldsymbol{r}, \boldsymbol{r}' \rangle} \kappa \cos(\theta_{\boldsymbol{r}} - \theta_{\boldsymbol{r}'}).$$
(54)

The XY rotor at any kagome lattice site \boldsymbol{r} has an orientation $\theta_{\boldsymbol{r}}$ and $L_{\boldsymbol{r}} = i \frac{\partial}{\partial \theta_{\boldsymbol{r}}}$. This XY Hamiltonian is invariant under global rotations generated by the total angular momentum $L = \sum_{\boldsymbol{r} \in \Lambda_K} L_{\boldsymbol{r}}$. A duality maps this model to a solid-on-solid like (\mathbb{Z}) gauge theory (similar to the one discussed in the main text) which now appears on the dice lattice Λ_D ,

$$H_{\mathsf{XYK}}^{D} = \sum_{\boldsymbol{l}\in\Lambda_{D}} \lambda(R_{\boldsymbol{l}} + R_{\boldsymbol{l}}^{\dagger}) + \sum_{\boldsymbol{p}\in\Lambda_{D}} \frac{1}{2I} \Big(\sum_{\boldsymbol{l}\in\partial\boldsymbol{p}} (-1)^{\boldsymbol{l}} X_{\boldsymbol{l}}\Big)^{2}.$$
(55)

Here, l are the oriented links of the dice lattice, p its plaquettes, and $(-1)^{l} = \pm 1$, with $(-1)^{l} = -1$ only if lviewed as a link in Λ_{D} has opposite orientation to that assigned by $l \in \partial p$ (∂p = oriented boundary of p). Similar to the examples that we discussed earlier, the exact dual possesses incomplete plaquette at the boundaries; this is seen in the lower panel of the figure. Under the duality transformation, the generator of the global rotational symmetry of the XY model maps onto a holographic symmetry,

$$L = \sum_{\boldsymbol{r} \in \Lambda_K} L_r \xrightarrow{\Phi_{\mathsf{d}}} \sum_{\boldsymbol{p} \in \Lambda_D} \sum_{\boldsymbol{l} \in \partial \boldsymbol{p}} (-1)^{\boldsymbol{l}} X_{\boldsymbol{l}} = X_{\boldsymbol{l}'}.$$
(56)

Here, $X_{l'}$ is a degree of freedom placed at the boundary link marked with an open circle. Any other X_l enters



FIG. 3. The quantum, U(1) symmetric XY model on the kagome lattice is holographically dual to a solid-on-solid-like gauge theory on the dice lattice.

the twice sum, with opposite sign. This is then an example of holography for a continuous symmetry in a highly frustrated system.

- * Electronic address: ecobaner@indiana.edu
- [1] F. J. Wegner, J. Math. Phys. **12**, 2259 (1971).
- [2] J. Bricmont and J. Frölich, Phys. Lett. B 122, 73 (1983).
- [3] K. Fredenhagen and M. Marcu, Phys. Rev. Lett. 56, 223 (1986); Comm. Math. Phys. 92, 81 (1983).
- [4] K. Gregor, D. A. Huse, R. Moessner, and S. L. Sondhi, New J.Phys. 13, 025009 (2011).
- [5] G. Ortiz, E. Cobanera, an Z. Nussinov, Nuclear Physics B 854, 780 (2011).
- [6] E. Cobanera, G. Ortiz, and Z. Nussinov, Phys. Rev. Lett. 104, 020402 (2010); Adv. Phys. 60, 679 (2011).
- [7] A. Kitaev, Ann. Phys. 303, 2 (2003).
- [8] M. D. Schulz, S. Dusuel, R. Orus, J. Vidal, and K. P. Schmidt, New J. Phys. 14, 025005 (2012).
- [9] I.S. Tupitsyn, A. Kitaev, N.V. Prokof'ev, and P.C.E. Stamp, Phys. Rev. B 82, 085114 (2010).