Comment on "Evidence of Non-Mean-Field-Like Low-Temperature Behavior in the Edwards-Anderson Spin-Glass Model"

Ref. [1] compares the low-temperature phase of the 3D Edwards-Anderson model (EA) to the Sherrington-Kirkpatrick model (SK), studying the overlap distributions $P_{\mathcal{J}}(q)$ and concluding that the two models behave differently. A similar analysis using state-of-the-art, larger data sets for EA (generated with Janus [2] in [3]) and for SK (from [4]) leads to a very clear interpretation of the results of [1], showing that EA behaves as predicted by the replica symmetry breaking (RSB) theory.

Ref. [1] studies $\Delta(\kappa, q_0)$, probability of finding in $P_{\mathcal{J}}(q)$ a peak greater than κ for $q < q_0$. In a RSB system, $\lim_{N\to\infty} \Delta(\kappa, q_0) = 1$. Fig. 5 of [1] shows that, at fixed q_0 and at the same T/T_c , Δ grows for SK, but seems to reach a plateau for EA. In the inset of Fig. 1 we show that, considering larger systems ($N \leq 32^3$ as opposed to $N \leq 12^3$ of [1]), Δ clearly grows with N also for EA. We use the same value of q_0 as in [1] and T = 0.703. Even this simple analysis is sufficient, when using state-of-theart lattice sizes, to show that Δ has the same qualitative behavior in both models.

Still, the choice of comparing data for different models at the same T/T_c and N does not have a strong basis. Indeed, according to the mean-field picture, the fluctuations of the $P_{\mathcal{J}}(q)$ are ruled by the shape of the averaged P(q) [5], so it is more appropriate to select T such that P(q) is similar for EA and SK. Now, it is universally accepted that the peak at $q = q_{\text{EA}}$ in P(q) grows with Nmore slowly for EA, so the simplest assumption that all the individual peaks for $q < q_{\text{EA}}$ scale at the same rate would already explain the results reported in [1].

According to RSB theory, in the large-N limit $P_{\mathcal{J}}(q) = \sum_{\gamma} W_{\gamma} \delta(q - q_{\gamma})$. Let us assume that for large but finite N the weight distribution is unchanged, but the delta functions are smoothed to a finite height H(N) [6]. The self-averaging peak at $q = q_{\text{EA}}$ will also be smoothed, so we can estimate $H(N) \sim P(q_{\text{EA}})$. $\Delta(\kappa, q_0)$ is the probability of finding a peak with weight $W_{\gamma} > \kappa/H(N)$, which, for small q_0 , is $\Delta(\kappa, q_0) \sim [\kappa/H(N)]^{-I(q_0)} \sim (P(q_{\text{EA}})/\kappa)^{I(q_0)}$, where $I(q_0) = \mathbb{P}(|q| < q_0)$ [5].

We show Δ at T = 0.4 for SK (top) and at T = 0.703for EA (middle), where the temperatures are such that P(0) are very similar (for the largest systems, q_0 ranges from 0.02 to 0.44). The curves show universal scaling for large N. The bottom panel compares Δ for SK and EA using similar effective sizes.

In short, the simple assumption that peaks for all values of q scale at the same rate is consistent with the numerical data and explains the slower growth of Δ with Nfor EA. Therefore, contrary to the claims in [1], we find no quantitative difference between EA and SK, as long as one is careful when comparing non-universal quantities and uses state-of-the-art system sizes.

We have been supported through research contracts nos. 247328 (ERC); FIS2012-35719-C02-01 and FIS2010-16587 (MICINN); and GR10158 (Junta de Extremadura). We thank the Janus Collaboration for granting us use of the EA data.

A. Billoire¹, L.A. Fernandez², A. Maiorano³, E. Marinari³, V. Martin-Mayor², G. Parisi³, F. Ricci-Tersenghi³, J.J. Ruiz-Lorenzo⁴, D. Yllanes³.

¹IPhT, CEA Saclay, 91191 Gif-sur-Yvette, France;

²Dep. Física Teórica I, UCM, 28040 Madrid, Spain;

³Dip. Fisica, La Sapienza Università di Roma, 00185 Roma, Italy;

⁴Dep. Física, Univ. Extremadura, 06071 Badajoz, Spain

- [1] B. Yucesoy, H.G. Katzgraber, and J. Machta, Phys. Rev. Lett. **109**, 177204 (2012) [preprint arXiv:1206.0783].
- [2] M. Baity-Jesi et al., Eur. Phys. J. Special Topics 210, 33 (2012) [preprint arXiv:1204.4134].
- [3] R.A. Baños et al., J. Stat. Mech. (2010) P06026 [preprint arXiv:1003.2569].
- [4] T. Aspelmeier, A. Billoire, E. Marinari, and M.A. Moore, J. Phys. A 41, 324008 (2008) [preprint arXiv:0711.3445].
- [5] G. Parisi, J. Stat. Phys. **72**, 857 (1993).
- [6] R. A. Baños et al., Phys. Rev. B 84, 174209 (2011) [preprint arXiv:1107.5772].




FIG. 1. (color online) $\Delta(\kappa, q_0)$ against $[P(q_{\rm EA})/\kappa]^{I(q_0)}$ for SK (top) and EA (middle). Inset: $\Delta(\kappa, q_0)$ for fixed q_0 for the EA model. Bottom: comparison of the EA and SK models for similar values of $P(q_{\rm EA})$.