

**EXACT CONSTANTS IN POINCARÉ TYPE
INEQUALITIES FOR FUNCTIONS WITH ZERO MEAN
BOUNDARY TRACES**

A. I. NAZAROV AND S. I. REPIN

ABSTRACT. In this paper, we investigate Poincaré type inequalities for the functions having zero mean value on the whole boundary of a Lipschitz domain or on a measurable part of the boundary. We find exact and easily computable constants in these inequalities for some basic domains (rectangles, cubes, and right triangles) and discuss applications of the inequalities to quantitative analysis of partial differential equations.

1. INTRODUCTION

Let Ω be a connected bounded domain in \mathbb{R}^d with Lipschitz boundary $\partial\Omega$. The classical Poincaré inequality reads

$$(1.1) \quad \|w\|_{2,\Omega} \leq C_P(\Omega) \|\nabla w\|_{2,\Omega}, \quad \forall w \in \tilde{H}^1(\Omega),$$

where

$$\tilde{H}^1(\Omega) := \{w \in H^1(\Omega) \mid \{ \! \! \{ w \} \! \! \} _{\Omega} = 0\}.$$

Here and later on $\{ \! \! \{ g \} \! \! \} _{\omega}$ denotes the mean value of g on the set ω while $\|\cdot\|_{2,\omega}$ stands for the norm in $L^2(\omega)$.

The inequality (1.1) was established by H. Poincaré ([12], [13]) for a certain class of domains. V.A. Steklov [16] found that the sharp constant in (1.1) is equal to $\lambda^{-\frac{1}{2}}$, where λ is the smallest positive eigenvalue of the problem

$$(1.2) \quad \begin{aligned} -\Delta u &= \lambda u & \text{in } \Omega; \\ \partial_{\mathbf{n}} u &= 0 & \text{on } \partial\Omega. \end{aligned}$$

L.E. Payne and H.F. Weinberger [11] proved that for convex domains in \mathbb{R}^d this constant satisfies the estimate $C_P(\Omega) \leq \frac{\text{diam}\Omega}{\pi}$. Estimates of constants in Poincaré type inequalities have been studied by many authors (see, e.g., [3, 4, 6, 7, 10] and other publications cited therein).

V.A. Steklov Institute of Mathematics in St. Petersburg, 191023, Fontanka 27, St.-Petersburg, Russia and St. Petersburg State University, 198504, Universitetskii pr. 28, St.-Petersburg, Russia.

V.A. Steklov Institute of Mathematics in St. Petersburg, 191023, Fontanka 27, St.-Petersburg, Russia and University of Jyväskylä, P.O. Box 35 (Agora), FIN-40014, Jyväskylä, Finland.

In this paper, we consider estimates similar to (1.1), for the functions having zero mean on a certain part of the boundary (or on the whole boundary), namely,

$$(1.3) \quad \|w\|_{2,\Omega} \leq C_1(\Omega, \Gamma) \|\nabla w\|_{2,\Omega}, \quad \forall w \in \tilde{H}^1(\Omega, \Gamma),$$

$$(1.4) \quad \|w\|_{2,\Gamma} \leq C_2(\Omega, \Gamma) \|\nabla w\|_{2,\Omega}, \quad \forall w \in \tilde{H}^1(\Omega, \Gamma),$$

where Γ is a measurable part of $\partial\Omega$ (we assume that $(d-1)$ -measure of Γ is positive),

$$\tilde{H}^1(\Omega, \Gamma) = \{w \in H^1(\Omega) \mid \llbracket w \rrbracket_\Gamma = 0\}.$$

Since the quantity $\|\nabla w\|_{2,\Omega} + |\int_\Gamma w ds|$ is a norm equivalent to the original norm of $H^1(\Omega)$, existence of the constants $C_1(\Omega, \Gamma)$ and $C_2(\Omega, \Gamma)$ is easy to prove.

We recall that the extremal function in (1.3) is an eigenfunction $u \in \tilde{H}^1(\Omega, \Gamma)$ of the boundary value problem

$$(1.5) \quad \begin{aligned} -\Delta u &= \lambda u && \text{in } \Omega; \\ \partial_{\mathbf{n}} u &= \mu \equiv -\frac{\lambda}{|\Gamma|} \int_\Omega u dx && \text{on } \Gamma; \\ \partial_{\mathbf{n}} u &= 0 && \text{on } \partial\Omega \setminus \Gamma \end{aligned}$$

corresponding to the least eigenvalue $\lambda > 0$. This fact follows from standard variational arguments.

Analogously, the extremal function in (1.4) is an eigenfunction $u \in \tilde{H}^1(\Omega, \Gamma)$ of the boundary value problem

$$(1.6) \quad \begin{aligned} \Delta u &= 0 && \text{in } \Omega; \\ \partial_{\mathbf{n}} u &= \lambda u && \text{on } \Gamma; \quad \partial_{\mathbf{n}} u = 0 && \text{on } \partial\Omega \setminus \Gamma, \end{aligned}$$

which corresponds to the least positive eigenvalue.

The problem (1.6) in the case $\Gamma = \partial\Omega$ was introduced by V. A. Steklov in [17]. We note also that for $n = 2$ ($n = 3$) and a particular choice of Γ , eigenvalues of the problem (1.6) give so-called *sloshing frequencies* of free oscillations of a liquid in a channel (container, respectively); see, for example, [8], ch. IX.

It is easy to show that the eigenfunctions of the problems (1.5) and (1.6) form complete orthogonal systems in $L^2(\Omega)$ and in $L^2(\Gamma)$, respectively, and the exact constants in (1.3) and (1.4) are equal to $\lambda^{-\frac{1}{2}}$, where λ is the minimal positive eigenvalue. For simply connected domains in \mathbb{R}^2 , eigenvalues of the problem (1.6) were estimated from above in [18, 5] (in fact these estimates are valid for a wider class of problems, which contain a positive weight function on Γ). In [2], it was shown that for doubly connected domains in \mathbb{R}^2 the symmetrization method yields certain estimates of the smallest positive eigenvalue from below.

Our goal is to find exact values of $C_1(\Omega, \Gamma)$ and $C_2(\Omega, \Gamma)$ for certain basic domains (rectangular domains and right triangles). Finding the constants amounts to finding the corresponding minimal positive eigenvalues. In some cases, this goal can be achieved by standard methods. For example, if Ω is a rectangle (parallelepiped) and Γ coincides with one side of Ω , then all the eigenfunctions of (1.5) and (1.6) can be constructed explicitly (by means of the separation of variables method).

However, these type arguments do not work in other cases because it is impossible to construct explicitly *all* eigenfunctions. The goal can be achieved if we find a suitable eigenfunction and prove that it indeed corresponds to the minimal positive eigenvalue. Proving this fact requires rather sophisticated argumentation, which combines analytical and geometrical analysis. Moreover, the proof strongly depends on the domain and boundary conditions so that arguments used for rectangular domains and $\Gamma = \partial\Omega$ considered in Section 2 differ from those used for triangles considered in Section 3.

Section 4 presents an example, which shows that the constants are valuable for quantitative analysis of differential equations. We consider two elliptic boundary value problems with different boundary conditions and source terms. The second problem is a certain simplification of the first one, namely, if the functions presenting source terms and Dirichlet or Neumann boundary conditions are complicated, then they are replaced by simple (piecewise constant or affine) functions. Such a simplified problem is more convenient for numerical methods because simplified boundary conditions can be exactly satisfied by simple approximations and elementwise integrals can be sharply computed by simple quadrature formulas. We deduce an easily computable bound of the difference between two exact solutions. It contains constants in (1.3), (1.4), and Poincaré inequalities. Computation of the bound is simple and needs only integration of known functions. In practice, this estimate is useful if a certain desired tolerance level is stated a priori. The estimate suggests possible simplifications of data and suitable meshes for which the difference is smaller than the tolerance level. Then, in computations (e.g., in the finite element method) we can use the simplified problem instead of the original one. Our analysis is performed for a particular linear elliptic equation, but similar arguments lead to similar estimates for other differential equations associated with the pair of conjugate operators grad and $-\text{div}$. Other applications of (1.3) and (1.4) are related to a posteriori error estimation methods for partial differential equations, where computable bounds between exact solutions and approximations often involve constants in Poincaré type inequalities (see [14]).

2. EXACT CONSTANTS FOR RECTANGLES AND PARALLELEPIPEDES

In this section, we deal with rectangles and parallelepipeds. First, we consider the simplest case where Γ coincides with one side of Ω .

Proposition 2.1. 1. *Let $\Omega = (0, h_1) \times (0, h_2)$ and $\Gamma = \{x_1 = 0, x_2 \in [0, h_2]\}$. Then the sharp constants in (1.3) and (1.4) are equal to $\frac{1}{\pi} \max\{2h_1; h_2\}$ and $\left(\frac{\pi}{h_2} \tanh\left(\frac{\pi h_1}{h_2}\right)\right)^{-\frac{1}{2}}$, respectively.*

2. *Let $\Omega = (0, h_1) \times (0, h_2) \times (0, h_3)$ and $\Gamma = \partial\Omega \cap \{x_1 = 0\}$. Then the sharp constants in (1.3) and (1.4) are equal to*

$$\frac{1}{\pi} \max\{2h_1; h_2; h_3\} \quad \text{and} \quad \left(\frac{\pi}{\max\{h_2; h_3\}} \tanh\left(\frac{\pi h_1}{\max\{h_2; h_3\}}\right)\right)^{-\frac{1}{2}},$$

respectively.

Proof. In this case, all the eigenfunctions of the problems (1.5) and (1.6) can be found by the separation of variables. For instance, the eigenfunctions of (1.6) in the rectangle are defined by the relation

$$u_k(x) = \cos\left(\frac{\pi k}{h_2} x_2\right) \cosh\left(\frac{\pi k}{h_2} (x_1 - h_1)\right), \quad k = 0, 1, 2, \dots$$

and evidently form a complete orthogonal system in $L^2(\Gamma)$. Therefore, the corresponding least eigenvalue of the problem (1.6) is

$$\lambda_1 = \frac{\pi}{h_2} \tanh\left(\frac{\pi h_1}{h_2}\right).$$

The proof of other statements is quite similar, and we omit it. \square

Now we turn to the case $\Gamma = \partial\Omega$. The problem in a rectangle is symmetric with respect to two axes. Therefore, it is convenient to select the coordinate system such that $\Omega = (-\frac{h_1}{2}, \frac{h_1}{2}) \times (-\frac{h_2}{2}, \frac{h_2}{2})$ (see Fig. 1).

Theorem 2.1. *Let $\Omega = (-\frac{h_1}{2}, \frac{h_1}{2}) \times (-\frac{h_2}{2}, \frac{h_2}{2})$ and $\Gamma = \partial\Omega$. Then the sharp constant in (1.3) is equal to $\frac{1}{\pi} \max\{h_1; h_2\}$.*

Proof. Due to the biaxial symmetry all the eigenfunctions of (1.5) and (1.6) can be chosen either even or odd with respect to the axes x_1 and x_2 . First, we consider the eigenfunctions of (1.5), which are odd with respect to x_1 . In this case, $\mu = 0$ and we arrive at the following problem:

$$(2.1) \quad \begin{aligned} -\Delta u &= \lambda u \quad \text{in} \quad \Omega^+ := \left(0, \frac{h_1}{2}\right) \times \left(-\frac{h_2}{2}, \frac{h_2}{2}\right), \\ u &= 0 \quad \text{on} \quad \{x_1 = 0\} \cap \Omega, \quad \partial_{\mathbf{n}} u = 0 \quad \text{on} \quad \partial\Omega^+ \setminus \{x_1 = 0\}. \end{aligned}$$

It is easy to see that the eigenfunctions of (2.1) are defined by the relations

$$\begin{aligned} u_{km}^{(1)}(x) &= \sin\left(\frac{\pi(2k+1)}{h_1} x_1\right) \cos\left(\frac{2\pi m}{h_2} x_2\right), \\ u_{km}^{(2)}(x) &= \sin\left(\frac{\pi(2k+1)}{h_1} x_1\right) \sin\left(\frac{\pi(2m+1)}{h_2} x_2\right), \quad k, m = 0, 1, \dots \end{aligned}$$

They form a system of orthogonal functions, which is complete in $L^2(\Omega^+)$. Hence, we conclude that the least eigenvalue of the problem (2.1) is $\lambda_{00}^{(1)} = \left(\frac{\pi}{h_1}\right)^2$.




FIGURE 1. Ω , Ω^+ , Ω^{++} , and Γ^+ .

All eigenfunctions of the problem (1.5), which are odd with respect to x_2 can be constructed quite similarly and the corresponding least eigenvalue is $\left(\frac{\pi}{h_2}\right)^2$.

It remains to consider the eigenfunctions, which are even with respect to both variables. They belong to the space $\tilde{H}^1(\Omega^{++}, \Gamma^+)$, where

$$\Omega^{++} := \left(0, \frac{h_1}{2}\right) \times \left(0, \frac{h_2}{2}\right) \quad \text{and} \quad \Gamma^+ = \Gamma \cap \partial\Omega^{++}.$$

In this case, we need to solve the problem

$$(2.2) \quad \begin{aligned} -\Delta u &= \lambda u \quad \text{in} \quad \Omega^{++}; \\ \partial_{\mathbf{n}} u &= \mu \quad \text{on} \quad \Gamma^+; \quad \partial_{\mathbf{n}} u = 0 \quad \text{on} \quad \partial\Omega^{++} \setminus \Gamma^+. \end{aligned}$$

The eigenvalues λ_k^e of the problem (2.2) (enumerated in the increasing order and repeated according to their multiplicity) are critical values of the Rayleigh quotient

$$(2.3) \quad Q[v] \equiv \frac{\|\nabla v\|_{2, \Omega^{++}}^2}{\|v\|_{2, \Omega^{++}}^2},$$

on the space $\tilde{H}^1(\Omega^{++}, \Gamma^+)$.

Consider the functional Q on the whole space $H^1(\Omega^{++})$. By the variational principle (see, e.g., [1], (1.15)), its critical values $\tilde{\lambda}_k^e$ enumerated in the increasing order and repeated in accordance with their multiplicity satisfy the relation¹ $\tilde{\lambda}_k^e \leq \lambda_k^e \leq \tilde{\lambda}_{k+1}^e$. Therefore, if there exists an eigenvalue of the problem (2.2) in the interval $(\tilde{\lambda}_0^e, \tilde{\lambda}_1^e)$, then it must be λ_0^e .

Note that $\tilde{\lambda}_k^e$ are eigenvalues of the conventional Neumann problem

$$-\Delta u = \lambda u \quad \text{in} \quad \Omega^{++}; \quad \partial_{\mathbf{n}} u = 0 \quad \text{on} \quad \partial\Omega^{++},$$

¹Note that $\tilde{H}^1(\Omega^{++}, \Gamma^+)$ has codimension 1 in $H^1(\Omega^{++})$.

and thus, $\tilde{\lambda}_0^e = 0$, $\tilde{\lambda}_1^e = \min\{(\frac{2\pi}{h_1})^2; (\frac{2\pi}{h_2})^2\}$.

Let us consider the equation

$$(2.4) \quad \frac{h_1}{2} \cot\left(\frac{\omega h_2}{2}\right) + \frac{h_2}{2} \cot\left(\frac{\omega h_1}{2}\right) + \frac{2}{\omega} = 0.$$

It is easy to see that the function in the left-hand side of (2.4) decreases from $+\infty$ to $-\infty$ in the interval $(0, \min\{\frac{2\pi}{h_1}; \frac{2\pi}{h_2}\})$, so that the equation has a unique solution ω_0 in this interval. The equation (2.4) arises in our analysis by the following reasons. Direct calculation shows that the function

$$v_0(x) = \frac{\cos(\omega_0 x_1)}{\sin(\frac{\omega_0 h_1}{2})} + \frac{\cos(\omega_0 x_2)}{\sin(\frac{\omega_0 h_2}{2})}$$

solves the problem (2.2) with $\lambda = \omega_0^2$. Then, the condition $\int_{\Gamma^+} v_0 ds = 0$ infers the equation (2.4).

Thus, we conclude that $\lambda_0^e = \omega_0^2$. However, it is easy to see that

$$\omega_0 > \min\left\{\frac{\pi}{h_1}; \frac{\pi}{h_2}\right\}.$$

Therefore, the least eigenvalue of the problem (1.5) is $\min\{(\frac{\pi}{h_1})^2; (\frac{\pi}{h_2})^2\}$, and the statement follows. \square

Theorem 2.2. *If $\Omega = (-\frac{h_1}{2}, \frac{h_1}{2}) \times (-\frac{h_2}{2}, \frac{h_2}{2})$ and $\Gamma = \partial\Omega$, then the smallest constant $C_2(\Omega, \Gamma)$ in (1.4) equals $\left(\frac{2z_0(\alpha_0)}{\sqrt{h_1 h_2}} \tanh(\frac{z_0(\alpha_0)}{\alpha_0})\right)^{-\frac{1}{2}}$, where $\alpha_0 = \sqrt{\frac{\max\{h_1; h_2\}}{\min\{h_1; h_2\}}}$ and $z_0(\alpha)$ is a unique root of the equation*

$$(2.5) \quad \tanh\left(\frac{z}{\alpha}\right) \tan(z\alpha) = 1,$$

such that $0 < z_0\alpha < \frac{\pi}{2}$.

Proof. First, we consider the eigenfunctions of (1.6), which are even with respect to both variables. They belong to the space $\tilde{H}^1(\Omega^{++}, \Gamma^+)$ and solve the following problem:

$$(2.6) \quad \begin{aligned} \Delta u &= 0 & \text{in } \Omega^{++}; \\ \partial_{\mathbf{n}} u &= \lambda u & \text{on } \Gamma^+; \quad \partial_{\mathbf{n}} u = 0 & \text{on } \partial\Omega^{++} \setminus \Gamma^+. \end{aligned}$$

Moreover, the eigenvalues Λ_k^e of the problem (2.6) (complemented by zero, enumerated in the increasing order, and repeated according to their multiplicity) are critical values of the Rayleigh quotient

$$\mathcal{Q}^+[v] \equiv \frac{\|\nabla v\|_{2, \Omega^{++}}^2}{\|v\|_{2, \Gamma^+}^2}$$

over the space $H^1(\Omega^{++})$. Consider another Rayleigh quotient

$$\tilde{\mathcal{Q}}[v] \equiv \frac{\|\nabla v\|_{2, \Omega^{++}}^2}{\|v\|_{2, \partial\Omega^{++}}^2}$$

on the same space. Since $\tilde{\mathcal{Q}}[v] \leq \mathcal{Q}^+[v]$, by the variational principle its critical values $\tilde{\Lambda}_k^e$ (which are also enumerated in the increasing order and repeated according to their multiplicity) satisfy the relation $\tilde{\Lambda}_k^e \leq \Lambda_k^e$. However, by scaling arguments $\tilde{\Lambda}_k^e = 2\lambda_k$, where λ_k is the k -th eigenvalue of the original problem (1.6). Therefore, an eigenfunction of (1.6) even with respect to both variables cannot generate the least eigenvalue².

Let us consider the eigenfunctions odd with respect to x_1 . They lead to the following problem:

$$(2.7) \quad \begin{aligned} \Delta u &= 0 & \text{in } \Omega^+; & \quad u = 0 & \text{on } \{x_1 = 0\}; \\ \partial_{\mathbf{n}} u &= \lambda u & \text{on } \partial\Omega^+ \setminus \{x_1 = 0\}. \end{aligned}$$

We claim that the eigenfunction of (2.7) corresponding to the least eigenvalue must be of constant sign in Ω^+ . Indeed, the function

$$v(x_1, x_2) = |u(|x_1|, x_2)| \cdot \text{sign}(x_1)$$

belongs to $\tilde{H}^1(\Omega, \Gamma)$ and provides the same value λ of the Rayleigh quotient

$$(2.8) \quad \mathcal{Q}[v] \equiv \frac{\|\nabla v\|_{2,\Omega}^2}{\|v\|_{2,\Gamma}^2}$$

as u . If λ coincides with the infimum of \mathcal{Q} on $\tilde{H}^1(\Omega, \Gamma)$, then v must be a solution of (1.6). By the maximum principle, v cannot vanish in Ω^+ , and the claim follows. Moreover, since the eigenfunctions of (2.7) are orthogonal in $L^2(\partial\Omega^+ \setminus \{x_1 = 0\})$, no eigenfunction except the first one can be positive in Ω^+ .

Now we consider the equation

$$(2.9) \quad \tan\left(\frac{\omega h_1}{2}\right) \tanh\left(\frac{\omega h_2}{2}\right) = 1,$$

which obviously has a unique solution ω_1 in the interval $(0, \frac{2\pi}{h_1})$. Direct calculation shows that the function

$$v_1(x) = \sin(\omega_1 x_1) \cosh(\omega_1 x_2)$$

²We can suggest another proof of this fact, which is interesting by itself. Let u be a solution of (2.6). We claim that at least one of sets $\varpi_{\pm} = \Omega^{++} \cap \{u \gtrless 0\}$ has a connected component which touches Γ^+ but does not touch one of the coordinate axes. Indeed, consider a connected component of ϖ_+ touching Γ^+ (in view of the condition $\int_{\Gamma^+} u = 0$, such a component exists). If this component touches both axes then any connected component of ϖ_- touching Γ^+ is separated either from $\{x_1 = 0\}$ or from $\{x_2 = 0\}$. To be definite, let ϖ be a connected component of ϖ_- which touches Γ^+ but does not touch $\{x_1 = 0\}$. Then the function

$$v(x_1, x_2) = u(|x_1|, |x_2|) \cdot \chi_{\varpi}(|x_1|, |x_2|) \cdot \text{sign}(x_1)$$

belongs to $\tilde{H}^1(\Omega, \Gamma)$ and provides the same value Λ of the Rayleigh quotient (2.8) as u . If Λ minimizes \mathcal{Q} on $\tilde{H}^1(\Omega, \Gamma)$ then v should be a solution of (1.6) which is impossible by the maximum principle. Unfortunately, this argument is purely 2-dimensional.

is positive in Ω^+ and solves the problem (2.7) with $\lambda = \omega_1 \tanh\left(\frac{\omega_1 h_2}{2}\right)$ (the equation (2.9) reflects the equality of $\partial_{\mathbf{n}}u/u$ on sides of rectangle). We substitute $z_0 = \frac{\omega_1}{2} \sqrt{h_1 h_2}$ and conclude that the least eigenvalue of (2.7) is equal to $\frac{2z_0(\sqrt{\kappa})}{\sqrt{h_1 h_2}} \tanh\left(\frac{z_0(\sqrt{\kappa})}{\sqrt{\kappa}}\right)$, where $z_0(\sqrt{\kappa})$ is the root of (2.5) with $\alpha = \sqrt{\kappa} := \sqrt{\frac{h_1}{h_2}}$.

The eigenfunctions of (1.6), which are odd with respect to x_2 , can be constructed in a similar way. The corresponding least eigenvalue is equal to $\frac{2z_0(\frac{1}{\sqrt{\kappa}})}{\sqrt{h_1 h_2}} \tanh\left(\sqrt{\kappa} z_0\left(\frac{1}{\sqrt{\kappa}}\right)\right)$, where $z_0\left(\frac{1}{\sqrt{\kappa}}\right)$ is the root of (2.5) with $\alpha = \frac{1}{\sqrt{\kappa}}$.

To complete the proof it suffices to show that the function $f(\alpha) = z_0 \tanh\left(\frac{z_0}{\alpha}\right)$ decreases on $(0, +\infty)$. We claim that, in fact, $\alpha f(\alpha)$ is a decreasing function. Indeed, differentiation of (2.5) after some transformations yields

$$\frac{d}{d\alpha} (\alpha f(\alpha)) = \frac{2z_0(1 - \tanh^4\left(\frac{z_0}{\alpha}\right))}{1 + \alpha^2 - \tanh^2\left(\frac{z_0}{\alpha}\right)(1 - \alpha^2)} \cdot \left[\frac{\tanh\left(\frac{z_0}{\alpha}\right)}{1 + \tanh^2\left(\frac{z_0}{\alpha}\right)} - z_0\alpha \right].$$

The fraction here is obviously positive. Further, (2.5) implies $z_0\alpha > \frac{\pi}{4}$. Therefore,

$$\frac{\tanh\left(\frac{z_0}{\alpha}\right)}{1 + \tanh^2\left(\frac{z_0}{\alpha}\right)} - z_0\alpha < \frac{1}{2} - \frac{\pi}{4} < 0,$$

and the claim follows. \square

Let us now consider three-dimensional generalizations of Theorems 2.1 and 2.2.

Theorem 2.3. *Let $\Omega = \left(-\frac{h_1}{2}, \frac{h_1}{2}\right) \times \left(-\frac{h_2}{2}, \frac{h_2}{2}\right) \times \left(-\frac{h_3}{2}, \frac{h_3}{2}\right)$ and let $\Gamma = \partial\Omega$. Then the exact constant in (1.3) is equal to $\frac{1}{\pi} \max\{h_1; h_2; h_3\}$.*

Proof. The proof is similar to the proof of Theorem 2.1. Instead of (2.4) we obtain the equation

$$\frac{h_1 h_2}{2} \cot\left(\frac{\omega h_3}{2}\right) + \frac{h_1 h_3}{2} \cot\left(\frac{\omega h_2}{2}\right) + \frac{h_2 h_3}{2} \cot\left(\frac{\omega h_1}{2}\right) + \frac{2}{\omega} (h_1 + h_2 + h_3) = 0.$$

The unique solution of this equation in $(0, \frac{2\pi}{h})$ (where $h = \max\{h_1; h_2; h_3\}$) is greater than $\frac{\pi}{h}$, and the statement follows. \square

Theorem 2.4. *Let Ω and $\Gamma = \partial\Omega$ be as in Theorem 2.3. To be definite, assume that $\max\{h_1; h_2; h_3\} = h_3$. Then the exact constant in (1.4) is equal to $\left(\frac{z_1}{h_1} \tanh(z_1)\right)^{-\frac{1}{2}}$, where (z_1, z_2) is a unique solution of the system*

$$(2.10) \quad \begin{aligned} \frac{z_1}{h_1} \tanh(z_1) &= \frac{z_2}{h_2} \tanh(z_2) = \\ &= \frac{z_1}{h_1} \sqrt{1 + \frac{\tanh^2(z_1)}{\tanh^2(z_2)}} \cdot \cot\left(\frac{z_1 h_3}{h_1} \sqrt{1 + \frac{\tanh^2(z_1)}{\tanh^2(z_2)}}\right), \end{aligned}$$

such that $0 < \frac{z_1 h_3}{h_1} \sqrt{1 + \frac{\tanh^2(z_1)}{\tanh^2(z_2)}} < \frac{\pi}{2}$.

Proof. By the same arguments as in the proof of Theorem 2.2, we conclude that the eigenfunction of (1.6) corresponding to the least eigenvalue must be odd with respect to one of the coordinate axes (e.g., with respect to x_3). This gives the following analog of (2.7):

$$(2.11) \quad \begin{aligned} \Delta u &= 0 & \text{in } \widehat{\Omega}^+ &:= \left(-\frac{h_1}{2}, \frac{h_1}{2}\right) \times \left(-\frac{h_2}{2}, \frac{h_2}{2}\right) \times \left(0, \frac{h_3}{2}\right); \\ u &= 0 & \text{on } \{x_3 = 0\}; \\ \partial_{\mathbf{n}} u &= \Lambda u & \text{on } \partial\widehat{\Omega}^+ \setminus \{x_3 = 0\}. \end{aligned}$$

Repeating the proof of Theorem 2.2, we find that the least eigenvalue is generated by the eigenfunction of (2.11), which is positive in $\widehat{\Omega}^+$.

It is easy to see that the equation

$$(2.12a) \quad \mu \tanh\left(\frac{\mu h_1}{2}\right) = \nu \tanh\left(\frac{\nu h_2}{2}\right)$$

defines an increasing function $\nu = \nu(\mu)$ on \mathbb{R}_+ , and the equation

$$(2.12b) \quad \mu \tanh\left(\frac{\mu h_1}{2}\right) = \sqrt{\mu^2 + \nu^2(\mu)} \cdot \cot\left(\sqrt{\mu^2 + \nu^2(\mu)} \frac{h_3}{2}\right)$$

has a unique solution μ_0 such that $\sqrt{\mu_0^2 + \nu^2(\mu_0)} \frac{h_3}{2} < \frac{\pi}{2}$. Direct calculation shows that the function

$$U_1(x) = \cosh(\mu_0 x_1) \cosh(\nu(\mu_0) x_2) \sin\left(\sqrt{\mu_0^2 + \nu^2(\mu_0)} x_3\right)$$

is positive in $\widehat{\Omega}^+$ and solves the problem (2.11) with $\Lambda = \mu_0 \tanh\left(\frac{\mu_0 h_1}{2}\right)$ (note that (2.12) reflects the boundary conditions on the sides of parallelepiped).

Substituting $z_1 = \frac{\mu_0 h_1}{2}$, $z_2 = \frac{\nu(\mu_0) h_2}{2}$, we conclude that the least eigenvalue of (2.11) equals $\frac{2z_1}{h_1} \tanh(z_1)$, where (z_1, z_2) is the solution of (2.10).

The eigenfunctions odd with respect to other coordinates can be constructed quite analogously. However, some additional calculations show that U_1 is the best eigenfunction provided that h_3 is the longest edge of Ω . Thus, the statement follows. \square

3. EXACT CONSTANTS FOR ISOSCELES RIGHT TRIANGLES

In this section, Ω is an isosceles right triangle. We find sharp constants in (1.3) and (1.4) for the following three cases: Γ is a leg, Γ is the union of two legs, and Γ is the hypotenuse. Finding exact constants in (1.3) and (1.4) for the case $\Gamma = \partial\Omega$ remains an open problem.

3.1. Case 1: Γ is a leg. It is convenient to select the coordinate system such that $\Omega = \{0 < x_2 < x_1 < h\}$ and $\Gamma = \{x_1 = h, x_2 \in [0, h]\}$ (see Fig. 2a).

Theorem 3.1. *The exact constant in (1.3) is equal to $\widetilde{z}_0^{-1} h$, where $\widetilde{z}_0 \approx 2.02876$ is a unique root of the equation*

$$(3.1) \quad z \cot(z) + 1 = 0$$




FIGURE 2. Right triangles

in the interval $(0, \pi)$.

Proof. We use the same arguments as in the proof of Theorem 2.1 and note that the eigenvalues λ_k^Δ of the problem (1.5) (enumerated in the increasing order and repeated according to their multiplicity) are critical values of the Rayleigh quotient $Q[v] \equiv \frac{\|\nabla v\|_{2,\Omega}^2}{\|v\|_{2,\Omega}^2}$ over the space $\tilde{H}^1(\Omega, \Gamma)$.

Let us consider $Q[v]$ on the whole space $H^1(\Omega)$. In accordance with the variational principle, the corresponding critical values $\tilde{\lambda}_k^\Delta$ (which are also enumerated in the increasing order and repeated according to their multiplicity) satisfy the relation $\tilde{\lambda}_k^\Delta \leq \lambda_k^\Delta \leq \tilde{\lambda}_{k+1}^\Delta$. Therefore, if the interval $(\tilde{\lambda}_0^\Delta, \tilde{\lambda}_1^\Delta)$ contains an eigenvalue of the problem (1.5), then it must be λ_0^Δ .

Note that $\tilde{\lambda}_k^\Delta$ are eigenvalues of the conventional Neumann problem

$$(3.2) \quad -\Delta u = \lambda u \quad \text{in } \Omega; \quad \partial_{\mathbf{n}} u = 0 \quad \text{on } \partial\Omega.$$

By even reflection with respect to the line $\{x_1 = x_2\}$, we conclude that any eigenfunction of (3.2) is an eigenfunction of the Neumann problem in the square $(0, h) \times (0, h)$. In particular, $\tilde{\lambda}_0^\Delta = 0$ corresponds to the eigenfunction $\tilde{u}_0 \equiv 1$, and $\tilde{\lambda}_1^\Delta = \left(\frac{\pi}{h}\right)^2$ corresponds to the eigenfunction $\tilde{u}_1(x) = \cos\left(\frac{\pi x_1}{h}\right) + \cos\left(\frac{\pi x_2}{h}\right)$.

Direct calculation shows that the function

$$\tilde{v}_0(x) = \cos\left(\frac{\tilde{z}_0 x_1}{h}\right) + \cos\left(\frac{\tilde{z}_0 x_2}{h}\right)$$

solves the problem (1.5) with $\lambda = \left(\frac{\tilde{z}_0}{h}\right)^2$ (the equation (3.1) is just the condition $\int_{\Gamma} \tilde{v}_0 ds = 0$). Thus, we conclude that $\lambda_0^\Delta = \left(\frac{\tilde{z}_0}{h}\right)^2$, and the statement follows. \square

Theorem 3.2. *The sharp constant in (1.4) is equal to $\left(\frac{\tilde{z}_0}{h} \tanh(\tilde{z}_0)\right)^{-\frac{1}{2}}$ where $\tilde{z}_0 \approx 2.3650$ is a unique root of the equation*

$$(3.3) \quad \tan(z) + \tanh(z) = 0$$

in the interval $(0, \pi)$.

Proof. Let v be the minimizer of the Rayleigh quotient $\mathcal{Q}[v] \equiv \frac{\|\nabla v\|_{2,\Omega}^2}{\|v\|_{2,\Gamma}^2}$ over the space $\tilde{H}^1(\Omega, \Gamma)$. Since the trace $v|_\Gamma$ is orthogonal to 1 in $L^2(\Gamma)$, its primitive $F(y) = \int_0^y v(h, t) dt$ vanishes at 0 and at h .

We claim that F has constant sign on $(0, h)$. Indeed, otherwise there exists $\alpha \in (0, 1)$ such that

$$(3.4) \quad \int_0^{\alpha h} v(h, x_2) dx_2 = \int_{\alpha h}^h v(h, x_2) dx_2 = 0.$$

We split Ω into two subdomains

$$\Omega_1 = \Omega \cap \{x_2 < \alpha x_1\} \quad \text{and} \quad \Omega_2 = \Omega \cap \{x_2 > \alpha x_1\}.$$

Let us compare the respective Rayleigh quotients

$$\mathcal{Q}_j[v] = \frac{\|\nabla v\|_{2,\Omega_j}^2}{\|v\|_{2,\Gamma_j}^2}, \quad j = 1, 2,$$

where $\Gamma_j := \Gamma \cap \partial\Omega_j$. Assume that $\mathcal{Q}_1[v] \leq \mathcal{Q}_2[v]$. We define the function $V(x_1, x_2) = v(x_1, \alpha x_2)$. Then $\int_\Gamma V dx_2 = 0$ by (3.4), and

$$\mathcal{Q}[V] = \frac{\int_\Omega (V_{x_1}^2 + V_{x_2}^2) dx_1 dx_2}{\int_\Gamma V^2 dx_2} = \frac{\int_{\Omega_1} (v_{x_1}^2 + \alpha^2 v_{x_2}^2) dx_1 dx_2}{\int_{\Gamma_1} v^2 dx_2} \leq \mathcal{Q}_1[v] \leq \mathcal{Q}[v].$$

Moreover, the first inequality in the above relations is strict. Indeed, the converse implies $v_{x_2} \equiv 0$, what is impossible since v is harmonic.

Analogously, if $\mathcal{Q}_2[v] \leq \mathcal{Q}_1[v]$, then we consider the function $V(x_1, x_2) = v(x_1, (1 - \alpha)x_2 + \alpha h)$ and again find that $\mathcal{Q}[V] < \mathcal{Q}[v]$. Thus, the claim follows.

Further, direct calculation shows that the function

$$\tilde{v}_1(x) = \cos\left(\frac{\hat{z}_0 x_1}{h}\right) \cosh\left(\frac{\hat{z}_0 x_2}{h}\right) + \cosh\left(\frac{\hat{z}_0 x_1}{h}\right) \cos\left(\frac{\hat{z}_0 x_2}{h}\right)$$

solves the problem (1.6) with $\lambda = \frac{\hat{z}_0}{h} \tanh(\hat{z}_0)$ (the equation (3.3) is just the condition $\int_\Gamma \tilde{v}_1 ds = 0$). Since $\frac{\pi}{2} < \hat{z}_0 < \pi$, $\tilde{v}_1|_\Gamma$ is monotone decreasing in x_2 and, in particular, the primitive of $\tilde{v}_1|_\Gamma$ has constant sign.

Finally, any other eigenfunction v of the problem (1.6) can be chosen orthogonal to 1 and to \tilde{v}_1 in $L^2(\Gamma)$. But if F (the primitive of $v|_\Gamma$) has a constant sign (e.g., positive), then we arrive at a contradiction because

$$\int_0^h v(h, t) \tilde{v}_1(h, t) dt = - \int_0^h F(h, t) \tilde{v}_1'(h, t) dt > 0.$$

Thus, \tilde{v}_1 should minimize $\mathcal{Q}[v]$ over $\tilde{H}^1(\Omega, \Gamma)$. The proof is complete. \square

3.2. Case 2: Γ is the union of two legs. Let (see Fig. 2b)

$$\Omega = \{0 < x_2 < x_1 < h\} \quad \text{and} \quad \Gamma = \{x_1 = h, x_2 \in [0, h]\} \cup \{x_2 = 0, x_1 \in [0, h]\}.$$

Theorem 3.3. *The sharp constant in (1.3) is equal to $\frac{h}{\pi}$.*

Proof. Making even reflection with respect to the line $\{x_1 = x_2\}$, we conclude that any eigenfunction of (1.5) is an eigenfunction of the same problem in the square domain $\Omega' = (0, h) \times (0, h)$ with $\Gamma = \partial\Omega'$. The corresponding eigenvalue problem has been studied in Theorem 2.1, which states that the least positive eigenvalue is equal to $(\frac{\pi}{h})^2$. The corresponding eigenspace has the dimension 2 and contains the function $\cos(\frac{\pi}{h}x_1) + \cos(\frac{\pi}{h}x_2)$ which solves the original problem in the triangle. \square

Theorem 3.4. *The sharp constant in (1.4) is equal to $(\frac{2z_0}{h} \tanh(z_0))^{-\frac{1}{2}}$, where $z_0 \approx 0.93755$ is a unique root of the equation (2.5) in $(0, \frac{\pi}{2})$ for $\alpha = 1$.*

Proof. Again, we use even reflection with respect to the line $\{x_1 = x_2\}$ and reduce our problem to the problem in the square domain $(0, h) \times (0, h)$. By Theorem 2.2, we conclude that the least positive eigenvalue is equal to $\frac{2z_0}{h} \tanh(z_0)$. The corresponding eigenspace has the dimension 2 and contains the function

$$\sin \xi_1 \cosh \xi_2 + \cosh \xi_1 \sin \xi_2,$$

where $\xi_1 = z_0 \left(\frac{2x_1}{h} - 1\right)$ and $\xi_2 = z_0 \left(\frac{2x_2}{h} - 1\right)$. It solves the original problem in the triangle. \square

3.3. Case 3: Γ is the hypotenuse. In this case, it is convenient to select the coordinate system such that (see Fig. 2c)

$$\Omega = \{0 < |x_2| < x_1 < h\} \quad \text{and} \quad \Gamma = \{x_1 = h, x_2 \in [-h, h]\}.$$

Theorem 3.5. *The sharp constant in (1.3) is equal to $\tilde{z}_0^{-1}h$, where \tilde{z}_0 is defined in Theorem 3.1.*

Proof. First, we consider the eigenfunctions of (1.5), which are odd with respect to x_2 . Then, $\mu = 0$ and we arrive at the following problem:

$$(3.5) \quad \begin{aligned} -\Delta u &= \lambda u & \text{in } \tilde{\Omega}^+ &:= \{0 < x_2 < x_1 < h\}, \\ u &= 0 & \text{on } \{x_2 = 0\}, \\ \partial_{\mathbf{n}} u &= 0 & \text{on } \partial\tilde{\Omega}^+ \setminus \{x_2 = 0\}. \end{aligned}$$

As in Theorems 3.3 and 3.4, we use even reflection with respect to the line $\{x_1 = x_2\}$ and reduce (3.5) to the problem in $(0, h) \times (0, h)$. Thus, we conclude that the least eigenvalue of the problem (3.5) is equal to $\frac{1}{2}(\frac{\pi}{h})^2$ and corresponds to the eigenfunction

$$\hat{u}_0(x) = \sin\left(\frac{\pi x_1}{2h}\right) \sin\left(\frac{\pi x_2}{2h}\right).$$

Next, we consider the eigenfunctions, which are even with respect to x_2 . Then, we arrive at the problem (1.5) in $\tilde{\Omega}^+$ which has been already solved in Theorem 3.1.

To complete the proof we compare $\frac{1}{2}(\frac{\pi}{h})^2$ and $(\frac{\tilde{z}_0}{h})^2$. It is easy to check that $\frac{\pi}{\sqrt{2}} \cdot \cot(\frac{\pi}{\sqrt{2}}) < -1$. Since $t \mapsto t \cdot \cot(t)$ is a decreasing function on $(0, \pi)$, this means that $\frac{\pi}{\sqrt{2}} > \tilde{z}_0$, and the statement follows. \square

Theorem 3.6. *The exact constant in (1.4) is equal to $h^{\frac{1}{2}}$.*

Proof. First, we consider eigenfunctions of the problem (1.6) even with respect to x_2 . Then we arrive at the problem (1.6) in $\tilde{\Omega}^+$ which is solved in Theorem 3.2.

Further, let us consider the eigenfunctions, which are odd with respect to x_1 . We arrive at the following problem in $\tilde{\Omega}^+$:

$$(3.6) \quad \begin{aligned} \Delta u &= 0 & \text{in } \tilde{\Omega}^+; & & \partial_{\mathbf{n}} u &= 0 & \text{on } \{x_1 = x_2\}; \\ u &= 0 & \text{on } \{x_2 = 0\}; & & \partial_{\mathbf{n}} u &= \lambda u & \text{on } \{x_1 = h\}. \end{aligned}$$

Direct calculation shows that the function $x_1 x_2$ is positive in $\tilde{\Omega}^+$ and solves the problem (3.6) with $\lambda = \frac{1}{h}$. By the same arguments as we used for the problem (2.7), it can be shown that this function corresponds to the least eigenvalue.

To complete the proof we compare $\frac{\hat{z}}{h} \tanh(\hat{z}_0)$ and $\frac{1}{h}$. Since $\hat{z}_0 > \frac{\pi}{2}$, we have $\hat{z}_0 \tanh(\hat{z}_0) > \frac{\pi}{2} \tanh(\frac{\pi}{2}) > 1$, and the statement follows. \square

4. AN APPLICATION OF THE ESTIMATES

Exact values of $C_1(\Omega, \Gamma)$ and $C_2(\Omega, \Gamma)$ for rectangular domains and right triangles presented in Theorems 2.1–2.2 and 3.1–3.5 yield guaranteed bounds of constants in analogous inequalities for arbitrary nondegenerate triangles and convex quadrilaterals. The corresponding constants are estimated by standard techniques based on affine equivalent transformations (see [9]). Estimates of the constant C_2 can be also obtained by monotonicity arguments. Indeed, if Ω_1 and Ω_2 are two Lipschitz domains such that $\Omega_1 \subset \Omega_2$ and Γ belongs to $\partial\Omega_1 \cap \partial\Omega_2$, then $C_2(\Omega_1, \Gamma) \geq C_2(\Omega_2, \Gamma)$.

Estimates (1.3) and (1.4) with known constants can be used in various problems arising in quantitative analysis of partial differential equations. For example, let Ω be a polygonal domain covered by a simplicial mesh \mathcal{T}_h , which cells T_i ($i = 1, 2, \dots, N$) have a character size h . Assume that a function $w_h \in H^1(\Omega)$ satisfies the condition $\{\{w_h\}\}_{E_{ij}} = 0$ on any edge $E_{ij} = \partial T_i \cap \partial T_j$. Then, we find that

$$\|w_h\|_{\Omega} \leq C \|\nabla w_h\|_{\Omega}, \text{ where } C = \max_{i=1,2,\dots,N} \min_{E_{ij} \subset \partial T_i} C_1(T_i, E_{ij}).$$

Estimates of this type contain explicitly known constants (which are proportional to h) and allow us to derive fully computable a posteriori estimates

of the accuracy of approximate solutions (in [9] this question is studied in the context of parabolic type equations).

However, a posteriori analysis of approximation errors is only one possible area of application. Below, we briefly discuss another example, in which the estimates are used to deduce a computable a priori estimate of the difference between exact solutions of two boundary value problems. Estimates of this type show when it is worth using a simplified mathematical model instead of the original (complicated) one.

Consider the following *Problem \mathcal{P}* : find $u \in H^1(\Omega)$ such that

$$u - u_0 \in V_0 := \{w \in H^1(\Omega) : w|_{\Gamma^D} = 0\}$$

and

$$(4.1) \quad \int_{\Omega} A \nabla u \cdot \nabla w \, dx = \int_{\Omega} f w \, dx + \int_{\Gamma^N} F w \, ds, \quad \forall w \in V_0,$$

where $\partial\Omega$ consists of two measurable nonintersecting parts Γ^D and Γ^N associated with Dirichlet and Neumann boundary conditions, respectively, $f \in L^2(\Omega)$, $F \in L^2(\Gamma^N)$, and the trace of function $u_0 \in H^1(\Omega)$ defines the Dirichlet boundary condition. We assume that the matrix A is symmetric, bounded, and satisfies the uniform ellipticity condition $A\xi \cdot \xi \geq c|\xi|^2$, where c is a positive constant and the dot stands for the scalar product of vectors.

Let Ω be divided into a finite set \mathcal{O} of "simple" nonoverlapping subdomains Ω_i (e.g., they can be cells of a certain polygonal mesh). Each Ω_i belongs to one of the following three subsets:

$$\begin{aligned} \mathcal{O}^D &:= \{\Omega_i \subset \Omega \mid \partial\Omega_i \cap \Gamma^D =: \Gamma_i^D \neq \emptyset\}, \\ \mathcal{O}^N &:= \{\Omega_i \subset \Omega \mid \partial\Omega_i \cap \Gamma^N =: \Gamma_i^N \neq \emptyset\}, \quad \mathcal{O}^I := \mathcal{O} \setminus (\mathcal{O}^D \cup \mathcal{O}^N). \end{aligned}$$

Here, \mathcal{O}^I contains interior subdomains, \mathcal{O}^D contains subdomains associated with Γ^D , and elements of \mathcal{O}^N are the subdomains associated with Γ^N . Then,

$$\overline{\Omega} = \overline{\Omega}^D \cup \overline{\Omega}^I \cup \overline{\Omega}^N,$$

where Ω^D , Ω^N , and Ω^I consist of Ω_i from \mathcal{O}^D , \mathcal{O}^N , and \mathcal{O}^I , respectively. In the general case, Ω_i may have common boundaries with Γ^N and Γ^D , but then we can subdivide it into two and obtain a decomposition of the above type, which subdomains form three sets: \mathcal{O}^I , \mathcal{O}^N , and \mathcal{O}^D (see Fig. 3).

Now, instead of \mathcal{P} we consider a modified *problem $\widehat{\mathcal{P}}$* : find $\widehat{u} \in H^1(\Omega)$ such that

$$\widehat{u} - \widehat{u}_0 \in V_0 := \{w \in H^1(\Omega) : w|_{\Gamma^D} = 0\}$$

and

$$(4.2) \quad \int_{\Omega} A \nabla \widehat{u} \cdot \nabla w \, dx = \int_{\Omega} \widehat{f} w \, dx + \int_{\Gamma^N} \widehat{F} w \, ds, \quad \forall w \in V_0,$$

where the functions \widehat{f} , \widehat{F} , and \widehat{u}_0 are simplified counterparts of f , F , and u_0 , respectively. Simplifications are done on subdomains and can be motivated




FIGURE 3.

by different reasons, e.g., they may be applied in order to avoid difficult integration procedures, which are necessary if the functions have a complicated behavior with many local details.

Our goal is to deduce an estimate of the difference between the exact solutions of these two problems in terms of the norm

$$\|u - \hat{u}\|^2 := \int_{\Omega} A \nabla(u - \hat{u}) \cdot \nabla(u - \hat{u}) \, dx,$$

which presents the *error of simplification*. If this error is insignificant, then the problem (4.2) can be successfully used instead of (4.1).

In the derivation, we use the Poincaré estimate (1.1) for cells in the set Ω_I with the respective constant $C_P(\Omega_i)$ (or its upper bound $\frac{\text{diam}\Omega_i}{\pi}$). For the cells in Ω_N , we use the estimate (1.4). If the cells are formed by triangles, then Γ_i^N may contain either one or two sides of triangles (see Fig. 3). Therefore, we can use estimates, which follow from Theorems 3.1 and 3.3. Analogously, for the cells in Ω_D , we use the estimate (1.3) and Theorems 3.2 and 3.4 related to the cases, in which functions have zero mean value either on one or on two sides of triangles.

We define the quantities

$$(4.3) \quad D_1^2 := \sum_{\Omega_i \in \mathcal{O}} \mathbb{C}_i^2 \|f - \hat{f}\|_{2, \Omega_i}^2 \quad \text{and} \quad D_2^2 := \sum_{\Omega_i \in \mathcal{O}^N} C_2(\Omega_i, \Gamma_i^N)^2 \|F - \hat{F}\|_{2, \Gamma_i^N}^2,$$

where

$$(4.4) \quad \mathbb{C}_i = \begin{cases} C_P(\Omega_i) & \text{if } \Omega_i \in \mathcal{O}^I \cup \mathcal{O}^N, \\ C_1(\Omega_i, \Gamma_i^D) & \text{if } \Omega_i \in \mathcal{O}^D. \end{cases}$$

Finding the quantities is a very easy task. It is reduced to integration of known functions and does not require solving a boundary value problem. The theorem below shows that a guaranteed majorant of $\|u - \hat{u}\|$ can be

expressed throughout D_1 , D_2 and easily computable quantities

$$(4.5) \quad \mathcal{I}_0 = \sum_{\Omega_i \in \mathcal{O}^D} \{ \{ u_0 - \widehat{u}_0 \} \}_{\Gamma_i^D} \int_{\Omega_i} (f - \widehat{f}) dx,$$

$$(4.6) \quad \mathcal{I}_1(\phi) = \int_{\Omega} (f - \widehat{f}) \phi dx, \quad \mathcal{I}_2(\phi) = \int_{\Gamma^N} (F - \widehat{F}) \phi ds,$$

where ϕ is an arbitrary function in $H^1(\Omega)$ such that $\phi = u_0 - \widehat{u}_0$ on Γ^D .

Theorem 4.1. *Let u and \widehat{u} be the solutions of (4.1) and (4.2), respectively, and*

$$(4.7) \quad \{ \{ f - \widehat{f} \} \}_{\Omega_i} = 0 \quad \forall \Omega_i \in \mathcal{O}^I \cup \mathcal{O}^N,$$

$$(4.8) \quad \{ \{ F - \widehat{F} \} \}_{\Gamma_i^N} = 0 \quad \forall \Omega_i \in \mathcal{O}^N.$$

Then

$$(4.9) \quad \|u - \widehat{u}\| \leq \rho_1 + \sqrt{\rho_2 + \rho_1^2},$$

where

$$(4.10) \quad 2\rho_1 = \frac{D_1 + D_2}{\sqrt{c}} + \|\phi\| \quad \text{and} \quad \rho_2 = \mathcal{I}_0 + \mathcal{I}_1(\phi) + \mathcal{I}_2(\phi).$$

Proof. From (4.1) and (4.2) it follows that

$$(4.11) \quad \int_{\Omega} A(\nabla u - \nabla \widehat{u}) \cdot \nabla w dx \\ = \int_{\Omega} (f - \widehat{f}) w dx + \int_{\Gamma^N} (F - \widehat{F}) w ds \quad \forall w \in V_0.$$

Since $w = u - \widehat{u} - \phi \in V_0$ we can use it as a test function. Then

$$(4.12) \quad \int_{\Omega} A \nabla(u - \widehat{u}) \cdot \nabla w dx = \|u - \widehat{u}\|^2 + \int_{\Omega} A \nabla(u - \widehat{u}) \cdot \nabla \phi dx \\ \geq \|u - \widehat{u}\|^2 - \|u - \widehat{u}\| \|\phi\|.$$

Consider the first term in the right-hand side of (4.11):

$$(4.13) \quad \int_{\Omega} (f - \widehat{f}) w dx \\ = \sum_{\Omega_i \in \mathcal{O}^I \cup \mathcal{O}^N} \int_{\Omega_i} (f - \widehat{f})(u - \widehat{u}) dx + \sum_{\Omega_i \in \mathcal{O}^D} \int_{\Omega_i} (f - \widehat{f})(u - \widehat{u}) dx + \mathcal{I}_1(\phi).$$

The terms of the first sum in (4.13) are estimated using (4.7) and (1.1):

$$\begin{aligned} \int_{\Omega_i} (f - \widehat{f})(u - \widehat{u}) dx &= \int_{\Omega_i} (f - \widehat{f})(u - \widehat{u} - \{ \{ u - \widehat{u} \} \}_{\Omega_i}) dx \\ &\leq C_P(\Omega_i) \|f - \widehat{f}\|_{2,\Omega_i} \|\nabla(u - \widehat{u})\|_{2,\Omega_i}, \quad \Omega_i \in \mathcal{O}^I \cup \mathcal{O}^N. \end{aligned}$$

The terms of the second sum are estimated with the help of (1.3) as follows:

$$\begin{aligned} \int_{\Omega_i} (f - \widehat{f})(u - \widehat{u}) dx &= \int_{\Omega_i} (f - \widehat{f})(u - \widehat{u} - \{ \{ u - \widehat{u} \} \}_{\Gamma_i^D}) dx \\ &+ \int_{\Omega_i} (f - \widehat{f}) \{ \{ u - \widehat{u} \} \}_{\Gamma_i^D} dx \leq C_1(\Omega_i, \Gamma_i^D) \|f - \widehat{f}\|_{2,\Omega_i} \|\nabla(u - \widehat{u})\|_{2,\Omega_i} \\ &+ \{ \{ u - \widehat{u} \} \}_{\Gamma_i^D} \int_{\Omega_i} (f - \widehat{f}) dx, \quad \Omega_i \in \mathcal{O}^D. \end{aligned}$$

Summing up these estimates and using (4.5) we obtain

$$\begin{aligned} (4.14) \quad \int_{\Omega} (f - \widehat{f})w dx &\leq \sum_{\Omega_i \in \mathcal{O}} \mathbb{C}_i \|f - \widehat{f}\|_{2,\Omega_i} \|\nabla(u - \widehat{u})\|_{2,\Omega_i} + \mathcal{I}_0 + \mathcal{I}_1(\phi) \\ &\leq \frac{D_1}{\sqrt{c}} \|u - \widehat{u}\| + \mathcal{I}_0 + \mathcal{I}_1(\phi). \end{aligned}$$

By means of (4.8) and (1.4) we find that

$$\int_{\Gamma_i^N} (F - \widehat{F})(u - \widehat{u}) ds \leq C_2(\Omega, \Gamma_i^N) \|F - \widehat{F}\|_{2,\Gamma_i^N} \|\nabla(u - \widehat{u})\|_{2,\Omega_i},$$

and (cf. (4.6))

$$(4.15) \quad \int_{\Gamma^N} (F - \widehat{F})w ds \leq \frac{D_2}{\sqrt{c}} \|u - \widehat{u}\| + \mathcal{I}_2(\phi).$$

Now (4.11), (4.13), (4.14), and (4.15) yield the estimate

$$(4.16) \quad \|u - \widehat{u}\|^2 \leq 2\rho_1 \|u - \widehat{u}\| + \rho_2,$$

where the quantities ρ_1 and ρ_2 are defined by (4.10).

The quadratic inequality (4.16) easily implies (4.9). \square

Remark 4.1. *It is worth noting that the quantity ρ_2 may be negative. However, the right hand side of (4.9) is always nonnegative. Moreover, it vanishes if and only if $u = \widehat{u}$ (i.e., if two problems are identical).*

Theorem 4.1 presents the most general form of the estimate. If $\widehat{u}_0 = u_0$, then it can be simplified. Indeed, in this case one can choose $\phi \equiv 0$, and

(4.9) is reduced to

$$\|u - \widehat{u}\| \leq \frac{D_1 + D_2}{\sqrt{c}}.$$

Moreover, in this case we can replace $C_1(\Omega_i, \Gamma_i^D)$ in (4.4) by a smaller constant $C_F(\Omega_i, \Gamma_i^D)$ such that

$$(4.17) \quad \|w\|_{2,\Omega} \leq C_F(\Omega_i, \Gamma_i^D) \|\nabla w\|_{2,\Omega}, \quad \forall w \in H^1(\Omega_i) : w|_{\Gamma_i^D} = 0.$$

For simple domains such as rectangles or isocles right triangles these constants are well-known.

Finally, we note that similar analysis can be performed for other differential equations associated with the pair of conjugate operators grad and $-\text{div}$.

5. ACKNOWLEDGMENTS

The first author was supported by RFBR grants 11-01-00825 and 14-01-00534 and by St.Peterburg University grant 6.38.670.2013. The second author was supported by RFBR grant 11-01-00531-a. We thank N. Filonov for useful comments. Also, we are grateful to the anonymous reviewer and to S. Krömer who found gaps in the previous proofs of Theorem 3.2.

REFERENCES

- [1] M.S. Birman, M.Z. Solomyak. *Quantitative analysis in Sobolev imbedding theorems and applications to spectral theory*. In: Proceed. of X Summer Mathematical School. Yu.A. Mitropolskii, A.F. Shestopal (Eds), 1974, 5–189 (in Russian). English transl.: AMS Translations, Series 2, **114**, AMS, Providence, R.I., 1980.
- [2] B. Dittmar and A.Yu. Solynin. *Mixed Stekloff eigenvalue problem and new extremal properties of the Grötzsch ring*. Zap. Nauchn. Sem. POMI, **270**, POMI, St. Petersburg, 2000, 51–79.
- [3] L. Esposito, C. Nitsch, C. Trombetti. *Best constants in Poincaré inequalities for convex domains*. Journal of Convex Analysis 20 (2013), No. 1, 253–264.
- [4] M. Fuchs. *Computable upper bounds for the constants in Poincaré-type inequalities for fields of bounded deformation*. Mathematical Methods in the Applied Sciences, **34**, Issue 15, 1920–1932, October 2011.
- [5] J. Hersch, L.E. Payne, M.M. Schiffer. *Some inequalities for Stekloff eigenvalues*. Arch. Rat. Mech. Anal., **57** (1974), 99–114.
- [6] B. Kawohl. *Rearrangements and convexity of level sets in PDE*. Springer Lecture Notes in Math., **1150** (1985).
- [7] F. Kikuchi, X. Liu. *Estimation of interpolation error constants for the P0 and P1 triangular finite elements*. Comput. Methods. Appl. Mech. Engrg. **196** (37–40) (2007), 3750–3758
- [8] H. Lamb. *Hydrodynamics*. Cambridge, University Press, 1932.
- [9] S. Matculevich and S. Repin. *Computable bounds of the distance to the exact solution of parabolic problems based on Poincaré type estimates*. Zapiski Nauchn. Semin. POMI, **425**, 2014 (in press). Preprint at <http://arxiv.org/abs/1407.6875>.
- [10] A.I. Nazarov, N.G. Kuznetsov, S.V. Poborchii. *V.A. Steklov and the Problem of Sharp (Exact) Constants in Inequalities of Mathematical Physics*. Preprint at <http://arxiv.org/abs/1307.8025>

- [11] L.E. Payne, H.F. Weinberger. *An optimal Poincaré inequality for convex domains.* Arch. Rat. Mech. Anal., **5** (1960), 286–292.
- [12] H. Poincaré. *Sur les équations aux dérivées partielles de la physique mathématique.* Am. J. Math., **12** (1890), 211–294.
- [13] H. Poincaré. *Sur les équations de la physique mathématique.* Rend. Circ. Mat. Palermo, **8** (1894), 57–156.
- [14] S. Repin. *A posteriori error estimates for partial differential equations.* Walter de Gruyter, Berlin, 2010.
- [15] S. Repin, S. Samrowski, S. Sauter. *Combined a posteriori modeling–discretization error estimate for elliptic problems with complicated interfaces.* ESAIM: Mathematical Modelling and Numerical Analysis, **46** (2012), 1389–1405.
- [16] V.A. Steklov, *On the expansion of a given function into a series of harmonic functions,* Communs Kharkov Math. Soc., Ser. 2, **5** (1896), 60–73. (In Russian.)
- [17] W. Stekloff. *Sur les problèmes fondamentaux de la physique mathématique (suite et fin).* Annales sci. ENS, Sér. 3, **19** (1902), 455–490.
- [18] R. Weinstock. *Inequalities for a classical eigenvalue problem.* J. Rational Mech. Anal., **3** (1954), 745–753.