

On Profinite Hyperbolicity and Diophantine Geometry

Arash Rastegar

September 3, 2018

Abstract

In this note, we explore the notion of hyperbolicity of topologically finitely generated profinite groups. Some applications to diophantine geometry are suggested and we try to reformulate certain problems in diophantine geometry in terms of hyperbolic profinite groups. Then, we introduce many occasions in which Galois groups are free profinite and try to explore implications of this condition in the world of diophantine geometry. In particular, we prove that, Grothendieck's "section conjecture" plus Shafarevich's "freeness conjecture" imply that hyperbolic curves have infinitely many solutions over the maximal abelian extension of a global field. This makes Moerdell's conjecture, which was proved by Faltings, more interesting.

Introduction

Gromov's theory of hyperbolic groups, studies finitely generated groups whose Cayley graphs are hyperbolic metric spaces, when equipped with word metric. This notion is independent of choices made for finitely many generators. We intend to explore the implications of extension of this theory to finitely generated profinite groups. The purpose of this paper is to imply that hyperbolicity of profinite groups has applications in finiteness and infiniteness results in diophantine geometry.

Given an elliptic curve E over \mathbb{Q} and fixing a prime ℓ , Tate associated a representation

$$\rho_\ell : Gal(\overline{\mathbb{Q}}/\mathbb{Q}) \rightarrow GL_2(\mathbb{Z}_\ell)$$

by taking the inverse limit over n of the groups of ℓ^n -torsion points of E as Galois modules. On the other hand, one can think of the ℓ^n -torsion points of E as the Galois group of the function field extension associated to the multiplication map $\times \ell^n : E \rightarrow E$. Since every Galois etale covering of an elliptic curve is again an elliptic curve, we can take a limit over all Galois etale covers $E' \rightarrow E$ of degree ℓ^n for some n , and obtain the same Galois module structure on $\mathbb{Z}_\ell \times \mathbb{Z}_\ell$ as part of the algebraic fundamental group of E .

For an arbitrary curve X defined over an algebraically closed field of characteristic zero, the algebraic fundamental group $\pi_1^{alg}(X)$ is defined by Grothendieck as $\lim Gal(K'/K)$ where K is the function field of X and K' runs over all Galois

extensions of K such that the corresponding curve X' is etale over X . For example, $\pi_1^{alg}(\mathbb{P}^1) = 1$ and for every elliptic curve E we have

$$\pi_1^{alg}(E) = \prod_{\ell} \mathbb{Z}_{\ell} \times \mathbb{Z}_{\ell}.$$

Grothendieck proved that, for a curve X of genus g , the algebraic fundamental group of X is isomorphic to the completion of the ordinary topological fundamental group of X over \mathbb{C} [Grt]. If X is defined over \mathbb{Q} one can induce an action of $Gal(\overline{\mathbb{Q}}/\mathbb{Q})$ on $\pi_1^{alg}(X)$.

One can also define the ℓ -adic algebraic fundamental group $\pi_{1,\ell}^{alg}(X)$ as the limit

$$\lim Gal(K'/K)$$

where K is the function field of X and K' runs over all degree ℓ^n Galois extensions of K such that the corresponding curve X' is etale over X . The group $\pi_{1,\ell}^{alg}(X)$ is isomorphic to the ℓ -adic completion of $\pi_1^{top}(X)$. If X is defined over \mathbb{Q} one can associate an ℓ -adic Galois representation to X which is a direct summand of the above mentioned representation

$$\rho_{\ell} : Gal(\overline{\mathbb{Q}}/\mathbb{Q}) \rightarrow Aut(\pi_{1,\ell}^{alg}(X)).$$

The Frattini subgroup $\Phi(G)$ of the pro- ℓ group $G = \pi_{1,\ell}^{alg}(X)$ is fixed by each automorphism [Rib-Zal]. Therefore we get a representation

$$\rho_{\ell} : Gal(\overline{\mathbb{Q}}/\mathbb{Q}) \rightarrow Aut(G/\Phi(G)).$$

Since $\Phi(G) = \overline{G^{\ell}[G, G]}$ we can think of $G/\Phi(G)$ as a finite dimensional vector space over the field \mathbb{F}_{ℓ} with ℓ elements [Rib-Zal]. This way we have recovered the mod- ℓ Galois representation associated to X via Tate module.

The classical approach to definition of an ℓ -adic Galois representation is to consider the action of Galois group on the Tate module associated to the Jacobian variety $Jac(X)$ as above. This representation can be reconstructed from the one we introduced above by abelianization. But lots of information is lost during this process. We believe the structure of fundamental group can help us to understand the Galois action better.

The notion of etale fundamental group introduced by Grothendieck, links the notion of Galois group with its function field counterpart, which is geometric fundamental group [Grt]. By "Grothendieck's anabelian conjecture" the arithmetic fundamental group of a hyperbolic algebraic curve completely determines the algebraic structure of the curve. More precisely, if X is an algebraic variety over a field k which is proper, connected, and geometrically reduced, we have the following exact sequence

$$0 \longrightarrow \pi_1(X_{\bar{k}}) \longrightarrow \pi_1(X_k) \longrightarrow Gal(\bar{k}/k) \longrightarrow 0$$

where $\pi_1(X_{\bar{k}})$ is called the geometric fundamental group and $\pi_1(X_k)$ is called the arithmetic fundamental group. This determines a homomorphism

$$\pi_1(X_k) \rightarrow \text{Aut}(\pi_1(X_{\bar{k}}))$$

by conjugating the group $\pi_1(X_{\bar{k}})$ by elements of $\pi_1(X_k)$ and then, after taking quotients, one obtains a homomorphism landing in the outer automorphism group

$$\rho_X : \text{Gal}(\bar{k}/k) \rightarrow \text{Out}(\pi_1(X_{\bar{k}}))$$

To consider the outer action of Galois group is equivalent to considering $\pi_1(X_k)$ as an extension of $\text{Gal}(\bar{k}/k)$.

By Grothendieck's "fundamental conjecture", if k is finitely generated over the prime field and if X is a hyperbolic algebraic variety, or in other words sufficiently non-abelian, ρ_X determines X over k . Grothendieck's "hom conjecture" states that if X and Y are algebraic varieties over a finitely generated field k , then the natural map between dominant morphisms from X to Y over k to Galois compatible open homomorphisms between the arithmetic fundamental groups

$$\text{Hom}(X, Y) \rightarrow \text{Hom}_{\text{Gal}(\bar{k}/k)}(\pi_1(X_k), \pi_1(Y_k))$$

induces a bijection if right hand side is considered up to composition with an inner automorphism of $\pi_1(Y_k)$. According to "section conjecture", splittings of the previously mentioned short exact sequence are in one to one correspondence with the k -rational points of the projectivized variety. Moreover, sections corresponding to points at infinity are group theoretically characterized. There are several more explicit versions of this general philosophy, similar to "fundamental conjecture", "hom conjecture" and the "section conjecture". We refer the reader to [Nak-Tam-Moc] for an exposition.

The known half of the section conjecture, tells us that to any point of X defined over k one can associate a splitting and this association is injective. Therefore, in order to prove finiteness of the rational points it is enough to show that $\pi_1(X_k)$ has only finitely many splittings as an extension of $\text{Gal}(\bar{k}/k)$ by $\pi_1(X_{\bar{k}})$. This brings us to the realm of profinite group theory. We suggest that hyperbolicity of $\pi_1(X_{\bar{k}})$ is enough to get such finiteness results. This leads us to the following conjecture which implies Mordell's conjecture proved by Faltings:

Conjecture 0.1 *Let H denote a finitely generated profinite hyperbolic group, and let G denote a split extension of the Galois group of the maximal extension unramified outside a set S of finitely many primes of a number field k by a hyperbolic finitely generated profinite group H*

$$0 \rightarrow H \rightarrow G \rightarrow \text{Gal}(\bar{k}_S/k) \rightarrow 0$$

Then, there are only finitely many H -conjugacy classes of splittings of the above short exact sequence.

In fact, we expect that $Gal(\bar{k}_S/k)$ be profinite hyperbolic and the conjecture to hold for Galois group replaced by any profinite hyperbolic group. Note that, we have the following finiteness result for restricted ramification (10.9.11 in [Neu-Sch-Wij]). For k a number field and S a finite set of primes in K , then $Gal_S(K)$ is topologically generated by conjugacy classes of finitely many elements.

We also try to imply that freeness of Galois groups has important implications in diophantine geometry. We will use the freeness of Galois groups to get infiniteness results. The following conjecture of Shafarevich states that certain Galois groups are profinite free:

Conjecture 0.2 (*Shafarevich*) *Let k be a global field and k^{cycl} denote its maximal abelian extension. Then $Gal(k^{cycl}/k)$ is a free profinite group.*

There are a number of evidences for Shafarevich's conjecture. Iwasawa proved that Galois group of the maximal solvable extension of k^{cycl} is profinite free, and Tate proved that $Gal(k^{cycl}/k)$ has cohomological dimension one [Ser] hence by a theorem of Gruenberg it is projective [Gru].

Shafarevich's conjecture was proved in the function field case by F. Pop [Pop1]. We shall see that Grothendieck's "anabelian" considerations plus Shafarevich's conjecture imply infiniteness of points defined over the maximal unramified extensions of number fields which brings Mordell's conjecture proved by Faltings to a new perspective.

Main Theorem 0.3 *Let k denote a global field and X a hyperbolic algebraic curve defined over k , which has a point over k . Then, Grothendieck's "section conjecture" plus Shafarevich's "freeness conjecture" imply that the number of points of X defined over k^{cycl} is infinite.*

Remark 0.4 *If Shafarevich's conjecture is true for any finitely generated field in characteristic zero we get the same result over such fields.*

Organization of the paper is as follows. In the first section, we review basic definition of hyperbolic metric spaces and hyperbolic groups. We review needed background in profinite groups and pro- p groups as a special class of profinite groups in the second section. In the third section, we define finitely generated hyperbolic profinite groups. The fourth section contains a profinite hyperbolic group theory formulation of diophantine geometry. In the fifth section, we review results on topological fundamental group and in the sixth section, we introduce the algebraic geometric formulation of the Poincare conjecture. Then we review the construction of outer representations. In next section, we show that outer representation lands in the completion of the mapping class group. In the ninth section, we review basic definition of free profinite groups and profinite free products. Then, in the tenth section, we review a few classical results in algebraic number theory which imply that certain Galois groups are profinite free. Afterwards, we explore connections with diophantine geometry and prove the main theorem in the final section.

1 Preliminaries on hyperbolic groups

Motivated by hyperbolic phenomena in the geometry of manifolds and spaces of negative curvature, Gromov defined the notion of hyperbolic finitely generated groups. Trying to make the formulation of this concept independent of curvature, he invented the notion of a quasi-isometry between metric spaces [Grm].

Definition 1.1 *Let (X, d) be a metric space. The Gromov product of points x and y of X with respect to a base point x_0 is defined to be the real number*

$$(x.y) = (x.y)_{x_0} = \frac{1}{2}(d(x, x_0) + d(y, x_0) - d(x, y))$$

The metric space X is called ‘Gromov hyperbolic’ or simply ‘hyperbolic’, if it is δ -hyperbolic for some $\delta \geq 0$ which means that

$$(x.y) \geq \min((x.z), (y.z)) - \delta$$

should hold for all $x, y, z \in X$ and every choice of basepoint x_0 . A geodesic metric space X is hyperbolic if and only if for some $\delta \geq 0$ every geodesic triangle is δ -thin, meaning that every side is contained in the δ -neighborhood of the union of two other sides.

Important examples of hyperbolic spaces are: the hyperbolic upper half-space \mathbb{H}^n and all simply-connected Riemannian manifolds with sectional curvature bounded above but away from zero, \mathbb{R} -trees and Cayley graphs of hyperbolic groups.

Definition 1.2 *A finitely generated group G is called hyperbolic, if for some finite generating set Γ the Cayley graph of G with respect to Γ is a hyperbolic space with word metric.*

In order to show that hyperbolicity is well-defined one has to use the notion of quasi-isometries between metric spaces and the fact that hyperbolicity is a quasi-isometric invariant. Important examples of hyperbolic groups are finite groups, free groups of finite rank, fundamental groups of compact Riemannian manifolds with negative sectional curvature, and groups properly discontinuously acting on hyperbolic spaces with compact quotient. Here are some properties of hyperbolic groups: A hyperbolic group is finitely presented. It can not contain a subgroup isomorphic to $\mathbb{Z} \oplus \mathbb{Z}$. A finite index subgroup is hyperbolic if and only if the ambient group is. A hyperbolic groups acts freely and transitively on a contractible finite dimensional locally finite polyhedral complex. As a consequence, it contains only finitely many conjugacy classes of torsion elements.

2 Preliminaries on profinite groups

Interest in profinite groups originates in the study of the Galois groups of infinite extensions of fields. Galois groups naturally carry ‘Krull’ topology under which

they become Hausdorff compact totally disconnected topological groups. These properties characterize profinite groups and indeed, profinite groups are precisely Galois groups.

Definition 2.1 *Let C denote a class of finite groups. A pro- C group G is an inverse limit of a surjective inverse system of groups G_i in C with discrete topology.*

$$G = \varprojlim G_i$$

We say that G is topologically generated by a finite subset Γ if its image in G_i generates G_i , or equivalently if the closure of the group generated by Γ is G . In this case, we say that G is the pro- C completion of the group generated by Γ .

In case C is the class of all finite groups, then G is called a profinite group and if C is the class of p -groups, then G is called a pro- p group. Since we intend to discuss hyperbolic profinite groups, we shall first recognize the class of free pro- C groups as the most important class of hyperbolic profinite groups, to be introduced in the next section.

Definition 2.2 *For any profinite space X there exists a unique pro- C group $F_C(X)$ containing X as a profinite subset which satisfies the following universal property: Any continuous map $X \rightarrow G$ to a pro- C group G whose image generates G , uniquely breaks through inclusion $X \hookrightarrow F$ and a continuous homomorphism $F \rightarrow G$. $F_C(X)$ is called the free pro- C group on the profinite basis X .*

In the case of finitely generated free pro- C groups, any generating set with the same cardinality as the basis is again a basis. In fact, pro- C completion of the free group generated by finite basis X is the free pro- C group on the basis X . Free pro- C groups are examples of free pro- C products.

Definition 2.3 *A family of homomorphisms $\rho_i : H_i \rightarrow H$ between pro- C groups, for $i \in I$, is called convergent to 1 if $\rho_i(H_i)$ is convergent to 1 which means that any open neighborhood of 1 contains almost all $\rho_i(H_i)$. The free pro- C product of a family of H_i is a pro- C group $H = *_{i \in I} H_i$ together with a convergent family of homomorphisms $\rho_i : H_i \rightarrow H$ which is universal among all such families targeting in any group H' . Free pro- C products are unique up to isomorphisms.*

Subgroups of finite index in free pro- C products are again free pro- C products, which could be an indication of the fact that free products being similar to free groups are usually hyperbolic groups. For example, free products of finitely many finitely generated hyperbolic groups are again hyperbolic groups.

3 Hyperbolic profinite groups

Now, we shall try to introduce a profinite version of the hyperbolic phenomena. In order to discuss hyperbolicity of a profinite group, we need to compensate existence

of a geometric object which replaces the role of Cayley graph of a hyperbolic group.

Definition 3.1 *Let G be a residually finite profinite group, which is isomorphic to inverse limit of all its finite images G_i of G , and let Γ denote a finite topological generating set. If Γ generates a hyperbolic finitely generated subgroup, we say that G is profinite hyperbolic.*

Proposition 3.2 *Free pro- C products of finitely many finitely hyperbolic pro- C groups G_i are again hyperbolic pro- C groups.*

Proof. Let Γ_i denote a finite generating set for G_i for $i \in I$. If the Cayley graphs of $\langle \Gamma_i \rangle \subset G_i$ with respect to Γ_i are δ_i -hyperbolic, then the Cayley graph of $*_{i \in I} G_i$ with respect to $\cup_i \Gamma_i$ will also be δ -hyperbolic for some δ . \square

4 Profinite hyperbolicity and Diophantine objects

We shall start with the following basic conjecture:

Conjecture 4.1 *For any number field k , the Galois group $Gal(\bar{k}_S/k)$ of the maximal extension unramified over a set S of finitely many primes of a number field k is a hyperbolic profinite group.*

There exists a pro- p version of the Grothendieck short exact sequence for a curve defined over any finitely generated field

$$0 \longrightarrow \pi_1^{(p)}(X_{\bar{k}}) \longrightarrow \pi_1^{(p)}(X_k) \longrightarrow Gal(\bar{k}/k)(p) \longrightarrow 0$$

and the philosophy of Grothendieck's section conjecture could be also implied in this case. In a forthcoming section, many examples are introduced where we get hyperbolicity of certain pro- p Galois groups for p outside a finite set of primes S . Together with hyperbolicity of the corresponding $\pi_1^{(p)}(X_{\bar{k}})$ one is lead to the following conjecture

Conjecture 4.2 *Let k denote a finitely generated field and X a hyperbolic algebraic curve defined over k , and assume that $Gal(\bar{k}_S/k)(p)$ is hyperbolic for the finite set of primes S of bad reduction for X , then the number of points of X defined over A_S is infinite, where A_S denotes the integral closure of \mathbb{Z} in the maximal extension of k unramified outside S .*

Remark 4.3 *Let k denote a finitely generated field and X a hyperbolic algebraic curve defined over k of genus ≥ 1 after compactification. We prove in [Ras] that the number of points of X defined over A_S is infinite, where A_S denotes the integral closure of \mathbb{Z} in the maximal extension of k unramified outside a finite set of places S .*

In a more abstract setting, one can also think of a profinite group Γ surjecting to the Galois group, as an "arithmetic" space

$$\Gamma \longrightarrow \text{Gal}(\bar{k}/k) \longrightarrow 0$$

and ask for pure algebraic hyperbolicity conditions implying the finiteness of splittings $\text{Gal}(\bar{K}/K) \rightarrow \Gamma$ of the corresponding short exact sequence. This would be a generalization of the Mordell conjecture. Using such algebraic methods one will have a proof of the following folklore more strong version of Mordell's conjecture:

Conjecture 4.4 *The number of rational points of any hyperbolic curve defined over a number field, is bounded above by a bound depending only on the genus and the number field.*

This result is proved in the function field case using Model theory. Up to now there exists no number theoretical proof of the Mordell's conjecture in the function field case. This is one of the main implications of the section conjecture.

5 The topological fundamental group revisited

The fundamental group plays a fundamental role in geometry. Let X be a compact surface of genus g . Hempel in 1976 proved that Poincaré conjecture holds iff there exists a unique epimorphism

$$\pi_1^{\text{top}}(X) \rightarrow F(g) \times F(g)$$

up to automorphisms of $\pi_1^{\text{top}}(X)$ and automorphisms of the components $F(g)$ which are free groups with g generators [Hem]. Existence of such an epimorphism is evident from the presentation

$$\pi_1^{\text{top}}(X) = \langle a_1, \dots, a_g, b_1, \dots, b_g \mid \prod_{i=1}^g [a_i, b_i] = 1 \rangle.$$

Having this presentation, one can show that the fundamental group is a torsion-free group with trivial center. We also have access on a necessary and sufficient condition for a one-relator group to be isomorphic to the fundamental group [Col-Zie].

For a free group F , we say that two surjective maps $\varphi_1, \varphi_2 : \pi_1^{\text{top}}(X) \rightarrow F$ are equivalent if there exist an automorphism of $\pi_1^{\text{top}}(X)$ and an automorphism of F which take φ_1 to φ_2 , i.e. there exists $\alpha \in \text{Aut}(\pi_1^{\text{top}}(X))$ and $\beta \in \text{Aut}(F)$ such that the following diagram is commutative

$$\begin{array}{ccc} \varphi_1 : \pi_1^{\text{top}}(X) & \longrightarrow & F \\ \alpha \downarrow & & \downarrow \beta \\ \varphi_2 : \pi_1^{\text{top}}(X) & \longrightarrow & F \end{array}$$

We say that φ_1 and φ_2 are strictly equivalent if there exist an automorphism α of $\pi_1^{\text{top}}(X)$ such that $\varphi_2 = \varphi_1 \circ \alpha$. Zieschang in 1964 proved that if rank of F is less than or equal to g there exists only one equivalence class of surjective maps $\pi_1^{\text{top}}(X) \rightarrow F$ [Zie]. With the same assumption, Kurchanov and Grigorchuk in 1989 proved that there exists only one strict equivalence class of such surjective maps [Kur-Gri]. Existence of such a surjection is an easy consequence of the presentation of the fundamental group. Nielsen in 1927 proved that every automorphism of $\pi_1^{\text{top}}(X)$ is induced by an automorphism of the corresponding free group which preserves the relation R [Nie].

Suppose H_0 and H_1 are subgroups of $\langle a_1, \dots, a_g, b_1, \dots, b_g \mid \prod_{i=1}^g [a_i, b_i] = 1 \rangle$ which are generated by a_1, \dots, a_g and b_1, \dots, b_g respectively. The subgroups L^+ and L^- defined as such

$$L^- = \{\varphi \in \text{Aut}(\pi_1^{\text{top}}(X)) : \varphi(H_0) = H_0\}$$

$$L^+ = \{\varphi \in \text{Aut}(\pi_1^{\text{top}}(X)) : \varphi(H_1) = H_1\}$$

are conjugate in $\text{Aut}(\pi_1^{\text{top}}(X))$. The group $\text{Aut}(\pi_1^{\text{top}}(X))$ is generated by L^- and L^+ .

Every automorphism of $\pi_1^{\text{top}}(X)$ induces an automorphism of the abelian group $\pi_1^{\text{top}}(X)/[\pi_1^{\text{top}}(X), \pi_1^{\text{top}}(X)]$ whose matrix coordinatizes a representation

$$\rho : \text{Aut}(\pi_1^{\text{top}}(X)) \longrightarrow \text{Sp}_{2g}(\mathbb{Z})$$

Here $\text{Sp}_{2g}(\mathbb{Z})$ is the group of matrices S of order $2g$ such that $SJS^t = \pm J$ where

$$J = \begin{pmatrix} 0 & I_g \\ -I_g & 0 \end{pmatrix}.$$

The representation ρ takes elements of L^- to lower triangular matrices

$$\begin{pmatrix} a & 0 \\ b & c \end{pmatrix}$$

and elements of L^+ go to upper triangular matrices

$$\begin{pmatrix} a & b \\ 0 & c \end{pmatrix}.$$

6 The algebro-geometric formulation of Poincaré conjecture

Providing information about the completed fundamental group is the key to a better understanding of the Galois representation. Let G denote the pro- p completion of the fundamental group. Taking pro- p completions from the surjective maps $\pi_1^{\text{top}}(X) \rightarrow F(g) \times F(g)$ and $\pi_1^{\text{top}}(X) \rightarrow F(2g)$ one gets surjections from G to the pro- p completions $F(g)_p \times F(g)_p$ and $F(2g)_p$. Also, every automorphism of $\pi_1^{\text{top}}(X)$ induces an automorphism of G . One can ask if the analogue of the Poincaré conjecture is true here.

Conjecture 6.1 *There is only one strict equivalence class of surjections*

$$G \longrightarrow F(g)_p \times F(g)_p.$$

The representation $\rho : Aut(\pi_1^{top}(X)) \longrightarrow Sp_{2g}(\mathbb{Z})$ induces a map

$$\widehat{\rho} : Aut(G) \longrightarrow Sp_{2g}(\mathbb{Z})_p$$

which after combining with the natural surjection

$$Sp_{2g}(\mathbb{Z})_p \longrightarrow Sp_{2g}(\mathbb{Z}_p)$$

induces a representation

$$\rho_X : Aut(G) \longrightarrow Sp_{2g}(\mathbb{Z}_p).$$

If we combine ρ with the Galois action on $\pi_1^{alg}(X)$ one has associated a p -adic Galois representation to the algebraic curve X :

$$\rho_p : Gal(\overline{\mathbb{Q}}/\mathbb{Q}) \rightarrow Sp_{2g}(\mathbb{Z}_p).$$

This is the same as the Galois representation associated to the Tate module of the Jacobian variety of X .

7 The Galois action on $\pi_1^{alg}(X)$

The outer automorphism group of $\pi_1^{alg}(X)$ as a quotient of $Aut(\pi_1^{alg}(X))$ accepts a representation from the Galois group

$$\rho_p : Gal(\overline{\mathbb{Q}}/\mathbb{Q}) \longrightarrow Aut(\pi_1^{alg}(X)) \longrightarrow Out(\pi_1^{alg}(X))$$

Mochizuki proved that if X and X' are curves over \mathbb{Q} of genus greater than or equal to 2, the map

$$Isom_{\mathbb{Q}}(X, X') \longrightarrow Out_{Gal(\overline{\mathbb{Q}}/\mathbb{Q})}(Out(\pi_1^{alg}(X)), Out(\pi_1^{alg}(X')))$$

is a one-to-one correspondence, where $Out_{Gal(\overline{\mathbb{Q}}/\mathbb{Q})}$ denotes Galois equivariant isomorphisms between the two groups [Moc]. In particular, the Galois representation on $Out(\pi_1^{alg}(X))$ completely determines the curve X over \mathbb{Q} .

The representation $\widehat{\rho}$ breaks through $Out(\pi_1^{alg}(X))$ because inner automorphisms are neutral when reduced to abelianization of a group

$$\widehat{\rho} : Aut(\pi_1^{alg}(X)) \longrightarrow Out(\pi_1^{alg}(X)) \longrightarrow Sp_{2g}(\mathbb{Z})_p.$$

Thus for each prime p one can associate a p -adic representation to $Out(\pi_1^{alg}(X))$ by considering the pro- p part of the outer automorphism group

$$\rho_X : Out(\pi_1^{alg}(X)) \longrightarrow Sp_{2g}(\mathbb{Z}_p).$$

and therefore the Galois representation

$$\rho_p : Gal(\overline{\mathbb{Q}}/\mathbb{Q}) \rightarrow Sp_{2g}(\mathbb{Z}_p)$$

could be associated to the Galois action on $Out(\pi_1^{alg}(X))$. This shows that the language of outer automorphisms is an appropriate one for studying the arithmetic of curves over \mathbb{Q} . To study the geometric Galois representation

$$\rho : Gal(\overline{\mathbb{Q}}/\mathbb{Q}) \rightarrow Out(\pi_1^{alg}(X))$$

the first step, is to study $Out(\pi_1^{alg}(X))$.

8 Preliminaries on free profinite groups

We shall first define the class of free pro- C groups via its universal property:

Definition 8.1 *For any profinite space X there exists a unique pro- C group $F_C(X)$ containing X as a profinite subset which satisfies the following universal property: Any continuous map $X \rightarrow G$ to a pro- C group G whose image generates G , uniquely breaks through inclusion $X \hookrightarrow F$ and a continuous homomorphism $F \rightarrow G$. $F_C(X)$ is called the free pro- C group on the profinite basis X .*

In the case of finitely generated free pro- C groups, any generating set with the same cardinality as the basis is again a basis. In fact, pro- C completion of the free group generated by finite basis X is the free pro- C group on the basis X . Free pro- C groups are examples of free pro- C products.

One can obtain free profinite groups by taking profinite free product of free profinite groups of rank one. This is because profinite free products of profinite free groups are again profinite free. Here we remind the notion of profinite free product.

Definition 8.2 *A family of homomorphisms $\rho_i : H_i \rightarrow H$ between pro- C groups, for $i \in I$, is called convergent to 1 if $\rho_i(H_i)$ is convergent to 1 which means that any open neighborhood of 1 contains almost all $\rho_i(H_i)$. The free pro- C product of a family of H_i is a pro- C group $H = *_{i \in I} H_i$ together with a convergent family of homomorphisms $\rho_i : H_i \rightarrow H$ which is universal among all such families targeting in any group H' . Free pro- C products are unique up to isomorphisms.*

9 Free Galois groups

This section contains no original results. All of the theorems mentioned are classical results of algebraic number theory. We have just collected a few examples of profinite free Galois groups to be used in the next section.

Let $K(C)$ denote the maximal pro- C extension of K and $Gal_K(C)$ denote the maximal pro- C quotient of Gal_K . For a global field K and a prime p , considering

a class of p -groups, we have an injection $Gal_{K_\beta}(p) \rightarrow Gal_K(p)$ where β is a prime of $K(p)$. If β_1, \dots, β_r are different primes of $K(p)$ then the natural map $*G_{K_{\beta_i}}(p) \rightarrow G_K(p)$ is injective (Theorem 9.3.2 in [Neu-Sch-Wij]). This could imply freeness in certain cases. Let us discuss the function field case first.

1. Let K be a function field of characteristic p and let $k \subset K$ be the finite constant field with algebraic closure \bar{k} . Let S be a finite set of places of K of cardinality n and $Gal_S(K) = Gal(K_S/K)$. We have the following exact sequence:

$$0 \rightarrow Gal_S(K\bar{k}) \rightarrow Gal_S(K) \rightarrow Gal_k \rightarrow 0$$

Gal_k is a free profinite group of rank 1 topologically generated by Frobenius automorphism. The group $Gal_S(K\bar{k})$ is trivial in case $S = \emptyset$ and zero genus $g(K) = 0$, and thus $Gal_S(K) = Gal_k \cong \hat{\mathbb{Z}}$. In case $S \neq \emptyset$, for every $l \neq p$ we have a pro- l presentation of $Gal_S(K\bar{k})(l)$ by $2g + n$ generators and one relation

$$Gal_S(K\bar{k})(l) \cong \langle x_1, \dots, x_{2g}, y_1, \dots, y_n \mid [x_1, x_2] \dots [x_{2g-1}, x_{2g}] y_1 \dots y_n = 1 \rangle.$$

which is a free pro- l group of rank $2g + n - 1$. In fact the following is true:

Riemann's Existence Theorem 9.1 (*Algebraic Form*) *Let K be function field of characteristic p with finite constant field k and let S be a non-empty set of places of K and $g(K) > 0$ then for every l different from p we have the following exact sequence*

$$0 \rightarrow *_{p \in S(K_\emptyset(p))} T_p(l) \rightarrow Gal_S(K\bar{k})(l) \rightarrow G_\emptyset(K\bar{k})(l) \rightarrow 0$$

where $*$ denotes the free pro- l product and $T_p \cong \mathbb{Z}_l(1)$ is the pro- l quotient of inertia at p and $S(K_\emptyset(p))$ denotes the set of primes above S in $K_\emptyset(p)$.

2. A number theoretical analogue of Riemann's existence theorem exists:

Riemann's Existence Theorem 9.2 (*Number Theoretical Form*) *Let K be number field and p be a prime, and let $S \supset T \supset S_p(K) \cup S_\infty$ be sets of places of K , then the canonical homomorphism*

$$*_{p \in S \setminus T(K_T(p))} T(K_p(p)/K_p) \rightarrow Gal(K_S(p)/K_T(p))$$

is an isomorphism where $*$ denotes the free pro- l product and $T(K_p(p)/K_p) \subset Gal(K_p(p)/K_p)$ is the inertia group at p and $S \setminus T(K_T(p))$ denotes the set of primes above $S \setminus T$ in $K_T(p)$.

This is again a freeness assertion. Note that, for primes $p \in S \setminus T(K)$ for which $N(p) \equiv 1 \pmod{p}$ we have $T(K_p(p)/K_p) \cong \mathbb{Z}_p(1)$ and otherwise we have $T(K_p(p)/K_p) = 1$. The following corollary is driven from this theorem in 10.5.3 of [Neu-Sch-Wij]

Corollary 9.3 *With assumptions of above theorem, there is a canonical isomorphism*

$$*_{p \in S \setminus T(K_T)} T_p(p) \rightarrow Gal(K_S/K_T)(p)$$

where T_p is the inertia group in the full local group G_{K_p} .

In the special case of $K = \mathbb{Q}$, by Kronecker-Weber theory, we have $\mathbb{Q}_{S_p \cup S_\infty}(p) = \mathbb{Q}_\infty$. Applying Riemann's existence theorem, one gets the following result of Neukirch (theorem 10.5.6 in [Neu-Sch-Wij]):

Theorem 9.4 *Let p be an odd prime, and let $S \supset S_p \cup S_\infty$ be a finite set of places of \mathbb{Q} , then the canonical homomorphism*

$$*_{p \in S \setminus S_p \cup S_\infty} T(\mathbb{Q}_p(p)/\mathbb{Q}_p) \rightarrow \text{Gal}(\mathbb{Q}_S(p)/\mathbb{Q}_\infty)$$

is an isomorphism.

Note that $\text{Gal}(\mathbb{Q}_\infty/\mathbb{Q}) \cong \mathbb{Z}_p$.

3. The following free product decomposition theorem is also proven 10.7.2 of [Neu-Sch-Wij].

Theorem 9.5 *Let $\delta = 1$ (or $\delta_v = 1$), if $\mu_p \subset K$ (or $\mu_p \subset K_v$ respectively). Suppose $\delta = \sum_v \delta_v$ and that for $S_0 \subseteq S^{\text{finite}}$ the dual group of*

$$\ker(H^1(\text{Gal}_S(K), \mu_p) \rightarrow \bigoplus_{v \in S_0} H^1(K_v, \mu_p))$$

vanishes, then there exists a set of primes $T \supseteq S$ such that the canonical homomorphism

$$*_{v \in S \setminus S_0} \text{Gal}(K_v(p)/K_v) * *_{v \in S \setminus S_0} \text{Gal}(K_v(p)/K_v)/T_v \rightarrow \text{Gal}_S(K)(p)$$

is an isomorphism, where T_v denotes the Inertia subgroup of v .

For $K = \mathbb{Q}$, p arbitrary or for $K = \mathbb{Q}(\mu_p)$, p regular with $S = S_p \cup S_\infty$, the assumptions on the kernel above are satisfied, and we get another freeness statement.

10 Diophantine results

It is time to prove our main result.

Proof(Main Theorem). By Shafarevich's conjecture $\text{Gal}(k^{\text{cycl}}/k)$ is a free profinite group. Let us call it F . The following exact sequence

$$0 \longrightarrow \pi_1(X_{\bar{k}}) \longrightarrow \pi_1(X_{k^{\text{cycl}}}) \longrightarrow \text{Gal}(\bar{k}/k^{\text{cycl}}) \longrightarrow 0$$

is uniquely associated to an outer representation

$$\text{Gal}(\bar{k}/k^{\text{cycl}}) \rightarrow \text{Out}(\pi_1(X_{\bar{k}})).$$

This is because $\pi_1(X_{\bar{k}})$ has trivial center (Look at [Eil-Mac] and chapter IV of [Bro]). The above exact sequence embeds in the following

$$0 \longrightarrow \pi_1(X_{\bar{k}}) \longrightarrow \pi_1(X_k) \longrightarrow \text{Gal}(\bar{k}/k) \longrightarrow 0$$

the quotient being F . Now, any point over k gives a splitting of the above short exact sequence, which in turn induces a splitting of the upper short exact sequence. Conjugating by inner automorphisms of $Gal(\bar{k}/k)$ which lift inner automorphisms of F , we can induce infinitely many splittings of

$$0 \longrightarrow \pi_1(X_{\bar{k}}) \longrightarrow \pi_1(X_{k^{cycl}}) \longrightarrow Gal(\bar{k}/k^{cycl}) \longrightarrow 0$$

which by section conjecture correspond to points defined over k^{cycl} . \square

Remark 10.1 *As mentioned before, there exists a pro- p version of the Grothendieck short exact sequence for a curve defined over any finitely generated field*

$$0 \longrightarrow \pi_1^{(p)}(X_{\bar{k}}) \longrightarrow \pi_1^{(p)}(X_k) \longrightarrow Gal(\bar{k}/k)(p) \longrightarrow 0$$

and the philosophy of Grothendieck's section conjecture could be also implied in this case. In the previous section, many examples are introduced where we get freeness of certain pro- p Galois groups for p outside a finite set of primes S . Together with hyperbolicity of the corresponding $\pi_1^{(p)}(X_{\bar{k}})$ one is lead to use Grothendieck's fundamental conjecture to prove Diophantine results out of a pro- p version of the Grothendieck's conjectures look at [Pop2].

Acknowledgements

I would like to thank Abdus Salam International Center for Theoretical physics (ICTP) and Institute des hautes etude scientifique (IHES) for hospitality and D.T. Le for constant encouragement in the process of preparation of this note. I would also like to thank F. Oort and M. Gromov for conversations which led indirectly to this result. I shall thank corrections to an earlier version suggested by M. Hadian and N. Nikolov.

References

- [Bro] Brown; *Cohomology of Groups*, GTM 87 Springer-Verlag.
- [Co-Zi] D. J. Collins, H. Zieschang; *Combinatorial Group Theory and Fundamental Groups* In Algebra VII by A.N. Parshin and I.R. Shafarevich; Encyclopedia of mathematical sciences, Vol. 58, Springer-Verlag 1993.
- [Eil-Mac] S. Eilenberg, S. MacLane; *Cohomology theory in abstract groups II. Group extensions with non-abelian kernel*, Annales of Math. Vol. 48, No. 2, April, 1947.
- [Grm] M. Gromov; *Hyperbolic Groups*, in 'Essays in Group Theory' editor M. Gersten, MSRI-publications 1987,75-264
- [Grt] A. Grothendieck; *SGA1*, Lecture notes in math. 224, Springer-Verlag 1971.

- [Gru] J.W. Gruenberg; *Projective profinite Groups*, J. London Math. Soc. 42(1967) 155-165
- [Hem] Hemple J.; *3-manifolds*, Ann. Math. Stud. 86. Zbl.345.57001.
- [Kur-Gri] Kurchanov P.F., Grigorchuck R.I.; *Classification of epimorphisms from the fundamental groups of a surface to free groups*, Int. Conf. in memory of A.I. Mal'tsev, Novosibirsk 1989, p. 72 (Russian). Zbl.725.20001.
- [Nak-Tam-Moc] H. Nakayama, A. Tamagawa, S. Mochizuki; *The crothendieck conjecture on the fundamental groups of curves*, Sugako Expositions 14(2001)31-53
- [Neu-Sch-Win] J. Neukirch, A. Schmidt, K. Wingberg; *Cohomology of number fields*, Springer 2000
- [Nie] Nielsen J.; *Om flytningsgruppen i den hyperbolske plan*, Mat. Tidsskrift B. 65-75, Jbuch 53, 543.
- [Pop1] F. Pop; *Etale Galois covers of affine smooth curves*, Invent. Math. 120(1995)555-578
- [Pop2] F. Pop; *Pro- ℓ birational anabelian geometry over algebraically closed fields I*, arXiv:math.AG/0307076
- [Rib-Zal] Ribes L., Zalesski P.A.; *Profinite Groups* Springer 2000.
- [Zie] H. Zieschang; *Alternierende Produkt in freien Gruppen*, Abh. Math. Semin. Univ. Ham. 27, 13-31. Zbl.135.418.