SELF-INTERSECTION OF OPTIMAL GEODESICS

FABIO CAVALLETTI AND MARTIN HUESMANN

ABSTRACT. Let (X, d, m) be a geodesic metric measure space. Consider a geodesic μ_t in the L^2 -Wasserstein space. Then as s goes to t the support of μ_s and the support of μ_t have to overlap, provided an upper bound on the densities holds. We give a more precise formulation of this self-intersection property. We consider for each t the set of times for which a geodesic belongs to the support of μ_t and we prove that t is a point of Lebesgue density 1 for this set, in the integral sense. Our result applies to spaces satisfying $CD(K, \infty)$. The non branching property is not needed.

1. Introduction

Let (X, d, m) be a complete and separable metric measure space with the additional property that

- X coincide with the support of m ;
- (X, d) is a geodesic space.

It is then well-known that the associated L^2 -Wasserstein space $(\mathcal{P}_2(X), W_2)$ is geodesic as well: so to any $\mu_0, \mu_1 \in \mathcal{P}_2(X)$ we can associate a geodesic $[0, 1] \ni t \mapsto \mu_t$ joining μ_0 to μ_1 . See [\[5\]](#page-2-0) for an overview (and much more) on optimal transportation.

Under some general assumption on the metric measure space (X, d, m) , like $CD(K, \infty)$, see [\[1\]](#page-2-1), [\[3\]](#page-2-2), [\[4\]](#page-2-3) for their definition, it is possible to prove that if μ_0 and μ_1 are both absolute continuous with respect to m with bounded densities, then the same property holds for the density of μ_t , in particular $\mu_t \ll m$ and its density in bounded uniformly in $t \in [0, 1]$, see for instance [\[2\]](#page-2-4).

Thanks to this uniform bound on the density, by means of standard arguments in measure theory, one can prove that the support of μ_t has to overlap with μ_0 as t goes to 0, otherwise to much "mass" would be present inside the support of μ_0 . The same property holds for another time s different from 0 as t goes to 0. This overlapping property is to our knowledge the only qualitative property of the support of μ_t that has been proved so far. Even in the Euclidean framework, examples have been constructed showing that even assuming for instance convexity of supp $|\mu_0|$ and supp $|\mu_1|$, then supp $|\mu_t|$ is not convex and in general one can expect supp $[\mu_t]$ to be "hardly" disconnected.

Our interest is to give a more careful analysis of this overlapping property and to prove a structure property of supp $[\mu_t]$. If we denote with $\mathcal{G}(X) \subset C([0,1];X)$ the subset of geodesics in (X,d) and with $\mathcal{P}(\mathcal{G}(X))$ the space of probability measures over it, it is again well-known that to each geodesic $t \mapsto \mu_t \in \mathcal{P}_2(X)$ it is possible to associate $\gamma \in \mathcal{P}(\mathcal{G}(X))$, so that

$$
(e_t)_\sharp \gamma = \mu_t, \qquad e_t : C([0,1];X) \to X, \quad e_t(\gamma) := \gamma_t.
$$

for all $t \in [0, 1]$, with e_t the evaluation map at time t. Our result will be stated in terms of the support of γ . So denote with $G \subset \mathcal{G}(X)$ the support of γ .

For each $t \in [0, 1]$ consider the set

$$
I_t(\gamma) := \{ \tau \in [0,1] : \gamma_\tau \in e_t(G) \},
$$

hence $I_t(\gamma)$ is the set of times for which γ remains inside the support of $e_t(G)$. So clearly $t \in I_t(\gamma)$. We will prove that if there exists a positive constant C so that

$$
\mu_{\tau} = \varrho_{\tau} m, \quad \varrho_{\tau} \leq C,
$$

for all τ in a neighborhood of $t \in (0,1)$, then t is a point of Lebesgue density 1 for $I_t(\gamma)$ in the $L^1(\gamma)$ -sense that is

$$
\lim_{\varepsilon \to 0} \frac{\mathcal{L}^1\big(I_t(\gamma) \cap (t - \varepsilon, t + \varepsilon)\big)}{2\varepsilon} = 1, \qquad \text{in } L^1(G, \gamma).
$$

MH gratefully acknowledges funding by SFB 611.

2. The Result

So let (X, d, m) be a metric measure space verifying the assumption stated before. Let $\mu_0, \mu_1 \in \mathcal{P}_2(X)$ and $t \mapsto \mu_t \in \mathcal{P}_2(X)$ a geodesic connecting them. Moreover $\gamma \in \mathcal{P}(\mathcal{G}(X))$ is the dynamical optimal plan associated to μ_t .

Theorem 2.1. Fix $t \in (0,1)$ and assume the existence of a positive constant C and a neighborhood $U_t \subset [0,1]$ of t such that $\mu_{\tau} = \varrho_{\tau} m$ with $\varrho_{\tau} \leq C$ for each $\tau \in U_t$. Then

(2.1)
$$
\lim_{\varepsilon \to 0} \frac{\mathcal{L}^1\big(I_t(\gamma) \cap (t - \varepsilon, t + \varepsilon)\big)}{2\varepsilon} = 1,
$$

in $L^1(G, \gamma)$.

Proof. Assume by contradiction [\(2.1\)](#page-1-0) doesn't hold. Therefore there exists a set $H \subset G$, with $\gamma(H) > 0$ so that for all $\gamma \in H$ we have two possibilities: or

$$
0 \leq \liminf_{\varepsilon \to 0} \frac{\mathcal{L}^1\big(I_t(\gamma) \cap (t-\varepsilon,t+\varepsilon)\big)}{2\varepsilon} < \limsup_{\varepsilon \to 0} \frac{\mathcal{L}^1\big(I_t(\gamma) \cap (t-\varepsilon,t+\varepsilon)\big)}{2\varepsilon} \leq 1.
$$

either the limit exists but is not one:

$$
\lim_{\varepsilon \to 0} \frac{\mathcal{L}^1\big(I_t(\gamma) \cap (t - \varepsilon, t + \varepsilon)\big)}{2\varepsilon} < 1.
$$

In the first part of the proof we show that in the first case, the lim sup must be equal to 1, neglecting a set of zero γ -measure. The same argument excludes immediately the second case.

Step 1. Suppose by contradiction the existence of a set $H \subset G$, with $\gamma(H) > 0$ so that for all $\gamma \in H$

$$
\limsup_{\varepsilon \to 0} \frac{\mathcal{L}^1(I_t(\gamma) \cap (t-\varepsilon,t+\varepsilon))}{2\varepsilon} < 1.
$$

and therefore possibly restricting H ,

$$
\liminf_{\varepsilon \to 0} \frac{\mathcal{L}^1\big(I_t^c(\gamma) \cap (t - \varepsilon, t + \varepsilon)\big)}{2\varepsilon} > \alpha,
$$

for some $\alpha > 0$. Let

$$
E := \{ (\gamma, s) \in H \times (0, 1) : t + s \in I_t(\gamma)^c \} = \{ (\gamma, s) \in H \times (0, 1) : d(\gamma_{t+s}, e_t(G)) > 0 \}.
$$

Then by Fubini's Theorem

$$
\gamma \otimes \mathcal{L}^1(E) = \int_{(0,1)} \mathcal{L}^1(E(\gamma)) \gamma(d\gamma), \qquad E(\gamma) := P_2\Big(E \cap (\{\gamma\} \times (0,1))\Big),
$$

where P_i denotes the projection map on the *i*-th component, for $i = 1, 2$. From Fatoú's Lemma

$$
\liminf_{\varepsilon \to 0} \frac{\gamma \otimes \mathcal{L}^1\Big(E \cap \big(H \times (t-\varepsilon,t+\varepsilon)\big)\Big)}{2\varepsilon} \ge \alpha \gamma(H),
$$

therefore

$$
\liminf_{\varepsilon \to 0} \frac{1}{2\varepsilon} \int_{(t-\varepsilon,t+\varepsilon)} \gamma(E(\tau)) \mathcal{L}^1(ds) \ge \alpha \gamma(H), \qquad E(\tau) := P_1\Big(E \cap \big(H \times \{\tau\}\big)\Big).
$$

So there must be a sequence of $\{s_n\}_{n\in\mathbb{N}}$ converging to 0 so that $\gamma(E(t+s_n)) \geq C$, for some $C > 0$. Then, since $e_{t+s_n}(G)$ converges to $e_t(G)$ in Hausdorff topology as s_n goes to 0, we have

$$
m(e_t(G)^{\varepsilon}) \ge m(e_t(G) \cup e_{t+s_n}(E(t+s_n))) \ge m(e_t(G)) + m(e_{t+s_n}(E(t+s_n))),
$$

where $e_t(G)^\varepsilon := \{z \in X : d(z, e_t(G)) \leq \varepsilon\}$. Since by assumption $\varrho_\tau \leq C$ on $e_\tau(G)$ for all $\tau \in U_t$, it follows that $m(e_{t+s_n}(E(t+s_n)))$ remains uniformly strictly positive as s_n goes to 0. Since

$$
m(e_t(G)) \ge \limsup_{\varepsilon \to 0} m(e_t(G)^{\varepsilon}),
$$

we have a contradiction. Hence we have shown that there exists H , γ -negligible, so that

$$
\limsup_{\varepsilon \to 0} \frac{\mathcal{L}^1(I_t(\gamma) \cap (t - \varepsilon, t + \varepsilon))}{2\varepsilon} = 1,
$$

for all $\gamma \in G \setminus H$. Using the same reasoning we can also prove a stronger statement: for any sequence $\varepsilon_n \to 0$ there exists H, γ -negligible and depending on the sequence ε_n , so that

$$
\limsup_{n \to \infty} \frac{\mathcal{L}^1(I_t(\gamma) \cap (t - \varepsilon_n, t + \varepsilon_n))}{2\varepsilon_n} = 1,
$$

for all $\gamma \in G \setminus H$. We now show that L^1 -convergence holds.

Step 2. Consider any sequence ε_n converging to 0. Then from the equality

$$
1 - \frac{\mathcal{L}^1(I_t^c(\gamma) \cap (t - \varepsilon_n, t + \varepsilon_n))}{2\varepsilon_n} = \frac{\mathcal{L}^1(I_t(\gamma) \cap (t - \varepsilon_n, t + \varepsilon_n))}{2\varepsilon_n},
$$

integrating over any set $K \subset G$ we get

$$
\gamma(K) - \limsup_{n \to \infty} \int_{K} \frac{\mathcal{L}^1(I_t^c(\gamma) \cap (t - \varepsilon_n, t + \varepsilon_n))}{2\varepsilon_n} \gamma(d\gamma)
$$

=
$$
\liminf_{n \to \infty} \int_{K} \frac{\mathcal{L}^1(I_t(\gamma) \cap (t - \varepsilon_n, t + \varepsilon_n))}{2\varepsilon_n} \gamma(d\gamma).
$$

Hence, if there exists $K \subset G$ with $\gamma(K) > 0$ so that

$$
\liminf_{n\to\infty}\int_K \frac{\mathcal{L}^1(I_t(\gamma)\cap(t-\varepsilon_n,t+\varepsilon_n))}{2\varepsilon_n}\gamma(d\gamma)<\gamma(K),
$$

then

$$
\limsup_{n \to \infty} \int_K \frac{\mathcal{L}^1(I_t^c(\gamma) \cap (t - \varepsilon_n, t + \varepsilon_n))}{2\varepsilon_n} \gamma(d\gamma) > 0.
$$

Using Fubini's Theorem as in Step 1. we could find a sequence of slices of E, say $E(t+s_n)$, with uniformly positive γ-measure and therefore we would get a contradiction. Reasoning in the same manner for the lim sup we get that for any $K \subset G$,

$$
\liminf_{n \to \infty} \int_{K} \frac{\mathcal{L}^1(I_t(\gamma) \cap (t - \varepsilon_n, t + \varepsilon_n))}{2\varepsilon_n} \gamma(d\gamma) = \gamma(K).
$$

Since ε_n was arbitrarily chosen

$$
\lim_{\varepsilon \to 0} \frac{\mathcal{L}^1(I_t(\gamma) \cap (t - \varepsilon, t + \varepsilon))}{2\varepsilon} = 1, \qquad \text{in } L^1(\gamma \sqcup_G).
$$

and the claim follows. \Box

REFERENCES

- [1] J. Lott and C. Villani. Ricci curvature for metric-measure spaces via optimal transport. Ann. of Math., 169(3):903–991, 2009.
- [2] T. Rajala. Interpolated measures with bounded density in metric spaces satisfying the curvature-dimension conditions of Sturm. J. Funct. Anal., 263:896–924, 2012.
- [3] K.T. Sturm. On the geometry of metric measure spaces.I. Acta Math., 196(1):65-131, 2006.
- [4] K.T. Sturm. On the geometry of metric measure spaces.II. Acta Math., 196(1):133-177, 2006.
- [5] C. Villani. Optimal transport, old and new. Springer, 2008.

RWTH, Department of Mathematics, Templergraben 64, D-52062 Aachen (Germany) E-mail address: cavalletti@instmath.rwth-aachen.de

UNIVERSITÄT BONN, INSTITUT FÜR ANGEWANDTE MATHEMATIK, ENDENICHER ALLEE 60, D-53115 BONN (GERMANY) E-mail address: huesmann@iam.uni-bonn.de