SQUARE FUNCTION AND MAXIMAL FUNCTION ESTIMATES FOR OPERATORS BEYOND DIVERGENCE FORM EQUATIONS

ANDREAS ROSÉN¹

ABSTRACT. We prove square function estimates in L_2 for general operators of the form $B_1D_1 + D_2B_2$, where D_i are partially elliptic constant coefficient homogeneous first order self-adjoint differential operators with orthogonal ranges, and B_i are bounded accretive multiplication operators, extending earlier estimates from the Kato square root problem to a wider class of operators. The main novelty is that B_1 and B_2 are not assumed to be related in any way. We show how these operators appear naturally from exterior differential systems with boundary data in $L₂$. We also prove non-tangential maximal function estimates, where our proof needs only off-diagonal decay of resolvents in L_2 , unlike earlier proofs which relied on interpolation and L_p estimates.

1. INTRODUCTION

In this paper, we generalize the square function estimates from the Kato square root problem, to a wider class of operators on $L_2(\mathbf{R}^n;\mathbf{C}^N)$, $n, N \geq 1$. Previously, estimates were known for perturbations of a homogeneous first order constant coefficients self-adjoint partial differential operator D , of the form DB or BD , with B being a bounded multiplication operator which is accretive on the range of D. Let us first recall how such operators appear in connection with divergence form equations. The celebrated Kato square root estimate

$$
\|\sqrt{-\text{div}A\nabla}u\|_2 \approx \|\nabla u\|_2
$$

for divergence form operators with general bounded accretive coefficients $A \in L_{\infty}(\mathbf{R}^n; \mathcal{L}(\mathbf{C}^n)),$ were proved in one dimension by Coifman, McIntosh and Meyer [\[9\]](#page-19-0) and in full generality by Auscher, Hofmann, Lacey, McIntosh and Tchamitchian [\[6\]](#page-19-1). It can be written $\|\sqrt{(BD)^2}[u, 0] \|_2 \approx \|BD[u, 0]\|_2$, with $D =$ $\begin{bmatrix} 0 & \mathrm{div} \end{bmatrix}$ $-\nabla \quad 0$ 1 and $B =$ $\begin{bmatrix} I & 0 \end{bmatrix}$ $0\quad A$ 1 , acting on vectors of dimension $N = 1 + n$. This estimate is in turn a consequence of square function estimates for the operator BD. This approach to the Kato square root estimate was developed in [\[7,](#page-19-2) [8\]](#page-19-3).

Operators of the form DB and BD appear not only in connection with divergence form operators on \mathbb{R}^n , but also in connection with divergence form equations on the half-space $\mathbf{R}^{1+n}_{+} := \{(t,x) : t > 0, x \in \mathbf{R}^n\}$, with $L_2(\mathbf{R}^n)$ or $H^1(\mathbf{R}^n)$ boundary data. We recall the following approach to boundary value problems from [\[5,](#page-19-4) [3\]](#page-19-5). Consider a divergence form equation

$$
\mathrm{div}_{t,x} A(t,x) \nabla_{t,x} u(t,x) = 0, \qquad t > 0, x \in \mathbf{R}^n,
$$

¹ Formerly Andreas Axelsson.

Supported by Grant 621-2011-3744 from the Swedish research council, VR.

2 ANDREAS ROSEN´

with coefficients $A =$ $\begin{bmatrix} a & b \\ c & d \end{bmatrix}$, splitting $\mathbf{C}^{1+n} = \mathbf{C} \oplus \mathbf{C}^n$. Write e_0, e_1, \ldots, e_n for the standard basis in \mathbf{R}^{1+n} , with coordinates $x_0 = t, x_1, \ldots, x_n$, and $f_{\perp} := e_0 \cdot f$ for the normal component and $f_{\parallel} := f - f_{\perp}e_0$ for the tangential part.

On the one hand, at the level of $\dot{H}^1(\mathbf{R}^n)$ boundary data $u|_{\mathbf{R}^n}$, we consider the conormal gradient

$$
f = \begin{bmatrix} f_{\perp} \\ f_{\parallel} \end{bmatrix} := \begin{bmatrix} a\partial_t u + b\nabla_x u \\ \nabla_x u \end{bmatrix}.
$$

In terms of f, the divergence form equation is $\partial_t f_\perp + \text{div}_x (c(a^{-1}f_\perp - a^{-1}bf_\parallel) + df_\parallel) = 0.$ The conormal gradient f , with the inward conormal derivative as normal component f_{\perp} , is in one-to-one correspondence with the potential u, modulo curl-freeness and constants. Written in terms of f, the curl-free condition is $\partial_t f_{\parallel} = \nabla_x (a^{-1} f_{\perp} - a^{-1} b f_{\parallel}),$ $\text{curl}_x f_{\parallel} = 0$. In vector notation this means that the divergence form equation for u is equivalent to the vector valued ordinary differential equation

$$
\partial_t f + DBf = 0
$$

for f under the constraint $f \in \overline{R(D)}$ for each $t > 0$, with $D := \begin{bmatrix} 0 & \text{div}_x \\ -\nabla_x & 0 \end{bmatrix}$ $-\nabla_x$ 0 1 acting along \mathbf{R}^n and $B := \begin{bmatrix} a^{-1} & -a^{-1}b \\ a^{-1} & d & a^{-1} \end{bmatrix}$ 1

 ca^{-1} d – $ca^{-1}b$ being a multiplication operator, which turns out to be accretive if and only if A is so.

On the other hand, at the level of $L_2(\mathbf{R}^n)$ boundary data $u|_{\mathbf{R}^n}$, we can write

$$
f_{\perp} = (A\nabla u)_{\perp} =: \text{div}_x v_{\parallel},
$$

with a tangential vector field v_{\parallel} , for each fixed t, assuming appropriate decay of u at infinity, since $\int_{\mathbf{R}^n} f_\perp dx = 0$ by the divergence theorem. Inserting this ansatz into the divergence form equation and commuting ∂_t and div_x yields div_x($\partial_t v_{\parallel} + (A\nabla_{t,x}u)_{\parallel}$) = 0. Since v_{\parallel} is only defined modulo tangential divergence free vector fields, we may choose it so that $\partial_t v_{\parallel} + (A\nabla_{t,x}u)_{\parallel} = 0$. In vector notation this means that the divergence form equation for u is equivalent to the vector valued ordinary differential equation

$$
\partial_t v + BDv = 0,
$$

for the vector field

$$
v = \begin{bmatrix} v_{\perp} \\ v_{\parallel} \end{bmatrix} := \begin{bmatrix} -u \\ \operatorname{div}_x^{-1}(a\partial_t u + b\nabla_x u) \end{bmatrix}
$$

.

One can view both $\partial_t f + DBf = 0$ and $\partial_t v + BDv = 0$ as generalized Cauchy– Riemann systems. In particular, the *n* components of v_{\parallel} should be viewed as some generalized harmonic conjugate functions.

Estimates of operators of the form DB or BD are by now well understood, see [\[8,](#page-19-3) [5,](#page-19-4) [4\]](#page-19-6). The aim of this paper is to prove fundamental estimates for more general operators of the form

$$
B_1D_1+D_2B_2,
$$

which appear for example when, similar to above, writing a more general exterior differential system as a vector valued ordinary differential equation in the variable transversal to the boundary. See Section [3.](#page-7-0) We assume that $D_1D_2 = 0$ but, unlike earlier results [\[4\]](#page-19-6), not that $D_2B_2B_1D_1=0$.

We next formulate our results in detail. Consider four operators D_1, D_2, B_1 and B_2 acting in the Hilbert space $L_2(\mathbf{R}^n; \mathbf{C}^N)$ with norm $\|\cdot\|_2$, where $n, N \geq 1$. We assume the following.

- The operators D_1 and D_2 are constant coefficient homogeneous first order differential operators which are self-adjoint and such that $R(D_2) \subset N(D_1)$. Assume the partial ellipticity estimates $||D_i f||_2 \ge ||f||_{\dot{H}^1(\mathbf{R}^n)}$ for all $f \in$ $R(D_i), i = 1, 2.$
- The operators B_1 and B_2 are bounded multiplication operators $B_i : f(x) \mapsto$ $B_i(x)f(x), x \in \mathbf{R}^n$, where $B_i(\cdot) \in L_\infty(\mathbf{R}^n; \mathcal{L}(\mathbf{C}^N)), i = 1, 2$. Assume the partial accretivity estimates $\text{Re}(B_i f, f) \gtrsim ||f||_2^2$ for all $f \in \mathsf{R}(D_i)$, $i = 1, 2$.

Denote by $\omega_i := \sup_{f \in \mathsf{R}(D_i) \setminus \{0\}} |\arg(B_i f, f)| < \pi/2$ the angle of accretivity for B_i on $R(D_i), i = 1, 2.$ For $0 \leq \alpha < \pi/2$, define the closed sectors $S_{\alpha+} := {\lambda \in \mathbb{C} \setminus \{0\}}$; $|\arg \lambda| \leq \alpha$ \cup {0} and $S_{\alpha-} := -S_{\alpha+}$, and bisectors $S_{\alpha} := S_{\alpha+} \cup S_{\alpha-}$, as well as the corresponding open sectors/bisectors $S^o_{\alpha\pm}$, S^o_{α} , being the interior of $S_{\alpha\pm}$, S_{α} respectively.

Denote by $R(\cdot)$, $N(\cdot)$ and $D(\cdot)$ the range, null space and domain of an operator. Define the operator

$$
T := B_1 D_1 + D_2 B_2, \qquad D(T) := \{ f \in D(D_1) \; ; \; B_2 f \in D(D_2) \}.
$$

The definition of operators $\psi(t)$, $\phi(t)$ in the functional calculus of T, is found in Section [2.](#page-4-0) For a function h on \mathbb{R}^{1+n}_+ , define the $(L_2$ Whitney averaged) nontangential maximal function

$$
\widetilde{N}_{*}h(x) := \sup_{t>0} \left(\iint_{W(t,x)} |h(s,y)|^2 \frac{dsdy}{|W(t,x)|} \right)^{1/2}, \qquad x \in \mathbf{R}^n,
$$

where $W(t, x)$ denotes a Whitney region around (t, x) , for example $W(t, x)$ = $B(x, t) \times (t/2, 2t)$.

Theorem 1.1. Under the above hypothesis, T is a closed and densely defined operator in $L_2(\mathbf{R}^n;\mathbf{C}^N)$, with spectrum $\sigma(T) \subset S_\omega$, where $\omega := \max(\omega_1, \omega_2)$, and resolvent estimates $\|(\lambda I - T)^{-1}\| \lesssim 1/\text{dist}(\lambda, \sigma(T)).$

Moreover, the following estimates of holomorphic functions of the operator T hold.

• We have square function estimates

$$
\int_{\mathbf{R}^n}\int_0^\infty |\psi(tT)f(x)|^2\frac{dtdx}{t}\lesssim \int_{\mathbf{R}^n}|f(x)|^2dx,\qquad f\in\overline{\mathcal{R}(T)},
$$

for any holomorphic symbol $\psi : S^o_\mu \to \mathbf{C}, \ \omega < \mu < \pi/2$, with estimates $|\psi(\lambda)| \lesssim \min(|\lambda|^s, |\lambda|^{-s})$ for some $s > 0$.

If furthermore $\psi|_{S_{\omega+}}$ and $\psi|_{S_{\omega-}}$ are not identically zero, then the reverse square function estimates \geq hold for all $f \in \overline{R(T)}$.

• We have non-tangential maximal function estimates

$$
\int_{\mathbf{R}^n} |\widetilde{N}_*(\phi(tT)f)(x)|^2 dx \approx \int_{\mathbf{R}^n} |f(x)|^2 dx, \qquad f \in \overline{R(T)},
$$

for any holomorphic symbol $\phi : S_\mu^o \to \mathbf{C}, \ \omega < \mu < \pi/2$, with estimates $|\phi(\lambda) - 1| \lesssim |\lambda|^s$ and $|\phi(\lambda)| \lesssim |\lambda|^{-s}$, for some $s > 0$.

4 ANDREAS ROSEN´

The estimates in Theorem [1.1](#page-2-0) go back to the techniques from the solution of the Kato square root problem by Auscher, Hofmann, Lacey, McIntosh and Tchamitchian [\[6\]](#page-19-1). The connection between the Kato square root problem and square function estimates for first order differential operators was developed by Auscher, McIntosh and Nahmod [\[7\]](#page-19-2). More directly, both the square function and non-tangential maximal function estimates in Theorem [1.1](#page-2-0) build on the author's joint work [\[3\]](#page-19-5) with Auscher and Hofmann.

So far, the Kato techniques have been applied to establish square function estimates for three main classes of first order differential operators.

- (1) Operators of the form DB and BD , with D being a self-adjoint constant coefficient homogeneous first order differential operator, and B being a bounded multiplication operator which is accretive on $R(D)$.
- (2) Operators of the form $\Gamma + B_1 \Gamma^* B_2$, with Γ being a nilpotent (that is $\Gamma^2 = 0$) constant coefficient homogeneous first order differential operator, and B_1, B_2 being bounded multiplication operators such that $\Gamma^* B_2 B_1 \Gamma^* = 0 = \Gamma B_1 B_2 \Gamma$, which are accretive on $R(\Gamma^*)$ and $R(\Gamma)$ respectively.
- (3) Operators of the form $B_1D_1 + D_2B_2$, with D_1, D_2 being self-adjoint constant coefficient homogeneous first order differential operators with orthogonal ranges, and B_1 , B_2 being bounded multiplication operators such that $D_2B_2B_1D_1 = 0$, which are accretive on $R(D_1)$ and $R(D_2)$ respectively.

Note that for all these classes of operators, there is essentially only one multiplication operator B as we are considering a small generalization of the case $B_2 = B_1^{-1}$. This is in constrast to Theorem [1.1,](#page-2-0) where the two multiplication operators B_1 and B_2 are independent of each other.

The main example of operators of type [\(2\)](#page-3-0) are Hodge–Dirac operators, with Γ being the exterior derivative acting on differential forms, see Section [3.](#page-7-0) For operators of type [\(2\)](#page-3-0), square function estimates were proved in [\[8,](#page-19-3) Thm. 2.7]. By a simple operator theoretic argument, square function estimates for operators of type [\(1\)](#page-3-1) follow from such estimates of operators of type (2) , as shown in [\[8,](#page-19-3) Thm. 3.1]. In fact this argument can be reversed. It was shown in [\[4,](#page-19-6) Sec. 8.1] that conversely square function estimates for operators of type [\(2\)](#page-3-0) follow from such estimates for operators of type [\(1\)](#page-3-1). Operators of type [\(3\)](#page-3-2) are nothing but a direct sum of two operators of type [\(1\)](#page-3-1), and hence square function estimates are immediate as shown in [\[4,](#page-19-6) Sec. 8.2]. The type [\(3\)](#page-3-2) operators first appeared, in disguise, in the work by Auscher, Axelsson and Hofmann [\[3\]](#page-19-5), where boundary value problems for Dirac equations of the form $(\Gamma + B_1 \Gamma^* B_2) f = 0$, Γ being the exterior derivative, were studied. Similarly to Section [3](#page-7-0) here, it was shown in [\[4,](#page-19-6) Sec. 8.3], that solving for the t-derivatives, this equation can be written $(\partial_t + (BD_1 + D_2B^{-1}))Uf = 0$, under suitable similarity transformation U . It should be noted that in [\[3\]](#page-19-5), B_1 and B_2 were related in exactly the way so that the associated operator $BD_1 + D_2B^{-1}$ is of type (3) and not of the more general form T considered in Theorem [1.1,](#page-2-0) where the two multiplication operators are independent.

Coming to the non-tangential maximal function estimates in Theorem [1.1,](#page-2-0) these build on the estimates by Auscher, Axelsson and Hofmann in [\[3,](#page-19-5) Prop. 2.56]. Although set in the framework with Dirac equations, what was actually proved there was non-tangential maximal function estimates for operators of type [\(1\)](#page-3-1), in the special case when the differential operator is of the form $D =$ $\begin{bmatrix} 0 & \text{div} \end{bmatrix}$ $-\nabla \quad 0$ Ť . Even for operators of type (1) with more general D, the non-tangential maximal function estimate in Theorem [1.1](#page-2-0) are new. Also the non-tangential maximal function estimates for general operators of type [\(2\)](#page-3-0) and [\(3\)](#page-3-2), which is a special case of Theorem [1.1,](#page-2-0) are new.

The key idea in the proof of the square function estimates in Theorem [1.1](#page-2-0) is to use a splitting of L_2 adapted to the operators B_1D_1 and D_2B_2 , in which the operator $B_1D_1 + D_2B_2$ is triangular due to the assumption $D_1D_2 = 0$. The proof of the non-tangential maximal function estimates in Theorem [1.1](#page-2-0) is much inspired by the proof of [\[3,](#page-19-5) Prop. 2.56]. The main difference is that our proof here is a pure L_2 proof, in that it only requires L_2 off-diagonal decay of resolvents. In [\[3,](#page-19-5) Prop. 2.56], interpolation theory to prove L_p off-diagonal decay, $p \approx 2$, was needed. Another novelty is a Caccioppoli type estimate, Lemma [5.2,](#page-17-0) for operators beyond divergence form equations.

The outline of this paper is as follows. In Section [3,](#page-7-0) we show how operators of the form $B_1D_1 + D_2B_2$ arise naturally in connection with exterior differential systems systems in \mathbb{R}^{1+n}_+ for differential forms. The special case of one-forms, that is vector fields, amounts to divergence form equations. In Section [2](#page-4-0) we prove the resolvent estimates for the operator T , in Section [4](#page-12-0) we prove the square function estimates for the operator T , and finally in Section [5](#page-16-0) we prove the non-tangential maximal function estimates for the operator T . The (roadmap to the) proof of Theorem [1.1](#page-2-0) is in Section [5.](#page-16-0)

2. Resolvent estimates

In this section, we establish the basic operator theoretical properties of the operator $T = B_1D_1 + D_2B_2$. A fundamental observation for the unperturbed operator $D_1 + D_2$ is the orthogonal splitting

$$
L_2(\mathbf{R}^n;\mathbf{C}^N)=(\mathsf{N}(D_1)\cap\mathsf{N}(D_2))\oplus\overline{\mathsf{R}(D_1)}\oplus\overline{\mathsf{R}(D_2)}=:\mathcal{H}_0\oplus\mathcal{H}_1\oplus\mathcal{H}_2.
$$

The natural perturbation of this splitting which is adapted to the operator T is

$$
L_2(\mathbf{R}^n; \mathbf{C}^N) = (\mathsf{N}(D_1) \cap \mathsf{N}(D_2B_2)) \oplus \overline{\mathsf{R}(B_1D_1)} \oplus \overline{\mathsf{R}(D_2)}.
$$

Proposition 2.1. We have a topological (but in general not orthogonal) splitting

$$
L_2(\mathbf{R}^n;\mathbf{C}^N)=\mathcal{H}_0^{B_2}\oplus B_1\mathcal{H}_1\oplus \mathcal{H}_2,
$$

where $\mathcal{H}_0^{B_2}$ B_2^B := (H₀ ⊕ H₂) ∩ B_2^{-1} (H₀ ⊕ H₁), $B_2^{-1}V$:= {f ∈ L₂ ; B₂f ∈ V} and $B_1V := \{B_1f \, ; \, f \in V\}.$

Proof. We observe that we have two topological splittings

$$
L_2=B_1\mathcal{H}_1\oplus(\mathcal{H}_0\oplus\mathcal{H}_2)
$$

and $L_2 = B_2^* \mathcal{H}_2 \oplus (\mathcal{H}_0 \oplus \mathcal{H}_1)$, since B_1 is accretive on \mathcal{H}_1 and B_2 , and hence B_2^* , is accretive on \mathcal{H}_2 . See for example [\[5,](#page-19-4) Prop. 3.3]. Taking orthogonal complements in the second splitting, we obtain a third topological splitting

$$
L_2=B_2^{-1}(\mathcal{H}_0\oplus \mathcal{H}_1)\oplus \mathcal{H}_2,
$$

since $B_2^{-1}(\mathcal{H}_0 \oplus \mathcal{H}_1) = (B_2^*\mathcal{H}_2)^\perp$ and $\mathcal{H}_2 = (\mathcal{H}_0 \oplus \mathcal{H}_1)^\perp$.

The result is now a consequence of Lemma [2.2](#page-5-0) below, with $X_1 = B_1 \mathcal{H}_1$, $X_2 =$ $\mathcal{H}_0 \oplus \mathcal{H}_2$, $X_3 = B_2^{-1}(\mathcal{H}_0 \oplus \mathcal{H}_1)$ and $X_4 = \mathcal{H}_2$.

Lemma 2.2. Assume that a Banach space X splits topologically in two ways

$$
X = X_1 \oplus X_2 = X_3 \oplus X_4,
$$

into closed subspaces such that $X_4 \subset X_2$. Then X splits topologically into three closed subspaces

$$
X = X_1 \oplus (X_2 \cap X_3) \oplus X_4.
$$

Proof. It is straightforward to verify that these three subspaces are closed and intersect pair wise only at 0. Also, given $x \in X$, we can write $x = x_1 + x_2$ and $x_2 = x_3 + x_4$ with $x_i \in X_i$, $i = 1, 2, 3, 4$. We have $x = x_1 + x_3 + x_4$, with $x_3 = x_2 - x_4 \in X_3 \cap X_2$, so the three subspaces span X.

Proposition 2.3. For the operator T, the null space is $N(T) = H_0^{B_2}$ $_0^{B_2}$, the range is $R(T) = B_1 R(D_1) + R(D_2)$ and the domain is

$$
D(T) = \{u_0 + u_1 + u_2 \in \mathcal{H}_0^{B_2} \oplus B_1\mathcal{H}_1 \oplus \mathcal{H}_2 ; u_1 \in D(D_1), u_1 + u_2 \in D(D_2B_2)\}.
$$

Proof. For the null space, we note that $f \in N(T)$ if and only if $B_1D_1f = -D_2B_2f$ Since $B_1\mathcal{H}_1 \cap \mathcal{H}_2 = \{0\}$, this is equivalent to $f \in N(D_1) \cap B_2^{-1}N(D_2)$.

Clearly $R(T) \subset B_1R(D_1) + R(D_2)$. For the converse implication, assume that $f_1 = B_1D_1u_1 \in B_1R(D_1)$ and $f_2 = D_2B_2u_2 \in R(D_2) = R(D_2B_2)$. Write $u_1 =$ $u_1^1 + u_1^2 \in \mathcal{H}_2 \oplus B_2^{-1}(\mathcal{H}_0 \oplus \mathcal{H}_1)$ and $u_2 = u_2^1 + u_2^2 \in \mathcal{H}_2 \oplus B_2^{-1}(\mathcal{H}_0 \oplus \mathcal{H}_1)$. Then $(B_1D_1+D_2B_2)(u_1^2+u_2^1)=B_1D_1u_1+D_2B_2u_2=f_1+f_2$, so $R(T)=B_1R(D_1)+R(D_2)$.

The result for the domain follows from the facts that $\mathcal{H}_0^{B_2} \subset D(D_1) \cap D(D_2B_2)$ and $\mathcal{H}_2 \subset D(D_1)$.

We now express the resolvents of T in terms of the resolvents

$$
R_t^1 := (I + itB_1D_1)^{-1}
$$
 and $R_t^2 := (I + itD_2B_2)^{-1}$

of B_1D_1 and D_2B_2 . It is known that $\sigma(B_1D_1) \subset S_{\omega_1} \cup \{0\}$ with resolvent estimates $||R_1^1|| \lesssim 1/(|t| \text{dist} (i/t, S_{\omega_1}))$, and that $\sigma(D_2B_2) \subset S_{\omega_2} \cup \{0\}$ with resolvent estimates $||R_t^2|| \lesssim 1/(|t| \text{dist}\left(\frac{i}{t}, S_{\omega_2}\right))$. See for example [\[5,](#page-19-4) Prop. 3.3].

Proposition 2.4. The operator T is closed and densely defined in $L_2(\mathbf{R}^n;\mathbf{C}^N)$. The spectrum is contained in the bisector $S_{\omega} \cup \{0\}$, $\omega = \max(\omega_1, \omega_2)$, and in the splitting $L_2(\mathbf{R}^n;\mathbf{C}^N)=\mathcal{H}_0^{B_2}\oplus B_1\mathcal{H}_1\oplus \mathcal{H}_2$, the resolvent has the expression

$$
(I + itT)^{-1} = \begin{bmatrix} I & 0 & 0 \\ 0 & R_t^1 & 0 \\ 0 & (R_t^2 - I)R_t^1 & R_t^2 \end{bmatrix}, \quad i/t \notin S_\omega \cup \{0\},\
$$

with estimates $||(I + itT)^{-1}|| \lesssim 1/(|t| \text{dist} (i/t, S_{\omega}))$.

Proof. Consider $(I + itT)u = f$ with $u \in D(T)$, and write $u = u_0 + u_1 + u_2$ and $f = f_0 + f_1 + f_2$ in the splitting from Proposition [2.1.](#page-4-1) Then

$$
u_0 = f_0, \qquad u_1 + itB_1D_1u_1 = f_1,
$$

$$
u_2 + itD_2B_2(u_1 + u_2) = f_2.
$$

Solving for u , we equivalently have

$$
u_0 = f_0, \qquad u_1 = (I + itB_1D_1)^{-1}f_1,
$$

$$
u_2 = (I + itD_2B_2)^{-1}f_2 - itD_2B_2(I + itD_2B_2)^{-1}(I + itB_1D_1)^{-1}f_1.
$$

This shows that $I+itT$ is injective with the stated resolvent expression and estimate.

To show surjectivity, given $f \in L_2(\mathbf{R}^n;\mathbf{C}^N)$, define $u := f_0 + (I + itD_2B_2)^{-1}f_2 +$ $(I + itD₂B₂)⁻¹(I + itB₁D₁)⁻¹f₁$. Then reversing the above calculation, shows that $u \in D(T)$ and $(I + itT)u = f$. It follows that $I + itT$ is surjective and that T is a closed operator. That T is densely defined, follows from the fact that $D(T)$ contains the dense subspace $D(D_2B_2) \cap R(D_2B_2)$.

Proposition 2.5. The adjoint of T is $T^* = B_2^* D_2 + D_1 B_1^*$, with domain $D(T^*) :=$ ${f \in D(D_2) ; B_1^* f \in D(D_1)}$.

Proof. It suffices to show that if

(1)
$$
D(T) \to \mathbf{C} : u \mapsto (Tu, v)
$$

is L_2 continuous, then $v \in D(D_2) \cap D(D_1B_1^*)$. The splitting for v analogous to Proposition [2.1](#page-4-1) for u , is

$$
v = v_0 + v_1 + v_2 \in ((\mathcal{H}_0 \oplus \mathcal{H}_1) \cap (B_1^*)^{-1}(\mathcal{H}_0 \oplus \mathcal{H}_2)) \oplus B_2^* \mathcal{H}_2 \oplus \mathcal{H}_1.
$$

We need to show $v_1 \in D(D_2)$ and $v_1 + v_2 \in D(D_1B_1^*)$. To this end, let $u = u_2 \in D(D_1B_1^*)$. $\mathcal{H}_2 \cap D(D_2B_2) \subset D(T)$ in [\(1\)](#page-6-0). Then

$$
|(D_2B_2u_2, v_1)| = |(Tu_2, v)| \lesssim ||u_2||_2.
$$

It follows that $v_1 \in D(D_2)$, since $(D_2B_2)^* = B_2^*D_2$. Therefore, for general $u \in D(T)$, we have

$$
(Tu, v) = (B_1D_1u_1, v) + (u, B_2^*D_2v_1),
$$

so $|(B_1D_1u_1, v_1 + v_2)| \lesssim ||u||_2$. Since $(B_1D_1)^* = D_1B_1^*$, it follows that $v_1 + v_2 \in$ $\mathsf{D}(D_1B_1^*$). \Box

We end this section with a short discussion of the definition of the functional calculus of bisectorial operators. For further details see [\[1\]](#page-19-7), where the corresponding theory for sectorial operators is readily adapted to bisectorial operators.

Given a bisectorial operator T in a Hilbert space H , that is a closed and densely defined operator T with $\sigma(T) \subset S_\omega$ for some $\omega < \pi/2$ and resolvent bounds

$$
\|(\lambda I - T)^{-1}\| \lesssim 1/\text{dist}\,(\lambda, S_{\omega}),
$$

there is a natural definition of $\phi(T)$ for any rational function $\phi(\lambda)$ which is bounded and without poles in S_{ω} . Useful such symbols in this paper are for example $1/(1 +$ $t^2\lambda^2$ and $t\lambda/(1+t^2\lambda^2)$, with scale parameter $t>0$.

If T is not injective, then there is a topological splitting $\mathcal{H} = \mathcal{H}_0 \oplus \mathcal{H}_1$, with $\mathcal{H}_0 = N(T)$ and $\mathcal{H}_1 = R(T)$. Indeed,

$$
I = (I + itT)^{-1} + itT(I + itT)^{-1},
$$

where the two terms converge strongly to the projections onto \mathcal{H}_0 and \mathcal{H}_1 respectively as $t \to \infty$. Thus $T = 0 \oplus T_1$ and $(\lambda I - T_1)^{-1} = \lambda^{-1} I \oplus (\lambda I - T_1)^{-1}$, where $T_1 := T|_{\mathcal{H}_1}$ is an injective bisectorial operator in \mathcal{H}_1 .

With the Dunford integral

$$
\psi(T) := \psi(0)I_{\mathcal{H}_0} \oplus \frac{1}{2\pi i} \int_{\gamma} \psi(\lambda) (\lambda I_{\mathcal{H}_1} - T_1)^{-1} d\lambda,
$$

the functional calculus is extended to all symbols $\psi : S^o_\mu \cup \{0\} \to \mathbb{C}$ which are holomorphic on an open bisector $S^o_\mu \supset S_\omega \setminus \{0\}$, with estimates $|\psi(\lambda)| \lesssim \min(|\lambda|^s, |\lambda|^{-s})$ for some $s > 0$, so that the integral is convergent in the operator norm on \mathcal{H}_1 . Here the curve $\gamma = \{te^{\pm i\theta} \, ; \, t \in \mathbf{R}\}, \, \omega < \theta < \mu$, is oriented counter clockwise around S_{ω} .

To obtain $\phi(T)$ as bounded operators on H for general bounded holomorphic symbols ϕ , without decay at 0 and ∞ , square function estimates

$$
\int_0^\infty \|\psi(tT)f\|_{\mathcal{H}_1}^2 \frac{dt}{t} \approx \|f\|_{\mathcal{H}_1}^2, \qquad f \in \mathcal{H}_1,
$$

are required. Usually, it suffices to show estimates \lesssim , as estimates \gtrsim for T follow from estimates \lesssim for T^* . Another basic result concerning square function estimates, is that if such hold for one symbol ψ , then they hold for all ψ such that $|\psi(\lambda)| \lesssim$ $\min(|\lambda|^s, |\lambda|^{-s})$ for some $s > 0$, and $\tilde{\psi}|_{S_{\omega} \pm} \neq 0$.

Given such square function estimates, it follows that $\psi_n(T)$ are uniformly bounded and converges strongly in $\mathcal{L}(\mathcal{H})$ whenever $\sup_{n\in\mathbf{Z}_+,\lambda\in S_\mu^o} |\psi_n(\lambda)| < \infty$ and $\psi_n(\lambda)$ converges for each $\lambda \in S^o_\mu \cup \{0\}$. Through such a limiting argument, we construct a bounded homomorphism

 $\phi \mapsto \phi(T),$

taking bounded symbols $\phi: S^o_\mu \cup \{0\} \to \mathbb{C}$ which are holomorphic on S^o_μ to bounded linear operators $\phi(T)$ on H.

3. Applications to exterior differential systems

In this section, we show how operators of the form $B_1D_1 + D_2B_2$ appear in connection with exterior differential systems for differential form. We first fix notation. Instead of writing $\{dx_0, dx_1, \ldots, dx_n\}$ for the basis one-forms, we shall keep the notation $\{e_0, e_1, \ldots, e_n\}$ from Section [1](#page-0-0) for the basis vectors, and we use the terminology k-vector field instead of k-form, in the euclidean space \mathbb{R}^{1+n} .

The space of k-vectors in \mathbb{R}^{1+n} we define to be the $\binom{1+n}{k}$ $\binom{+n}{k}$ dimensional complex linear space

 $\wedge^k \mathbf{R}^{1+n} := \text{span}_{\mathbf{C}} \{ e_{s_1} \wedge \ldots \wedge e_{s_k} \; ; \; 0 \le s_1 < s_2 < \ldots s_k \le n \}, \qquad k = 2, \ldots, n+1,$

and we let $\wedge^0 {\bf R}^{1+n} := {\bf C} \wedge^1 {\bf R}^{1+n} := {\bf C}^{1+n}$ and $\wedge^k {\bf R}^{1+n} := \{0\}$ if $k \notin \{0, 1, \ldots, n+1\}.$ Given a vector $v = \sum_{j=0}^{n} v_j e_j \in \wedge^1 \mathbb{R}^{1+n}$ and a k-vector $w = \sum_{0 \le s_1 < ... s_k \le n} w_s e_s \in$ $\wedge^k \mathbf{R}^{1+n}$, writing $s = \{s_1, \ldots, s_k\}$ and $e_s := e_{s_1} \wedge \ldots \wedge e_{s_k}$, we have in particular the exterior product $v \wedge w \in \wedge^{k+1} \mathbb{R}^{1+n}$ and the (left) interior product $v \wedge w \in \wedge^{k-1} \mathbb{R}^{1+n}$ defined bilinearly using

$$
e_j\wedge e_s:=\begin{cases}\epsilon(j,s)e_{\{j\}\cup s},&j\notin s,\\0,&j\in s,\end{cases}\hspace{0.5cm}e_j\mathbin{\lrcorner} e_s:=\begin{cases}0,&j\notin s,\\ \epsilon(j,s\setminus\{j\})e_{s\setminus\{j\}},&j\in s,\end{cases}
$$

where the permutation sign is $\epsilon(j, s) := (-1)^{|\{s_i : j>s_i\}|}$. Defining inner products on $\wedge^k \mathbf{R}^{1+n}$, $k = 0, 1, \ldots, n+1$, so that the standard bases above are ON-bases, we have that

$$
v \wedge (\cdot) : \wedge^k \mathbf{R}^{1+n} \to \wedge^{k+1} \mathbf{R}^{1+n}
$$
 and $v \cup (\cdot) : \wedge^{k+1} \mathbf{R}^{1+n} \to \wedge^k \mathbf{R}^{1+n}$,

are adjoint multiplication operators if v is a vector, that is $v \in \wedge^1 \mathbb{R}^{1+n}$, with real coefficients. The corresponding differential operators are the exterior derivative operator

$$
\nabla_{t,x} \wedge f := \sum_{j=0}^n e_j \wedge \partial_j f,
$$

mapping k-vector fields $f: \mathbf{R}^{1+n} \to \wedge^k \mathbf{R}^{1+n}$ to $k+1$ -vector fields, and the interior derivative operator

$$
\nabla_{t,x}\,\lrcorner\, g:=\sum_{j=0}^n e_j\,\lrcorner\,\partial_j g,
$$

mapping $k+1$ -vector fields $g: \mathbf{R}^{1+n} \to \wedge^{k+1} \mathbf{R}^{1+n}$ to k-vector fields. As special cases of these operators, we have the gradient and curl, being the exterior derivative acting on scalars and vectors $(k = 0 \text{ and } k = 1 \text{ respectively})$, and the divergence being the interior derivative acting on vectors. We also note the duality $\int \int (\nabla_{t,x} \wedge f, g) dt dx =$ $-\iiint (f, \nabla_{t,x} \, \lrcorner \, g) dt dx$ for compactly supported fields.

The basic exterior differential system in \mathbb{R}^{1+n} that we want to consider is

(2)
$$
\begin{cases} \nabla_{t,x} \,\underline{\quad} \tilde{f}_{k+1} = \nabla_{t,x} \wedge \tilde{f}_{k-1}, \\ \nabla_{t,x} \wedge \tilde{f}_{k+1} = 0 = \nabla_{t,x} \,\underline{\quad} \tilde{f}_{k-1}, \end{cases}
$$

for a $k + 1$ -vector field f_{k+1} and a $k - 1$ -vector field f_{k-1} . Two important special cases are the following. If $k = 0$, then the system reads $\text{div}_{t,x} \tilde{f}_1 = 0 = \text{curl}_{t,x} \tilde{f}_1$, since $\tilde{f}_{-1} = 0$. This is nothing but the Laplace equation, written for the gradient as in Section [1.](#page-0-0) If $k = 1$, then the system reads $\nabla_{t,x} \,\underline{\mathcal{J}} \, \tilde{f}_2 = \nabla_{t,x} \tilde{f}_0, \, \nabla_{t,x} \wedge f_2 = 0.$ This equation is the Stokes' system of linearized hydrostatics, written for the vorticity \tilde{f}_2 and the the pressure f_0 .

Consider next a bilipschitz map $\rho : \mathbf{R}^{1+n}_+ \to \Omega \subset \mathbf{R}^{1+n}$, and the system [\(2\)](#page-8-0) in Ω. We want to pull back this system of equations to \mathbf{R}^{1+n}_+ , and recall therefore the following facts from differential geometry. At a fixed point in \mathbb{R}^{1+n}_+ , denote by ρ the Jacobian matrix of all partial derivatives of ρ. Extend this linear map as a \wedge -homomorphism to $\wedge^k \mathbf{R}^{1+n}$, letting

$$
\underline{\rho}(e_{s_1} \wedge \ldots \wedge e_{s_k}) := (\underline{\rho}e_{s_1}) \wedge \ldots \wedge (\underline{\rho}e_{s_k}).
$$

Given a k-vector field $f: \Omega \to \wedge^k \mathbf{R}^{1+n}$, we define the pullback of f by ρ to be the k-vector field

$$
\rho^* f(t,x) := \underline{\rho}_{(t,x)}^* (f(\rho(t,x)))
$$

in \mathbf{R}^{1+n}_+ , where ρ^*_{α} $_{(t,x)}^*$ is the adjoint of the Jacobian matrix at (t, x) . A fundamental well known result is that

(3) $\nabla_{t,x} \wedge (\rho^* f) = \rho^* (\nabla_{t,x} \wedge f).$

Less commonly used is the equivalent dual result that

(4)
$$
\nabla_{t,x} \mathop{\lrcorner} (J_{\rho} \rho_*^{-1} g) = J_{\rho} \rho_*^{-1} (\nabla_{t,x} \mathop{\lrcorner} g),
$$

where J_{ρ} is the Jacobian determinant of ρ and

$$
\rho_*^{-1}g(t,x):=\underline{\rho}_{(t,x)}(g(\rho(t,x)))
$$

10 ANDREAS ROSEN´

is the push forward of g by ρ^{-1} . Applying [\(3\)](#page-8-1) and [\(4\)](#page-8-2), we find that [\(2\)](#page-8-0) in Ω is equivalent to

(5)
$$
\begin{cases} \nabla_{t,x} \, \mathbf{1}(A_{k+1}(t,x) f_{k+1}) = A_k(t,x) (\nabla_{t,x} \wedge f_{k-1}), \\ \nabla_{t,x} \wedge f_{k+1} = 0 = \nabla_{t,x} \, \mathbf{1}(A_{k-1}(t,x) f_{k-1}) \end{cases}
$$

in \mathbf{R}^{1+n}_+ , where $f_j := \rho^* \tilde{f}_j$ and $A_j := J_\rho(\rho^* \rho_*)^{-1}$ is the Jacobian determinant times the inverse of the metric tensor $G = \rho^* \rho_*$, extended as a ∧-homomorphism to $\wedge^j \mathbf{R}^{1+n}$.

We now show, analogous to the case $k = 0$ in the introduction, how [\(5\)](#page-9-0) is equivalent to a vector valued ordinary differential equation $\partial_t f_t + T f_t = 0$, for general bounded measurable and accretive coefficients $A_j(t,x) \in \mathcal{L}(\wedge^j \mathbf{R}^{1+n}), j =$ $k-1, k, k+1$, with an infinitesimal generator T of the form $T = B_1D_1 + D_2B_2$. We use the natural identifications

$$
\wedge^k \mathbf{R}^n \oplus \wedge^{k+1} \mathbf{R}^n = \wedge^{k+1} \mathbf{R}^{1+n} : \overline{f}_k \oplus \overline{f}_{k+1} \approx e_0 \wedge \overline{f}_k + \overline{f}_{k+1} = f_{k+1},
$$

$$
\wedge^{k-2} \mathbf{R}^n \oplus \wedge^{k-1} \mathbf{R}^n = \wedge^{k-1} \mathbf{R}^{1+n} : \overline{f}_{k-2} \oplus \overline{f}_{k-1} \approx e_0 \wedge \overline{f}_{k-2} + \overline{f}_{k-1} = f_{k-1},
$$

with corresponding splittings of the coefficient matrices so that

$$
A_{k+1}f_{k+1} = e_0 \wedge (a_{k+1}\overline{f}_k + b_{k+1}\overline{f}_{k+1}) + (c_{k+1}\overline{f}_k + d_{k+1}\overline{f}_{k+1}),
$$

$$
A_{k-1}f_{k-1} = e_0 \wedge (a_{k-1}\overline{f}_{k-2} + b_{k-1}\overline{f}_{k-1}) + (c_{k-1}\overline{f}_{k-2} + d_{k-1}\overline{f}_{k-1}),
$$

and similarly for A_k .

Let \mathcal{H}^j_\wedge denote the closure of the range of $\nabla_x \wedge (\cdot) : L_2(\mathbf{R}^n; \wedge^{j-1}\mathbf{R}^n) \to L_2(\mathbf{R}^n; \wedge^j\mathbf{R}^n)$, and let \mathcal{H}_{\perp}^{j} be the closure of the range of $\nabla_x \Box(\cdot) : L_2(\mathbf{R}^n; \wedge^{j+1}\mathbf{R}^n) \to L_2(\mathbf{R}^n; \wedge^j\mathbf{R}^n)$. Fundamental results are that \mathcal{H}^j_\wedge is the null space of $\nabla_x \wedge (\cdot) : L_2(\mathbb{R}^n; \wedge^j \mathbb{R}^n) \to$ $L_2(\mathbf{R}^n; \wedge^{j+1}\mathbf{R}^n)$, $\mathcal{H}^j_{\mathfrak{I}}$ is the nullspace of $\nabla_x \mathfrak{I}(\cdot) : L_2(\mathbf{R}^n; \wedge^j\mathbf{R}^n) \to L_2(\mathbf{R}^n; \wedge^{j-1}\mathbf{R}^n)$, and we have an orthogonal Hodge splitting

$$
L_2(\mathbf{R}^n; \wedge^j \mathbf{R}^n) = \mathcal{H}^j_\wedge \oplus \mathcal{H}^j_\omega.
$$

Proposition 3.1. Assume that $A_j \in L_\infty(\mathbf{R}^n; \mathcal{L}(\wedge^{j-1}\mathbf{R}^n \oplus \wedge^j\mathbf{R}^n))$ are t-independent and accretive on $\mathcal{H}^{j-1}_{\lambda} \oplus \mathcal{H}^{j}_{\lambda}$, $j = k - 1, k, k + 1$. Define the $\wedge^{k-2} \mathbb{R}^n \oplus \wedge^{k-1} \mathbb{R}^n \oplus$ $\wedge^k {\bf R}^n \oplus \wedge^{k+1} {\bf R}^n$ valued function \hat{f} by

$$
\hat{f} = \begin{bmatrix} \hat{f}_{k-2} \\ \hat{f}_{k-1} \\ \hat{f}_{k} \\ \hat{f}_{k+1} \end{bmatrix} := \begin{bmatrix} a_{k-1} \overline{f}_{k-2} + b_{k-1} \overline{f}_{k-1} \\ \overline{f}_{k-1} \\ a_{k+1} \overline{f}_{k} + b_{k+1} \overline{f}_{k+1} \end{bmatrix}.
$$

Then the exterior differential system [\(5\)](#page-9-0) for the $\wedge^{k-1} \mathbf{R}^{1+n} \oplus \wedge^{k+1} \mathbf{R}^{1+n}$ valued function $f_{k-1} \oplus f_{k+1}$ is equivalent to the vector valued ordinary differential equation

$$
\partial_t \hat{f} + (B_1 D_1 + D_2 B_2) \hat{f} = 0,
$$

together with the constraint $\hat{f} \in \overline{R(B_1D_1 + D_2B_2)}$ for each t, where

$$
B_1 := \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & a_k^{-1} & -a_k^{-1}b_k & 0 \\ 0 & c_ka_k^{-1} & d_k - c_ka_k^{-1}b_k & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix},
$$

\n
$$
D_1 := \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & \nabla_{x} \mathbf{1}(\cdot) & 0 \\ 0 & -\nabla_{x} \wedge (\cdot) & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix},
$$

\n
$$
D_2 := \begin{bmatrix} 0 & \nabla_{x} \mathbf{1}(\cdot) & 0 & 0 \\ -\nabla_{x} \mathbf{1}(\cdot) & 0 & 0 & 0 \\ 0 & 0 & 0 & \nabla_{x} \mathbf{1}(\cdot) \\ 0 & 0 & -\nabla_{x} \mathbf{1}(\cdot) & 0 \end{bmatrix} \quad \text{and}
$$

\n
$$
B_2 := \begin{bmatrix} a_{k-1}^{-1} & -a_{k-1}^{-1}b_{k-1} & 0 & 0 \\ c_{k-1}a_{k-1}^{-1} & d_{k-1} - c_{k-1}a_{k-1}^{-1}b_{k-1} & 0 & 0 \\ 0 & 0 & a_{k+1}^{-1} & -a_{k+1}^{-1}b_{k+1} \\ 0 & 0 & c_{k+1}a_{k+1}^{-1} & d_{k+1} - c_{k+1}a_{k+1}^{-1}b_{k+1} \end{bmatrix}
$$

satisfy the hypothesis in Theorem [1.1.](#page-2-0)

Proof. The equation $\nabla_{t,x} \wedge f_{k+1} = 0$ is equivalent to

$$
\begin{cases} \partial_t \overline{f}_{k+1} - \nabla_x \wedge \overline{f}_k = 0, \\ \nabla_x \wedge \overline{f}_{k+1} = 0. \end{cases}
$$

The equation $\nabla_{t,x} \mathcal{A}_{k-1} f_{k-1} = 0$ is equivalent to

$$
\begin{cases} \nabla_x \mathbf{1} \left(a_{k-1} \overline{f}_{k-2} + b_{k-1} \overline{f}_{k-1} \right) = 0, \\ \partial_t (a_{k-1} \overline{f}_{k-2} + b_{k-1} \overline{f}_{k-1}) + \nabla_x \mathbf{1} \left(c_{k-1} \overline{f}_{k-2} + d_{k-1} \overline{f}_{k-1} \right) = 0. \end{cases}
$$

The equation $\nabla_{t,x} \mathcal{A}(A_{k+1}f_{k+1}) = A_k(\nabla_{t,x} \wedge f_{k-1})$ is equivalent to

$$
\begin{cases}\n-\nabla_{x} \mathop{\lrcorner} (a_{k+1}\overline{f}_k + b_{k+1}\overline{f}_{k+1}) = a_k(\partial_t \overline{f}_{k-1} - \nabla_{x} \wedge \overline{f}_{k-2}) + b_k(\nabla_x \wedge \overline{f}_{k-1}), \\
\partial_t (a_{k+1}\overline{f}_k + b_{k+1}\overline{f}_{k+1}) + \nabla_{x} \mathop{\lrcorner} (c_{k+1}\overline{f}_k + d_{k+1}\overline{f}_{k+1}) \\
= c_k(\partial_t \overline{f}_{k-1} - \nabla_x \wedge \overline{f}_{k-2}) + d_k(\nabla_x \wedge \overline{f}_{k-1}).\n\end{cases}
$$

Written in terms of \hat{f} , the four evolution equations are

$$
\begin{cases} \partial_t \hat{f}_{k-2} + \nabla_x \mathop{\lrcorner} (c_{k-1} \overline{f}_{k-2} + d_{k-1} \overline{f}_{k-1}) = 0, \\ \partial_t \hat{f}_{k-1} - \nabla_x \wedge \overline{f}_{k-2} + a_k^{-1} b_k (\nabla_x \wedge \overline{f}_{k-1}) + a_k^{-1} \nabla_x \mathop{\lrcorner} (a_{k+1} \overline{f}_k + b_{k+1} \overline{f}_{k+1}) = 0, \\ \partial_t \hat{f}_k + \nabla_x \mathop{\lrcorner} (c_{k+1} \overline{f}_k + d_{k+1} \overline{f}_{k+1}) - c_k (\partial_t \hat{f}_{k-1} - \nabla_x \wedge \overline{f}_{k-2}) - d_k (\nabla_x \wedge \overline{f}_{k-1}) = 0, \\ \partial_t \hat{f}_{k+1} - \nabla_x \wedge \overline{f}_k = 0, \end{cases}
$$

and the remaining two equations give the constraints $\nabla_{\underline{x}} \wedge \hat{f}_{k+1} = 0 = \nabla_{x} \cup \hat{f}_{k-2}$. We next write the evolution equations in terms of \hat{f} , using $\overline{f}_{k-2} = a_{k-1}^{-1}$ $_{k-1}^{-1}(\hat{f}_{k-2}-b_{k-1}\hat{f}_{k-1})$ and $\overline{f}_k = a_{k+1}^{-1}(\hat{f}_k - b_{k+1}\hat{f}_{k+1})$. The tangential derivatives in the evolution equations

that appear with coefficients to the right are

$$
\begin{bmatrix}\n\nabla_{x} \mathbf{1} (c_{k-1} a_{k-1}^{-1} (\hat{f}_{k-2} - b_{k-1} \hat{f}_{k-1}) + d_{k-1} \hat{f}_{k-1}) \\
-\nabla_{x} \wedge (a_{k-1}^{-1} (\hat{f}_{k-2} - b_{k-1} \hat{f}_{k-1})) \\
\nabla_{x} \mathbf{1} (c_{k+1} a_{k+1}^{-1} (\hat{f}_{k} - b_{k+1} \hat{f}_{k+1}) + d_{k+1} \hat{f}_{k+1}) \\
-\nabla_{x} \wedge (a_{k+1}^{-1} (\hat{f}_{k} - b_{k+1} \hat{f}_{k+1}))\n\end{bmatrix} = D_{2} B_{2} \hat{f}.
$$

The tangential derivatives in the evolution equations that appear with coefficients to the left are

$$
\begin{bmatrix} 0 \\ a_k^{-1}b_k \nabla_x \wedge \hat{f}_{k-1} + a_k^{-1} \nabla_x \wedge \hat{f}_k \\ c_k (a_k^{-1}b_k \nabla_x \wedge \hat{f}_{k-1} + a_k^{-1} \nabla_x \wedge \hat{f}_k) - d_k \nabla_x \wedge \hat{f}_{k-1} \\ 0 \end{bmatrix} = B_1 D_1 \hat{f}.
$$

This shows that the evolution equation for \hat{f} is $\partial_t \hat{f} + (B_1 D_1 + D_2 B_2) \hat{f} = 0$. To show that the constraint $\nabla_x \wedge \hat{f}_{k+1} = 0 = \nabla_x \sqcup \hat{f}_{k-2}$ is equivalent to $f \in \overline{R(B_1D_1 + D_2B_2)}$, we note that

$$
\overline{\mathsf{R}(B_1D_1+D_2B_2)}=B_1\overline{\mathsf{R}(D_1)}\oplus \overline{\mathsf{R}(D_2)}
$$

= $B_1(\mathcal{H}_{\wedge}^k\oplus \mathcal{H}_{\perp}^{k-1})\oplus ((\mathcal{H}_{\perp}^k\oplus \mathcal{H}_{\wedge}^{k-1})\oplus (\mathcal{H}_{\wedge}^{k+1}\oplus \mathcal{H}_{\perp}^{k-2}))$
= $L_2(\mathbf{R}^n; \wedge^{k-1}\mathbf{R}^n\oplus \wedge^k\mathbf{R}^n)\oplus (\mathcal{H}_{\wedge}^{k+1}\oplus \mathcal{H}_{\perp}^{k-2}),$

by Proposition [2.3](#page-5-1) and a L_2 Hodge splitting of $L_2(\mathbf{R}^n; \wedge^{k-1}\mathbf{R}^n \oplus \wedge^k\mathbf{R}^n)$ adapted to B_1 .

Given Theorem [1.1](#page-2-0) and Proposition [3.1,](#page-9-1) we can proceed as in [\[5,](#page-19-4) Thm. 2.3], where the case $k = 0$ was treated, to represent solutions to the exterior differential system [\(5\)](#page-9-0) with functional calculus as outlined in Section [2.](#page-4-0) To this end, define symbols

$$
e^{-t\lambda}\chi^+(\lambda) := \begin{cases} e^{-t\lambda}, & \text{Re }\lambda > 0, \\ 0, & \text{Re }\lambda \le 0, \end{cases} \qquad t \ge 0,
$$

$$
e^{-t\lambda}\chi^-(\lambda) := \begin{cases} 0, & \text{Re }\lambda \ge 0, \\ e^{-t\lambda}, & \text{Re }\lambda < 0, \end{cases} \qquad t \le 0.
$$

For $t = 0$, we obtain bounded spectral projections $\chi^{\pm}(T)$, with $\chi^{+}(T) + \chi^{-}(T)$ being the projection onto $\overline{R(T)}$ along $N(T)$. The following result roughly states that the spectral subspace $\chi^+(T)L_2 := \overline{R(\chi^+(T))}$ is a Hardy type subspace containing traces of solutions to [\(5\)](#page-9-0) in \mathbf{R}^{1+n}_+ , whereas the spectral subspace $\chi^-(T)L_2 := \mathsf{R}(\chi^-(T))$ is a Hardy type subspace containing traces of solutions to (5) in \mathbb{R}^{1+n}_{-} , and the operators $e^{-tT}\chi^{\pm}(T)$ are Cauchy integral type operators, giving the value of the function at (t, \cdot) from the boundary trace.

Theorem 3.2. Consider the exterior differential system (5) , with bounded, t-independent, accretive coefficients A_{k-1}, A_k, A_{k+1} , and the associated operator $T = B_1D_1 + D_2B_2$ as in Proposition [3.1.](#page-9-1) Given $\hat{f}_0^{\pm} \in \chi^{\pm}(T) L_2$, the function $f \approx \hat{f}$ defined by

$$
\hat{f}^{\pm}(t,x) := (e^{-tT} \chi^{\pm}(T)) \hat{f}_0^{\pm}(x), \qquad (t,x) \in \mathbf{R}_{\pm}^{1+n},
$$

is a solution to [\(5\)](#page-9-0), with limits $\lim_{t\to 0^{\pm}} || \hat{f}_t^{\pm} - \hat{f}_0^{\pm} ||_2 = 0$ and $\lim_{t\to \pm \infty} || \hat{f}_t^{\pm} ||_2 = 0$. Conversely, any solution $f^{\pm} \approx \hat{f}^{\pm}$ to [\(5\)](#page-9-0) in \mathbf{R}_{\pm}^{1+n} with estimates $\sup_{t>0} \int_{t < \pm s < 2t} ||f_s||_2^2 ds$ ∞ is of this form, and in particular has the stated limits at $t = 0$, for some $\hat{f}_0^{\pm} \in \chi^{\pm}(T)L_2$, and $t = \infty$.

These solutions have square function, non-tangential maximal function and $L^t_{\infty}L^x_2$ estimates

$$
\int_{\pm t>0} \|\partial_t \hat{f}_t^{\pm}\|_2^2 t dt \approx \|\widetilde{N}_*(\hat{f}^{\pm})\|_2^2 \approx \sup_{\pm t>0} \|\hat{f}_t^{\pm}\|_2^2 \approx \|\hat{f}_0^{\pm}\|_2^2.
$$

The idea of proof is found in [\[5,](#page-19-4) Thm. 3.2] and [\[2,](#page-19-8) Thm. 8.2]. In particular, the estimates follow from Theorem [1.1,](#page-2-0) using the symbol $\psi(\lambda) = \lambda e^{-\lambda} \chi^{\pm}(\lambda)$ for the square function estimates, and the symbol $\phi(\lambda) = e^{-\lambda} \chi^{\pm}(\lambda)$ for the non-tangential maximal function estimates and the $L^t_{\infty} L^x_2$ estimates. We omit the details.

4. Square function estimates

In this section, we prove the square function estimates for the operator T in Theorem [1.1.](#page-2-0) We start by simplifying the problem with Lemma [4.1,](#page-12-1) and we use the following operators.

$$
P_t^1 := (I + t^2 (B_1 D_1)^2)^{-1} = \frac{1}{2} (R_t^1 + R_{-t}^1),
$$

\n
$$
Q_t^1 := t B_1 D_1 (I + t^2 (B_1 D_1)^2)^{-1} = \frac{i}{2} (R_t^1 - R_{-t}^1),
$$

\n
$$
P_t^2 := (I + t^2 (D_2 B_2)^2)^{-1} = \frac{1}{2} (R_t^2 + R_{-t}^2),
$$

\n
$$
Q_t^2 := t D_2 B_2 (I + t^2 (D_2 B_2)^2)^{-1} = \frac{i}{2} (R_t^2 - R_{-t}^2).
$$

It is known that these operators are uniformly bounded for $t > 0$ and that square function estimates

$$
\int_0^\infty \|Q_t^1 f\|_2^2 \frac{dt}{t} + \int_0^\infty \|Q_t^2 f\|_2^2 \frac{dt}{t} \lesssim \|f\|_2^2, \qquad f \in L_2(\mathbf{R}^n; \mathbf{C}^N),
$$

hold. See for example [\[5,](#page-19-4) Thm. 3.4].

Lemma 4.1. Let

$$
\Theta_t:=Q_t^2R_t^1B_1.
$$

Assume that we have square function estimates

$$
\int_0^\infty \|\Theta_t f\|_2^2 \frac{dt}{t} \lesssim \|f\|_2^2, \qquad \text{for all } f \in \mathcal{H}_1.
$$

Then we have square function estimates $\int_0^\infty \|\psi(tT)f\|_2^2 \frac{dt}{t} \lesssim \|f\|_2^2$, $f \in L_2(\mathbf{R}^n; \mathbf{C}^N)$, for ψ as in Theorem [1.1.](#page-2-0)

Proof. It is known, see [\[1\]](#page-19-7), that it suffices to prove the square function estimate for $\psi(\lambda) = \lambda/(1 + \lambda^2)$. For this ψ , we see from Proposition [2.4](#page-5-2) that

$$
\psi(tT) = \frac{i}{2} \left((I + itT)^{-1} - (I - itT)^{-1} \right) = \begin{bmatrix} 0 & 0 & 0 \\ 0 & Q_t^1 & 0 \\ 0 & (R_{-t}^2 - I)Q_t^1 + Q_t^2 R_t^1 & Q_t^2 \end{bmatrix}, \qquad t > 0,
$$

by writing

$$
\frac{1}{2} \left(\frac{tD_2B_2}{I + itD_2B_2} \frac{1}{I + itB_1D_1} + \frac{tD_2B_2}{I - itD_2B_2} \frac{1}{I - itB_1D_1} \right)
$$
\n
$$
= \frac{itD_2B_2}{I - itD_2B_2} \frac{tB_1D_1}{I + t^2(B_1D_1)^2} + \frac{tD_2B_2}{I + t^2(D_2B_2)^2} \frac{1}{I + itB_1D_1}.
$$

Since R_{-t}^2 are uniformly bounded and since we have square function estimates for Q_t^1 and Q_t^2 , it suffices to prove square function estimates for $Q_t^2 R_t^1$ on $B_1 \mathcal{H}_1$ as claimed. \square

We now prove the square function estimates for Θ_t using techniques from the proof of the Kato square root estimate, following [\[3,](#page-19-5) Sec. 4].

Definition 4.2. Let $\mathcal{D} = \bigcup_{j \in \mathbf{Z}} \mathcal{D}_{2^{-j}}$ denote the dyadic cubes in \mathbb{R}^n , with

$$
\mathcal{D}_t := \{ 2^{-j}(0,1)^n + 2^{-j}k \; ; \; k \in \mathbb{Z}^n \}, \qquad 2^{-j-1} < t \le 2^{-j}, j \in \mathbb{Z}.
$$

Write $\ell(Q)$ for side length and |Q| for measure of a cube Q. Given $Q \in \mathcal{D}$, write

$$
A_0(Q) := Q, \qquad A_k(Q) := (2^k Q) \setminus (2^{k-1} Q), \quad k \ge 1,
$$

for the dyadic annuli around Q, where aQ denote the cube with same center as Q but with $\ell(aQ) = a\ell(Q)$.

The key tool in the proof of the square function estimates, as well as for the nontangential maximal function estimates, are the following L_2 off-diagonal estimates.

Proposition 4.3. For any $m < \infty$, there exists $C_m < \infty$ such that

$$
\|\Theta_t f\|_{L_2(F)} \lesssim C_m (t/\text{dist}(E, F))^m \|f\|_2,
$$

for all $f \in L_2(\mathbf{R}^n;\mathbf{C}^N)$ with supp $f \subset E$, and any closed subsets $E, F \subset \mathbf{R}^n$ such that dist $(E, F) := \inf\{|x - y| : x \in E, y \in F\} > 0.$

Proof. These estimates are known to hold for R_t^j t_i^j , and therefore for R_{-}^j $\frac{j}{-t}$, P_t^j and Q_t^j $_t^j,$ $j = 1, 2$, see for example [\[4,](#page-19-6) Sec. 5]. From this, the estimates for $\Theta_t := Q_t^2 (I - R_t^1) B_1$ follow as in $[3, \text{ Lem. } 2.26]$.

These L_2 off-diagonal estimates enable us to approximate the family of operators ${\{\Theta_t\}}_{t>0}$ by a family of multiplication operators ${\{\gamma_t\}}_{t>0}$, where formally $\gamma_t = \Theta_t 1$. More precisely, we let

$$
\gamma_t(x)v := \sum_{k=0}^{\infty} \Theta_t(v\chi_{A_k(Q)})(x), \qquad x \in Q \in \mathcal{D}_t, v \in \mathbf{C}^N,
$$

where $\chi_{A_k(Q)}$ denotes the characteristic function of the dyadic annulus $A_k(Q)$. From Proposition [4.3,](#page-13-0) we have the estimate

(6)
$$
\|\gamma_t\|_{L_2(Q)} \lesssim \sum_{k=0}^{\infty} 2^{-km} 2^{kn/2} \leq C
$$

uniformly for all $Q \in \mathcal{D}_t$, $t > 0$, by choosing $m > n/2$.

We also need the following Sobolev–Poincaré inequality, see [\[10,](#page-19-9) Sec. 7.8].

Lemma 4.4. Let $1 < r < n$, or $r = n = 1$, and $1/r^* = 1/r - 1/n$. Assume that $1 \leq q \leq r^*$ and $r \leq p \leq \infty$. Then there exists $C < \infty$ such that for all $u \in H^1_{loc}(\mathbb{R}^n)$ and $0 < r \leq R < \infty$, we have the estimate

$$
||u - u_S||_{L_q(\Omega)} \leq C |\Omega|^{1/q - 1/p + 1/n} R^n |S|^{-1} ||\nabla u||_{L_p(\Omega)},
$$

for any convex set Ω with diameter R and measure $|\Omega|$ and any measurable subset $S \subset \Omega$ with measure $|S|$.

Proposition 4.5. We have the estimate

$$
\int_0^\infty \|\Theta_t f - \gamma_t E_t f\|_2^2 \frac{dt}{t} \lesssim \|f\|_2^2, \quad \text{for all } f \in \mathcal{H}_1,
$$

where E_t denotes the dyadic averaging operator

$$
E_t f(x) = E_Q f := \frac{1}{|Q|} \int_Q f(y) dy, \qquad x \in Q \in \mathcal{D}_t.
$$

Proof. Let P_t, Q_t denote the unpertubed operators P_t^1, Q_t^1 , that is

$$
P_t := (I + t^2 D_1^2)^{-1}, \qquad Q_t := t D_1 (I + t^2 D_1^2)^{-1}.
$$

Write

 $\Theta_t f - \gamma_t E_t f = \Theta_t (I - P_t) f + (\Theta_t - \gamma_t E_t) P_t f + (\gamma_t E_t) E_t (P_t - I) f =: I + II + III.$

For the first term we have

$$
||I||_2 = ||Q_t^2(I - R_t^1)Q_t f||_2 \lesssim ||Q_t f||_2,
$$

since $(I + itB_1D_1)^{-1}B_1(t^2D_1^2(I + t^2D_1^2)) = (I + itB_1D_1)^{-1}(tB_1D_1)Q_t$, and square function estimates for Q_t give the desired estimate.

For the term III we note from [\(6\)](#page-13-1) that $\|\gamma_t E_t\|_{L_2\to L_2} \leq C$. Square function estimates for $E_t(P_t - I)$ can be proved as in [\[8,](#page-19-3) Prop. 5.7], replacing Π there by the operator D_1 .

The term II we write

$$
(\Theta_t - \gamma_t E_t) P_t f = \sum_{k \ge 0} \Theta_t ((P_t f - E_Q(P_t f)) \chi_{A_k(Q)}), \quad \text{on } Q \in \mathcal{D}_t.
$$

Proposition [4.3](#page-13-0) and Poincaré's inequality in Lemma [4.4](#page-13-2) yields

$$
||II||_2^2 \lesssim \sum_{Q \in \mathcal{D}_t} \left(\sum_{k=0}^{\infty} 2^{-km} ||P_t f - E_Q(P_t f)||_{L_2(A_k(Q))} \right)^2
$$

$$
\lesssim \sum_{Q \in \mathcal{D}_t} \sum_{k=0}^{\infty} 2^{-km} ||P_t f - E_Q(P_t f)||_{L_2(A_k(Q))}^2
$$

$$
\lesssim \sum_{Q \in \mathcal{D}_t} \sum_{k=0}^{\infty} 2^{-km+2k(n+1)} ||t \nabla P_t f||_{L_2(A_k(Q))}^2 \approx ||t \nabla P_t f||_2^2,
$$

if we choose m sufficiently large. Since D_1 is elliptic on $R(D_1)$, the square function estimate for II follows from that for Q_t . .

To prove square function estimates for the remaining paraproduct term $\gamma_t E_t f$, we use the following test functions. For a small fixed parameter $\epsilon > 0$, we define for all dyadic cubes $Q \in \mathcal{D}$ and unit vectors $v \in \mathbb{C}^N$, the test function

$$
f_Q^v := (I + (\epsilon \ell(Q) D_1 B_1)^2)^{-1} (\eta_Q v),
$$

where $\eta_Q = 1$ on $2Q$ and supp $\eta_Q \subset (3Q)$, $\|\nabla \eta_Q\|_{\infty} \lesssim 1/\ell(Q)$. The parameter ϵ is chosen so that the accretivity condition

$$
\operatorname{Re}\left(v,\int_{Q}f_{Q}^{v}dx\right)\geq|Q|/2
$$

holds. This is possible since it is known that we have the estimate

$$
\left|\frac{1}{|Q|}\int_{Q} f_{Q}^{v} dx - v\right| \lesssim \sqrt{\epsilon}.
$$

This can be proved by applying [\[8,](#page-19-3) Lem. 5.6] with the operator D_1 , similar to [8, Lem. 5.10].

Proposition 4.6. We have the estimate

$$
\int_0^\infty \int_{\mathbf{R}^n} |E_t f(x)|^2 |\gamma_t(x)|^2 \frac{dxdt}{t} \lesssim ||f||_2^2, \quad \text{for all } f \in L_2(\mathbf{R}^n; \mathbf{C}^N).
$$

Proof. By Carleson's embedding theorem, it suffices to show that

$$
\int_0^{\ell(Q)} \int_Q |\gamma_t(x)|^2 \frac{dxdt}{t} \lesssim |Q|, \quad \text{for all } Q \in \mathcal{D}.
$$

Following the proof of the Kato square root problem, see for example [\[8,](#page-19-3) Sec. 5.3], we now do (1) a sufficiently fine sectorial decomposition of $\mathcal{L}(\mathbb{C}^N)$, run (2) a stopping time argument to construct an large sawtooth sub region of the Carleson box $Q \times$ $(0, \ell(Q))$ where the test function f_Q^v is paraaccretive, and make (3) a John–Nirenberg bootstrapping argument for Carleson measure, to show that it suffices to prove the estimate

$$
\int_0^{\ell(Q)} \int_Q |\gamma_t(x) E_t f_Q^v|^2 \frac{dxdt}{t} \lesssim |Q|
$$

for all $Q \in \mathcal{D}$ and unit vectors $v \in \mathbb{C}^N$.

We first note that

$$
\|\Theta_t f_Q^v\|_2 = \|Q_t^2(B_1 f_Q^v) - itQ_t^2 R_t^1 B_1 (D_1 B_1)(I + (\epsilon \ell(Q)D_1 B_1)^2)^{-1} (\eta_Q v)\|_2
$$

\$\lesssim \|Q_t^2(B_1 f_Q^v)\|_2 + t/\epsilon \ell(Q)\sqrt{|Q|}\$,

since Q_t^2 and R_t^1 and $\epsilon \ell(Q)D_1B_1(I+(\epsilon \ell(Q)D_1B_1)^2)^{-1}$ are uniformly bounded. Therefore

$$
\int_0^{\ell(Q)} \|\Theta_t f_Q^v\|^2 \frac{dt}{t} \lesssim \|B_1 f_Q^v\|_2^2 + |Q| \lesssim |Q|,
$$

so it suffices to prove

$$
\int_0^{\ell(Q)} \int_Q |(\Theta_t - \gamma_t E_t) f_Q^v|^2 \frac{dxdt}{t} \lesssim |Q|.
$$

Although $f_Q^v \notin \mathcal{H}_1$ in general, we have that $f_Q^v - \eta_Q v = -\epsilon \ell(Q) D_1 B_1 Q_{\epsilon \ell(Q)}^1(\eta_Q v) \in$ $R(D_1)$. Thus Proposition [4.5](#page-14-0) applies to $f = f_Q^v - \eta_Q v$, and it remains to show

$$
\int_0^{\ell(Q)} \int_Q |(\Theta_t - \gamma_t E_t)(\eta_Q v)|^2 \frac{dxdt}{t} \lesssim |Q|.
$$

We have on Q that

$$
(\Theta_t - \gamma_t E_t)(\eta_Q v) = \sum_{k \ge 0} \Theta_t \big(v(\eta_Q - 1)\chi_{A_k(Q)} \big) = \sum_{k \ge 2} \Theta_t \big(v(\eta_Q - 1)\chi_{A_k(Q)} \big),
$$

and Proposition [4.3](#page-13-0) gives

$$
\|(\Theta_t - \gamma_t E_t)(\eta_Q v)\|_{L_2(Q)} \lesssim \sum_{k=2}^{\infty} (t/2^k \ell(Q))^m (2^{nk} |Q|)^{1/2} \lesssim (t/\ell(Q))^m |Q|^{1/2},
$$

if m is chosen large enough. We obtain

$$
\int_0^{\ell(Q)} \int_Q |(\Theta_t - \gamma_t E_t)(\eta_Q v)|^2 \frac{dxdt}{t} \lesssim \int_0^{\ell(Q)} (t/\ell(Q))^{2m} |Q| \frac{dt}{t} \approx |Q|
$$

and the proof is complete.

5. Non-tangential maximal function estimates

In this section, we prove the non-tangential maximal function estimates for the operator T in Theorem [1.1.](#page-2-0) We first consider the operator D_2B_2 , that is the operator T in the special case when $D_1 = 0$. We prove the following result.

Theorem 5.1. We have non-tangential maximal function estimates

 $\|\widetilde{N}_*(\phi(tD_2B_2)f)\|_2 \lesssim \|f\|_2, \qquad f \in \mathcal{H}_2,$

for ϕ as in Theorem [1.1.](#page-2-0)

This result was proved in [\[3,](#page-19-5) Prop. 2.56] for operators DB with D of the form $D =$ $\begin{bmatrix} 0 & \mathrm{div} \end{bmatrix}$ $-\nabla \begin{array}{cc} 0 \\ 0 \end{array}$ 1 , which appear in connection with boundary value problems for divergence form elliptic system. Below we give a simplified proof for general operators of the form DB. Before doing so, we complete the proof of Theorem [1.1](#page-2-0) using Theorem [5.1.](#page-16-1)

Proof of Theorem [1.1.](#page-2-0) Proposition [2.4](#page-5-2) proves that T is closed, densely defined and has the stated estimates of spectrum and resolvents. The square function estimates for $\psi(t)$ follows from Propositions [4.5](#page-14-0) and [4.6,](#page-15-0) using Lemma [4.1.](#page-12-1) Using Proposition [2.5,](#page-6-1) the reverse square function estimates follow by duality.

To prove the non-tangential maximal function estimates, we first note that estimates \geq hold since $\|\phi(tT)f - f\|_2 \to 0$ as $t \to 0$ and

$$
\sup_{t>0} t^{-1} \int_t^{2t} \|\phi(sT)f\|_2^2 ds \lesssim \|\widetilde{N}_*(\phi(tT)f)\|_2^2,
$$

as proved in [\[2,](#page-19-8) Lem. 5.3]. For the estimate \lesssim , it suffices to consider the resolvents $(I + itT)^{-1}$, since

$$
\|\widetilde N_*(\phi(tT)f-(I+i tT)^{-1}f)\|_2^2\lesssim \int_0^\infty \|\phi(tT)f-(I+i tT)^{-1}f\|_2^2\frac{dt}{t}\lesssim \|f\|^2,
$$

where a proof of the first estimate can be found in [\[2,](#page-19-8) Lem. 5.3] and the second estimate follows from square function estimates with $\psi(\lambda) := \phi(\lambda) - (1 + i\lambda)^{-1}$. By Proposition [2.4,](#page-5-2) it remains to prove

(7) $\|\widetilde{N}_*(R_t^2 f)\|_2 \lesssim \|f\|_2, \quad f \in \mathcal{H}_2,$

(8)
$$
\|\widetilde{N}_*(R_t^1f)\|_2 \lesssim \|f\|_2, \quad f \in B_1\mathcal{H}_1,
$$

(9)
$$
\|\widetilde{N}_*(R_t^2 R_t^1 f)\|_2 \lesssim \|f\|_2, \quad f \in B_1 \mathcal{H}_1.
$$

Estimate [\(7\)](#page-16-2) follows from Theorem [5.1.](#page-16-1) For [\(8\)](#page-16-3), we estimate

$$
\|\widetilde{N}_*(R_t^1B_1v)\|_2 = \|\widetilde{N}_*(B_1(I + itD_1B_1)^{-1}v)\|_2 \lesssim \|v\|_2 \lesssim \|B_1v\|_2, \qquad v \in \mathcal{H}_1,
$$

using that B_1 is a bounded multiplication operator and Theorem [5.1,](#page-16-1) with D_2B_2 replaced by D_1B_1 , for the first estimate, and that B_1 is accretive on \mathcal{H}_1 for the second estimate.

To prove [\(9\)](#page-16-4), we introduce the auxiliary non-tangential maximal functions

$$
\widetilde{N}_*^k u(x) := \sup_{t>0} \left(2^{-kn} t^{-(1+n)} \iint_{B(x, 2^k t) \times (t/2, t)} |u(s, y)|^2 ds dy \right)^{1/2}, \qquad k \ge 0.
$$

From L_2 off-diagonal estimates for R_t^2 , as in Proposition [4.3,](#page-13-0) we get that

$$
\widetilde{N}_*^0(R_t^2 R_t^1 f)(x) \lesssim \sum_{k=0}^{\infty} 2^{-km} \widetilde{N}_*^k(R_t^1 f)(x).
$$

We claim that

$$
\|\widetilde{N}^k_*(R^1_tf)\|_2 \lesssim 2^{kn/2} \|\widetilde{N}^0_*(R^1_tf)\|_2.
$$

Thus, choosing m large, we obtain the desired estimates

$$
\|\widetilde{N}_*^0(R_t^2 R_t^1 f)\|_2 \lesssim \|\widetilde{N}_*^0(R_t^1 f)\|_2 \lesssim \|f\|_2.
$$

To prove the claim, assume that $N_*^k g(x) > \lambda$. It follows that there exists $x_1 \in$ $B(x_0, 2^k t)$ such that $t^{-(1+n)} \iint_{B(x_1,t/2)\times(t/2,t)} |g|^2 \gtrsim \lambda^2$. We conclude that for all $y \in$ $B(x_1, t/2)$ we have $\overline{N}_{*}^0 g(y) \gtrsim \lambda$, and therefore

$$
|\{x \; ; \; \widetilde{N}_*^k g(x) > \lambda\}| \le |\{x \; ; \; M(\chi_{\{y \; ; \; \widetilde{N}_*^0 g(y) \gtrsim \lambda\}})(x) \gtrsim 2^{-kn}\}| \lesssim 2^{kn} |\{y \; ; \; \widetilde{N}_*^0 g(y) \gtrsim \lambda\}|,
$$
using the way L , boundedness of the Hardy–L it
lewood maximal function M from

using the weak L_1 boundedness of the Hardy–Littlewood maximal function M , from which the claim follows,. \Box

We now turn to the proof of Theorem [5.1,](#page-16-1) where we write $D := D_2$ and $B := B_2$ to simplify notation. We use the following version of the Caccioppoli estimate for equations of the form $\partial_t v + BDv = 0$.

Lemma 5.2. Let $\eta \in C_0^{\infty}(\mathbf{R}^{1+n})$ and consider a function v solving $\partial_t v + BDv = 0$ on a neighbourhood of supp η. Then

$$
\int \int |Dv(t,x)|^2 |\eta(t,x)|^2 dt dx \lesssim \int \int |v(t,x)|^2 |\nabla \eta(t,x)|^2 dt dx.
$$

Proof. Integrating by part twice, using anti-symmetry of ∂_t and symmetry of D, we obtain

$$
\iint (BDv, Dv)\eta^2 = -\iint (\partial_t v, Dv)\eta^2
$$

=
$$
\iint (v, \partial_t Dv)\eta^2 + 2\iint (v, Dv)\eta \partial_t \eta = -\iint (v, DBDv)\eta^2 + 2\iint (v, Dv)\eta \partial_t \eta
$$

=
$$
-\iint (D(\eta^2 v), BDv) + 2\iint (v, Dv)\eta \partial_t \eta.
$$

We get the estimate

$$
\operatorname{Re} \iint (BDv, Dv) \eta^2 \lesssim \iint |v| |Dv| |\eta| |\nabla \eta|.
$$

For each $t \in \mathbf{R}$, we have by the accretivity of B that

$$
\int |D(\eta v)|^2 dx \lesssim \text{Re} \int (BD(\eta v), D(\eta v)) dx.
$$

Integrating both sides with respect to t and using the product rule for the derivatives, we get

$$
\int \int |Dv|^2 \eta^2 dt dx \lesssim \int \int (|Dv|\eta)(|v||\nabla \eta|) dt dx + \int \int |v|^2 |\nabla \eta|^2 dt dx.
$$

Using the absorption inequality, we obtain the stated estimate. \Box

Proof of Theorem [5.1.](#page-16-1) To prove the non-tangential maximal function estimates, it suffices to estimate the semigroups $e^{-tDB}\chi^+(DB)$ and $e^{tDB}\chi^-(DB)$, since

$$
\begin{aligned} \|\widetilde{N}_*(\phi(tDB)f - e^{-tDB}\chi^+(DB)f - e^{tDB}\chi^-(DB)f)\|_2^2 \\ &\lesssim \int_0^\infty \|\phi(tT)f - e^{-tDB}\chi^+(DB)f - e^{tDB}\chi^-(DB)f\|_2^2 \frac{dt}{t} \lesssim \|f\|^2, \end{aligned}
$$

where the second estimate follows from square function estimates for DB with $\psi(\lambda) := \phi(\lambda) - e^{-|\lambda|}$. Moreover, by a limiting argument, we may assume that $f \in R(D)$.

Consider first $e^{-tDB}\chi^+(DB)f$ and a Whitney region $W = B(x_0, t_0) \times (t_0/2, t_0)$. Write $\chi^+(DB)f = Dv$ with $v \in \chi^+(BD)L_2 \cap D(D)$. Let P be the orthogonal projection onto $\overline{R(D)}$, and denote by $[Pv] := |B(x_0, t_0)|^{-1} \int_{B(x_0, t_0)} Pv(x) dx$ the average of $Pv \in \overline{R(D)}$. Note that $Dv = D(Pv)$ since P projects along the null space N(D). Define the function

$$
w(t,x) := t\psi_1(tBD)BDv(x) + \phi_1(tBD)(Pv - [Pv])(x), \qquad (t,x) \in \mathbf{R}^{1+n}_+,
$$

where

$$
\psi_1(\lambda) := \lambda^{-1} \big(e^{-\lambda} - (1 - \lambda) / (1 + \lambda^2) \big) \chi^+(\lambda),
$$

$$
\phi_1(\lambda) := (1 - \lambda) / (1 + \lambda^2).
$$

In the second term, $[Pv]$ is regarded as a constant function. Since $\phi_1(tBD)$ is a linear combination of resolvents, it extends to a bounded operator $L_{\infty}(\mathbf{R}^n) \to L_2(B(x_0, t_0))$ as in the estimate [\(6\)](#page-13-1). For the first term, the decay of ψ_1 at $\lambda = 0$ and ∞ shows that we have square function estimates for $\psi_1(tBD)$. We calculate for $t > 0$ that

$$
Dw = De^{-tBD}v = e^{-tDB}\chi^+(DB)f,
$$

\n
$$
\partial_t w = -BDe^{-tBD}v,
$$

using $D\phi_1(tBD)[Pv] = \partial_t \phi_1(tBD)[Pv] = 0$. Since $\partial_t w + BDw = 0$, we get from Lemma [5.2](#page-17-0) that

$$
\iint_W |e^{-tDB}\chi^+(DB)f|^2 dt dx = \iint_W |Dw|^2 dt dx \lesssim \iint_{\widetilde{W}} |t^{-1}w|^2 dt dx,
$$

with a slightly enlarged Whitney region $\widetilde{W} = B(x_0, 2t_0) \times (t_0/4, 2t_0)$. Using square function estimates, we obtain

$$
\|\widetilde{N}_*(\psi_1(tBD)BDv)\|_2^2 \lesssim \int_0^\infty \|\psi_1(tBD)BDv\|_2^2 \frac{dt}{t} \lesssim \|BDv\|_2^2 \lesssim \|f\|_2^2,
$$

so it remains to estimate $\phi_1(tBD)(Pv - [Pv])$. To this end, for fixed $t_0/4 < t < 2t_0$, use the L_2 off-diagonal estimates for the resolvents of BD to get

$$
\|\phi_1(tBD)(Pv - [Pv])\|_{L_2(B(x_0, 2t_0))}^2 \lesssim \sum_{k=0}^{\infty} 2^{-km} \|Pv - [Pv]\|_{L_2(B(x_0, 2^k 2t_0))}^2
$$

$$
\lesssim \sum_{k=0}^{\infty} 2^{k(-m+3n-2n/p+2)} t_0^{n-2n/p+2} \|\nabla v\|_{L_p(B(x_0, 2^k 2t_0))}^2,
$$

using the Sobolev–Poincaré inequality from Lemma [4.4](#page-13-2) with suitable $1 \leq p < 2 = q$, $\Omega = B(x_0, 2^k 2t_0)$ and $S = B(x_0, 2t_0)$. Integrating with respect to t, we get

$$
t_0^{-1-n} \iint_{\widetilde{W}} |t^{-1}\phi_1(tBD)(Pv - [Pv])|^2 dt dx
$$

$$
\lesssim \sum_{k=0}^{\infty} 2^{k(-m+3n+2)} M(|\nabla Pv|^p)(x_0)^{2/p} \lesssim M(|\nabla Pv|^p)(x_0)^{2/p},
$$

choosing the parameter m large. Therefore, boundedness of the Hardy–Littlewood maximal function M on $L_{2/p}$ gives

$$
\|\widetilde{N}_*(e^{-tDB}\chi^+(DB)f)\|_2^2 \lesssim \|f\|_2^2 + \|\nabla Pv\|_2^2 \lesssim \|f\|_2^2 + \|DPv\|_2^2 \lesssim \|f\|_2^2,
$$

by the ellipticity of D on $R(D)$.

The estimate of the semigroup $e^{tDB}\chi^{-}(DB)$ follows from the above estimate upon replacing D by $-D$, since $e^{-t(-DB)}\chi^+(-DB) = e^{tDB}\chi^-(DB)$. This proves the nontangential maximal function estimate for $\phi(tDB)$.

REFERENCES

- [1] Albrecht, D., Duong, X., and McIntosh, A. Operator theory and harmonic analysis. In Instructional Workshop on Analysis and Geometry, Part III (Canberra, 1995), vol. 34 of Proc. Centre Math. Appl. Austral. Nat. Univ. Austral. Nat. Univ., Canberra, 1996, pp. 77–136.
- [2] Auscher, P., and Axelsson, A. Weighted maximal regularity estimates and solvability of non-smooth elliptic systems I. Invent. Math. 184, 1 (2011), 47–115.
- [3] AUSCHER, P., AXELSSON, A., AND HOFMANN, S. Functional calculus of Dirac operators and complex perturbations of Neumann and Dirichlet problems. J. Funct. Anal. 255, 2 (2008), 374–448.
- [4] Auscher, P., Axelsson, A., and McIntosh, A. On a quadratic estimate related to the Kato conjecture and boundary value problems. Contemp. Math. 205 (2010), 105–129.
- [5] Auscher, P., Axelsson, A., and McIntosh, A. Solvability of elliptic systems with square integrable boundary data. Ark. Mat. 48 (2010), 253-287.
- [6] AUSCHER, P., HOFMANN, S., LACEY, M., MCINTOSH, A., AND TCHAMITCHIAN, P. The solution of the Kato square root problem for second order elliptic operators on \mathbb{R}^n . Ann. of Math. (2) 156, 2 (2002), 633–654.
- [7] Auscher, P., McIntosh, A., and Nahmod, A. The square root problem of Kato in one dimension, and first order elliptic systems. Indiana Univ. Math. J. 46, 3 (1997), 659–695.
- [8] AXELSSON, A., KEITH, S., AND MCINTOSH, A. Quadratic estimates and functional calculi of perturbed Dirac operators. Invent. Math. 163, 3 (2006), 455–497.
- [9] COIFMAN, R. R., MCINTOSH, A., AND MEYER, Y. L'intégrale de Cauchy définit un opérateur borné sur L^2 pour les courbes lipschitziennes. Ann. of Math. (2) 116, 2 (1982), 361–387.
- [10] GILBARG, D., AND TRUDINGER, N. Elliptic partial differential equations of second order. No. 224 in Grundlehren der Mathematischen Wissenschaften. Springer-Verlag, Berlin, 1983.

ANDREAS ROSÉN, MATEMATISKA INSTITUTIONEN, LINKÖPINGS UNIVERSITET, 581 83 LINKÖPING, **SWEDEN**

E-mail address: andreas.rosen@liu.se