

Multiplication of conjugacy classes, colligations, and characteristic functions of matrix argument

YURY A. NERETIN¹

We extend the classical construction of operator colligations and characteristic functions. Consider the group G of finitary block unitary matrices of size $\alpha + \infty + \dots + \infty$ (m times). Consider the subgroup $K = U(\infty)$, which consists of block diagonal unitary matrices with a block 1 of size α and a matrix $u \in U(\infty)$ repeated m times. It appears that there is a natural multiplication on the conjugacy classes $G//K$. We construct 'spectral data' of conjugacy classes, which visualize the multiplication and are sufficient for a reconstruction of a conjugacy class.

MSC 22E66, 47A48, 15A72, 16W22

1 Formulation of the problem

1.1. Notation. By $1 = 1_n$ we denote the unit matrix of order n , by A^t the transposed matrix. Let $\text{Mat}(n)$ be the space of complex matrices of size $n \times n$, $\text{GL}(n, \mathbb{C})$ be the group of invertible matrices of order n , $U(n)$ be the unitary subgroup in $\text{GL}(n, \mathbb{C})$.

Denote by $\text{MAT}(\infty)$ the space of all infinite matrices g such that $g - 1$ has only finite number of non-zero elements. By $\text{GL}(\infty, \mathbb{C})$, we denote the group of invertible elements of $\text{MAT}(\infty)$. By $U(\infty)$ we denote its subgroup consisting of unitary matrices.

Denote $\text{Mat}(\infty)$ the space of all infinite matrices having only finite number of non-zero matrix elements, i.e. $g \in \text{Mat}(\infty) \Leftrightarrow g + 1 \in \text{MAT}(\infty)$.

Let G be a group, $H \subset G$ a subgroup. By $G//H$ we denote the space of conjugacy classes G with respect to H , by $H \backslash G/H$ the double cosets.

1.2. Colligations. The following objects arose by independent reasons in spectral theory of non-selfadjoint operators, system theory, and representation theory of infinite-dimensional groups (see [12], [13], [29], [2], [3], [4], [7], [23], [24], [15]).

Fix a non-negative integer α . Denote by G the group $U(\infty)$, by K its subgroup consisting of matrices of the form $\begin{pmatrix} 1_\alpha & 0 \\ 0 & h \end{pmatrix}$. Consider the space $G//K$ of conjugacy classes of G with respect to K , i.e., elements of G defined up to the equivalence

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \sim \begin{pmatrix} 1_\alpha & 0 \\ 0 & h \end{pmatrix} \begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} 1_\alpha & 0 \\ 0 & h \end{pmatrix}^{-1}.$$

There is a natural multiplication on the set $G//K$ defined in the following way. Let $\mathfrak{g}, \mathfrak{h}$ be conjugacy classes, let g, h be their representatives, let g, h actually

¹Supported by the grants FWF, P22122, P25142.

be contained in $U(\alpha + N)$, i.e., they are given by block $(\alpha + N + \infty)$ -matrices of the form

$$g = \begin{pmatrix} a_1 & b_1 & 0 \\ c_1 & d_1 & 0 \\ 0 & 0 & 1_\infty \end{pmatrix}, \quad g = \begin{pmatrix} a_2 & b_2 & 0 \\ c_2 & d_2 & 0 \\ 0 & 0 & 1_\infty \end{pmatrix}.$$

We define the product $\mathfrak{g} \circ \mathfrak{h}$ as the conjugacy class containing

$$\begin{aligned} & \begin{pmatrix} a_1 & b_1 & 0 \\ c_1 & d_1 & 0 \\ 0 & 0 & 1_\infty \end{pmatrix} \circ \begin{pmatrix} a_2 & b_2 & 0 \\ c_2 & d_2 & 0 \\ 0 & 0 & 1_\infty \end{pmatrix} := \\ & := \begin{pmatrix} a_1 & b_1 & 0 & 0 \\ c_1 & d_1 & 0 & 0 \\ 0 & 0 & 1_N & 0 \\ 0 & 0 & 0 & 1_\infty \end{pmatrix} \begin{pmatrix} a_2 & 0 & b_2 & 0 \\ 0 & 1_N & 0 & 0 \\ c_2 & 0 & d_2 & 0 \\ 0 & 0 & 0 & 1_\infty \end{pmatrix} = \begin{pmatrix} a_1 a_2 & b_1 & a_1 b_2 & 0 \\ c_1 a_2 & d_1 & c_1 b_2 & 0 \\ c_2 & 0 & d_2 & 0 \\ 0 & 0 & 0 & 1_\infty \end{pmatrix}. \end{aligned} \tag{1.1}$$

This product is a well-defined associative operation on $G//K$.

This operation admits the following visualization: To a matrix $g = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ we assign the *characteristic function* as a function on \mathbb{C} taking values in $\alpha \times \alpha$ -matrices given by

$$\chi(z) = a + zb(1 - zd)^{-1}c.$$

This function depends only on a conjugacy class \mathfrak{g} and not on a matrix g , so we can write $\chi_{\mathfrak{g}}(z)$.

Next, denote by $\eta_{\mathfrak{g}}$ the set of eigenvalues of d that are contained in the circle $|z| = 1$.

The following statement is well-known (see, e.g., [3], in this work the complete unitary group instead of $U(\infty)$ is considered).

Theorem 1.1 a) $\chi_{\mathfrak{g} \circ \mathfrak{h}}(z) = \chi_{\mathfrak{g}}(z)\chi_{\mathfrak{h}}(z)$.

b) If $|z| \leq 1$, then $\|\chi(z)\| \leq 1$; if $|z| = 1$, then $\chi(z) \in U(n)$.

c) A conjugacy class \mathfrak{g} is uniquely determined by the characteristic function $\chi_{\mathfrak{g}}(z)$ and the set $\eta_{\mathfrak{g}}$.

A similar object is a semigroup of conjugacy classes of $Q := GL(\infty, \mathbb{C})$ by a subgroup H consisting of matrices $\begin{pmatrix} 1_\alpha & 0 \\ 0 & h \end{pmatrix}$, for a further discussion see [4], [9], [14].

The \circ -multiplication is a representative of a wide class of operations on sets of conjugacy classes and double cosets. Such operations arise in a natural way in representation theory of infinite-dimensional classical groups, see [16], [17], [18]. The purpose of this paper is to give a way for a visualization of such operations. We consider a minimal example but our receipt works in the wider generality of [17], we explain this in Section 4.

1.3. Product of conjugacy classes. Fix integers $\alpha \geq 0$ and $m \geq 1$. Let $N > 0$. Consider the space $\text{Mat}(\alpha + mN)$. We write its elements as block matrices

$$g = \begin{pmatrix} a & b_1 & \dots & b_m \\ c_1 & d_{11} & \dots & d_{1m} \\ \vdots & \vdots & \ddots & \vdots \\ c_m & d_{m1} & \dots & d_{mm} \end{pmatrix} \quad (1.2)$$

of size $\alpha + N + \dots + N$. For an element $u \in \text{GL}(N, \mathbb{C})$ we denote by $\iota(u)$ the following matrix

$$\iota(u) = \begin{pmatrix} 1_\alpha & 0 & 0 & \dots & 0 \\ 0 & u & 0 & \dots & 0 \\ 0 & 0 & u & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \dots & u \end{pmatrix}. \quad (1.3)$$

Denote by

$$\mathcal{M}_N = \mathcal{M}_N^{\alpha, m} := \text{Mat}(\alpha + mN) // \text{GL}(N, \mathbb{C})$$

the set of conjugacy classes of $\text{Mat}(\alpha + mN)$ by $\text{GL}(N, \mathbb{C})$, i.e.,

$$g \sim \iota(u)g\iota(u)^{-1}. \quad (1.4)$$

We call elements of this set by *colligations*.

Usually, we omit superscripts α, m from $\mathcal{M}_N^{\alpha, m}$.

Next, we formulate the definition of a colligation in another form. Consider the space $V = \mathbb{C}^m$ and $Z_N = \mathbb{C}^N$. Then an operator g can be regarded as an operator

$$\mathbb{C}^\alpha \oplus (V \otimes Z_N) \rightarrow \mathbb{C}^\alpha \oplus (V \otimes Z_N)$$

defined up to a conjugation by an element of $\text{GL}(N, \mathbb{C})$ acting in Z_N .

We wish to define a canonical multiplication

$$\mathcal{M}_{N_1} \times \mathcal{M}_{N_2} \rightarrow \mathcal{M}_{N_1+N_2} \quad (1.5)$$

(α, m are fixed). First, we consider the case $m = 2$. The multiplication is given

by

$$\begin{aligned}
g \circ h &= \begin{pmatrix} a & b_1 & b_2 \\ c_1 & d_{11} & d_{12} \\ c_2 & d_{21} & d_{22} \end{pmatrix} \circ \begin{pmatrix} p & q_1 & q_2 \\ c_1 & d_{11} & d_{12} \\ c_2 & d_{21} & d_{22} \end{pmatrix} = \\
&= \begin{pmatrix} a & b_1 & 0 & b_2 & 0 \\ c_1 & d_{11} & 0 & d_{12} & 0 \\ 0 & 0 & 1_{N_2} & 0 & 0 \\ c_2 & d_{21} & 0 & d_{22} & 0 \\ 0 & 0 & 0 & 0 & 1_{N_2} \end{pmatrix} \begin{pmatrix} p & 0 & q_1 & 0 & q_2 \\ 0 & 1_{N_2} & 0 & 0 & 0 \\ c_1 & 0 & d_{11} & 0 & d_{12} \\ 0 & 0 & 0 & 1_{N_2} & 0 \\ c_2 & 0 & d_{21} & 0 & d_{22} \end{pmatrix} = \\
&= \begin{pmatrix} aa' & | & b_1 & ab'_1 & & b_1 & ab'_1 \\ - & + & - & - & - & - & - \\ c_1 a' & | & d_{11} & c_1 b'_1 & & d_{12} & c_1 b'_2 \\ c'_1 & | & 0 & d'_{11} & & 0 & d'_{12} \\ \vdots & & & & & & \\ c_2 a' & | & d_{21} & c_2 b'_1 & & d_{22} & c_2 b'_2 \\ c'_2 & | & 0 & d'_{21} & & 0 & d'_{22} \end{pmatrix}
\end{aligned}$$

and we take the conjugacy class containing the latter matrix.

Pass from $m = 2$ to an arbitrary m is evident. However we formulate a formal definition, which is valid for all m . We extend the operator g to an operator \tilde{g} in

$$\mathbb{C}^\alpha \oplus (V \otimes Z_{N_1+N_2}) = \left[\mathbb{C}^\alpha \oplus (V \otimes Z_{N_1}) \right] \oplus \left[V \otimes Z_{N_2} \right]$$

This operator acts as g on the first summand and as 1 on the second summand. Next, we extend h to an operator \hat{h} acting in the same way in the space

$$\mathbb{C}^\alpha \oplus (V \otimes Z_{N_1+N_2}) = \left[\mathbb{C}^\alpha \oplus (V \otimes Z_{N_2}) \right] \oplus \left[V \otimes Z_{N_1} \right],$$

and set

$$g \circ h := \tilde{g} \cdot \hat{h}.$$

Proposition 1.2 *The \circ -multiplication is associative, i.e., for any N_1, N_2, N_3 and $g_1 \in \mathcal{M}_{N_1}, g_2 \in \mathcal{M}_{N_2}, g_3 \in \mathcal{M}_{N_3}$ the following equality holds:*

$$(g_1 \circ g_2) \circ g_3 = g_1 \circ (g_2 \circ g_3).$$

This is more or less obvious (see also [16]).

1.4. Infinite-dimensional version. Denote by \mathcal{M}_∞ the set of conjugacy classes on $\text{MAT}(\infty)$ with respect to $\text{GL}(\infty, \mathbb{C})$. Then \mathcal{M}_∞ is a semigroup with respect to \circ -multiplication.

1.5. Variants. Denote

$$\begin{aligned}
\mathcal{GL}_N &= \mathcal{GL}_N^{\alpha, m} = \text{GL}(\alpha + mN, \mathbb{C}) // \text{GL}(N, \mathbb{C}) \\
\mathcal{U}_N &= \mathcal{U}_N^{\alpha, m} = \text{U}(\alpha + mN) // \text{U}(N),
\end{aligned}$$

in all cases we consider the equivalence (1.4), in the unitary case we suppose that u is unitary. Thus we get the operations

$$\begin{aligned}\mathcal{GL}_{N_1}^{\alpha,m} \times \mathcal{GL}_{N_2}^{\alpha,m} &\rightarrow \mathcal{GL}_{N_1+N_2}^{\alpha,m}; \\ \mathcal{GL}_{\infty}^{\alpha,m} \times \mathcal{GL}_{\infty}^{\alpha,m} &\rightarrow \mathcal{GL}_{\infty}^{\alpha,m}; \\ \mathcal{U}_{N_1}^{\alpha,m} \times \mathcal{U}_{N_2}^{\alpha,m} &\rightarrow \mathcal{U}_{N_1+N_2}^{\alpha,m}; \\ \mathcal{U}_{\infty}^{\alpha,m} \times \mathcal{U}_{\infty}^{\alpha,m} &\rightarrow \mathcal{U}_{\infty}^{\alpha,m}.\end{aligned}$$

In this section α, m are fixed and we will omit upper superscripts.

1.6. The origin of \circ -multiplication. The spaces $\mathcal{M}_{\infty}^{\alpha,1}$ and $\mathcal{U}_{\infty}^{\alpha,1}$ were discussed above in Subsection 1.2. In 80s a family of operations of this type arose in a natural way in representation theory of infinite-dimensional classical and infinite symmetric groups, see [22], [23], [15], Section IX.4. For instance, there is a semigroup structure on double cosets

$$\lim_{n \rightarrow \infty} \mathrm{O}(n) \backslash \mathrm{U}(\alpha + n) / \mathrm{O}(n).$$

Moreover, this semigroup acts in the space of $\mathrm{O}(\infty)$ -fixed vectors of unitary representations of $\mathrm{U}(\alpha + \infty)$. Big zoo of operations on $K \backslash G / K$ (with 'small' subgroups K) arose in [16], [17], [19], [5], [21]; our $\mathrm{U}(\alpha + m\infty) // \mathrm{U}(\infty)$ is inside this zoo. For another explanation, see [18] (the \circ -product is an infinite-dimensional limit of convolutions of δ -measures on conjugacy classes).

1.7. The purpose of the work. In [16] there was obtained an analog of the characteristic function for the semigroup $\mathcal{U}_{\infty}^{\alpha,m}$ (see also p -adic case in [20]). The purpose of this work is to obtain spectral data of matrices visualizing the \circ -multiplications and separating points of $\mathcal{U}_{\infty}^{\alpha,m}$ and of $\mathcal{GL}_{\infty}^{\alpha,m}$. In fact, for each conjugacy class we assign a rational functions $\chi^{(j)} : \mathrm{Mat}(jm) \rightarrow \mathrm{Mat}(j\alpha)$ and a divisors $\xi^{(j)}$ in $\mathrm{Mat}(jm)$. The main statements are Theorems 2.10, 2.11.

Notice that the spectral data for collections of matrices were widely discussed, see different approaches in surveys [1], [10]. It seems that our approach produces another kind of spectral data. On the other hand, it extends classical approach of operator theory existing for $m = 1$.

1.8. Structure of the paper. Section 2 contains a construction of 'spectral data', and main Theorems 2.10, 2.11. Proofs are contained in Section 3. Some extensions of our construction are discussed in Section 4.

2 Spectral data

2.1. Categorical quotient. Notice that the spaces \mathcal{M}_N are non-Hausdorff. There are many ways to construct Hausdorff spaces from set-theoretical quotients. We will use the following approach (see, e.g., [25]).

Let a reductive group G act on an affine algebraic variety X . Consider the algebra $\mathbb{C}[X]^G$ of G -invariant regular functions on X . The *categorical quotient* $[X/G]$ is the set of maximal ideals of $\mathbb{C}[X]^G$.

We consider categorical quotients $[\mathcal{M}_N]$ of $\text{Mat}(\alpha + mN)$ by $\text{GL}(N, \mathbb{C})$ and $[\mathcal{GL}_N]$ of $\text{GL}(\alpha + mN, \mathbb{C})$ by $\text{GL}(N, \mathbb{C})$.

Notice that regular functions on $\text{Mat}(\alpha + mN)$ are polynomials. The algebra of regular functions on $\text{GL}(\alpha + mN, \mathbb{C})$ is generated by polynomials and $\det(g)^{-1}$.

Proposition 2.1 a) *The natural map $[\mathcal{M}_N] \rightarrow [\mathcal{M}_{N+1}]$ is injective.*

b) *The natural map $[\mathcal{GL}_N] \rightarrow [\mathcal{GL}_{N+1}]$ is injective, and $[\mathcal{GL}_N] = [\mathcal{M}_N] \setminus \{g : \det g = 0\}$.*

c) *The natural map $\mathcal{U}_N \rightarrow \mathcal{U}_{N+1}$ is injective.*

The statement is proved in Subsection 3.4.

This allows to define the infinite-dimensional 'categorical quotient' $[\mathcal{M}_\infty]$ as the inductive limit. On the other hand this shows that the topological quotient \mathcal{U}_∞ is Hausdorff.

Proposition 2.2 *The \circ -multiplication determines a map of categorical quotients,*

$$[\mathcal{M}_{N_1}^{\alpha, m}] \times [\mathcal{M}_{N_2}^{\alpha, m}] \rightarrow [\mathcal{M}_{N_1+N_2}^{\alpha, m}].$$

The statement is proved in Subsection 3.5.

2.2. Characteristic functions. Fix $g \in \text{Mat}(\alpha + mN)$ given by (1.2). Let S range in $\text{Mat}(m)$. We write the following equation:

$$\begin{pmatrix} q \\ x_1 \\ \vdots \\ x_m \end{pmatrix} = \begin{pmatrix} a & b_1 & \dots & b_m \\ c_1 & d_{11} & \dots & d_{1m} \\ \vdots & \vdots & \ddots & \vdots \\ c_m & d_{m1} & \dots & d_{mm} \end{pmatrix} \begin{pmatrix} p \\ s_{11}x_1 + \dots + s_{1m}x_m \\ \vdots \\ s_{m1}x_1 + \dots + s_{mm}x_m \end{pmatrix}. \quad (2.1)$$

Eliminating variables x_1, \dots, x_m , we get a relation

$$q = \chi_g(S)p,$$

where $\chi_g(S)$ is a rational matrix-valued function, $\text{Mat}(m) \rightarrow \text{Mat}(\alpha)$. We call χ_g by *the characteristic function* of g .

The following statements were obtained in [16].

Proposition 2.3 *The characteristic function of g depends only on a $\text{GL}(N, \mathbb{C})$ -conjugacy class of g (and not on g itself).*

Proposition 2.4 *Consider the natural map $I_N : \mathcal{M}_N \rightarrow \mathcal{M}_{N+1}$. Then*

$$\chi_{I_N g}(S) = \chi_g(S).$$

This is obvious.

Theorem 2.5

$$\chi_{g \circ h}(S) = \chi_g(S)\chi_h(S).$$

Theorem 2.6 *Let g be unitary. Then*

- a) *If $\|S\| \leq 1$, then $\|\chi_g(S)\| \leq 1$.*
- b) *If S is unitary, then $\chi_g(S)$ is unitary.*

REMARK. The space B_n of $n \times n$ matrices with norm < 1 is a well-known object, it is an Hermitian symmetric space $U(n, n)/U(n) \times U(n)$, also they are called homogeneous Cartan domains of the first kind. The unitary group $U(n)$ is a Shilov boundary of this domain. Characteristic functions are holomorphic maps $B_m \rightarrow B_\alpha$, which send the Shilov boundary to the Shilov boundary. For $m = 1$ such functions ('inner functions') are well-known topic of function theory, see [6], [26]. \diamond

Denoting $\tilde{S} := S \otimes 1_N$, we represent (2.1) as

$$\begin{aligned} q &= ap + b\tilde{S}x \\ x &= cp + d\tilde{S}x \end{aligned}$$

Therefore

$$x = (1 - d\tilde{S})^{-1}cp$$

and

$$\chi_g(S) = a + b\tilde{S}(1 - d\tilde{S})^{-1}c. \quad (2.2)$$

2.3. The language of Grassmannians. Now we reformulate the definition of characteristic function. For a linear space W denote by $\text{Gr}(W)$ the set of all subspaces in W . For an even-dimensional space denote by $\text{Gr}^{1/2}(W)$ the set of subspaces having dimension $\frac{1}{2} \dim W$.

Consider the Grassmannian $\text{Gr}^{1/2}(V \oplus V)$ in the linear space $V \oplus V$. For any $L \in \text{Gr}^{1/2}(V \oplus V)$ consider the subspace

$$L \otimes Z_N \subset (V \oplus V) \otimes Z_N = (V \otimes Z_N) \oplus (V \otimes Z_N)$$

We write the equation

$$\begin{pmatrix} q \\ x_1 \\ \vdots \\ x_m \end{pmatrix} = \begin{pmatrix} a & b_1 & \dots & b_m \\ c_1 & d_{11} & \dots & d_{1m} \\ \vdots & \vdots & \ddots & \vdots \\ c_m & d_{m1} & \dots & d_{mm} \end{pmatrix} \begin{pmatrix} p \\ y_1 \\ \vdots \\ y_m \end{pmatrix}. \quad (2.3)$$

Consider the set $\mathcal{X}_g(L)$ of all (q, p) such that there are $(x, y) \in L \otimes Z_N$ satisfying (2.3). The set $\mathcal{X}_g(L)$ is a subspace in $\mathbb{C}^\alpha \oplus \mathbb{C}^\alpha$. The following statement is straightforward.

Proposition 2.7 *Let L be a graph of an operator $S : V \rightarrow V$. Assume that S is a non-singular point of χ_g . Then $\mathcal{X}_g(L)$ is the graph of the operator $\chi_g(S) : \mathbb{C}^\alpha \rightarrow \mathbb{C}^\alpha$.*

Thus \mathcal{X}_g is a rational map $\text{Gr}^{1/2}(V \oplus V) \rightarrow \text{Gr}^{1/2}(\mathbb{C}^\alpha \oplus \mathbb{C}^\alpha)$.

2.4. The distinguished divisor. Generally, the characteristic function does not determine an element of \mathcal{U}_N (and, also of the space $[\mathcal{M}_N]$). Indeed, consider the following matrix of size $\alpha + (k+l) + (k+l)$:

$$\begin{pmatrix} a & b_1 & 0 & b_2 & 0 \\ c_1 & d_{11} & 0 & d_{12} & 0 \\ 0 & 0 & e_{11} & 0 & e_{12} \\ c_2 & d_{21} & 0 & d_{22} & 0 \\ 0 & 0 & e_{21} & 0 & e_{22} \end{pmatrix} \in \mathcal{U}_{k+l}^{\alpha,2}.$$

Then its characteristic functions does not depend on the matrix $\begin{pmatrix} e_{11} & e_{12} \\ e_{21} & e_{22} \end{pmatrix}$.

Therefore additional invariants are necessary.

Let $g \in \text{Mat}(\alpha + mN)$. We write the equation

$$\begin{pmatrix} x_1 \\ \vdots \\ x_m \end{pmatrix} = \begin{pmatrix} d_{11} & \dots & d_{1m} \\ \vdots & \ddots & \vdots \\ d_{m1} & \dots & d_{mm} \end{pmatrix} \begin{pmatrix} s_{11}x_1 + \dots + s_{1m}x_m \\ \vdots \\ s_{m1}x_1 + \dots + s_{mm}x_m \end{pmatrix} \quad (2.4)$$

and consider the set ξ_g of all S such that this equation has a nonzero solution.

Next, we reformulate the definition in a more precise form. Consider the polynomial p_g on $\text{Mat}(m)$ determined by the equation

$$\begin{aligned} p_g(S) &= \det(1 - d\tilde{S}) = \\ &= \det \left(1_{mN} - \begin{pmatrix} d_{11} & \dots & d_{1m} \\ \vdots & \ddots & \vdots \\ d_{m1} & \dots & d_{mm} \end{pmatrix} \begin{pmatrix} s_{11} \cdot 1_N & \dots & s_{1m} \cdot 1_N \\ \vdots & \ddots & \vdots \\ s_{m1} \cdot 1_N & \dots & s_{mm} \cdot 1_N \end{pmatrix} \right) \end{aligned}$$

The set ξ_g is the set of zeros of the polynomial p_g . It is more natural to consider the set ξ_g as a *divisor* (see, e.g. [8]), i.e. we decompose $p_g(S)$ as a product of irreducible factors

$$p_g(S) = \prod_i h_i(S)^{v_i},$$

where h_i are pairwise distinct. Then we consider the collection of hypersurfaces² $h_i(S) = 0$ with assigned multiplicities v_i .

Evidently,

$$p_{g \circ h}(S) = p_g(S) p_h(S).$$

Equivalently,

$$\xi_{g \circ h} = \xi_g + \xi_h.$$

The sign '+' means that we consider the union of hypersurfaces taking in accounts their multiplicities.

²of complex codimension 1.

Proposition 2.8 *The following identity holds*

$$\det \chi_g(S) = \frac{\det \begin{pmatrix} a & -b\tilde{S} \\ c & 1 - d\tilde{S} \end{pmatrix}}{\det(1 - d\tilde{S})}. \quad (2.5)$$

In particular, the denominator of the rational function $\det \chi_g(S)$ is a divisor of $p_g = \det(1 - \tilde{S})$.

PROOF. Applying the formula for the determinant of a block matrix

$$\det \begin{pmatrix} A & B \\ C & D \end{pmatrix} = \det(D) \det(A - BD^{-1}C),$$

we get

$$\det \begin{pmatrix} a & -b\tilde{S} \\ c & 1 - d\tilde{S} \end{pmatrix} = \det(1 - d\tilde{S}) \cdot \det[a + b\tilde{S}(1 - d\tilde{S})^{-1}c].$$

By (2.2), the expression in the square brackets is the characteristic function. \square

Let us pass to the language of Grassmannians. Consider another matrix coordinate $\Lambda = S^{-1}$ on Grassmannian. Then the equation for the divisor passes to the form

$$\det(d - \Lambda) = \left(\begin{pmatrix} d_{11} & \dots & d_{1m} \\ \vdots & \ddots & \vdots \\ d_{m1} & \dots & d_{mm} \end{pmatrix} - \begin{pmatrix} \lambda_{11} \cdot 1_N & \dots & \lambda_{1m} \cdot 1_N \\ \vdots & \ddots & \vdots \\ \lambda_{m1} \cdot 1_N & \dots & \lambda_{mm} \cdot 1_N \end{pmatrix} \right) = 0.$$

Two equations

$$\det(1 - dS) = 0, \quad \det(d - \Lambda) = 0$$

determine a divisor Ξ_g in $\text{Gr}^{1/2}(V \oplus V)$. Indeed, we have two charts in $\text{Gr}^{1/2}(V \oplus V)$, one consists of graphs of operators $V \oplus 0 \rightarrow 0 \oplus V$, another from graphs of operators $0 \oplus V \rightarrow V \oplus 0$. These charts do not cover the whole Grassmannian, but the complement of their union has codimension 2. Therefore any hypersurface in $\text{Gr}^{1/2}(V \oplus V)$ has an intersection with at least one of the charts. In fact, all hypersurfaces in $\text{Gr}^{1/2}(V \oplus V)$ are observable in the chart S , except the hypersurface $\det \Lambda = 0$ (it is a complement to the chart S).

The degree of the polynomial p_g for a generic g is mN . If the divisor Ξ_g contains the component $\det \Lambda = 0$ with multiplicity l , then the degree of p_g is $m(N - l)$.

2.5. Characteristic functions for \mathcal{M}_∞ . We have a natural embedding

$$I_N : \mathcal{M}_N \rightarrow \mathcal{M}_{N+1}$$

induced by the embedding $Z_N \rightarrow Z_{N+1}$. Evidently,

$$\chi_{I_N g}(S) = \chi_g(S).$$

Therefore, we can define a characteristic function for elements of $\mathcal{M}_\infty = \cup_N \mathcal{M}_N$. On the other hand, we can repeat the definition of characteristic function for $N = \infty$, this produces the same result.

Next,

$$p_{I_N g}(S) = p_g(S) \cdot \det(1 - S).$$

Denoting by δ the divisor $\det(1 - S) = 0$ we get

$$\Xi_{I_N g} = \Xi_g + \delta.$$

Thus we can define a 'divisor' $\Xi_g \subset \text{Gr}^{1/2}(V \oplus V)$ for $g \in \mathcal{M}_\infty$. Notice that its component δ has multiplicity ∞ . Except δ , we have finite number of components of finite multiplicity.

REMARK. In the same way, $\xi_{I_N g} = \xi_g + \delta$. We can define ξ_g for $g \in \mathcal{M}_\infty$. However, acting in this way we loss information concerning the multiplicity of the divisor $\det \Lambda = 0$. \square

2.6. Multiple characteristic functions.

Proposition 2.9 *For fixed α, m and sufficiently large N the Taylor coefficients of the characteristic function χ_g and of the polynomial p_g at zero do not generate the algebra of $\text{GL}(N)$ -invariant polynomials on $\text{Mat}(\alpha + mN)$.*

Proof is given in Subsection 3.6.

For this reason, we introduce additional invariants. Let $j = 1, 2, \dots$. For $g \in \text{GL}(\infty, \mathbb{C})$ we consider the direct sum

$$g^{[j]} := g \oplus \dots \oplus g$$

of j copies of g . It acts in the space

$$\begin{aligned} \left[\mathbb{C}^\alpha \oplus (V \otimes Z_N) \right] \oplus \dots \oplus \left[\mathbb{C}^\alpha \oplus (V \otimes Z_N) \right] &= \mathbb{C}^{j\alpha} \oplus ((V \oplus \dots \oplus V) \otimes Z_N) = \\ &= \mathbb{C}^{j\alpha} \oplus ((V \otimes \mathbb{C}^j) \otimes Z_N). \end{aligned}$$

Thus we get an embedding

$$\mathcal{M}_N^{\alpha, m} \rightarrow \mathcal{M}_N^{j\alpha, jm}.$$

It is compatible with \circ -multiplication.

Now for a given $g \in \mathcal{M}_N^{\alpha, m}$ we get the collection of characteristic functions

$$\chi_g, \chi_{g^{[2]}}, \chi_{g^{[3]}}, \dots; \quad \chi_{g^{[j]}} : \text{Mat}(jm) \rightarrow \text{Mat}(j\alpha) \quad (2.6)$$

and the collection of divisors

$$\Xi_g, \Xi_{g^{[2]}}, \Xi_{g^{[3]}}, \dots; \quad \Xi_{g^{[j]}} \subset \text{Gr}((V \otimes \mathbb{C}^j) \oplus (V \otimes \mathbb{C}^j)). \quad (2.7)$$

Theorem 2.10 a) For any α, m, N the characteristic functions (2.6) and divisors (2.7) uniquely determine an element of the categorical quotient $[\mathcal{M}_N^{\alpha, m}]$

b) The characteristic functions (2.6) and divisors (2.7) uniquely determine an element of the categorical quotient $[\mathcal{GL}_N^{\alpha, m}]$

Proof is contained in Subsections 3.1–3.2.

Theorem 2.11 a) The characteristic functions (2.6) and divisors (2.7) uniquely determine an element of the space $\mathcal{U}_N^{\alpha, m}$ of conjugacy classes.

b) The characteristic functions (2.6) and divisors (2.7) uniquely determine an element of the space $\mathcal{U}_\infty^{\alpha, m}$ of conjugacy classes.

Proof is contained in Subsection 3.3.

3 Invariants

3.1. Invariants for $\mathcal{M}_N^{\alpha, m}$. Denote

$$a^{[j]} = a \otimes 1_j, \quad b^{[j]} = b \otimes 1_j, \quad c^{[j]} = c \otimes 1_j, \quad d^{[j]} = d \otimes 1_j.$$

Denote by $b_\beta[k]$ the k -th row of the matrix b_β , by $c_\gamma[l]$ the l -th column of the matrix c_γ . Let us regard $\text{Mat}(\alpha + mN)$ as a linear space with action of $\text{GL}(N, \mathbb{C})$. A point of the space is a collection of m^2 matrices d_{ij} , αm vectors $c_i[l]$, and αm of covectors $b_j[k]$. Generators of the algebra of invariants are known (see, e.g., [28], Section 11.8.1). The algebra is generated by the following polynomials

$$\text{tr } d_{\varphi_1 \psi_1} d_{\varphi_2 \psi_2} \dots d_{\varphi_n \psi_n}, \quad (3.1)$$

$$b_\beta[k] d_{\varphi_1 \psi_1} d_{\varphi_2 \psi_2} \dots d_{\varphi_n \psi_n} c_\gamma[l], \quad (3.2)$$

$$a_{\sigma\tau}. \quad (3.3)$$

We wish to show that all the generators can be expressed in the terms of Taylor coefficients of $\chi_{g^{[j]}}(S)$ and $p_{g^{[j]}}(S)$ at zero.

First, consider the expression

$$\ln p_{g^{[j]}}(S) = \ln \det(1 - d^{[j]} \tilde{S}), \quad (3.4)$$

where S is an operator in $V \otimes \mathbb{C}^j$.

It is convenient to think that matrix elements $s_{\varphi\psi}^{\mu\nu}$ of S depend on 4 indexes: indices $\varphi, \psi \leq m$ are responsible for an operator in the space \mathbb{C}^m , and $\mu, \nu \leq j$ for operators in the space \mathbb{C}^j . In a neighborhood of $S = 0$ we have the following

expansion

$$\begin{aligned}
\ln \det(1 - d^{[j]} \tilde{S}) &= \sum_{n>0} \frac{(-1)^n}{n} \operatorname{tr}(d^{[j]} \tilde{S})^n = \\
&= \sum_{\substack{n \geq 0, \\ \varphi_1, \dots, \varphi_n, \psi_1, \dots, \psi_n \leq m, \\ \mu_1, \dots, \mu_n \leq j}} \frac{(-1)^n}{n} s_{\varphi_1, \psi_1}^{\mu_1, \mu_2} s_{\varphi_2, \psi_2}^{\mu_2, \mu_3} \dots s_{\varphi_n, \psi_n}^{\mu_n, \mu_1} \times \\
&\quad \times \operatorname{tr} d_{\psi_1, \varphi_2} d_{\psi_2, \varphi_3} \dots d_{\psi_n, \varphi_1}. \quad (3.5)
\end{aligned}$$

Generally, Taylor coefficients (in $s_{\varphi\psi}^{\mu\nu}$) of this series are sums of several traces. However, assume that all elements of the sequence

$$\mu_1, \dots, \mu_n$$

are pairwise distinct. Then the coefficient at

$$s_{\varphi_2, \psi_2}^{\mu_2, \mu_3} \dots s_{\varphi_n, \psi_n}^{\mu_n, \mu_1}$$

is

$$\operatorname{tr} d_{\psi_1, \varphi_2} d_{\psi_2, \varphi_3} \dots d_{\psi_n, \varphi_1}.$$

We observe that all the invariants (3.1) are contained in the set of Taylor coefficients of (3.4).

Next we consider the characteristic function

$$\chi_{g^{[j]}}(S) = a^{[j]} + b^{[j]} \tilde{S}(1 - d^{[j]} \tilde{S})^{-1} c^{[j]}.$$

Substituting $S = 0$, we get invariants (3.3). Next, expand the matrix

$$X := b^{[j]} \tilde{S}(1 - d^{[j]} \tilde{S})^{-1} c^{[j]} = \sum_{n=0}^{\infty} b^{[j]} \tilde{S}(d^{[j]} \tilde{S})^n c^{[j]} \quad (3.6)$$

in a Taylor series. The operator X acts in $\mathbb{C}^m \otimes \mathbb{C}^j$. We enumerate its matrix elements as $x_{kl}^{\gamma\delta}$, where $k, l \leq \alpha$ and $\gamma, \delta \leq j$. The matrix elements are

$$\begin{aligned}
x_{kl}^{\gamma\delta} &= \sum_{n \geq 0} s_{\varphi_1, \psi_1}^{\gamma, \mu_2} s_{\varphi_2, \psi_2}^{\mu_2, \mu_3} \dots s_{\varphi_{n+1}, \psi_{n+1}}^{\mu_n, \delta} \times \\
&\quad \varphi_1, \dots, \varphi_{n+1} \leq m \\
&\quad \psi_1, \dots, \psi_{n+1} \leq m \\
&\quad \mu_2, \dots, \mu_n \leq j \\
&\quad \times b_{\varphi_1} [k] d_{\psi_1, \varphi_2} d_{\psi_2, \varphi_3} \dots d_{\psi_n, \varphi_{n+1}} c_{\psi_{n+1}} [l]
\end{aligned}$$

Again, consider a coefficient of the Taylor series at

$$s_{\varphi_1, \psi_1}^{\gamma, \mu_2} s_{\varphi_2, \psi_2}^{\mu_2, \mu_3} \dots s_{\varphi_{n+1}, \psi_{n+1}}^{\mu_n, \delta}$$

with pairwise distinct μ_2, \dots, μ_n that do not equal γ, δ . The factors d_{\dots} in the coefficient and their order are uniquely determined.

This proves Theorem 2.10.a.

3.2. Invariants for invertible matrices. Now consider the space $\mathcal{GL}_N^{\alpha, m}$. Invariant regular functions on this space have the form $p(g)/\det(g)^k$, where $p(g)$ is a polynomial satisfying

$$p(\iota(u)g\iota(u)^{-1}) = p(g), \quad \text{where } u \in \text{GL}(N, \mathbb{C}),$$

$\iota(u)$ is given by (1.3). Thus the algebra of invariant functions is generated by the same invariants and $\det(g)^{-1}$.

3.3. Invariants for unitary group. Consider invariants of $U(N)$ on $U(\alpha + mN)$. Consider the algebra \mathcal{A} of functions on $U(n)$ generated by matrix elements and $\det(g)^{-1}$. By the Peter-Weyl theorem it is dense in the algebra of continuous functions.

Consider two orbits $\mathcal{O}_1, \mathcal{O}_2 \subset U(\alpha + mN)$ of $U(N)$, consider an invariant continuous function φ separating these orbits. Consider a function $\psi \in \mathcal{A}$ approximating φ , it satisfies a condition of the form

$$\psi|_{\mathcal{O}_1} \leq a < b \leq \psi|_{\mathcal{O}_2}.$$

We have

$$\psi(g) = \frac{p(g)}{(\det g)^k},$$

where p is a polynomial. The average

$$f(g) = \int_{U(N)} \psi(\iota(h)g\iota(h)^{-1}) dh$$

has the same form $\frac{q(g)}{(\det g)^k}$ with a polynomial q . Therefore f has an invariant regular continuation to the whole group $\text{GL}(\alpha + mN, \mathbb{C})$. If $N > \alpha m$, then all such functions are polynomials in Taylor coefficients of $\chi_g(S)$ and $p_g(S)$.

3.4. Change of N . Next, we wish to show that the map $I_N : [\mathcal{M}_N^{\alpha, m}] \rightarrow [\mathcal{M}_{N+1}^{\alpha, m}]$ is an embedding (Proposition 2.1). The map I_N replaces

$$\begin{aligned} a &\rightarrow a; \\ b_i &\rightarrow (b_i \ 0); \\ c_l &\rightarrow \begin{pmatrix} c_l \\ 0 \end{pmatrix}; \\ d_{\varphi\psi} &\rightarrow \begin{pmatrix} d_{\varphi\psi} & 0 \\ 0 & 0 \end{pmatrix}, \quad i \neq j; \\ d_{\varphi\varphi} &\rightarrow \begin{pmatrix} d_{\varphi\varphi} & 0 \\ 0 & 1 \end{pmatrix}. \end{aligned}$$

Restrict generators (3.1)–(3.3) of invariants on $\text{Mat}(\alpha + m(N + 1))$ to $\text{Mat}(\alpha + mN)$. We get the same expressions in all the cases except traces of the form

$$\text{tr } d_{\varphi_1 \varphi_1} d_{\varphi_2 \varphi_2} \dots d_{\varphi_n \varphi_n},$$

they are shifted by 1. Therefore, restricting a collection of generators of $\text{GL}(N + 1, \mathbb{C})$ -invariants of $\text{Mat}(\alpha + m(N + 1))$ we get a collection of generators for algebra of $\text{GL}(N, \mathbb{C})$ -invariants on $\text{Mat}(\alpha + mN)$.

Thus we proved the desired statement.

3.5. Categorical quotient. Proposition 2.2 follows from Theorem 2.10 (invariants of \circ -products are uniquely determined by invariants of factors).

3.6. Proof of Proposition 2.9. Consider the algebra of $U(N)$ -invariants on $\text{Mat}(\alpha + mN)$. The number of the standard generators of the degree l has asymptotics $\sim \text{const} \cdot (m^2)^l$ as $l \rightarrow \infty$.

The Taylor coefficients of the characteristic function are polynomials of generators. Number of the Taylor coefficients equals to $\alpha^2 C_{m^2+l-1}^{l-1}$, it is a polynomial in l of degree m^2 .

The minimal (graded) degree of a relation between standard generators (3.1) is $N + 1$, see [27], Theorem 4.3.

It remains to choose a sufficiently large N such that number of independent monomials of the independent generators (3.1) of degree $\leq N$ is larger than number of monomials of the Taylor coefficients of degree $\leq N$.

4 Variations

Let A_1, \dots, A_l be square matrices. By $\text{Diag}(A_1, \dots, A_n)$ we denote the block-diagonal matrix with blocks A_1, \dots, A_l .

4.1. Several matrices determined up to a common conjugation. Let $G = U(\infty)$, and K be the stabilizer of the first α basis vectors as in Subsection 1.2. Consider the product $G^n = G \times \dots \times G$ of n copies of G and the subgroup K embedded to $G \times \dots \times G$ by diagonal, $h \mapsto (h, \dots, h)$. Consider the space $G^n // K$ of conjugacy classes

$$(g_1, \dots, g_n) \sim (hg_1h^{-1}, \dots, hg_nh^{-1}).$$

It is equipped with the component-wise product

$$(g_1, \dots, g_n) * (r_1, \dots, r_n) := (g_1 \circ r_1, \dots, g_n \circ r_n), \quad (4.1)$$

where \circ -multiplication of matrices is given by the formula (1.1).

To be definite, set $n = 2$. Consider a pair of matrices

$$\begin{pmatrix} a_1 & b_1 \\ c_1 & d_1 \end{pmatrix}, \quad \begin{pmatrix} a_2 & b_2 \\ c_2 & d_2 \end{pmatrix}$$

determined up to a common conjugation by a matrix $\begin{pmatrix} 1 & 0 \\ 0 & u \end{pmatrix}$. We compose a matrix

$$\begin{pmatrix} a_1 & 0 & b_1 & 0 \\ 0 & a_2 & 0 & b_2 \\ c_1 & 0 & d_1 & 0 \\ 0 & c_2 & 0 & d_2 \end{pmatrix}.$$

It is determined up to a conjugation by a matrix

$$\text{Diag}(1_\alpha, 1_\alpha, u, u) = \text{Diag}(1_{2\alpha}, u, u).$$

Thus we embedded the semigroup $G^n//K$ to the semigroup $U_\infty^{\alpha n, n}$. We know that multiple characteristic functions and distinguished divisors separate elements of $U_\infty^{\alpha n, n}$ and therefore they separate elements of $G^n//K$. We get the analog of Theorem 2.11. Analog of theorem 2.10 also holds but it requires a separate proof.

4.2. Another example with conjugacy classes. Now consider the group $Q_{\alpha, m, m}$ of finitary unitary matrices of size

$$\alpha + \underbrace{\infty + \dots + \infty}_{m \text{ times}} + \underbrace{\infty + \dots + \infty}_{m \text{ times}}$$

determined up to conjugations by the group H of unitary matrices of the form

$$\text{Diag}(1, \underbrace{u, \dots, u}_{m \text{ times}}, \underbrace{v, \dots, v}_{m \text{ times}}),$$

i.e., $H \simeq U(\infty) \times U(\infty)$. The formula for multiplication on $Q_{\alpha, m, m}/H$ is the same as in Subsection 1.3.

Now we explain how to obtain a characteristic function. To be definite, set $m = 2$. We fix two $m \times m$ matrices, T, S , set

$$\Lambda = T^{-1}, \quad K = S^{-1},$$

and write the equation

$$\begin{pmatrix} q \\ x_1 \\ x_2 \\ y_1 \\ y_2 \end{pmatrix} = \begin{pmatrix} a_1 & b_1 & b_2 & b_3 & b_4 \\ c_1 & d_{11} & d_{12} & d_{13} & d_{14} \\ c_2 & d_{21} & d_{22} & d_{23} & d_{24} \\ c_1 & d_{31} & d_{32} & d_{33} & d_{34} \\ c_1 & d_{41} & d_{42} & d_{43} & d_{44} \end{pmatrix} \begin{pmatrix} p \\ s_{11}x_1 + s_{12}x_2 \\ s_{21}x_1 + s_{22}x_2 \\ t_{11}y_1 + t_{12}y_2 \\ t_{21}y_1 + t_{22}y_2 \end{pmatrix} \quad (4.2)$$

(big matrix is an element $g \in Q_{\alpha, m, m}$). We eliminate variables s_{ij}, t_{ij} and get a relation

$$q = \chi_g(S, T) p.$$

Thus we get a rational function $\text{Mat}(m) \times \text{Mat}(m) \rightarrow \text{Mat}(\alpha)$, it sends $B_m \times B_m$ to B_α and $U(m) \times U(m) \rightarrow U(\alpha)$. As above, a product of conjugacy classes

corresponds to a point-wise product of functions. See several proofs of similar statements in [16].

We also define a distinguished divisor

$$\Xi_g \in \mathrm{Gr}^{1/2}(\mathbb{C}^m \oplus \mathbb{C}^m) \times \mathrm{Gr}^{1/2}(\mathbb{C}^m \oplus \mathbb{C}^m).$$

In the coordinates, $(T, S) \in \mathrm{Mat}(n) \times \mathrm{Mat}(n)$ it is determined by the equation

$$\det \left[1 - \begin{pmatrix} d_{11} & d_{12} & d_{13} & d_{14} \\ d_{21} & d_{22} & d_{23} & d_{24} \\ d_{31} & d_{32} & d_{33} & d_{34} \\ d_{41} & d_{42} & d_{43} & d_{44} \end{pmatrix} \begin{pmatrix} s_{11} & s_{12} & 0 & 0 \\ s_{21} & s_{22} & 0 & 0 \\ 0 & 0 & t_{11} & t_{12} \\ 0 & 0 & t_{21} & t_{22} \end{pmatrix} \right] = 0.$$

In coordinates $(\Lambda, K) \in \mathrm{Mat}(m) \times \mathrm{Mat}(m)$ we get the equation

$$\det \left[\begin{pmatrix} \lambda_{11} & \lambda_{12} & 0 & 0 \\ \lambda_{21} & \lambda_{22} & 0 & 0 \\ 0 & 0 & \varkappa_{11} & \varkappa_{12} \\ 0 & 0 & \varkappa_{21} & \varkappa_{22} \end{pmatrix} - \begin{pmatrix} d_{11} & d_{12} & d_{13} & d_{14} \\ d_{21} & d_{22} & d_{23} & d_{24} \\ d_{31} & d_{32} & d_{33} & d_{34} \\ d_{41} & d_{42} & d_{43} & d_{44} \end{pmatrix} \right] = 0.$$

These two charts cover the product of Grassmannians up to a subvariety of codimension 2. This is sufficient to define a divisor.

As in Subsection 2.6, we consider elements

$$\mathfrak{g}^{[j]} = \underbrace{\mathfrak{g} \oplus \cdots \oplus \mathfrak{g}}_{j \text{ times}} \in Q_{j\alpha, jm, jm} // H.$$

Theorem 4.1 *The collection of characteristic functions $\chi_{\mathfrak{g}^{[j]}}$ and the distinguished divisors $\Xi_{\mathfrak{g}^{[j]}}$ uniquely determine a conjugacy class $\mathfrak{g} \in Q_{\alpha, m, m} // U(\infty) \times U(\infty)$.*

A proof repeats considerations of Section 3.

4.3. Example with double cosets. Consider the group $G = U(\infty)$ and its subgroup L consisting of matrices $\begin{pmatrix} 1_\alpha & 0 \\ 0 & u \end{pmatrix}$, where u is real orthogonal. Consider the group $G^n = G \times \cdots \times G$ (n times) and the subgroup L embedded to G^n by diagonal. Consider double cosets $L \backslash G/L$, i.e., we consider tuples

$$\left(\begin{pmatrix} a_1 & b_1 \\ c_1 & d_1 \end{pmatrix}, \dots, \begin{pmatrix} a_n & b_n \\ c_n & d_n \end{pmatrix} \right)$$

up to the equivalence

$$\begin{aligned} & \left(\begin{pmatrix} a_1 & b_1 \\ c_1 & d_1 \end{pmatrix}, \dots, \begin{pmatrix} a_n & b_n \\ c_n & d_n \end{pmatrix} \right) \simeq \\ & \simeq \left(\begin{pmatrix} 1_\alpha & 0 \\ 0 & u \end{pmatrix} \begin{pmatrix} a_1 & b_1 \\ c_1 & d_1 \end{pmatrix} \begin{pmatrix} 1_\alpha & 0 \\ 0 & v \end{pmatrix}, \dots, \begin{pmatrix} 1_\alpha & 0 \\ 0 & u \end{pmatrix} \begin{pmatrix} a_n & b_n \\ c_n & d_n \end{pmatrix} \begin{pmatrix} 1_\alpha & 0 \\ 0 & v \end{pmatrix} \right), \end{aligned}$$

- [2] Brodskij, V.M. *On operator nodes and their characteristic functions*. Sov. Math., Dokl. 12, 696-700 (1971).
- [3] Brodskij, M.S. *Unitary operator colligations and their characteristic functions*. Russ. Math. Surv. 33, No.4, 159–191 (1978).
- [4] Dym, H. *Linear algebra in action*. American Mathematical Society, Providence, RI, 2007.
- [5] Gaifullin A.A., Neretin Yu.A. *Infinite symmetric group and bordisms of pseudomanifolds*. Preprint, arXiv:1501.04062
- [6] Garnett, J. B. *Bounded analytic functions*. Academic Press, Inc. New York-London, 1981.
- [7] Gohberg, I., Goldberg, S., Kaashoek, M. A. *Classes of linear operators. Vol. II*. Birkhauser, Basel, 1993.
- [8] Griffiths, Ph.; Harris, J. *Principles of algebraic geometry*. 2nd ed. Wiley Classics Library. New York, NY: John Wiley & Sons (1994).
- [9] Hazewinkel, M. *Lectures on invariants, representations and Lie algebras in systems and control theory*. Sémin. d’algèbre P. Dubreil et M.-P. Malliavin, 35ème Année, Proc., Paris 1982, Lect. Notes Math. 1029, 1-36 (1983).
- [10] Hitchin, N. *Riemann surfaces and integrable systems*, in *Integrable systems*, Clarendon Press, Oxford, 1999, 11–52.
- [11] Ismagilov, R.S., *Elementary spherical functions on the groups $SL(2, P)$ over a field P , which is not locally compact with respect to the subgroup of matrices with integral elements*. Math. USSR-Izvestiya, 1967, 1:2, 349–380.
- [12] Livshits, M.S. *On a certain class of linear operators in Hilbert space*. Mat. Sb., N. Ser. 19(61), 239-262 (1946); English transl. in Amer. Math. Soc. Transl. (Ser. 2), Vol. 13, 61-83 1960.
- [13] Livshits, M. S. *On spectral decomposition of linear nonself-adjoint operators*. Mat. Sbornik N.S. 34(76), (1954). 145–199. English transl. in Amer. Math. Soc. Transl. (Ser. 2), Vol 5, 1957, 67–114.
- [14] Martin, C.; Hermann, R. *Applications of algebraic geometry to systems theory: the McMillan degree and Kronecker indices of transfer functions as topological and holomorphic system invariants*. SIAM J. Control Optimization 16, 743-755 (1978).
- [15] Neretin, Yu. A. *Categories of symmetries and infinite-dimensional groups*. Oxford University Press, New York, 1996.
- [16] Neretin, Yu. A. *Multi-operator colligations and multivariate spherical functions*. Anal. and Math. Physics, Anal. Math. Phys. 1 (2011), no. 2-3, 121-138.

- [17] Neretin, Yu. A. *Sphericity and multiplication of double cosets for infinite-dimensional classical groups*. *Funct. Anal. Appl.*, 2011, 45:3, 225-239.
- [18] Neretin, Yu.A. *On degeneration of convolutions of double cosets at infinite-dimensional limit*. Preprint, <http://arxiv.org/abs/1211.6149>.
- [19] Neretin, Yu. *Infinite tri-symmetric group, multiplication of double cosets, and checker topological field theories*. *Int. Math. Res. Not. IMRN* 2012, no. 3, 501523.
- [20] Neretin, Yu.A. *Infinite-dimensional p -adic groups, semigroups of double cosets, and inner functions on Bruhat-Tits buildings*. *Izvestiya: Mathematics*, 2015, 79:3, 512–553
- [21] Neretin, Yu.A. *Infinite symmetric groups and combinatorial constructions of topological field theory type*. *Russ. Math. Surv.*, 2015, 70:4, 715–773
- [22] Olshanski, G. I. *Unitary representations of (G, K) -pairs that are connected with the infinite symmetric group $S(\infty)$* . *Leningrad Math. J.* 1 (1990), no. 4, 983-1014
- [23] Olshanski, G. I. *Unitary representations of infinite dimensional pairs and the formalism of R. Howe* in *Representation of Lie Groups and Related Topics*, Gordon and Breach, New York, 1990, 269–463.
- [24] Olshanskii, G. I. *Caractères généralisés du groupe $U(\infty)$ et fonctions intérieures*,— *C. R. Acad. Sci. Paris. Sér. 1*, 313:1 (1991), 9-12.
- [25] Popov, V.L.; Vinberg E.B. *Invariant Theory in Algebraic Geometry IV*, *Encyclopaedia of Math. Sci.*, v. 55, Springer-Verlag, 1994, 137-314.
- [26] Potapov, V. P. *The multiplicative structure of J -contractive matrix functions*. *Trudy Moskov. Mat. Obshchestva.* 4 (1955), 125–236; English transl. *Amer. Math. Soc. Transl.* (2) 15 (1960) 131–243.
- [27] Procesi, C. *The invariant theory of $n \times n$ matrices*. *Advances in Math.* 19 (1976), no. 3, 306-381.
- [28] Procesi, C. *Lie groups. An approach through invariants and representations*, NY: Springer, 2007.
- [29] Sz.-Nagy, B.; Foias, C. *Harmonic analysis of operators on Hilbert space*. Budapest: Akademiai Kiado (1970).

Math.Dept., University of Vienna,
 Nordbergstrasse, 15, Vienna, Austria
 &
 Institute for Theoretical and Experimental Physics,
 Bolshaya Cheremushkinskaya, 25, Moscow 117259, Russia
 &

Mech.Math. Dept., Moscow State University,
Vorob'evy Gory, Moscow
e-mail: neretin(at) mccme.ru
URL:www.mat.univie.ac.at/~neretin
wwwth.itep.ru/~neretin