

ON ABELIAN GROUP REPRESENTABILITY OF FINITE GROUPS

ELDHO K. THOMAS, NADYA MARKIN, FRÉDÉRIQUE OGGIER

ABSTRACT. A set of quasi-uniform random variables X_1, \dots, X_n may be generated from a finite group G and n of its subgroups, with the corresponding entropic vector depending on the subgroup structure of G . It is known that the set of entropic vectors obtained by considering arbitrary finite groups is much richer than the one provided just by abelian groups. In this paper, we start to investigate in more detail different families of non-abelian groups with respect to the entropic vectors they yield. In particular, we address the question of whether a given non-abelian group G and some fixed subgroups G_1, \dots, G_n end up giving the same entropic vector as some abelian group A with subgroups A_1, \dots, A_n , in which case we say that (A, A_1, \dots, A_n) represents (G, G_1, \dots, G_n) . If for any choice of subgroups G_1, \dots, G_n , there exists some abelian group A which represents G , we refer to G as being abelian (group) representable for n . We completely characterize dihedral, quasi-dihedral and dicyclic groups with respect to their abelian representability, as well as the case when $n = 2$, for which we show a group is abelian representable if and only if it is nilpotent. This problem is motivated by understanding non-linear coding strategies for network coding, and network information theory capacity regions.

1. INTRODUCTION

Let X_1, \dots, X_n be a collection of n jointly distributed discrete random variables over some alphabet of size N . We denote by \mathcal{A} a subset of indices from $\mathcal{N} = \{1, \dots, n\}$, and $X_{\mathcal{A}} = \{X_i, i \in \mathcal{A}\}$. The *entropic vector* corresponding to X_1, \dots, X_n is the vector

$$h = (H(X_1), \dots, H(X_1, X_2), \dots, H(X_1, \dots, X_n))$$

which collects all the joint entropies $H(X_{\mathcal{A}})$, $\mathcal{A} \subseteq \mathcal{N}$. Consider the $(2^n - 1)$ -dimensional Euclidean space, with coordinates labeled by all possible non-empty subsets $\mathcal{A} \subseteq \{1, \dots, n\}$. The region formed by all entropic vectors h is denoted by Γ_n^* . Determining Γ_n^* is notoriously difficult. However, it is also the key to understanding the capacity region of any arbitrary multi-source multi-sink acyclic network, which can be computed as a linear function of the entropy vector over Γ_n^* , under linear constraints [7, 5].

1991 *Mathematics Subject Classification.* 20D15 ; 94A17 .

Key words and phrases. nilpotent group, group representability, entropic vector.

E. K. Thomas was supported by Nanyang Technological University under Research Grant M58110049, N. Markin was supported by the Singapore National Research Foundation under Research Grant NRF-CRP2-2007-03, F. Oggier was partially funded by Nanyang Technological University under Research Grant M58110049.

It is known [1] that the closure $\bar{\Gamma}_n^*$ of the set of all entropic vectors is equal to the convex closure (the smallest closed convex set containing the set) of all entropic vectors generated by quasi-uniform random variables.

Definition 1. A probability distribution over a set of n random variables X_1, \dots, X_n is said to be *quasi-uniform* if for any $\mathcal{A} \subseteq \mathcal{N}$, $X_{\mathcal{A}}$ is uniformly distributed over its support $\lambda(X_{\mathcal{A}})$:

$$P(X_{\mathcal{A}} = x_{\mathcal{A}}) = \begin{cases} 1/|\lambda(X_{\mathcal{A}})| & \text{if } x_{\mathcal{A}} \in \lambda(X_{\mathcal{A}}), \\ 0 & \text{otherwise.} \end{cases}$$

By definition, the entropy of a quasi-uniform distribution is

$$H(X_{\mathcal{A}}) = - \sum_{x_{\mathcal{A}} \in \lambda(X_{\mathcal{A}})} P(X_{\mathcal{A}} = x_{\mathcal{A}}) \log P(X_{\mathcal{A}} = x_{\mathcal{A}}) = \log |\lambda(X_{\mathcal{A}})|.$$

We now turn to the question of characterizing quasi-uniform random variables. Let G be a finite group with n subgroups G_1, \dots, G_n , and let X be a random variable uniformly distributed over G , that is, $P(X = g) = 1/|G|$ for any $g \in G$. Define a new random variable $X_i = XG_i$, whose support is $[G : G_i]$ cosets of G_i in G . Then $P(X_i = gG_i) = |G_i|/|G|$ since $X_i = gG_i$ whenever $g \in G_i$, and $P(X_i = gG_i, i \in \mathcal{A}) = |\cap_{i \in \mathcal{A}} G_i|/|G|$. This shows that quasi-uniform random variables may be obtained from finite groups, more precisely:

Theorem 2. [1, 2] *For any finite group G and any subgroups G_1, \dots, G_n of G , there exists a set of n jointly distributed quasi-uniform discrete random variables X_1, \dots, X_n such that for all non-empty subsets \mathcal{A} of \mathcal{N} , $H(X_{\mathcal{A}}) = \log [G : G_{\mathcal{A}}]$, where $G_{\mathcal{A}} = \cap_{i \in \mathcal{A}} G_i$ and $[G : G_{\mathcal{A}}]$ is the index of $G_{\mathcal{A}}$ in G .*

Thus, given a group G and n subgroups G_1, \dots, G_n , one obtains n quasi-uniform random variables. The subgroup structure of G determines the correlation among the n random variables, and in turn the corresponding entropic vector.

Definition 3. Let X_1, \dots, X_n be n jointly distributed discrete random variables. The corresponding entropic vector h is said to be *group representable*, if there exists a group G , with subgroups G_1, \dots, G_n such that $H(X_{\mathcal{A}}) = \log [G : G_{\mathcal{A}}]$ for all \mathcal{A} . If in addition, the group G is abelian, we say that h is *abelian group representable*.

Interestingly, considering only quasi-uniform distributions coming from finite groups is in fact enough to understand $\bar{\Gamma}_n^*$:

Theorem 4. [3] *Let Υ_n be the region of all group representable entropic vectors, and let $\overline{\text{con}}(\Upsilon_n)$ be its convex closure. Then*

$$\overline{\text{con}}(\Upsilon_n) = \bar{\Gamma}_n^*.$$

It is also shown in [1] that the collection of all abelian group representable entropic vectors Υ_n^{ab} gives a non-trivial inner bound for $\bar{\Gamma}_n^*$ when n is at least 4.

Theorem 5. [1] *When $n \geq 4$, $\overline{\text{con}}\Upsilon_n^{ab} \neq \bar{\Gamma}_n^*$.*

Thus entropic vectors which are abelian group representable are a proper subset of entropic vectors coming from quasi-uniform random variables. This addresses a natural question of classifying groups with respect to the entropic vectors which they induce. In particular, we want to understand which groups belong to the same class as abelian groups with respect to this classification. We make this precise in the definitions below.

Definition 6. Let G be a group and let G_1, \dots, G_n be fixed subgroups of G . Suppose there exists an abelian group A with subgroups A_1, \dots, A_n such that for every non-empty $\mathcal{A} \subseteq \mathcal{N}$, $[G : G_{\mathcal{A}}] = [A : A_{\mathcal{A}}]$, where $G_{\mathcal{A}} = \bigcap_{i \in \mathcal{A}} G_i$, $A_{\mathcal{A}} = \bigcap_{i \in \mathcal{A}} A_i$. Then we say that (A, A_1, \dots, A_n) represents (G, G_1, \dots, G_n) .

Definition 7. If for every choice of subgroups G_1, \dots, G_n of G , there exists an abelian group A such that (A, A_1, \dots, A_n) represents (G, G_1, \dots, G_n) , we say that G is *abelian (group) representable for n* .

Note that the abelian group A may vary for different choices of subgroups G_1, \dots, G_n . Sometimes, however, it is possible to find an abelian group that works for all choices G_1, \dots, G_n .

Definition 8. Suppose there exists an abelian group A such that for every choice of subgroups $G_1, \dots, G_n \leq G$, there exist subgroups $A_1, \dots, A_n \leq A$ such that (A, A_1, \dots, A_n) represents (G, G_1, \dots, G_n) . Then we will say that G is *uniformly abelian (group) representable for n* . (Alternatively, A *uniformly represents G* .)

When G is abelian group representable, the quasi-uniform random variables X_1, \dots, X_n corresponding to subgroups G_i can also be obtained using an abelian group A and its subgroups A_1, \dots, A_n . If we choose the subgroup $G_i = G$, then $\log[G : G] = 0$, that is $H(X_i) = 0$, which implies X_i is actually taking values deterministically. If $G_i = \{1\}$, then $\log[G : \{1\}] = \log |G|$. Thus the entropy chain rule yields

$$H(X_i, X_{\mathcal{A}}) = H(X_i) + H(X_{\mathcal{A}}|X_i)$$

for every \mathcal{A} such that $i \notin \mathcal{A}$. Since $H(X_i) = \log |G|$ and $H(X_i, X_{\mathcal{A}}) = \log[G : \{1\} \cap G_{\mathcal{A}}] = \log |G|$, we conclude that

$$H(X_{\mathcal{A}}|X_i) = 0.$$

That is, given X_i , all the $n - 1$ other random variables are functions of X_i and we will consequently require that each subgroup G_1, \dots, G_n is non-trivial and proper. Hence n is at most the number of non-trivial proper subgroups of G .

The contributions of this paper are to introduce the notion of abelian group representability for an arbitrary finite group G , and to characterize the abelian group representability of several classes of groups:

- Dihedral, quasi-dihedral and dicyclic groups are shown to be abelian group representable for every $n \geq 2$ if (Section 2) and only if (Section 4) they are 2-groups. When they are abelian group representable, they are furthermore uniformly abelian group representable for every n (Section 2).
- p -groups: p -groups are shown to be uniformly abelian group representable for $n = 2$ in Section 3.
- Nilpotent groups: in Section 4 we show that representability of nilpotent groups is completely determined by representability of p -groups. The set of nilpotent groups is shown to contain the set of abelian representable groups for any n in Section 4, the two coincide for $n = 2$.

2. ABELIAN GROUP REPRESENTABILITY OF CLASSES OF 2-GROUPS

In this section we establish uniform abelian group representability of dihedral, quasi-dihedral and dicyclic 2-groups for any n . We begin with a general lemma showing how abelian group representability of a group H may imply abelian group representability of a group G .

Lemma 9. *Let*

$$\psi : G \rightarrow H$$

be a bijective map, which is additionally subgroup preserving, i.e., for any subgroup $G_i \leq G$, the set $\psi(G_i)$ is a subgroup of H . Suppose that H is abelian (resp. uniformly abelian) group representable. Then so is G . In particular, if H itself is abelian, then G is abelian group representable.

Proof. We want to show that given subgroups $G_1, \dots, G_n \leq G$, there exists an abelian group A with subgroups A_1, \dots, A_n such that for any subset $\mathcal{A} \subseteq \{1, \dots, n\}$ the intersection subgroup $G_{\mathcal{A}}$ has index $[G : G_{\mathcal{A}}] = [A : A_{\mathcal{A}}]$.

Since H is abelian group representable, $(H, \psi(G_1), \dots, \psi(G_n))$ can be represented by some (A, A_1, \dots, A_n) . We claim that (A, A_1, \dots, A_n) will also represent (G, G_1, \dots, G_n) .

Since ψ is bijective, for any $G_i \leq G$, the subgroup $\psi(G_i)$ has the same size and index in H as G_i has in G . In particular, $[A : A_i] = [H : \psi(G_i)] = [G : G_i]$. This takes care of 1-intersection, i.e., when $|\mathcal{A}| = 1$.

Now we want to show that in fact $[G : G_{\mathcal{A}}] = [A : A_{\mathcal{A}}]$ for any $\mathcal{A} \subseteq \mathcal{N}$. When considering intersections $G_{\mathcal{A}}$, let us first consider 2-intersections $G_{12} = G_1 \cap G_2 \leq G$. First observe that $\psi(G_1 \cap G_2) = \psi(G_1) \cap \psi(G_2)$. The containment $\psi(G_1 \cap G_2) \subseteq \psi(G_1) \cap \psi(G_2)$ is immediate. To see the containment $\psi(G_1 \cap G_2) \supseteq \psi(G_1) \cap \psi(G_2)$ observe that if $y = \psi(g_1) = \psi(g_2)$ for some $g_1 \in G_1, g_2 \in G_2$ then bijectivity of ψ implies that $g_1 = g_2$ and $y \in \psi(G_1 \cap G_2)$.

Now, recalling that $|G| = |H|$, we see that

$$[G : G_1 \cap G_2] = [H : \psi(G_1) \cap \psi(G_2)] = [A : A_1 \cap A_2].$$

More generally for arbitrary intersection $G_{\mathcal{A}}$, we have $[G : G_{\mathcal{A}}] = [A : A_{\mathcal{A}}]$: this follows by induction on the number of subgroups involved in the intersection. We conclude that $(A, A_i)_{i \in \mathcal{N}}$ represents $(G, G_i)_{i \in \mathcal{N}}$. If H was uniformly abelian group representable, then A was chosen independently of subgroups $\psi(G_1), \dots, \psi(G_n)$ and it follows that G is also uniformly abelian group representable. \square

2.1. Dihedral and Quasi-dihedral 2-Groups. We will establish the abelian representability for dihedral and quasi-dihedral 2-groups. We define the dihedral group D_m for $m \geq 3$ to be the symmetry group of the regular m -sided polygon. The group D_m is of order $2m$, with a well known description in terms of generators and relations:

$$D_m = \langle r, s \mid r^m = s^2 = 1, rs = sr^{-1} \rangle.$$

Each element of D_m is uniquely represented as $r^a s^j$ where $1 \leq a \leq m, j = 0, 1$.

Note that the generator s acts on r by conjugation sending r to an element $srs^{-1} = r^{-1}$. More generally, consider other possibilities for $srs^{-1} = r^z$. When we apply this map twice, we see that since s has order 2, it must be that $z^2 \equiv 0 \pmod{2^m}$. In case $m = 2^k$, there are 4 such choices modulo 2^k for $z \in \mathbb{Z}_{2^k}^\times$, i.e., $z \equiv \pm 1, 2^{k-1} \pm 1$. The choice $z = 1$ will result in an abelian group, $z = -1$ gives the dihedral group D_{2^k} above. Now we cover the remaining two choices. In either case, the subgroup structure is similar to that of a dihedral group, which will eventually allow us to conclude that these groups are abelian representable via Lemma 9.

Define two quasi-dihedral groups, each of order 2^{k+1} :

$$\mathcal{QD}_{2^k}^{-1} = \langle r, s \mid r^{2^k} = s^2 = 1, rs = sr^{2^{k-1}-1} \rangle,$$

$$\mathcal{QD}_{2^k}^{+1} = \langle r, s \mid r^{2^k} = s^2 = 1, rs = sr^{2^{k-1}+1} \rangle.$$

Let us make some brief observations about the structure of subgroups of $D_{2^k}, \mathcal{QD}_{2^k}^{-1}, \mathcal{QD}_{2^k}^{+1}$. First of all, note that a subgroup generated by $\langle r^{j_1}s, r^{j_2}s \rangle$ can also be generated by $\langle r^{j_1}s, r^{j_2}s(r^{j_1}s)^{-1} \rangle = \langle r^{j_1}s, r^{j_2-j_1} \rangle$ which in turn can be expressed as $\langle r^{2^i}, r^{j_1}s \rangle$ for some i . We conclude that only one generator of the form $r^j s$ is required. Trivially, only one generator of the form r^j is required as well, since the cyclic subgroup $\langle r \rangle$ contains only cyclic subgroups $\langle r^{2^i} \rangle$.

Hence any subgroup can be expressed in terms of at most 2 generators of the form $r^{2^i}, r^j x$. When the subgroup is given by $\langle r^{2^i}, r^j x \rangle$, without loss of generality we can further assume that $j < 2^i$: this is achieved by premultiplying the second generator $r^j s$ repeatedly with r^{-2^i} . This gives us 3 basic subgroup types of dihedral and quasi-dihedral groups.

Remark 10. *Subgroups of $D_{2^k}, \mathcal{QD}_{2^k}^{-1}, \mathcal{QD}_{2^k}^{+1}$ can be only of the following types:*

- (1) $\langle r^{2^i} \rangle, 0 \leq i < k,$
- (2) $\langle r^a s \rangle, 0 \leq a < 2^k,$
- (3) $\langle r^{2^i}, r^c s \rangle, 0 \leq i < k, 0 \leq c < 2^i.$

2.1.1. *Dihedral 2-groups.* We now establish abelian group representability of dihedral 2-groups for all n by applying the lemma above. We start by showing that subgroups of D_{2^k} map to subgroups of an abelian group.

Proposition 11. *Let $A = \bigoplus_{i=0}^k \mathbb{Z}_2$ be generated by the standard basis $\{e_0, \dots, e_k\}$ over integers \mathbb{Z}_2 modulo 2. There exists a subgroup preserving bijection from the dihedral group D_{2^k} to A .*

Proof. For each element $r^a s^j$ we use the base 2 representation of exponent $a = \sum_{i=0}^{k-1} a_i 2^i$ to define the map $\psi : D_{2^k} \rightarrow A$:

$$\psi : r^a s^j \mapsto \sum_{i=0}^{k-1} a_i e_i + j e_k.$$

We show that the map ψ is a subgroup preserving bijection.

First note that since each element is uniquely represented as $r^a s^j$ for some a, j , the map ψ is indeed well-defined and clearly bijective. Hence we only need to verify that ψ is subgroup preserving. In other words, if $H \leq D_{2^k}$ is a subgroup, then the image $\psi(H)$ is a subgroup of A . Since each element in A is its own inverse, and $\psi(H)$ contains the identity, we only need to check the closure property of $\psi(H)$ in order to prove that it is a subgroup.

To that end, we investigate the subgroups of D_{2^k} . All subgroups of D_{2^k} are of the form (see Remark 10)

- (1) $\langle r^{2^i} \rangle = \{r^a : a \equiv 0 \pmod{2^i}\}, 0 \leq i < k,$
- (2) $\langle r^a s \rangle = \{1, r^a s\}, 0 \leq a < 2^k,$
- (3) $\langle r^{2^i}, r^c s \rangle = \{r^a, r^b s : a \equiv 0 \pmod{2^i}, b \equiv c \pmod{2^i}\},$
 $0 \leq i < k, 0 \leq c < 2^k.$

Case 1: $H = \langle r^{2^i} \rangle = \{r^a : a \equiv 0 \pmod{2^i}\}$. Note that $a < 2^k$ is a multiple of 2^i if and only if $a = \sum_{j=i}^{k-1} a_j 2^j$, i.e., the terms a_0, \dots, a_{i-1} of the binary expansion of a are 0. In other words, the image

$$\psi(H) = \left\{ \sum_{j=i}^{k-1} a_j e_j \right\}.$$

Clearly this set is closed under addition.

Case 2: $H = \langle r^a s \rangle = \{1, r^a s\}$. Then the image

$$\psi(H) = \left\{ \sum_{i=0}^{k-1} a_i e_i + e_k, 0 \right\}.$$

Clearly this set of size 2 is a subgroup of A .

Case 3: $H = \langle r^{2^i}, r^c s \rangle$, $0 \leq c < 2^i$.

Indeed $H = \{r^{2^i h}, r^{c+2^i h} s : h \in \mathbb{Z}\}$.

We verify directly that $\psi(H)$ is closed under addition by showing that

$$x, y \in H \implies \psi(x) + \psi(y) \in H.$$

This involves considering 3 cases:

- $x, y \in \langle r^{2^i} \rangle$
- $x \in \langle r^{2^i} \rangle, y = r^{c+2^i h} s$
- both x, y are of the form $r^{c+2^i h} s$.

(1) $x, y \in \langle r^{2^i} \rangle$ implies that $\psi(x) + \psi(y) \in \psi(H)$. This follows identically to **Case 1**.

(2) $x \in \langle r^{2^i} \rangle, y = r^{c+2^i h'} s$ implies that $\psi(x) + \psi(y) \in \psi(H)$.

We show that $\psi(r^{2^i h}) + \psi(r^{2^i h'+c} s) = \psi(r^{2^i h'+c} s) \in \psi(H)$. To see this, observe using the binary expansion of exponents, that when $c < 2^i$ we can actually factor $\psi(r^{2^i h'+c}) = \psi(r^{2^i h'}) + \psi(r^c)$. Hence

$$\begin{aligned} \psi(r^{2^i h}) + \psi(r^{2^i h'+c} s) &= \psi(r^{2^i h}) + \psi(r^{2^i h'}) + \psi(r^c) + \psi(s) = \\ &= \psi(r^{2^i h''}) + \psi(r^c) + \psi(s) = \psi(r^{2^i h''+c} s) \in \psi(H). \end{aligned}$$

(3) both x, y of the form $r^{c+2^i h} s$ implies that $\psi(x) + \psi(y) \in \psi(H)$.

$$\begin{aligned} \text{Now } \psi(r^{2^i h+c} s) + \psi(r^{2^i h'+c} s) &= \psi(r^{2^i h+c}) + \psi(s) + \psi(r^{2^i h'+c}) + \psi(s) = \\ &= \psi(r^{2^i h}) + \psi(r^c) + \psi(r^{2^i h'}) + \psi(r^c) = \psi(r^{2^i h''}) \in \psi(H). \end{aligned}$$

□

Proposition 12. *The dihedral group D_{2^k} is uniformly abelian group representable for all n .*

Proof. This follows from Proposition 11 and Lemma 9. The uniform part comes from the fact that A only depends on the size of D_{2^k} and not on the choice of subgroups of D_{2^k} .

□

We next use abelian representability of dihedral groups to derive abelian representability of quasi-dihedral groups. The idea is that these two classes of groups have the same subgroups.

2.1.2. Quasi-dihedral 2-Groups. We already know by Remark 10 what subgroups of quasi-dihedral groups look like in terms of generators. However, for the purpose of defining a subgroup preserving map ψ (which will not be a homomorphism!), we want to know exactly what elements these subgroups consist of. The next lemma describes what the subgroups of quasi-dihedral groups look like.

Lemma 13. *Let $G = \mathcal{QD}_{2^k}^{-1}$ or $\mathcal{QD}_{2^k}^{+1}$ be a quasi-dihedral group of size 2^{k+1} . Then all the subgroups of G are of the form*

- (1) $\langle r^{2^i} \rangle = \{r^a : a \equiv 0 \pmod{2^i}\}$, $0 \leq i \leq k-1$,
- (2) $\langle r^{2^i}, r^j s \rangle = \{r^a, r^b s : a \equiv 0 \pmod{2^i}, b \equiv j \pmod{2^i}\}$, $0 \leq i \leq k-1$, $0 \leq j \leq 2^i - 1$,
- (3) • When $G = \mathcal{QD}_{2^k}^{-1}$, then $\langle r^j s \rangle = \begin{cases} \{1, r^j s\} & j \equiv 0 \pmod{2} \\ \{1, r^j s, r^{2^{k-1}}, r^{2^{k-1}+j} s\} & j \equiv 1 \pmod{2} \end{cases}$
- When $G = \mathcal{QD}_{2^k}^{+1}$, then $\langle r^j s \rangle = \begin{cases} \langle r^{2^j}, r^j s \rangle \text{ is of type (2)} & j \neq 0 \\ \{1, s\} & j = 0. \end{cases}$

Proof. First assume that $G = \mathcal{QD}_{2^k}^{-1}$.

- (1) $\langle r^{2^i} \rangle = \{r^a : a \equiv 0 \pmod{2^i}\}$
This case is obvious.
- (2) $\langle r^{2^i}, r^j s \rangle = \{r^a, r^b s : a \equiv 0 \pmod{2^i}, b \equiv j \pmod{2^i}\}$;
 $0 \leq i \leq k-1$, $0 \leq j \leq 2^i - 1$.
For this we observe that $r^j s r^{2^i} = r^{j+2^i(2^{k-1}-1)} s$.
Additionally note that

$$r^j s r^j s = r^{j+j(2^{k-1}-1)} s^2 = r^{j(2^{k-1})} = \begin{cases} 1 & j \equiv 0 \pmod{2} \\ r^{2^{k-1}} & j \equiv 1 \pmod{2} \end{cases}$$

- (3) $\langle r^j s \rangle = \begin{cases} \{1, r^j s\} & j \equiv 0 \pmod{2} \\ \{1, r^j s, r^{2^{k-1}}, r^{2^{k-1}+j} s\} & j \equiv 1 \pmod{2} \end{cases}$

In particular when j is odd, the subgroup $\langle r^j s \rangle$ can be expressed in form

- (2) as $\langle r^{2^{k-1}}, r^j s \rangle$.

Now let $G = \mathcal{QD}_{2^k}^{+1}$. The cases (1), (2) follow identically to $\mathcal{QD}_{2^k}^{-1}$.

For case (3), the case $j = 0$ is obvious, for $j > 0$, observe

$$r^j s r^j s = r^j r^{j(2^{k-1}+1)} = r^{j(2^{k-1}+2)} = r^{2j(2^{k-2}+1)}.$$

But $\langle r^{2j(2^{k-2}+1)} \rangle = \langle r^{2j} \rangle$ and hence we have

$$\langle r^j s \rangle = \langle r^{2j}, r^j s \rangle.$$

Setting $2^i || 2j$ be the highest power of 2 dividing j , this subgroup is in fact of type (2) and consists of elements $\{r^a, r^{j'} s : a \equiv 0 \pmod{2^i}, j' \equiv j \pmod{2^i}\}$. \square

Proposition 14. *The quasi dihedral group $\mathcal{QD}_{2^k}^{-1}$ and $\mathcal{QD}_{2^k}^{+1}$ are uniformly abelian group representable for all n .*

Proof. To prove the result we construct a subgroup-preserving bijection from quasi-dihedral to dihedral groups. By Lemma 9 and the fact that dihedral groups are uniformly abelian group representable, the result follows.

Let $G = \mathcal{QD}_{2^k}^{-1}$ or $\mathcal{QD}_{2^k}^{+1}$.

Elements of G can be uniquely written as

$$\{r^i s^j : 0 \leq i < 2^k, j = 0, 1\}.$$

Also all the elements in D_{2^k} can be uniquely written as

$$\{r^i s^j : 0 \leq i < 2^k, j = 0, 1\},$$

keeping in mind that $r, s \in D_{2^k}$ are not the same as $r, s \in G$, as the group law for the groups G and D_{2^k} is not the same.

Define a subgroup preserving bijection

$$\begin{aligned} \psi : G &\rightarrow D_{2^k} \\ \psi : r^i s^j &\mapsto r^i s^j. \end{aligned}$$

The map ψ is well-defined and clearly bijective. It remains to show that ψ is subgroup preserving.

Now Lemma 13 describes which elements subgroups of G consist of. The proof of Proposition 11 gives a similar description for subgroups of D_{2^k} . Verifying that the two coincide via ψ gives the result. As an example, consider a subgroup of form (2). We have

$$\begin{aligned} \psi(\langle r^{2^i}, r^j s \rangle) &= \psi(\{r^a, r^b s : a \equiv 0 \pmod{2^i}, b \equiv j \pmod{2^i}\}) = \\ &= \{r^a, r^b s : a \equiv 0 \pmod{2^i}, b \equiv j \pmod{2^i}\} \leq D_{2^k}. \end{aligned}$$

Cases (1) and (3) follow immediately as well. Hence ψ is a subgroup preserving bijection and since D_{2^k} is abelian group representable by Proposition 12, Lemma 9 implies that $\mathcal{QD}_{2^k}^{-1}$ and $\mathcal{QD}_{2^k}^{+1}$ are uniformly abelian group representable as well. \square

2.2. Dicyclic 2-Groups. Next we consider the case of dicyclic groups, another well studied class of non-abelian groups. The results are similar to that of dihedral groups. A dicyclic group DiC_m of order $4m$ is generated by two elements a, x as follows:

$$DiC_m = \langle a, x : a^{2m} = 1, x^2 = a^m, xa = a^{-1}x \rangle.$$

Every element of DiC_m can be uniquely presented as $a^i x^j$, where $0 \leq i < 2m$, $j = 0, 1$.

A *generalized quaternion group* is a dicyclic group with m a power of 2. We now study the subgroups of DiC_{2^k} . From the definition, we know that $|DiC_{2^k}| = 2^{k+2}$. All the elements of DiC_{2^k} are of the form $a^i, a^i x$, $1 \leq i \leq 2^{k+1}$ where $x^2 = a^{2^k}$ and $x^3 = a^{2^k} x$.

As is the case with dihedral groups, any subgroup $\langle a^{j_1} x, a^{j_2} x \rangle = \langle a^{j_1 - j_2}, a^{j_2} x \rangle$, which in turn can be represented as $\langle a^{2^i}, a^{j_2} x \rangle$, since $\langle a^{j_1 - j_2} \rangle = \langle a^{2^i} \rangle$ for some i . Hence we need at most one generator of the form $a^j x$. Trivially, since $\langle a \rangle$ is cyclic, we only need one generator of the form a^{2^i} as well.

Now consider the subgroup $\langle a^j x \rangle$. Its elements are $\{a^j x, x^2 = a^{2^k}, a^{2^k + j} x, 1\}$, so it can be written in the form $\langle a^{2^k}, a^j x \rangle$.

We conclude that subgroups of DiC_{2^k} are of types

- (1) $\langle a^{2^i} \rangle$, $0 \leq i \leq k$
- (2) $\langle a^{2^i}, a^j x \rangle$, $0 \leq i \leq k$, $0 \leq j \leq 2^i - 1$.

Proposition 15. *The dicyclic group $DiC_{2^{k-1}}$ is uniformly abelian group representable for all n .*

Proof. This proof is an application of Lemma 9, which allows us to use the fact that D_{2^k} was already shown to be abelian group representable, to conclude that so is $DiC_{2^{k-1}}$.

To apply Lemma 9, we must define a subgroup preserving bijection

$$\psi : DiC_{2^{k-1}} \rightarrow D_{2^k}.$$

Since the elements in $DiC_{2^{k-1}}$ can be uniquely written as

$$\{a^i x^j : 0 \leq i < 2^k, j = 0, 1\}$$

while all the elements in D_{2^k} can be uniquely written as

$$\{r^i s^j : 0 \leq i < 2^k, j = 0, 1\},$$

the map

$$\psi : a^i x^j \mapsto r^i s^j$$

is well-defined and clearly bijective. It remains to show that ψ is subgroup preserving. But from the discussion above, we see that the subgroups of $DiC_{2^{k-1}}$ are

- (1) $\langle a^{2^i} \rangle$, $0 \leq i < k$,
- (2) $\langle a^{2^i}, a^j x \rangle$, $0 \leq i < k$, $0 \leq j \leq 2^i - 1$.

The images of both kinds of subgroups under ψ do indeed form subgroups of D_{2^k} :

- (1) $\psi(\langle a^{2^i} \rangle) = \psi(\{a^{2^i h} : h \in \mathbb{Z}\}) = \{r^{2^i h} : h \in \mathbb{Z}\}$,
- (2) $\psi(\langle a^{2^i}, a^j x \rangle) = \psi(\{a^{2^i h}, a^{2^i h+j} x : h \in \mathbb{Z}\}) = \{r^{2^i h}, r^{2^i h+j} s : h \in \mathbb{Z}\}$.

Hence ψ is a subgroup preserving bijection and since by Proposition 12 the dihedral group D_{2^k} is abelian group representable, Lemma 9 implies that $DiC_{2^{k-1}}$ is uniformly abelian group representable as well. \square

Propositions 12 and 15 generalize Proposition 3 of [6].

The results we have obtained on representability of 2-groups rely on the use of a subgroup preserving bijection ψ and Lemma 9. The map ψ defined for dihedral groups in Proposition 11 does not generalize to p -groups, as we can no longer trivially claim closure under inverses. We employ different methods to deal with abelian representability of odd order p -groups in the following section.

3. ABELIAN GROUP REPRESENTABILITY p -GROUPS

We start with a simple lemma which establishes a necessary condition for abelian representability. In the next section this lemma is used to exclude the class of all non-nilpotent groups.

Lemma 16. *Let G be a group which is abelian group representable. Then for any subgroups G_1, G_2 with intersection G_{12} and corresponding indices i_1, i_2, i_{12} in G , it must be the case that*

$$i_{12} \mid i_1 i_2.$$

Proof. Let (A, A_1, A_2) represent (G, G_1, G_2) . Since A is abelian, the subgroups A_1, A_2, A_{12} are all normal in A and the quotient group A/A_{12} is abelian of order i_{12} .

Next note that the subgroups A_1/A_{12} and A_2/A_{12} of A/A_{12} are disjoint and normal. Hence A/A_{12} contains the subgroup $(A_1/A_{12})(A_2/A_{12})$ whose order divides the order of A/A_{12} :

$$|A_1/A_{12}||A_2/A_{12}| \mid i_{12}.$$

Observing that $|A_1/A_{12}| = \frac{|A/A_{12}|}{|A/A_1|}$ we obtain

$$\frac{|A/A_{12}|}{|A/A_1|} \frac{|A/A_{12}|}{|A/A_2|} = \frac{i_{12}}{i_1} \frac{i_{12}}{i_2} \mid i_{12}.$$

Equivalently, $i_{12} \mid i_1 i_2$. □

The following proposition proves abelian representability of p -groups for $n = 2$ by establishing a sufficient condition for $n = 2$ and showing that all p -groups in fact satisfy it.

Proposition 17. *Let G be a p -group. Then G is uniformly abelian group representable for $n = 2$.*

Proof. Let p^m be the order of G . Consider some subgroups $G_1, G_2, G_{12} = G_1 \cap G_2$ of orders p^i, p^j, p^k respectively. We show that the exponents i, j, k, m obey an inequality, which is sufficient to guarantee abelian representability of (G, G_1, G_2) .

Claim 1. Inequality $i + j - k \leq m$ holds for any p -group G .

To that end, consider the subset $S_{G_1}S_{G_2} = \{g_1g_2 \mid g_1 \in G_1, g_2 \in G_2\}$. Note that $S_{G_1}S_{G_2}$ is only a subgroup when one of G_1, G_2 is normal in G . Counting the number of elements in $S_{G_1}S_{G_2}$, we have

$$|S_{G_1}S_{G_2}| = \frac{|G_1||G_2|}{|G_{12}|} = \frac{p^i p^j}{p^k} = p^{i+j-k}.$$

Now since $S_{G_1}S_{G_2} \subseteq G$, we conclude that $i + j - k \leq m$.

Claim 2. Sufficiency of condition $i + j - k \leq m$.

We show that $i + j - k \leq m$ implies that (G, G_1, G_2) can be represented by some abelian (A, A_1, A_2) . Define A to be the elementary abelian p -group $A = C_p^m$.

We can express A as the following direct product:

$$C_p^k \times C_p^{i-k} \times C_p^{j-k} \times C_p^{m-(i+j-k)}.$$

Note that we needed the inequality $i + j - k \leq m$ in order for the exponent $m - (i + j - k)$ to be nonnegative. Now we define subgroups

$$\begin{aligned} A_1 &= C_p^k \times C_p^{i-k} \times \{1\} \times \{1\}, \\ A_2 &= C_p^k \times \{1\} \times C_p^{j-k} \times \{1\}. \end{aligned}$$

As the orders of A, A_1, A_2, A_{12} are same as those of G, G_1, G_2, G_{12} , clearly (A, A_1, A_2) represents (G, G_1, G_2) .

Since the choice of G_1, G_2 was arbitrary, we conclude that G is abelian group representable for $n = 2$. Moreover, G is uniformly abelian group representable, since A was chosen independently of G_1, G_2 . \square

Claim 2 of the above proof establishes a sufficient condition for abelian group representability for $n = 2$, namely the exponents i, j, k, m of orders of G_1, G_2, G_{12}, G obeying the inequality $i + j - k \leq m$. Lemma 16 on the other hand implies that this condition is also necessary: If (G, G_1, G_2) is representable, then $i_{12} \mid i_1 i_2$. But

$$\begin{aligned} i_{12} \mid i_1 i_2 &\iff \frac{p^m}{p^k} \mid \frac{p^m}{p^i} \frac{p^m}{p^j} \iff \frac{p^i p^j}{p^k} \mid p^m \\ &\iff i + j - k \leq m. \end{aligned}$$

Hence we have a numerical necessary and sufficient condition for abelian group representability of (G, G_1, G_2) in terms of exponents i, j, k, m of orders of subgroups G_1, G_2, G_{12}, G :

$$i + j - k \leq m.$$

Certainly we can use similar methods as in the proof of Claim 2, Proposition 17 to guarantee abelian representability whenever a similar ‘‘inclusion-exclusion’’ inequality holds for higher n . It will no longer, however, be a necessary condition. More specifically for $G_1, G_2, G_3 \leq G$ of orders $|G_{\mathcal{A}}| = p^{j_{\mathcal{A}}}$: $\mathcal{A} \subseteq \{1, 2, 3\}$, $|G| = p^m$, the similar inequality on exponents of orders

$$j_1 + j_2 + j_3 - (j_{12} + j_{13} + j_{23}) + j_{123} \leq m$$

while guaranteeing representability of (G, G_1, G_2, G_3) , need no longer hold for groups which are abelian representable. For example, we take the group of quaternions Q_8 with subgroups $\langle i \rangle, \langle j \rangle, \langle k \rangle$, the exponents of whose orders violate the inequality for $n = 3$: $2 + 2 + 2 - (1 + 1 + 1) + 1 = 4 \not\leq 3$.

The following question arises:

Remark 18. *Can we establish a necessary and sufficient numerical condition for abelian group representability for $n > 2$.*

We finish the section by giving a class of uniformly abelian representable p -groups for $n = 3$.

Proposition 19. *If p is a prime, a non-abelian group G of order p^3 is abelian group representable for $n = 3$.*

Proof. We show that the group $C_{p^2} \times C_p$ uniformly represents G . Since G is a non-trivial p -group, it has a non-trivial center, $Z(G)$.

If $|Z(G)| = p^2$, $G/Z(G)$ is cyclic and hence G is abelian. Thus we have $|Z(G)| = p$.

If G_1, G_2 and G_3 are three subgroups of G of order p^2 , then they are normal in G having non-trivial intersection with $Z(G)$, implies that $Z(G) \subseteq G_1, G_2, G_3$ (Theorem 1, Chapter 6, [4]). So the pairwise intersection of all subgroups of G of index p is $Z(G)$. Then the possible combinations of indices of $G_1, G_2, G_3, G_{12}, G_{13}, G_{23}, G_{123}$ are

- (1) $p, p, p, p^2, p^2, p^2, p^2$
- (2) $p, p, p^2, p^2, p^2, p^2, p^2$
- (3) $p, p, p^2, p^2, p^2, p^3, p^3$
- (4) $p, p, p^2, p^2, p^3, p^3, p^3$

- (5) $p, p^2, p^2, p^2, p^2, p^3, p^3$
- (6) $p, p^2, p^2, p^2, p^3, p^3, p^3$
- (7) $p, p^2, p^2, p^3, p^3, p^3, p^3$
- (8) $p^2, p^2, p^2, p^3, p^3, p^3, p^3$

The corresponding abelian group representation is given uniformly by the group $A = C_{p^2} \times C_p$, where $C_{p^2} = \langle g | g^{p^2} = 1 \rangle$, $C_p = \langle r | r^p = 1 \rangle$. For each combination of the indices above, let the subgroups G_1, G_2, G_3 be represented by A_1, A_2, A_3 respectively:

- (1) $A_1 = \langle g^p \rangle \times \langle r \rangle$, $A_2 = \langle g \rangle \times 1$, $A_3 = \langle (g, r) \rangle$, $A_{12} = \langle g^p \rangle \times 1$, $A_{13} = \langle g^p \rangle \times 1$, $A_{23} = \langle g^p \rangle \times 1$, $A_{123} = \langle g^p \rangle \times 1$
- (2) $A_1 = \langle g^p \rangle \times \langle r \rangle$, $A_2 = \langle g \rangle \times 1$, $A_3 = \langle g^p \rangle \times 1$, $A_{12} = \langle g^p \rangle \times 1$, $A_{13} = \langle g^p \rangle \times 1$, $A_{23} = \langle g^p \rangle \times 1$, $A_{123} = \langle g^p \rangle \times 1$
- (3) $A_1 = \langle g^p \rangle \times \langle r \rangle$, $A_2 = \langle g \rangle \times 1$, $A_3 = 1 \times \langle r \rangle$, $A_{12} = \langle g^p \rangle \times 1$, $A_{13} = 1 \times \langle r \rangle$, $A_{23} = 1 \times 1$, $A_{123} = 1 \times 1$
- (4) $A_1 = \langle (g, r) \rangle$, $A_2 = \langle g \rangle \times 1$, $A_3 = 1 \times \langle r \rangle$, $A_{12} = \langle g^p \rangle \times 1$, $A_{13} = 1 \times 1$, $A_{23} = 1 \times 1$, $A_{123} = 1 \times 1$
- (5) $A_1 = \langle g^p \rangle \times \langle r \rangle$, $A_2 = \langle g^p \rangle \times 1$, $A_3 = 1 \times \langle r \rangle$, $A_{12} = \langle g^p \rangle \times 1$, $A_{13} = 1 \times \langle r \rangle$, $A_{23} = 1 \times 1$, $A_{123} = 1 \times 1$
- (6) $A_1 = \langle g \rangle \times 1$, $A_2 = \langle g^p \rangle \times 1$, $A_3 = \langle (g^p, r) \rangle$, $A_{12} = \langle g^p \rangle \times 1$, $A_{13} = 1 \times 1$, $A_{23} = 1 \times 1$, $A_{123} = 1 \times 1$
- (7) $A_1 = \langle (g, r) \rangle$, $A_2 = \langle (g^p, r) \rangle$, $A_3 = 1 \times \langle r \rangle$, $A_{12} = 1 \times 1$, $A_{13} = 1 \times 1$, $A_{23} = 1 \times 1$, $A_{123} = 1 \times 1$
- (8) $A_1 = \langle g^p \rangle \times 1$, $A_2 = 1 \times \langle r \rangle$, $A_3 = \langle (g^p, r) \rangle$, $A_{12} = 1 \times 1$, $A_{13} = 1 \times 1$, $A_{23} = 1 \times 1$, $A_{123} = 1 \times 1$.

We obtain that A, A_1, A_2, A_3 represents G, G_1, G_2, G_3 and since the group A is independent of subgroups G_1, G_2, G_3 , the representation of G is uniform. \square

4. ABELIAN GROUP REPRESENTABILITY OF NILPOTENT GROUPS

A finite nilpotent group is a direct product of its Sylow $_p$ subgroups. In this section we obtain a complete classification of abelian representable groups for $n = 2$, as well as show that non nilpotent groups are never abelian group representable for general n . We start with an easy proposition which allows us to build new abelian representable groups from existing ones.

Proposition 20. *Let G and H be groups of coprime orders. Suppose G, H are abelian group representable, then their direct product $G \times H$ is abelian group representable.*

Proof. Since the orders of G, H are coprime, any subgroup $K_i \leq G \times H$ is in fact a direct product $K_i = G_i \times H_i$, where $G_i \leq G$, $H_i \leq H$. To prove abelian group representability for n , consider n subgroups $G_1 \times H_1, \dots, G_n \times H_n$ of $G \times H$.

Since G is abelian group representable, there exists an abelian group A with subgroups A_1, \dots, A_n which represents (G, G_1, \dots, G_n) . Similarly some (B, B_1, \dots, B_n) represents (H, H_1, \dots, H_n) .

It is easy to see that $(A \times B, A_1 \times B_1, \dots, A_n \times B_n)$ represents $G_1 \times H_1, \dots, G_n \times H_n$. To this end, consider an arbitrary intersection group $G_{\mathcal{A}} \times K_{\mathcal{A}} \leq G \times H$ for $\mathcal{A} \subseteq \{1, \dots, n\}$:

$$\begin{aligned} [G \times K : (G_{\mathcal{A}} \times K_{\mathcal{A}})] &= [G : G_{\mathcal{A}}][H : H_{\mathcal{A}}] = \\ &= [A : A_{\mathcal{A}}][B : B_{\mathcal{A}}] = [A \times B : (A_{\mathcal{A}} \times B_{\mathcal{A}})]. \end{aligned}$$

□

The previous proposition allows us to prove the following theorem.

Theorem 21. *All nilpotent groups are abelian group representable for $n = 2$.*

Proof. We know that finite nilpotent groups are direct products of their Sylow _{p} subgroups. That is, if G is a finite nilpotent group, then $G \cong S_{p_1} \times S_{p_2} \dots \times S_{p_r}$ where S_{p_i} is the Sylow _{p_i} subgroup. Since each S_{p_i} is abelian group representable by Proposition 17, the previous result implies that G is abelian group representable. □

Conversely, we show that the class of abelian representable groups must be contained inside the class of nilpotent groups for all n .

Proposition 22. *A non-nilpotent group G is not abelian group representable for all values of n .*

Proof. Recall that, a group is nilpotent if and only if all of its Sylow subgroups are normal. Since G is by assumption non-nilpotent, at least one of its Sylow _{p} subgroup S_p is not normal for a prime $p \mid r = |G|$. We know that $|S_p| = p^a$, the highest power of p dividing r , that is, $r = p^a m$ with $p \nmid m$. Since S_p is not normal, G has another Sylow _{p} subgroup $S_p^x = \{xS_p x^{-1} \mid x \in G, s \in S_p\}$ of order p^a with $S_p \neq S_p^x$ for some x .

Say their intersection $S_p \cap S_p^x$ is of order p^t . Since S_p and S_p^x are distinct, $a > t$ and thus $[G : S_p \cap S_p^x] = mp^{a-t} > m$. Thus we have subgroups $G_1 = S_p, G_2 = S_p^x, G_{12} = S_p \cap S_p^x$ of indices m, m, mp^{a-t} , respectively, in G . But this contradicts Lemma 16, which implies that $mp^{a-t} \mid m^2$. □

In particular, we have completely classified groups which are abelian group representable for $n = 2$.

Theorem 23. *A group G is abelian group representable for $n = 2$ if and only if it is nilpotent.*

The following corollary generalizes Proposition 4 of [6].

Corollary 24. *If m is not a power of 2, then the dihedral group D_m , the quasi-dihedral groups \mathcal{QD}_m^{-1} and \mathcal{QD}_m^{+1} , and the dicyclic group DiC_m are not abelian group representable for any $n > 1$.*

Proof. Let G_m denote either $D_m, \mathcal{QD}_m^{-1}, \mathcal{QD}_m^{+1}$, or DiC_m . Since subgroups of nilpotent groups are nilpotent, it suffices to show that $G_p \leq G_m$ is not nilpotent for a prime $p \mid m, p \geq 3$.

In case when $G_m = D_m, \mathcal{QD}_m^{-1}, \mathcal{QD}_m^{+1}$ is (quasi-)dihedral, clearly G_p is not nilpotent, as its Sylow₂ subgroup $\{1, s\}$ is not normal. Similarly, a Sylow₂ subgroup $\langle x \rangle \leq DiC_p$ is not normal, which shows that DiC_p is not nilpotent.

By Proposition 22 the result follows. □

5. CONCLUSIONS

In this paper we propose a classification of finite groups with respect to the quasi-uniform variables induced by the subgroup structure. In particular, we study which finite groups belong to the same class as abelian groups with respect to this classification, that is, which finite groups can be represented by abelian groups. We provide an answer to this question when the number n of quasi-uniform variables is 2: it is the class of nilpotent groups. For general n , we show that nilpotent groups are abelian representable if and only p -groups are, while non-nilpotent groups do not afford abelian representation. Hence the question of classifying abelian representable finite groups is completely reduced to answering the question for p -groups.

We demonstrate how some classes of p -groups afford abelian representation for all n , opening various interesting questions for further work. What other classes of p -groups can be shown to be abelian group representable? Is there a generalization of numerical criterion given for $n = 2$ providing a necessary and sufficient condition for abelian representability? It would be extremely interesting to show whether p -groups are indeed abelian group representable. If not - what is the grading with respect to representability within p -groups (and, consequently, nilpotent groups)? Finally, beyond the nilpotent case, the classification of groups with respect to the quasi-uniform distributions is completely open, e.g. - what are the finite groups which induce the same quasi-uniform variables as solvable groups?

REFERENCES

- [1] T.H. Chan, "Aspects of Information Inequalities and its Applications", M.Phil Thesis, Dept. of Information Engineering, The Chinese University of Hong Kong, September 1998.
- [2] T. H. Chan, "Group characterizable entropy functions," *2007 IEEE International Symposium on Information Theory (ISIT 2007)*, Nice, France, 24-29 June 2007.
- [3] T. H. Chan and R. W. Yeung, "On a relation between information inequalities and group theory," *IEEE Trans. on Information Theory*, Vol. 48, pp.1992-1995, July 2002.
- [4] David S.Dummit and Richard M.Foote, "Abstract Algebra (3rd edition) ", *John Wiley and Sons*, 2004.
- [5] B. Hassibi and S. Shadbakht, "Normalized entropy vectors, network information theory and convex optimization," *2007 Information Theory Workshop (ITW 2007)*, July 2007.
- [6] E. Thomas and F. Oggier, "A Note on Quasi-Uniform Distributions and abelian Group Representability", *2012 International Conference on Signal Processing and Communications (SPCOM 2012)*, Bangalore, India, July 21-25, 2012.
- [7] X. Yan, R. Yeung, and Z. Zhang, "The capacity for multi-source multi-sink network coding", *2007 International Symposium on Information Theory (ISIT 2007)*, Nice, France, June 24-29, 2007.

DIVISION OF MATHEMATICAL SCIENCES, SCHOOL OF PHYSICAL AND MATHEMATICAL SCIENCES,
NANYANG TECHNOLOGICAL UNIVERSITY, SINGAPORE.

E-mail address: ELDH01@e.ntu.edu.sg, NMarkin@ntu.edu.sg, frederique@ntu.edu.sg