

# HOMOLOGICAL LOCALISATION OF MODEL CATEGORIES

DAVID BARNES AND CONSTANZE ROITZHEIM

ABSTRACT. One of the most useful methods for studying the stable homotopy category is localising at some spectrum  $E$ . For an arbitrary stable model category we introduce a candidate for the  $E$ -localisation of this model category. We study the properties of this new construction and relate it to some well-known categories.

## INTRODUCTION

The stable homotopy category is spectacularly complicated yet of fundamental importance to homotopy theorists. A standard and highly successful method of dealing with this complexity is to “filter out” some of this information via a Bousfield localisation. In return we obtain a more structured category with useful and interesting patterns.

More precisely, we choose some homology theory  $E_*$  and replace the stable homotopy category  $\mathrm{Ho}(\mathcal{S})$  with  $\mathrm{Ho}(L_E\mathcal{S})$ , the full subcategory of  $\mathrm{Ho}(\mathcal{S})$  with objects the  $E$ -local spectra. This means that in the passage from  $\mathrm{Ho}(\mathcal{S})$  to  $\mathrm{Ho}(L_E\mathcal{S})$ , the  $E_*$ -isomorphisms are formally inverted. Bousfield’s paper [Bou79] is the original source of this idea.

There are a number of other categories that share of the properties  $\mathrm{Ho}(\mathcal{S})$ . It would be advantageous if we could generalise the specialised form of “homological” Bousfield localisation to these other contexts, namely arbitrary stable model categories. A comprehensive general study of Bousfield localisation of model categories is given in [Hir03], but we are specifically interested in the construction of a **homological localisation** of a stable model category.

The main motivation comes again from the study of the stable homotopy category. In order to understand spectra,  $\mathrm{Ho}(\mathcal{S})$  and its various  $E$ -localisations it is necessary to relate  $\mathcal{S}$  and  $L_E\mathcal{S}$  to other stable model categories  $\mathcal{C}$ . For example, one can study to what extent there is a stable model category  $\mathcal{C}$  whose homotopy category “models”  $\mathrm{Ho}(L_E\mathcal{S})$  and how similar  $\mathcal{C}$  is to  $L_E\mathcal{S}$  in terms of higher homotopy behaviour. To make those links it would be a desirable tool to have the corresponding  $E$ -localisations of  $\mathcal{C}$  in order to compare  $E$ -local spectra to other counterparts related to  $\mathcal{C}$ .

A stable model category  $\mathcal{C}$  is a model category whose associated homotopy category  $\mathrm{Ho}(\mathcal{C})$  is triangulated via the construction of [Hov99, Section 7]. Lenhardt proved in [Len12] that  $\mathrm{Ho}(\mathcal{C})$  is a module over  $\mathrm{Ho}(\mathcal{S})$  whenever  $\mathcal{C}$  is a stable model category. Hence we have a tensor product

$$-\wedge^L -: \mathrm{Ho}(\mathcal{C}) \times \mathrm{Ho}(\mathcal{S}) \longrightarrow \mathrm{Ho}(\mathcal{C})$$

and an enrichment of  $\mathrm{Ho}(\mathcal{C})$  in  $\mathrm{Ho}(\mathcal{S})$ . This technique is called **stable frames**.

---

*Date:* 8<sup>th</sup> December 2012.

The first author was supported by EPSRC grant EP/H026681/1, the second author by EPSRC grant EP/G051348/1.

Using this action on the homotopy category of a stable model category one could try to make a new model structure on  $\mathcal{C}$  such that the weak equivalences are the “ $E_*$ -isomorphisms”: those maps  $f: X \rightarrow Y$  in  $\mathcal{C}$  such that

$$f \wedge^L \text{Id}: X \wedge^L E \rightarrow Y \wedge^L E$$

is an isomorphism in  $\text{Ho}(\mathcal{C})$ . Such a model structure would deserve the name  $L_E\mathcal{C}$ . But it seems particularly difficult to check that this construction exists for general  $\mathcal{C}$ . For spectra, the argument appears in [EKMM97, Section VIII.1] and requires numerous unpleasant cardinality arguments.

For well-behaved stable model categories  $\mathcal{C}$  we are going to produce a new model structure  $\mathcal{C}_E$  that avoids such set-theoretic awkwardness. This  $\mathcal{C}_E$  is a good replacement for the  $E$ -localisation of  $\mathcal{C}$  because of the following universal property:  $\mathcal{C}_E$  is the “closest” model category to  $\mathcal{C}$  such that any Quillen adjunction from spectra to  $\mathcal{C}$

$$\mathcal{S} \rightleftarrows \mathcal{C}$$

gives rise to a Quillen adjunction

$$L_E\mathcal{S} \rightleftarrows \mathcal{C}_E$$

from  $E$ -local spectra to  $\mathcal{C}_E$ . We are also able to give another description of  $\mathcal{C}_E$  in terms of pushouts of model categories, which shows how strong the universal property of this new model structure is.

We are also able to give an improvement of [BR11, Theorem 9.5]: we can show that for all  $E$ , the homotopy information of  $E$ -local spectra is entirely encoded in the  $\text{Ho}(\mathcal{S})$ -module structure on the  $E$ -local stable homotopy category. This was previously only possible with the strong restriction that  $E$  is smashing. Hence we have the following, which appears as Theorem 7.1.

**Theorem.** *Let  $\mathcal{C}$  be a stable model category. Assume we have an equivalence of triangulated categories*

$$\Phi: \text{Ho}(L_E\mathcal{S}) \longrightarrow \text{Ho}(\mathcal{C})$$

*then  $L_E\mathcal{S}$  and  $\mathcal{C}$  are Quillen equivalent if and only if  $\Phi$  is an equivalence of  $\text{Ho}(\mathcal{S})$ -model categories.*

**Organisation.** Firstly, we recall some definitions and conventions regarding Bousfield localisation and stable frames. We also re-introduce the concept of **stably  $E$ -familiar model categories**: in [BR11] we studied those  $\mathcal{C}$  such that the action of  $\text{Ho}(\mathcal{S})$  factors over the functor  $\text{Ho}(\mathcal{S}) \rightarrow \text{Ho}(L_E\mathcal{S})$ . In particular the homotopy category of such a model category has an enrichment in the more structured category  $\text{Ho}(L_E\mathcal{S})$ . We called such categories stably  $E$ -familiar.

We then turn to the question of altering a model structure on a given category so as to obtain a stably  $E$ -familiar model category. In Section 3 we consider the simpler case of spectral model categories: such a model category is defined in a similar way to a simplicial model category, but with simplicial sets replaced by the model category of symmetric spectra. We construct the **stable  $E$ -familiarisation** of a spectral model category in this section.

In Section 4 we extend our results to more general stable model categories. We prove that the **stable  $E$ -familiarisation** of a model category  $\mathcal{C}$  is the closest stably  $E$ -familiar

model category to  $\mathcal{C}$  in the following sense. The result below also implies that our construction has the universal property we described earlier.

**Theorem.** *Let  $\mathcal{C}$  be a stable, proper and cellular model category such that the domains of the generating cofibrations of  $\mathcal{C}$  are cofibrant. Then there is a model structure  $\mathcal{C}_E$  on  $\mathcal{C}$  such that*

- (1)  $\mathcal{C}_E$  is stably  $E$ -familiar,
- (2) if  $F: \mathcal{C} \rightarrow \mathcal{E}$  is a left Quillen functor and  $\mathcal{E}$  is stably  $E$ -familiar, then  $F$  factors over  $\mathcal{C} \rightarrow \mathcal{C}_E$ .

Section 5 consists of several examples of  $\mathcal{C}_E$  for some  $E$  and  $\mathcal{C}$  involving algebraic model categories, chromatic localisations and module categories over a ringoid spectrum.

In Section 6 we rephrase the universal property of  $\mathcal{C}_E$  in terms of homotopy pushouts of model categories.

Finally, we prove a full version of the **modular rigidity** theorem that all homotopy information of  $E$ -local spectra is governed by the  $\mathrm{Ho}(\mathcal{S})$ -action on  $\mathrm{Ho}(L_E\mathcal{S})$  given by framings.

## 1. BOUSFIELD LOCALISATION

We begin with an introduction to Bousfield localisation at a homology theory  $E$ . Throughout the paper when we refer to spectra, we mean symmetric spectra [HSS00] unless stated otherwise.

Let  $E$  be a spectrum. Then  $E$  corepresents a homology functor  $E_*$  on the category of spectra via

$$E_*(X) = \pi_*(E \wedge X).$$

Bousfield used this to construct a homotopy category of spectra where maps which induce isomorphisms on  $E_*$ -homology are isomorphisms [Bou79]. We recap some of the definitions from this work. We denote homotopy classes of maps of spectra by  $[-, -]$ .

**Definition 1.1.** *A map  $f: X \rightarrow Y$  of spectra is an  $E$ -equivalence if  $E_*(f)$  is an isomorphism. A spectrum  $Z$  is  $E$ -local if  $f^*: [Y, Z] \rightarrow [X, Z]$  is an isomorphism for all  $E$ -equivalences  $f: X \rightarrow Y$ . A spectrum  $A$  is  $E$ -acyclic if  $[A, Z] = 0$  for all  $E$ -acyclic  $Z$ . An  $E$ -equivalence from  $X$  to an  $E$ -local object  $Z$  is called an  $E$ -localisation.*

Bousfield localisation of spectra gives a homotopy theory that is particularly sensitive towards  $E_*$  and  $E$ -local phenomena. The  $E$ -local homotopy theory is obtained from the category of spectra by formally inverting the  $E$ -equivalences.

This can be seen as a special case of a more general result by Hirschhorn. For  $X, Y \in \mathcal{C}$ , we let  $\mathrm{map}_{\mathcal{C}}(X, Y) \in \mathrm{sSet}$  denote the homotopy function object, see [Hir03, Chapter 17] and Section 2.

**Definition 1.2.** *Let  $S$  be a set of maps in  $\mathcal{C}$ . Then an object  $Z \in \mathcal{C}$  is  $S$ -local if*

$$\mathrm{map}_{\mathcal{C}}(s, Z) : \mathrm{map}_{\mathcal{C}}(B, Z) \longrightarrow \mathrm{map}_{\mathcal{C}}(A, Z)$$

*is a weak equivalence in simplicial sets for any  $s: A \rightarrow B$  in  $S$ . A map  $f: X \rightarrow Y \in \mathcal{C}$  is an  $S$ -equivalence if*

$$\mathrm{map}_{\mathcal{C}}(f, Z) : \mathrm{map}_{\mathcal{C}}(Y, Z) \longrightarrow \mathrm{map}_{\mathcal{C}}(X, Z)$$

is a weak equivalence for any  $S$ -local  $Z \in \mathcal{C}$ . An object  $W \in \mathcal{C}$  is  $S$ -acyclic if

$$\mathrm{map}_{\mathcal{C}}(W, Z) \simeq *$$

for all  $S$ -local  $Z \in \mathcal{C}$ .

A **left Bousfield localisation** of a model category  $\mathcal{C}$  with respect to a class of maps  $S$  is a new model structure  $L_S\mathcal{C}$  on  $\mathcal{C}$  such that

- the weak equivalences of  $L_S\mathcal{C}$  are the  $S$ -equivalences,
- the cofibrations of  $L_S\mathcal{C}$  are the cofibrations of  $\mathcal{C}$ ,
- the fibrations of  $L_S\mathcal{C}$  are those maps that have the right lifting property with respect to cofibrations that are also  $S$ -equivalences.

Hirschhorn proves that if  $S$  is a set and  $\mathcal{C}$  is left proper and cellular then  $L_S\mathcal{C}$  exists. (We will give rough definitions of these two terms below.) In the case of localising spectra at a homology theory one wants to invert the class of  $E_*$ -isomorphisms. Since this is not a set, one cannot use Hirschhorn’s result directly. In [EKMM97, Section VIII.1] it is shown that there is a set  $S$  whose  $S$ -equivalences are exactly the  $E_*$ -isomorphisms. Hence, the key to proving the existence of homological localisations is to find a set giving the correct notion of equivalence. We shall encounter this idea again when constructing  $\mathcal{C}_E$ .

A model category is **left proper** if the pushout of a weak equivalence along a cofibration is a weak equivalence. A model category is **right proper** if the pullback of a weak equivalence along a fibration is a weak equivalence. If a model category is both left and right proper, we say that it is **proper**.

We also need a stronger version of cofibrantly generated, which forces cell complexes to be better behaved. The actual definition is technical and not particularly illuminating, so we shall simply say that a model category is **cellular** if it is cofibrantly generated by sets  $I$  and  $J$ , and the domains and codomains of  $I$  and  $J$  satisfy some nice cardinality conditions. We leave the details to [Hir03, Definition 12.1.1].

## 2. STABLE FRAMINGS

Framings are a powerful tool that describe and classify Quillen functors from simplicial sets or spectra to arbitrary model categories. They were first developed by Hovey in [Hov99, Section 5.2]. For a model category  $\mathcal{C}$ , he investigates cosimplicial and simplicial resolutions of objects in  $\mathcal{C}$ . These are called “frames”. In more detail, a frame of an object  $A \in \mathcal{C}$  is a cofibrant replacement of the constant cosimplicial object  $A \in \mathcal{C}^\Delta$  in the Reedy model category of cosimplicial objects in  $\mathcal{C}$ . From these notions one obtains bifunctors

$$\begin{aligned} - \otimes - &: \mathcal{C} \times \mathrm{sSet} \longrightarrow \mathcal{C}, \\ \mathrm{map}_l(-, -) &: \mathcal{C}^{op} \times \mathcal{C} \longrightarrow \mathrm{sSet}, \\ (-)^{(-)} &: \mathcal{C} \times \mathrm{sSet}^{op} \longrightarrow \mathcal{C}, \\ \mathrm{map}_r(-, -) &: \mathcal{C}^{op} \times \mathcal{C} \longrightarrow \mathrm{sSet} \end{aligned}$$

satisfying certain adjunction properties. The notation  $\otimes$  stems from the fact that  $A \otimes \mathbb{S}^0 \simeq A$ . However, this set-up does not equip  $\mathcal{C}$  with the structure of a simplicial model category because the “mapping spaces”  $\mathrm{map}_l(X, Y)$  and  $\mathrm{map}_r(X, Y)$  only agree up to a zig-zag of weak equivalences for  $X, Y \in \mathcal{C}$  [Hov99, Proposition 5.4.7]. But their derived functors agree, leaving us with the following [Hov99, Theorem 5.5.3].

**Theorem 2.1** (Hovey). *Let  $\mathcal{C}$  be any model category. Then its homotopy category  $\mathrm{Ho}(\mathcal{C})$  is a closed  $\mathrm{Ho}(\mathrm{sSet})$ -module category.*

In particular, this equips any model category with the notion of a homotopy mapping space. Moreover, framings satisfy the following important properties.

- If  $\mathcal{C}$  carries the structure of a simplicial model category [Hov99, Definition 4.2.18], then the two  $\mathrm{Ho}(\mathrm{sSet})$ -module structures coming from either framings or the simplicial structure agree [Hov99, Theorem 5.6.2].
- If  $F : \mathrm{sSet} \rightarrow \mathcal{C}$  is a left Quillen functor with  $F(\mathbb{S}^0) = A$ , then the left derived functors of  $F$  and of the framing functor  $A \otimes - : \mathrm{sSet} \rightarrow \mathcal{C}$  agree. Thus, every left Quillen functor from simplicial sets to any model category can be described, up to homotopy, by a frame.

The second property follows from the fact that the category of cosimplicial objects  $\mathcal{C}^\Delta$  is equivalent to the category of adjunctions  $\mathrm{sSet} \rightleftarrows \mathcal{C}$ . A cosimplicial object  $A^\bullet$  corresponds to a Quillen adjunction under this equivalence if and only if it is a frame, that is  $A^\bullet$  is cofibrant and homotopically constant, [BR11, Proposition 3.2].

In [Len12] Fabian Lenhardt described an analogous set-up for spectra and stable model categories. Now let  $\mathcal{C}$  be a stable model category. First, Lenhardt shows that the category of adjunctions between spectra and a stable model category  $\mathcal{C}$  is equivalent to the category of “ $\Sigma$ -cospectra”  $\mathcal{C}^\Delta(\Sigma)$ . An object in  $\mathcal{C}^\Delta(\Sigma)$  consists of a sequence of cosimplicial objects  $X_n \in \mathcal{C}^\Delta$  together with structure maps

$$\Sigma X_{n+1} \longrightarrow X_n.$$

He then characterises those  $\Sigma$ -cospectra that give rise to Quillen adjunctions under this equivalence, calling them stable frames. These give rise to bifunctors  $-\wedge-$  and  $\mathrm{Map}(-, -)$  satisfying the expected adjunction properties.

As in the unstable case, this is not rigid enough to equip any stable model category  $\mathcal{C}$  with the structure of a spectral model category. However, the above bifunctors give rise to the following [Len12, Theorem 6.3].

**Theorem 2.2** (Lenhardt). *Let  $\mathcal{C}$  be a stable model category. Then  $\mathrm{Ho}(\mathcal{C})$  is a closed  $\mathrm{Ho}(\mathcal{S})$ -module category.*

Note that unlike in the stable case, certain technical subtleties only allow the construction of a  $\mathrm{Ho}(\mathcal{S})$ -module structure on  $\mathrm{Ho}(\mathcal{C})$  rather than a closed module structure.

As expected, this satisfies the following key properties.

- If  $\mathcal{C}$  is already a spectral model category, then the  $\mathrm{Ho}(\mathcal{S})$ -module structure derived from the spectral structure agrees with the  $\mathrm{Ho}(\mathcal{S})$ -module structure coming from stable frames [BR11, Example 6.7].
- By construction, every left Quillen functor  $F : \mathcal{S} \rightarrow \mathcal{C}$  is, up to homotopy, of the form  $X \wedge - : \mathcal{S} \rightarrow \mathcal{C}$  for some fibrant-cofibrant  $X \in \mathcal{C}$ .
- In particular, for any fibrant-cofibrant  $X \in \mathcal{C}$  there is a left Quillen functor  $\mathcal{S} \rightarrow \mathcal{C}$  that sends the sphere spectrum to  $X$ .
- Any stable frame and thus any Quillen functor  $\mathcal{S} \rightarrow \mathcal{C}$  is, up to homotopy, entirely determined by its image on the sphere.

As we have already mentioned, the homotopy theory of  $L_E\mathcal{S}$  is often much better understood than  $\mathcal{S}$ . So it is worth asking if some stable model categories have more in common with  $L_E\mathcal{S}$  than  $\mathcal{S}$ . We answer this question and obtain several useful results using this idea in [BR11]. We give the fundamental definitions below.

**Definition 2.3.** *We say that a stable frame  $X \in \mathcal{C}^\Delta(\Sigma)$  is an  **$E$ -local frame** if it gives rise to a Quillen functor pair*

$$X \wedge - : L_E\mathcal{S} \rightleftarrows \mathcal{C} : \text{Map}(X, -)$$

A stable model category  $\mathcal{C}$  is **stably  $E$ -familiar** if every stable frame is an  $E$ -local frame.

This is [BR11, Definition 7.1]. This generalises the notion of an  $L_E\mathcal{S}$ -model category in the following sense: if  $\mathcal{C}$  is already an  $L_E\mathcal{S}$ -model category, then the  $\text{Ho}(\mathcal{C})$ -module structure on  $\text{Ho}(\mathcal{C})$  agrees with the  $\text{Ho}(L_E\mathcal{S})$ -module structure given by  $E$ -local frames [BR11, Proposition 7.6]. We can further characterise stably  $E$ -familiar model categories as follows [BR11, Theorem 7.8].

**Theorem 2.4.** *Let  $\mathcal{C}$  be a stable model category. Then  $\mathcal{C}$  is stably  $E$ -familiar if and only if the homotopy mapping spectrum  $\mathbb{R}\text{Map}_{\mathcal{C}}(X, Y)$  is an  $E$ -local spectrum for all  $X, Y \in \mathcal{C}$ .*

We can use the theory of  $E$ -local framings to study algebraic model categories. An algebraic model category is a  $\text{Ch}(\mathbb{Z})$ -model category in the sense of [Hov99, Definition 4.2.18]. Thus a  $\text{Ch}(\mathbb{Z})$ -model category is enriched, tensored and cotensored over chain complexes and satisfies the  $\text{Ch}(\mathbb{Z})$ -analogue of the compatibility axiom (SM7). This implies that the homomorphism spectra obtained via framings are products of Eilenberg-MacLane spectra [GJ99, Proposition II.2.20], [DS07, Section 2.6]. Using the computations of Gutiérrez in [Gut10] one can draw the following conclusions [BR11, Section 9].

- For  $n \geq 1$  and there are no algebraic stably  $K(n)$ -familiar model categories, where  $n$  denotes the  $n^{\text{th}}$  Morava- $K$ -theory.
- Let  $E(n)$  denote the  $n^{\text{th}}$  chromatic Johnson-Wilson spectrum. An algebraic model category is stably  $E(n)$ -familiar if and only if it is rational.
- The category  $L_E\mathcal{S}$  of  $E$ -local spectra is algebraic if and only if  $E = H\mathbb{Q}$ .

Now we turn to the question of whether any model category can be made stably  $E$ -familiar in some natural way.

### 3. $E$ -FAMILIARISATION OF SPECTRAL MODEL CATEGORIES

For any homology theory  $E$  we can consider the category of  $E$ -local spectra,  $L_E\mathcal{S}$ . Hence we would like to know if a reasonable notion of  $E$ -localisation exists for an arbitrary stable model category  $\mathcal{C}$ .

Intuitively, a promising definition would be a Bousfield localisation  $L_E\mathcal{C}$  of  $\mathcal{C}$  where one localises at the class of “ $E$ -equivalences” given by

$$\{f : X \rightarrow Y \in \mathcal{C} \mid f \wedge^L E : X \wedge^L E \rightarrow Y \wedge^L E \text{ is a weak equivalence in } \mathcal{C}\}$$

where the action  $\wedge$  of a spectrum on an element of  $\mathcal{C}$  is defined via stable frames. However, showing the existence of Bousfield localisations at a class of maps is set-theoretically awkward. The standard method to circumvent this difficulty is to find a set of maps  $S$  such that the  $S$ -equivalences are precisely the  $E$ -equivalences. This is an extremely difficult

task, see [EKMM97, Section VIII.1], so it is not clear if a good notion of  $E$ -localisation exists for general model categories.

Instead, we will construct the *stable  $E$ -familiarisation* of  $\mathcal{C}$  which is the “closest”  $E$ -familiar model category to  $\mathcal{C}$ . We will then draw some conclusions about its properties which will show that this construction is the right choice for an analogue of  $E$ -localisation for general stable  $\mathcal{C}$ . For example, the first theorem will show that every Quillen adjunction

$$\mathcal{S} \rightleftarrows \mathcal{C}$$

will give rise to a Quillen adjunction

$$L_E\mathcal{S} \rightleftarrows \mathcal{C}_E.$$

The first question to answer is: what kind of maps do we want to invert in order to construct  $\mathcal{C}_E$ ? In a stably  $E$ -familiar model category  $\mathcal{D}$  any map of the form

$$\mathrm{Id} \wedge^L j : X \wedge^L A \rightarrow X \wedge^L B$$

for  $j : A \rightarrow B$  an  $E$ -equivalence of spectra and  $X \in \mathcal{D}$  is a weak equivalence. Hence we could try to localise  $\mathcal{C}$  at this class of maps. So we must find some set of maps  $S$  such that the  $S$ -equivalences equals this class.

We need a couple of technical results first. For this section we shall work with  $\mathcal{S}$ -model categories in the sense of [Hov99, Definition 4.2.18], where  $\mathcal{S}$  again denotes the model category of symmetric spectra. Such a model category  $\mathcal{D}$  is enriched, tensored and cotensored over symmetric spectra in simplicial sets and satisfies the appropriate analogue of Quillen’s SM7 axiom for simplicial model categories. We shall refer to  $\mathcal{D}$  as being a **spectral model category**. We may also talk about  $L_E\mathcal{S}$ -model categories, where we use the  $E$ -local model structure on  $\mathcal{S}$ . A spectral model category is in particular stable and simplicial, see [SS03, Lemma 3.5.2]. We will see later that the restriction to spectral model categories is not as big a restriction as it might seem at first.

We denote the pushout-product of two maps by  $\square$ , so for  $f : X \rightarrow Y$  and  $g : A \rightarrow B$  the pushout-product of  $f$  and  $g$  is

$$f \square g : X \wedge B \coprod_{X \wedge A} Y \wedge A \rightarrow Y \wedge B.$$

Recall that a set of maps  $S$  in a stable model category  $\mathcal{D}$  is said to be **stable** if the class of  $S$ -local objects is closed under suspension. By [BR12, Proposition 3.6] if  $\mathcal{D}$  and  $S$  are stable then so is  $L_S\mathcal{D}$ .

**Proposition 3.1.** *Let  $\mathcal{D}$  be a left proper, cellular and spectral model category. Let  $S$  be a stable set of maps in  $\mathcal{D}$ . Then  $L_S\mathcal{D}$  is also a spectral model category.*

*Proof.* Since  $\mathcal{D}$  is left proper and cellular,  $L_S\mathcal{D}$  exists. We must prove that if  $i$  is a cofibration of  $L_S\mathcal{D}$  and  $j$  is a cofibration of  $\mathcal{S}$  then  $i \square j$  is a cofibration of  $L_S\mathcal{D}$  that is a weak equivalence (in  $L_S\mathcal{D}$ ) if either of  $i$  or  $j$  is. Since  $\mathcal{D}$  is spectral and the cofibrations are unchanged by left Bousfield localisation, we know that  $i \square j$  is a cofibration whenever  $i$  and  $j$  are. Furthermore if  $j$  is an acyclic cofibration of symmetric spectra, then  $i \square j$  is a weak equivalence in  $\mathcal{D}$  and hence it is also an  $S$ -equivalence.

The third case is where  $i$  is an acyclic cofibration of  $L_S\mathcal{D}$  and  $j$  is a cofibration of symmetric spectra. We must show that  $i \square j$  is an  $S$ -equivalence. By [Hov99, Lemma

4.2.4] it suffices to prove this for  $j$  a generating cofibration of symmetric spectra and  $i$  a generating acyclic cofibration of  $L_S\mathcal{D}$ . Hence we may assume that  $j$  has form

$$F_n K \rightarrow F_n L$$

where  $F_n$  is the left adjoint to evaluation at level  $n$ , and  $K$  and  $L$  are simplicial sets. By [Hir03, Proposition 4.5.1] the domain of  $j$  is cofibrant, so it follows that both the domain and codomain of  $i\Box j$  are cofibrant. The set  $S$  is stable, so the class of  $S$ -equivalences in  $\text{Ho}(\mathcal{D})$  is closed under suspension and desuspension. Thus  $i\Box j$  is an  $S$ -equivalence if and only if

$$\Sigma^n(i\Box j) \cong i\Box \Sigma^n j$$

is an  $S$ -equivalence for all  $n$ . Note that we do not need to use the derived functor of  $\Sigma$  in that statement since all the terms are cofibrant.

We know that  $\Sigma^n F_n K$  is weakly equivalent to  $F_0 K$  in  $\mathcal{S}$ . Hence for any cofibrant  $X \in \mathcal{D}$ ,

$$X \wedge \Sigma^n F_n K \rightarrow X \wedge F_0 K$$

is a weak equivalence of  $\mathcal{D}$ . We also know that the domains of the maps  $i\Box \Sigma^n j$  and  $i\Box(F_0 K \rightarrow F_0 L)$  are pushouts of cofibrations between cofibrant objects. It follows that  $i\Box \Sigma^n j$  is weakly equivalent to the map  $i\Box(F_0 K \rightarrow F_0 L)$ . The bifunctor

$$- \wedge F_0 - : \mathcal{D} \times \text{sSet} \rightarrow \mathcal{D}$$

gives  $\mathcal{D}$  the structure of a simplicial model category. We may now use [Hir03, Theorem 4.1.1], which states that since  $\mathcal{D}$  is simplicial, so is  $L_S\mathcal{D}$ . Consequently we see that  $i\Box(F_0 K \rightarrow F_0 L)$  is an  $S$ -equivalence. Hence  $i\Box j$  is also an  $S$ -equivalence and  $L_S\mathcal{D}$  is a spectral model category.  $\square$

**Proposition 3.2.** *Let  $\mathcal{D}$  be a left proper, cellular and spectral model category with generating cofibrations  $I_{\mathcal{D}}$  and generating acyclic cofibrations  $J_{\mathcal{D}}$ . Let  $J_E$  be the set of generating cofibrations for  $L_E\mathcal{S}$ . Define*

$$S = I_{\mathcal{D}}\Box J_E = \{i\Box j \mid i \in I_{\mathcal{D}}, j \in J_E\}.$$

*Then  $L_S\mathcal{D}$  is an  $L_E\mathcal{S}$ -model category and hence is stably  $E$ -familiar.*

*Proof.* The set  $J_E$  is closed under desuspension in the sense that for any element  $j \in J_E$  there is an element  $j'$  with  $\Sigma j' \simeq j$ . It follows that the same holds for  $S$ , so it is stable in the sense of [BR12, Definition 3.2]. Thus  $L_S\mathcal{D}$  is also a stable model category. By Lemma 3.1 it is also a  $\mathcal{S}$ -model category.

To see that it is an  $L_E\mathcal{S}$ -model category we only need check that if  $i$  is a cofibration of  $L_S\mathcal{D}$  and  $j$  is an acyclic cofibration of  $L_E\mathcal{S}$  then  $i\Box j$  is an  $S$ -equivalence. By [Hov99, Lemma 4.2.4] it suffices to prove this for  $i \in I_{\mathcal{D}}$  and  $j \in J_E$ . But then  $i\Box j$  is an element of  $S$  and hence is an  $S$ -equivalence.  $\square$

We now show that this set  $S$  has the correct homotopical behaviour in terms of  $E$ -familiarity by giving another description of the weak equivalences of  $L_S\mathcal{D}$ .

**Proposition 3.3.** *Let  $\mathcal{D}$  be a left proper, cellular, spectral model category, such that the domains of the generating cofibrations of  $\mathcal{D}$  are cofibrant. Let  $T$  be the class of maps*

$$T = \{X \wedge^L f \mid X \in \mathcal{D}, f \text{ is an } E\text{-equivalence of spectra}\}.$$

*Then the class of  $T$ -equivalences is equal to the class of  $S$ -equivalences.*



*Proof.* Take some cofibrant  $X \in \mathcal{D}$ . Then the functor

$$X \wedge - : L_E \mathcal{S} \rightarrow L_S \mathcal{D}$$

is a left Quillen functor by Proposition 3.2. Hence  $X \wedge -$  takes  $E$ -equivalences between cofibrant spectra to  $S$ -equivalences. Thus every element of  $T$  is a weak equivalence in  $L_S \mathcal{D}$ .

Now we will show that every element of  $S$  is also a  $T$ -equivalence. Consider  $i \square j \in S$  for  $i: X \rightarrow Y$  a generating cofibration of  $\mathcal{D}$  and  $j: A \rightarrow B$  a generating acyclic cofibration for  $L_E \mathcal{S}$ . Since  $X, Y, A$  and  $B$  are all cofibrant, the maps  $X \wedge j$  and  $Y \wedge j$  are in the class  $T$ . Let  $P$  be the domain of  $i \square j$ , then by [Hir03, Lemma 3.4.2], the map from  $Y \wedge A \rightarrow P$  is also a  $T$ -equivalence. It follows by the two-out-of-three property that  $i \square j$  is a  $T$ -equivalence.  $\square$

If the category  $\mathcal{C}$  is already stably  $E$ -familiar then the class  $T$  is already contained in the category of weak equivalences. Hence so is the set  $S$ , and  $\mathcal{C}$  is in fact an  $L_E \mathcal{S}$ -model category.

**Corollary 3.4.** *Let  $\mathcal{D}$  be a left proper, cellular, spectral model category that is stably  $E$ -familiar. Assume that domain of the generating cofibrations of  $\mathcal{C}$  are cofibrant. Then  $\mathcal{C}$  is an  $L_E \mathcal{S}$ -model category.  $\square$*

#### 4. $E$ -FAMILIARISATION OF STABLE MODEL CATEGORIES

We now want to consider more model categories that are not necessarily spectral. Consider a proper and cellular stable model category  $\mathcal{C}$ . By [BR12, Theorem 7.2]  $\mathcal{C}$  is Quillen equivalent to a spectral model category, namely the category  $\mathcal{D} = \mathcal{S}^\Sigma(s\mathcal{C})$  of symmetric spectra in simplicial objects in  $\mathcal{C}$  equipped with a non-standard model structure. Hence there is a Quillen equivalence which by abuse of notation we call

$$\Sigma^\infty : \mathcal{C} \rightleftarrows \mathcal{D} = \mathcal{S}^\Sigma(s\mathcal{C}) : \Omega^\infty$$

This model category  $\mathcal{D}$  is also proper and cellular. Furthermore, if the generating cofibrations for  $\mathcal{C}$  have cofibrant domains, then so do the generating cofibrations for  $\mathcal{D}$ .

**Theorem 4.1.** *Let  $\mathcal{C}$  be a stable, proper and cellular model category, such that the domains of the generating cofibrations of  $\mathcal{C}$  are cofibrant. Then define  $\mathcal{C}_E$  to be the left Bousfield localisation of  $\mathcal{C}$  at the set of maps  $\Omega^\infty \hat{f} S$ , where  $\hat{f}$  denotes fibrant replacement in  $\mathcal{S}^\Sigma(s\mathcal{C})$  and where  $S = I_{\mathcal{D}} \square J_E$  as above. Then*

- (1)  $\mathcal{C}_E$  is stably  $E$ -familiar,
- (2) the weak equivalences of  $\mathcal{C}_E$  are the  $T'$ -equivalences, for  $T'$  the class below

$$T' = \{X \wedge^L f \mid X \in \mathcal{C}, f \text{ is an } E\text{-equivalence of spectra}\}$$

- (3) if  $F: \mathcal{C} \rightarrow \mathcal{E}$  is a left Quillen functor and  $\mathcal{E}$  is stably  $E$ -familiar, then  $F$  factors over  $\mathcal{C} \rightarrow \mathcal{C}_E$ .

*Proof.* The model categories  $\mathcal{C}$  and  $\mathcal{D} = \mathcal{S}^\Sigma(s\mathcal{C})$  are Quillen equivalent. Hence [Hir03, Theorem 3.3.20] tell us that the adjunction

$$\Sigma^\infty : \mathcal{C} \rightleftarrows \mathcal{D} : \Omega^\infty.$$

induces a Quillen equivalence between  $L_{\Omega^\infty \hat{f} S} \mathcal{C}$  and  $L_S \mathcal{D}$ . Thus by [BR12, Lemma 7.10]  $\mathcal{C}_E = L_{\Omega^\infty \hat{f} S} \mathcal{C}$  is stably  $E$ -familiar.

We may also conclude that the left derived functor of  $\Sigma^\infty$  induces a bijection between the weak equivalences of  $\mathcal{C}_E$  (considered as a class in  $\text{Ho}\mathcal{C}$ ) and the  $S$ -equivalences of  $\text{Ho}\mathcal{D}$ . Proposition 3.3 tells us that the class of  $S$ -equivalences in  $\mathcal{D}$  is equal to the class of  $T$ -equivalences where

$$T = \{X \wedge^L f \mid X \in \mathcal{D}, f \text{ is an } E\text{-equivalence of spectra}\}$$

Consider the class of maps

$$T' = \{X \wedge^L f \mid X \in \mathcal{C}, f \text{ is an } E\text{-equivalence of spectra}\}$$

Let  $\mathbb{L}\Sigma^\infty$  and  $\mathbb{R}\Omega^\infty$  denote the left and right derived functors of  $\Sigma^\infty$  and  $\Omega^\infty$  respectively. By [Len12, Theorem 6.3]

$$\mathbb{L}\Sigma^\infty(X \wedge^L f) = (\mathbb{L}\Sigma^\infty X) \wedge^L f.$$

Hence  $\mathbb{L}\Sigma^\infty$  takes elements of  $T'$  to elements of  $T$ . Consider some element  $Y \wedge^L f$  of  $T$ . This is weakly equivalent to

$$(\mathbb{L}\Sigma^\infty \mathbb{R}\Omega^\infty Y) \wedge^L f$$

and hence is in  $\mathbb{L}\Sigma^\infty T'$ . Thus the derived functor of  $\Sigma^\infty$  induces a bijection between the class  $T'$  and the class  $T$  up to weak equivalence. As a consequence the derived functor of  $\Sigma^\infty$  induces a bijection between the class of  $T'$ -equivalences and the class of  $T$ -equivalences. It follows that the  $T'$ -equivalences must be the class of weak equivalences of  $\mathcal{C}_E$ .

For the final point, let  $F: \mathcal{C} \rightarrow \mathcal{E}$  be a left Quillen functor. If  $\mathcal{E}$  is stably  $E$ -familiar, then the left derived functor of  $F$  takes the  $T'$ -equivalences to weak equivalences of  $\mathcal{E}$ . Hence  $F: \mathcal{C}_E \rightarrow \mathcal{E}$  is also a left Quillen functor.  $\square$

**Corollary 4.2.** *Let  $\mathcal{C}$  be a stable, proper and cellular model category, such that the domains of the generating cofibrations of  $\mathcal{C}$  are cofibrant. Then  $\mathcal{C}_E$  is the closest stably  $E$ -familiar model category to  $\mathcal{C}$ .*  $\square$

In particular a model category  $\mathcal{C}$  is stably  $E$ -familiar if and only if  $\mathcal{C}_E = \mathcal{C}$ .

**Remark 4.3.** *The assumptions on  $\mathcal{C}$  are more reasonable than they might seem in practice. Since we want to perform a left Bousfield localisation, we will have to assume that  $\mathcal{C}$  is left proper and cellular. To assume that  $\mathcal{C}$  is also right proper is not too much of a restriction on the kinds of model categories we are able to deal with.*

*We also need another assumption: that the domains of the generating cofibrations of  $\mathcal{C}$  are cofibrant. This is a subtle assumption that occurs elsewhere in the literature, for example in [Hov01]. We note that this assumption holds for almost all of the cofibrantly generated model categories that arise naturally.*

It is easy to check that the homotopy mapping spectra for  $\mathcal{C}_E$  are given by the formula below, where  $Y_E$  is the fibrant replacement of  $Y$  in  $\mathcal{C}_E$ .

$$\mathbb{R}\text{Map}_{\mathcal{C}_E}(X, Y) = \mathbb{R}\text{Map}_{\mathcal{C}}(X, Y_E)$$

In particular, this mapping spectrum is  $E$ -local. We can use this to draw some immediate consequences of  $E$ -familiarisation.

For example, the chromatic Johnson-Wilson theories  $E(n)$  satisfy

$$L_{E(n-1)}L_{E(n)} = L_{E(n)}$$

[Rav92, 7.5.3]. Thus,

**Corollary 4.4.** *For a proper and cellular stable model category we have*

$$(\mathcal{C}_{E(n)})_{E(n-1)} = \mathcal{C}_{E(n-1)}.$$

□

We can further use our knowledge of stably  $E$ -familiar algebraic model categories described at the end of Section 2 to read off the following corollaries.

**Corollary 4.5.** *Let  $\mathcal{C}$  be an algebraic model category and  $K(n)$  the  $n^{\text{th}}$  Morava- $K$ -theory for  $n \geq 1$ . Then  $\mathcal{C}_{K(n)}$  is trivial.* □

**Corollary 4.6.** *Let  $\mathcal{C}$  be an algebraic model category and let  $E(n)$  denote the  $n^{\text{th}}$  chromatic Johnson-Wilson spectrum. Then  $\mathcal{C}_{E(n)} = \mathcal{C}_{H\mathbb{Q}}$ .* □

If we assume that localisation at  $E$  is smashing, we can obtain a nicer description of the weak equivalences of  $\mathcal{C}_E$ : in the smashing case  $\mathcal{C}_E$  is precisely the “naive” localisation of  $\mathcal{C}$  at  $L_E S^0$  as motivated in the introduction of this section. Thus, we also obtain that

$$\mathcal{C}_E = \mathcal{C}_{L_E S^0}.$$

However, for a general model category  $\mathcal{C}$  and smashing  $E$  it is unclear whether this also implies that the model categories  $L_E \mathcal{C}$  and  $L_{L_E S^0} \mathcal{C}$  are Quillen equivalent.

**Lemma 4.7.** *In addition to the assumptions of Theorem 4.1, assume that localisation at  $E$  is smashing. Then a map  $f$  in  $\mathcal{C}_E$  is a weak equivalence if and only if  $f \wedge^L L_E S^0$  is a weak equivalence in  $\mathcal{C}$ . Hence the weak equivalences of  $\mathcal{C}_E$  are precisely the  $L_E S^0$ -equivalences.*

*Proof.* We first show the statement for a spectral model category  $\mathcal{D}$ . Recall the model category  $L_S \mathcal{D}$  for  $S$  the set  $I_{\mathcal{C}} \square J_E$  from the previous section. We will show that the  $S$ -equivalences are precisely the  $L_E S^0$ -equivalences of  $\mathcal{D}$ .

Every map in the set  $S$  is an  $L_E S^0$ -equivalence, hence every  $S$ -equivalence is a  $L_E S^0$ -equivalence. Now take some  $L_E S^0$ -equivalence  $f: X \rightarrow Y$  in  $\mathcal{D}$ . The map

$$X \rightarrow X \wedge^L L_E S^0$$

is a  $T$ -equivalence and hence an  $S$ -equivalence. Thus the commutative square

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ \downarrow & & \downarrow \\ X \wedge^L L_E S^0 & \xrightarrow{f \wedge^L L_E S^0} & Y \wedge^L L_E S^0 \end{array}$$

shows that  $f$  is  $S$ -equivalent to a weak equivalence in  $\mathcal{D}$ . Weak equivalences in  $\mathcal{D}$  are in particular  $S$ -equivalences, so by the 2-out-of-3 axiom of model categories,  $f$  must be an  $S$ -equivalence.

To move this result from  $\mathcal{D}$  to  $\mathcal{C}$  we use a similar argument to that of the second point of Theorem 4.1. The Quillen equivalence  $(\Sigma^\infty, \Omega^\infty)$  takes the  $L_E S^0$ -equivalences of  $\mathcal{D}$  bijectively to the  $L_E S^0$ -equivalences of  $\mathcal{C}$ . It follows that the  $L_E S^0$ -equivalences of  $\mathcal{C}$  are precisely the weak equivalences of  $\mathcal{C}_E$ . □

The following corollary shows that stable  $E$ -familiarisation restricts to  $E$ -localisation in the case of spectra. This shows that the notion of  $\mathcal{C}_E$  is indeed a good candidate for an analogue of  $E$ -localisation of a general  $\mathcal{C}$ .

**Corollary 4.8.** *Consider the category of modules over a ring spectrum  $R$ . Then*

$$(R\text{-mod})_E = L_E(R\text{-mod})$$

and in particular

$$\mathcal{S}_E = L_E\mathcal{S}.$$

□

We can now give a simple proof that stable  $E$ -familiarisation preserves Quillen equivalences.

**Proposition 4.9.** *Let  $\mathcal{C}$  and  $\mathcal{E}$  be proper, cellular and stable model categories such that the domains of their generating cofibrations are cofibrant. Let*

$$F : \mathcal{C} \rightleftarrows \mathcal{E} : G$$

be a Quillen equivalence. Then there is a Quillen equivalence between the  $E$ -familiarised model categories

$$F : \mathcal{C}_E \rightleftarrows \mathcal{E}_E : G.$$

*Proof.* Composing  $F$  with the identity on  $\mathcal{E}$  gives us a left Quillen functor

$$F : \mathcal{C} \rightarrow \mathcal{E}_E$$

and  $\mathcal{E}_E$  is of course stably  $E$ -familiar. Hence by the universal property of  $\mathcal{C}_E$  proved in Theorem 4.1 we have a left Quillen functor  $F : \mathcal{C}_E \rightarrow \mathcal{E}_E$ . We now need to show that gives us a Quillen equivalence. We do so using Proposition 3.3 and the method of the second part of the proof of Theorem 4.1.

Let  $T$  be the class of maps

$$T = \{A \wedge^L f \mid A \in \mathcal{C}, f \text{ is an } E\text{-equivalence of spectra}\}.$$

Similarly, let  $T'$  be the class of maps

$$T' = \{B \wedge^L f \mid B \in \mathcal{E}, f \text{ is an } E\text{-equivalence of spectra}\}.$$

Then  $\mathcal{C}_E = L_T\mathcal{C}$  and  $\mathcal{E}_E = L_{T'}\mathcal{E}$ . Let  $\mathbb{L}F$  and  $\mathbb{R}G$  denote the left and right derived functors of  $F$  and  $G$  respectively. By [Hir03, Theorem 3.3.20], the adjunction  $(F, G)$  induces a Quillen equivalence between  $L_T\mathcal{C}$  and  $L_{\mathbb{L}F(T)}\mathcal{E}$ . But the set  $\mathbb{L}F(T)$  is isomorphic in  $\text{Ho}\mathcal{E}$  to the set  $T'$  because Quillen equivalences induce equivalences of  $\text{Ho}(\mathcal{S})$ -module categories [Len12, Theorem 6.3]. □

## 5. EXAMPLES

Let  $\mathcal{C}$  be a spectral model category, such that the domains of its generating cofibrations are cofibrant. (Recall from [BR12, Theorem 7.2] that any stable, proper and cellular model category is Quillen equivalent to a spectral one.) Assume that  $\mathcal{C}$  has a set of compact generators for its homotopy category, [SS03, Definition 2.1.2]. Schwede and Shipley prove in the above-mentioned paper that any such  $\mathcal{C}$  is Quillen equivalent to a category  $\text{mod-}\mathcal{E}$  where  $\mathcal{E}$  can be thought of as a “ring spectrum with several objects”. In the case of  $\mathcal{C}$  having a single compact generator,  $\mathcal{E}$  is simply a ring spectrum.

Let us briefly recap some of the definitions and constructions of that result. Let  $\mathcal{G}$  denote the set of generators of  $\mathcal{C}$ . Then the  $\mathcal{S}$ -enriched category  $\mathcal{E}$  is simply defined as

the full  $\mathcal{S}$ -enriched subcategory of  $\mathcal{C}$  with objects  $\mathcal{G}$ . An object  $M \in \text{mod-}\mathcal{E}$  consists of a spectrum  $M(G)$  for each  $G \in \mathcal{G}$  plus morphisms of spectra

$$M(G) \wedge \mathcal{E}(G', G) \longrightarrow M(G') \text{ for } G, G' \in \mathcal{G}$$

satisfying certain coherence conditions. By adjunction, such an  $M$  is the same as a contravariant spectral functor from  $\mathcal{E}$  to  $\mathcal{S}$ . The model structure on  $\text{mod-}\mathcal{E}$  has weak equivalences and fibrations defined objectwise [SS03, Theorem A.1.1]: meaning that a natural transformation  $f : M \longrightarrow N$  is a weak equivalence or a fibration if and only if

$$f_G : M(G) \longrightarrow N(G)$$

is so for each  $G \in \mathcal{G}$ . Theorem 3.9.3 of [SS03] then describes a Quillen equivalence

$$\text{Hom}(\mathcal{G}, -) : \mathcal{C} \xleftrightarrow{\quad} \text{mod-}\mathcal{E} : - \wedge_{\mathcal{E}} \mathcal{G}$$

for spectral  $\mathcal{C}$ .

This is a highly useful description of a stable model category and we would like to obtain a description of the  $E$ -familiarisation  $\mathcal{C}_E$  of  $\mathcal{C}$  in terms of  $\text{mod-}\mathcal{E}$ . We note that this is a rather special case as not every stable model category has a set of compact generators [HS99, Corollary B.13].

By Proposition 4.9 we know that  $\mathcal{C}_E$  and  $(\text{mod-}\mathcal{E})_E$  are Quillen equivalent, so we shall find another description of  $(\text{mod-}\mathcal{E})_E$ .

The generating cofibrations of this model category have the form

$$\text{hom}(-, g) \wedge (i \square j)$$

where  $g$  is a cofibrant and fibrant replacement of one of the compact generators for  $\mathcal{C}$ ,  $i$  is a generating cofibration for  $\mathcal{S}$  and  $j$  is a generating acyclic cofibration for  $L_E\mathcal{S}$ .

We can make another model structure on  $\text{mod-}\mathcal{E}$  by taking the same cofibrations as before, but taking the generating set of acyclic cofibrations to be those maps of form

$$\text{hom}(-, g) \wedge j$$

for  $g$  a generator and  $j$  a generating acyclic cofibration for  $L_E\mathcal{S}$ . We shall call this set of maps  $K$ . One can either check directly that these sets give a model structure or one can alter [SS03, Theorem A.1.1] to use  $L_E\mathcal{S}$  instead of  $\mathcal{S}$ .

We claim that this model structure equals the model structure of  $(\text{mod-}\mathcal{E})_E$ . An element of  $K$  can be described as

$$\text{hom}(-, g) \wedge ((* \rightarrow S^0) \square j)$$

Hence every element of  $K$  is an acyclic cofibration of  $(\text{mod-}\mathcal{E})_E$ . Conversely, since  $E$ -acyclic cofibrations of spectra are closed under pushout along cofibrations, it follows that the set of acyclic cofibrations generated by the set  $K$  is precisely the set of acyclic cofibrations of  $(\text{mod-}\mathcal{E})_E$ . Thus we have shown the following.

**Proposition 5.1.** *The model category  $(\text{mod-}\mathcal{E})_E$  is the category of contravariant spectral functors from  $\mathcal{E}$  to  $L_E\mathcal{S}$ , equipped with the model structure where fibrations and weak equivalences are defined objectwise. Thus the fibrant objects are those functors  $M$  such that  $M(g)$  is fibrant in  $\mathcal{S}$  and  $E$ -local for all  $g \in \mathcal{E}$ .  $\square$*

Consider the case where  $\mathcal{C}$  has a single compact generator. Following the above we can replace this by a category of functors to  $\mathcal{S}$ . Indeed, [SS03, Theorem 3.1.1] states that  $\mathcal{C}$  is Quillen equivalent to the category of  $R$ -modules,  $\text{mod-}R$ , for some ring spectrum  $R$ . Hence we are essentially in the same situation as Corollary 4.8. Our work above recovers the well-known result that  $(\text{mod-}R)_E$  is the category of  $R$ -modules, with weak equivalences the underlying  $E$ -equivalences of spectra.

## 6. $E$ -FAMILIARISATION AND HOMOTOPY PUSHOUTS

We want to give another description of  $\mathcal{C}_E$  via a universal property. We will relate  $\mathcal{C}_E$  to a pushout of model categories. While the pullback of model categories is well-understood, [Ber11], the pushout is more complicated and is not often used. Roughly speaking, the homotopy pushout of a corner diagram of Quillen adjunctions

$$\mathcal{C} \begin{array}{c} \longleftarrow \\ \rightleftarrows \\ \longrightarrow \end{array} \mathcal{D} \begin{array}{c} \longrightarrow \\ \rightleftarrows \\ \longleftarrow \end{array} \mathcal{E}$$

is supposed to be a model category  $\mathcal{P}$  that satisfies a universal property analogous to the pushout of a diagram within a category. Unfortunately, the homotopy-theoretic pushout construction is rather delicate and its existence and description not always clear.

However there is a special case where we can construct pushouts of model categories and verify that they have the correct universal property. By working in a particular context, we avoid the general question of whether homotopy pushouts of model categories exist in general.

Let  $\mathcal{M}_2$  be a left Bousfield localisation of  $\mathcal{M}_1$  at a class of maps  $W$ . Without loss of generality we assume that the maps in  $W$  are morphisms between cofibrant objects. In particular, this gives us a Quillen pair

$$\text{Id} : \mathcal{M}_1 \begin{array}{c} \longleftarrow \\ \rightleftarrows \\ \longrightarrow \end{array} \mathcal{M}_2 = L_W \mathcal{M}_1 : \text{Id}$$

Assume that we have a Quillen adjunction

$$F : \mathcal{M}_1 \begin{array}{c} \longrightarrow \\ \rightleftarrows \\ \longleftarrow \end{array} \mathcal{N}_1 : G$$

We are now going to discuss the homotopy pushout of the corner diagram below for this special case

$$L_W \mathcal{M}_1 = \mathcal{M}_2 \begin{array}{c} \longleftarrow \\ \rightleftarrows \\ \longrightarrow \end{array} \mathcal{M}_1 \begin{array}{c} \longrightarrow \\ \rightleftarrows \\ \longleftarrow \end{array} \mathcal{N}_1.$$

**Definition 6.1.** *The homotopy pushout of the above diagram is defined, if it exists, as the Bousfield localisation  $L_{FW} \mathcal{N}_1$  of  $\mathcal{N}_1$ .*


To justify this definition we need to see that  $\mathcal{N}_2 = L_{FW} \mathcal{N}_1$  (provided it exists) satisfies the desired properties that a homotopy pushout is supposed to have. First we note that by [Hir03, Theorem 3.3.20]  $F$  and  $G$  induce a Quillen adjunction

$$F : \mathcal{M}_2 \begin{array}{c} \longrightarrow \\ \rightleftarrows \\ \longleftarrow \end{array} \mathcal{N}_2 : G$$


Assume that there is a model category  $\mathcal{D}$  with Quillen adjunctions

$$\begin{array}{ccc} \mathcal{M}_2 & \begin{array}{c} \longrightarrow \\ \rightleftarrows \\ \longleftarrow \end{array} & \mathcal{D} \\ F' : \mathcal{N}_1 & \begin{array}{c} \longrightarrow \\ \rightleftarrows \\ \longleftarrow \end{array} & \mathcal{D} : G' \end{array}$$

such that in the diagram below, the two different composites of left adjoints from  $\mathcal{M}_1$  to  $\mathcal{D}$  agree up to natural isomorphism.



Because the vertical functors in the square below are simply identity functors it follows immediately that we may add  $\mathcal{N}_2$  and obtain a commutative diagram of adjoint pairs.



We must check that the adjunction below is Quillen adjunction.

$$F' : \mathcal{N}_2 \rightleftarrows \mathcal{D} : G'$$

The model category  $\mathcal{N}_2$  is the Bousfield localisation of  $\mathcal{N}_1$  with respect to the class of maps  $Ff$  where  $f$  is a weak equivalence between cofibrant objects of  $\mathcal{M}_2$ . Thus  $(F' \circ F)(f)$  is a weak equivalence in  $\mathcal{D}$ . This means that  $F'$  uniquely factors over  $\mathcal{N}_2$ . Furthermore, by construction,  $\mathcal{N}_2$ , if it exists, is unique up to Quillen equivalence.

Recall that the stable  $E$ -familiarisation  $\mathcal{C}_E$  satisfies the following universal property. Given a left Quillen functor  $F : \mathcal{C} \rightarrow \mathcal{D}$  with  $\mathcal{D}$  stably  $E$ -familiar,  $F$  also gives rise to a left Quillen functor  $\mathcal{C}_E \rightarrow \mathcal{D}$  via

$$\begin{array}{ccc} \mathcal{C} & \xrightarrow{F} & \mathcal{D} \\ \text{id} \downarrow & \nearrow & \\ \mathcal{C}_E & & \end{array}$$

This fact allows us to relate  $\mathcal{C}_E$  and certain homotopy pushouts. Let  $X \in \mathcal{C}$  be fibrant and cofibrant. Then we have a Quillen adjunction

$$X \wedge - : \mathcal{S} \rightleftarrows \mathcal{C} : \text{Map}(X, -).$$

Using Definition 6.1 we can read off the following for a proper and cellular stable model category  $\mathcal{C}$ .

**Lemma 6.2.** *The homotopy pushout  $\mathcal{P}_X$  of the diagram*

$$L_E \mathcal{S} \rightleftarrows \mathcal{S} \rightleftarrows \mathcal{C}$$

exists and is the Bousfield localisation of  $\mathcal{C}$  with respect to the set of maps below, where  $J_E$  is the set of generating acyclic cofibrations of  $L_E\mathcal{S}$ .

$$X \wedge^L J_E = \{X \wedge^L j \mid j \in J_E\}$$

□

So in particular we know that this homotopy pushout exists. Because  $\mathcal{C}_E$  is stably  $E$ -familiar we have a commutative square of Quillen adjunctions

$$\begin{array}{ccc} \mathcal{S} & \rightleftarrows & \mathcal{C} \\ \updownarrow & & \updownarrow \\ L_E\mathcal{S} & \rightleftarrows & \mathcal{C}_E \end{array}$$

By the universal property of  $\mathcal{P}_X$ , there is a Quillen adjunction  $\mathcal{P}_X \rightleftarrows \mathcal{C}_E$  for each  $X$ . We can show that  $\mathcal{C}_E$  is the “closest” model category to those pushouts in the following sense.

**Theorem 6.3.** *The Quillen adjunction*

$$\mathcal{C} \rightleftarrows \mathcal{C}_E$$

*factors over*

$$\mathcal{P}_X \rightleftarrows \mathcal{C}_E$$

*for all fibrant-cofibrant  $X \in \mathcal{C}$ . If there is any other stable  $\mathcal{D}$  with a Quillen adjunction*

$$F : \mathcal{C} \rightleftarrows \mathcal{D} : G$$

*that factors over*

$$\mathcal{P}_X \rightleftarrows \mathcal{D}$$

*for all fibrant-cofibrant  $X$ , then  $(F, G)$  also factors over  $\mathcal{C}_E$ .*

*Proof.* The pushout  $\mathcal{P}_X$  is defined as the Bousfield localisation of  $\mathcal{C}$  at the set of maps  $X \wedge j$  with  $j \in J_E$ . By Proposition 3.3 we know that  $\mathcal{C}_E$  is the localisation of  $\mathcal{C}$  at the class of maps of form  $X \wedge f$  for  $f$  and  $E$ -equivalence of spectra. Thus we see that for every  $X \in \mathcal{C}$  the identity gives us a Quillen adjunction

$$\text{Id} : \mathcal{P}_X \rightleftarrows \mathcal{C}_E : \text{Id}$$

because every weak equivalence in  $\mathcal{P}_X$  is also a weak equivalence in  $\mathcal{C}_E$ .

If the given Quillen adjunction  $(F, G)$  induces a Quillen adjunction

$$F : \mathcal{P}_X \rightleftarrows \mathcal{D} : G,$$

then  $F$  sends all morphisms of the form  $X \wedge^L j$  to weak equivalences in  $\mathcal{D}$ . Thus it induces a Quillen adjunction

$$\mathcal{C}_E \rightleftarrows \mathcal{D}$$

which is what we wanted to prove. □



7. MODULAR RIGIDITY FOR  $E$ -LOCAL SPECTRA

We can show that stable frames encode all homotopical information of the  $E$ -local stable homotopy category. The triangulated structure of  $\mathrm{Ho}(L_E\mathcal{S})$  alone is not sufficient for this: given just a triangulated equivalence

$$\Phi : \mathrm{Ho}(L_E\mathcal{S}) \longrightarrow \mathrm{Ho}(\mathcal{C})$$

for a stable model category  $\mathcal{C}$  does not imply in general that  $L_E\mathcal{S}$  and  $\mathcal{C}$  are Quillen equivalent. In fact, Quillen equivalence can only be deduced from a triangulated equivalence of homotopy categories in some very special cases. To this date, the only nontrivial cases known of this ‘rigidity’ are the stable homotopy category itself [Sch07] and the case  $E = K_{(2)}$  [Roi07]. However, if do not only have a triangulated equivalence as above but also assume that this equivalence is a  $\mathrm{Ho}(\mathcal{S})$ -module equivalence, we can show that  $L_E\mathcal{S}$  and  $\mathcal{C}$  are Quillen equivalent.

For the case of  $E$  smashing, the result below appeared as [BR11, Theorem 9.5], but with a slight modification it would also work for all  $E$  such that  $\mathrm{Ho}(L_E\mathcal{S})$  has one compact generator, e.g.  $E = K(n)$ . The proof relied on the result of Schwede and Shipley [SS03, Theorem 3.1.1] that every stable model category  $\mathcal{C}$  such that  $\mathrm{Ho}(\mathcal{C})$  has one compact generator is Quillen equivalent to the category of modules over some ring spectrum. However, for some  $E$  the  $E$ -local stable homotopy category does not possess any compact objects at all [HS99, Corollary B.13], let alone a set of compact generators. But in fact a more general result is true.

**Theorem 7.1.** *Let  $\mathcal{C}$  be a stable model category. Assume we have an equivalence of triangulated categories*

$$\Phi : \mathrm{Ho}(L_E\mathcal{S}) \longrightarrow \mathrm{Ho}(\mathcal{C})$$

*then  $L_E\mathcal{S}$  and  $\mathcal{C}$  are Quillen equivalent if and only if  $\Phi$  is an equivalence of  $\mathrm{Ho}(\mathcal{S})$ -model categories.*

*Proof.* The “only if” part is true by [Len12, Theorem 6.3]: a Quillen equivalence induces a  $\mathrm{Ho}(\mathcal{S})$ -module equivalence.

Now let us assume that we have a  $\mathrm{Ho}(\mathcal{S})$ -module equivalence

$$\Phi : \mathrm{Ho}(L_E\mathcal{S}) \longrightarrow \mathrm{Ho}(\mathcal{C})$$

It follows that  $\Phi^{-1}$  induces a weak equivalence of homotopy mapping spectra

$$\Phi^{-1} : \mathbb{R}\mathrm{Map}_{\mathcal{C}}(X, Y) \longrightarrow \mathbb{R}\mathrm{Map}_{\mathcal{C}}(\Phi^{-1}X, \Phi^{-1}Y)$$

for  $X, Y \in \mathcal{C}$ . The right-hand-side is an  $E$ -local spectrum as  $L_E\mathcal{S}$  is stably  $E$ -familiar. Hence every homotopy mapping spectrum of  $\mathcal{C}$  is  $E$ -local, so  $\mathcal{C}$  is stably  $E$ -familiar by [BR11, Theorem 7.8].

Thus for fibrant and cofibrant  $X \in \mathcal{C}$ , the Quillen functor

$$X \wedge - : \mathcal{S} \longrightarrow \mathcal{C}$$

factors over  $L_E\mathcal{S}$  as a Quillen functor

$$X \wedge - : L_E\mathcal{S} \longrightarrow \mathcal{C}.$$

Now we consider  $X$  a cofibrant-fibrant replacement of  $\Phi(S^0)$ . Because  $\Phi$  is a  $\text{Ho}(\mathcal{S})$ -module equivalence we see that

$$X \wedge^L (-) = \Phi(L_E S^0) \wedge^L (-) = \Phi(L_E S^0 \wedge^L -) = \Phi(-).$$

This means that  $\Phi$  is derived from a Quillen functor. This Quillen functor must therefore be a Quillen equivalence, which is what we wanted to prove.  $\square$

## REFERENCES

- [Ber11] J.E. Bergner. Homotopy fiber products of homotopy theories. *Israel J. Math.*, 185:389–411, 2011.
- [Bou79] A. K. Bousfield. The localization of spectra with respect to homology. *Topology*, 18(4):257–281, 1979.
- [BR11] D. Barnes and C. Roitzheim. Local framings. *New York J. Math.*, 17:513–552, 2011.
- [BR12] D. Barnes and C. Roitzheim. Stable left and right bousfield localisations. To appear in *Glasgow Mathematical Journal*, 2012.
- [DS07] D. Dugger and B. Shipley. Enriched model categories and an application to additive endomorphism spectra. *Theory Appl. Categ.*, 18:400–439 (electronic), 2007.
- [EKMM97] A. D. Elmendorf, I. Kriz, M. A. Mandell, and J. P. May. *Rings, modules, and algebras in stable homotopy theory*, volume 47 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 1997. With an appendix by M. Cole.
- [GJ99] P.G. Goerss and J.F. Jardine. *Simplicial homotopy theory*, volume 174 of *Progress in Mathematics*. Birkhäuser Verlag, Basel, 1999.
- [Gut10] J.J. Gutiérrez. Homological localizations of Eilenberg-MacLane spectra. *Forum Math.*, 22(2):349–356, 2010.
- [Hir03] P. S. Hirschhorn. *Model categories and their localizations*, volume 99 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2003.
- [Hov99] M. Hovey. *Model categories*, volume 63 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 1999.
- [Hov01] M. Hovey. Spectra and symmetric spectra in general model categories. *J. Pure Appl. Algebra*, 165(1):63–127, 2001.
- [HS99] M. Hovey and N.P. Strickland. Morava  $K$ -theories and localisation. *Mem. Amer. Math. Soc.*, 139(666):viii+100, 1999.
- [HSS00] M. Hovey, B. Shipley, and J. Smith. Symmetric spectra. *J. Amer. Math. Soc.*, 13(1):149–208, 2000.
- [Len12] F. Lenhardt. Stable frames in model categories. *J. Pure Appl. Algebra*, 216(5):1080–1091, 2012.
- [Rav92] D.C. Ravenel. *Nilpotence and periodicity in stable homotopy theory*, volume 128 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, 1992. Appendix C by Jeff Smith.
- [Roi07] C. Roitzheim. Rigidity and exotic models for the  $K$ -local stable homotopy category. *Geom. Topol.*, 11:1855–1886, 2007.
- [Sch07] S. Schwede. The stable homotopy category is rigid. *Ann. of Math. (2)*, 166(3):837–863, 2007.
- [SS03] S. Schwede and B. Shipley. Stable model categories are categories of modules. *Topology*, 42(1):103–153, 2003.

D. BARNES, DEPARTMENT OF PURE MATHEMATICS, UNIVERSITY OF SHEFFIELD, HICKS BUILDING, HOUNSFIELD ROAD, SHEFFIELD S3 7RH, UK

C. ROITZHEIM, UNIVERSITY OF KENT, SCHOOL OF MATHEMATICS, STATISTICS AND ACTUARIAL SCIENCE, CORNWALLIS BUILDING, CANTERBURY, KENT, CT2 7NF, UK