

A MODULAR DESCRIPTION OF $ER(2)$

ROMIE BANERJEE

ABSTRACT. We give a description of the maximally unramified extension of completed second Real Johnson-Wilson theory using supersingular elliptic curves with $\Gamma_0(3)$ -level structures.

1. INTRODUCTION

1.1. Real Johnson-Wilson theories. Complex conjugation acts on the complex K -theory spectrum KU and the homotopy fixed points of this action is KO . In fact the complex orientation $MU \rightarrow KU$ is equivariant with respect to this C_2 -action. Localizing at the prime 2, there is a C_2 -equivariant orientation

$$BP \rightarrow KU_{(2)}$$

with kernel $\langle v_i, i > 1 \rangle$. Hu and Kriz ([7],[8]) have realized this as a map of honest C_2 -equivariant spectra

$$B\mathbb{P}\mathbb{R} \rightarrow K\mathbb{R}$$

where $K\mathbb{R}$ is Atiyah's *real* K -theory ([1]) localized at the prime 2. They have further generalized this to a map of equivariant spectra

$$B\mathbb{P}\mathbb{R} \rightarrow E\mathbb{R}(n)$$

which is an equivariant refinement of the orientation $BP \rightarrow E(n)$ with kernel $\langle v_i, i > n \rangle$. Here $E(n)$ is the $2(2^n - 1)$ -periodic Johnson-Wilson theory. It has coefficients $\mathbb{Z}_{(2)}[v_1, \dots, v_n^{\pm 1}]$ with $|v_i| = 2(2^i - 1)$.

The underlying nonequivariant spectrum of $E\mathbb{R}(n)$ is $E(n)$ and the homotopy fixed points with respect to the complex conjugation action is denoted by $ER(n)$. This is $2^{n+2}(2^n - 1)$ periodic. The spectrum $ER(1)$ is $KO_{(2)}$. Kitchloo and Wilson have done extensive computations with the spectrum $ER(2)$ ([12]).

The spectrum $ER(n)$ is homotopy commutative, but its completion is an E_∞ -ring spectrum. In fact Averett ([2]) has shown that after completion the fixed points inclusion map $ER(n) \rightarrow E(n)$ is a higher chromatic generalization of the C_2 -Galois extension (in the sense of [16]) $KO \rightarrow KU$.

More elaborately, let E_n be the n -th Morava E -theory spectrum associated to the Lubin-Tate space of deformations of the Honda formal group Γ_n over \mathbb{F}_{2^n} . This Lubin-Tate space is noncanonically isomorphic to $\mathrm{Spf} \pi_0(E_n) = \mathrm{Spf} W(\mathbb{F}_{2^n})[[u_1, \dots, u_{n-1}]]$ where $W(k)$ denotes the Witt ring of k . Hopkins-Miller theory gives a unique E_∞ -ring structure on E_n , with coefficients

$$\pi_* E_n = \pi_0(E_n)[u^{\pm 1}]$$

1991 *Mathematics Subject Classification.* 55N20, 55N91, 14K10, 14H52.

Key words and phrases. Real Johnson-Wilson theories, Lubin-Tate spaces, topological modular forms, algebraic stacks.

where $|u| = 2$. The n -th Morava stabilizer group $\mathbb{S}_n = \mathcal{O}_{D_{\frac{1}{n}}, \mathbb{Q}_2}^\times$ (the group of units in the maximal order of the division algebra over \mathbb{Q}_2 with Hasse invariant $\frac{1}{n}$) and the Galois group $Gal = \text{Gal}(\mathbb{F}_{2^n}/\mathbb{F}_2)$ act on the Lubin-Tate space. By Hopkins-Miller theory this lifts to an action of the extended stabilizer group $\mathbb{S}_n \rtimes Gal$ on the spectrum E_n through E_∞ -ring maps, upto contractible choices.

There is a subgroup $H(n) = \mathbb{F}_{2^n}^\times \rtimes Gal$ of the extended stabilizer and a central element $[-1]_{\Gamma_n} \in \mathbb{S}_n$ corresponding to the formal inverse of the Honda formal group law. Averett shows that the standard equivalence $L_{K(n)}E(n) \simeq E_n^{hH(n)}$ is equivariant with respect to the C_2 action coming from $[-1]_{\Gamma_n}$. In other words there is an equivalence

$$(1) \quad L_{K(n)}ER(n) \simeq E_n^{h(C_2 \times H(n))}.$$

1.2. Main result. In this paper we want to show that the Real Johnson-Wilson theory $ER(2)$ arises from certain modular curves in the same way the spectrum TMF of topological modular forms arises from the moduli stack of smooth elliptic curves [6].

The stabilizer group \mathbb{S}_2 has the maximal finite subgroup $\widehat{A}_4 = Q_8 \rtimes C_3$ where C_3 acts on Q_8 by conjugation, and $H(2) = C_3 \rtimes Gal \subset \widehat{A}_4 \rtimes Gal$.

Completion of TMF at the chromatic height 2 admits the description (for the primes 2 and 3)

$$(2) \quad L_{K(2)}TMF = \left(\prod_{x \in \text{Ell}^{ss}(\overline{\mathbb{F}}_p)} E_2^{h \text{Aut}(x)} \right)^{h Gal(\overline{\mathbb{F}}_p/\mathbb{F}_p)}.$$

Here Ell^{ss} is the locus of supersingular elliptic curves at the prime p and E_2 is the Hopkins-Miller spectrum associated to the Lubin-Tate space of deformations of the formal group law of x over $\overline{\mathbb{F}}_p$. Since there is a unique isomorphism class of elliptic curves x at these primes and $\text{Aut}(x)$ is the maximal finite subgroup of \mathbb{S}_2 , the right hand side is the higher real K -theory $EO_2^{h Gal(\overline{\mathbb{F}}_2/\mathbb{F}_2)}$.

We want to describe completed $ER(2)$ in terms of modular curves. This raises the question: *Given $p = 2$, does there exist an étale cover of \mathcal{X} of Ell^{ss} so that there exists $x \in \mathcal{X}$ for which $\text{Aut}(x)$ is the subgroup $C_2 \times C_3$ of \mathbb{S}_2 ?*

We state the main theorem here.

Theorem 1.1. *For a p -local commutative S -algebra A , let $A(\mu_{\infty, p})$ denote the S -algebra obtained by adjoining all the roots of unity of order prime to p . Then $L_{K(2)}ER(2)(\mu_{\infty, 2})$ is an algebra over TMF .*

There is a $K(2)$ -local C_2 -Galois extension

$$L_{K(2)}TMF_0(3)(\mu_{\infty, 2}) \rightarrow L_{K(2)}TMF_1(3)(\mu_{\infty, 2})$$

which is isomorphic to the extension $L_{K(2)}ER(2)(\mu_{\infty, 2}) \rightarrow L_{K(2)}E(2)(\mu_{\infty, 2})$.

It is worth pointing out that Behrens and Hopkins ([4]) have answered in great detail the following question: Given a *maximal* finite subgroup G of \mathbb{S}_n at the prime p , is the associated higher real K -theory EO_n a summand of the $K(n)$ -localization of a TAF spectrum associated to a unitary similitude group of type $(1, n-1)$? This paper is concerned with a similar question where $p = 2$, $n = 2$ and G is not maximal.

It is evident from the calculations of Mahowld and Rezk ([14]) that the spectrum $TMF_1(3)$ is an example of a *generalized $E(2)$* (see [3]). Presumably there is a notion of a generalized $E\mathbb{R}(n)$ so that $TMF_0(3)$ is an example of a generalized $ER(2)$.

In the next section we reformulate our main theorem (as Theorem 2.3) in algebraic-geometric language and review the various moduli spaces appearing in the formulation. The final section contains the proof of the main theorem.

Acknowledgements. The author has benefited greatly from discussions with and comments received from Jack Morava, Niko Naumann and Andrew Salch.

2. SETUP

2.1. Stacks from ring spectra. Let BP denote the p -local Brown-Peterson spectrum. We can consider the associated flat Hopf algebroid.

$$V = BP_* \simeq \mathbb{Z}_{(p)}[v_1, \dots]$$

$$W = BP_*BP \simeq BP_*[t_1, \dots]$$

Denote the stack associated to (V, W) by \mathcal{M} . This is an algebraic stack ([13, 4.1],[15]). Let $BP^{\wedge \bullet+1}$ denote the usual cosimplicial ring spectrum, then \mathcal{M} is the stack associated to the simplicial affine scheme $\text{Spec } \pi_*BP^{\wedge \bullet+1}$.

$$\mathcal{M} = \text{colim}_{\Delta^{op}} \text{Spec } \pi_*BP^{\wedge \bullet+1}$$

The colimit is taken in the 2-category of stacks [13, 12.4].

The algebraic stack \mathcal{M} is closely related to the moduli stack of one-dimensional commutative formal groups. A formal group over a scheme S is a commutative group object in the category of formal schemes over S that is *fpqc*-locally isomorphic to $(\hat{\mathbb{A}}^1, 0)$ as schemes. Let's denote this stack by $X_{1,fg}$. Given a ring R , the groupoid $X_{1,fg}(R)$ comprises formal groups G/R over $\text{Spec } R$ and their isomorphisms.

The stack $X_{1,fg}$ carries a canonical line bundle ω . For every R we can construct the locally free rank one R -module $\omega_{G/R}$ of invariant 1-forms of G over $\text{Spec } R$, and its formation is compatible with base change and therefore defines a line bundle ω over X .

The stack $\mathcal{M} = \mathcal{M}_{(V,W)}$ is a \mathbb{G}_m torsor over $X_{1,fg} \otimes \mathbb{Z}_{(p)}$. Its points can be described as follows: For a $\mathbb{Z}_{(p)}$ -algebra R , the groupoid $\mathcal{M}(R)$ consists of pairs $(G/R, \alpha : \omega_{G/R} \simeq R)$ of a formal group and a trivialization of ω section, and isomorphisms of G that respect the trivializations.

Definition 2.1. *Let X be a p -local homotopy commutative ring spectrum.*

Consider the cosimplicial ring spectrum $BP^{\wedge \bullet+1} \wedge X$ and the associated simplicial affine scheme $\text{Spec } \pi_(BP^{\wedge \bullet+1} \wedge X)$. Define*

$$\mathcal{M}_X = \text{colim}_{\Delta^{op}} \text{Spec } \pi_*(BP^{\wedge \bullet+1} \wedge X)$$

By definition, \mathcal{M}_X is an algebraic stack over $\text{Spec } \mathbb{Z}_{(p)}$.

We now make clear the relation of \mathcal{M}_X with formal groups. The map of cosimplicial ring spectra $BP^{\wedge \bullet+1} \rightarrow BP^{\wedge \bullet+1} \wedge X$ gives a map of algebraic stacks $\mathcal{M}_X \rightarrow \mathcal{M}$. There is an algebraic stack M_X over $X_{1,fg} \otimes \mathbb{Z}_{(p)}$ along with a line bundle $\omega_X : M_X \rightarrow B\mathbb{G}_m$, so that \mathcal{M}_X is a \mathbb{G}_m -torsor over M_X . The R -points of \mathcal{M}_X can then be identified with pairs $(P \in M_X(R), \alpha : \omega_X(P/R) \simeq R)$ of objects of M_X over $\text{Spec } R$ and trivializations of their ω_X sections.

If X is a homotopy BP -algebra then \mathcal{M}_X is an affine scheme.

Lemma 2.1. *If the map $\mathcal{M}_X \rightarrow \mathcal{M}$ is flat and Y is another p -local homotopy commutative ring spectrum, then the stack associated to the smash product product $X \wedge Y$ can be identified with the pullback.*

$$\mathcal{M}_{X \wedge Y} \simeq \mathcal{M}_X \times_{\mathcal{M}} \mathcal{M}_Y$$

Proof. Consider the affine case first. Suppose E and F are BP -algebras and the map $\mathcal{M}_E \rightarrow \mathcal{M}$ is flat. This means that E is a Landweber exact cohomology theory and there is an isomorphism $E_*F \simeq E_* \otimes_{BP_*} BP_*BP \otimes_{BP_*} F_*$. In terms of stacks this is,

$$\mathrm{Spec} E_*F \simeq \mathrm{Spec} E_* \times_{\mathcal{M}} \mathrm{Spec} F_*.$$

The rest of the argument follows via descent. Suppose \mathcal{X} and \mathcal{Y} are algebraic stacks with maps to \mathcal{M} . Let \mathcal{V}_\bullet and \mathcal{W}_\bullet be simplicial affine schemes such that $\mathcal{X} = \mathrm{colim}_{\Delta^{op}} \mathcal{V}_\bullet$ and $\mathcal{Y} \simeq \mathrm{colim}_{\Delta^{op}} \mathcal{W}_\bullet$. Then the pullback can be given an atlas by the bisimplicial affine (since \mathcal{M} is algebraic) scheme $\mathcal{X} \times_{\mathcal{M}} \mathcal{Y} \simeq \mathrm{colim}_{\Delta^{op} \times \Delta^{op}} \mathcal{V}_\bullet \times_{\mathcal{M}} \mathcal{W}_\bullet$.

If $\mathcal{M}_X \rightarrow \mathcal{X}$ is flat, it follows from the proof in the affine case that there is an equivalence of bisimplicial affine schemes $\mathrm{Spec} \pi_* BP^{\wedge \bullet+1} \wedge X \times_{\mathcal{M}} \mathrm{Spec} \pi_* BP^{\wedge \bullet+1} \wedge Y \simeq \mathrm{Spec} \pi_* BP^{\wedge \bullet+1} \wedge X \wedge BP^{\wedge \bullet+1} \wedge Y$. The totalization of the bicosimplicial ring spectrum $BP^{\wedge \bullet+1} \wedge X \wedge BP^{\wedge \bullet+1} \wedge Y$ is $X \wedge Y$. \square

The notion of height of a formal group gives a filtration of the moduli stack $X_{1,fg}$ and this canonically lifts to a filtration of the \mathbb{G}_m -torsor \mathcal{M} . One can give an explicit construction of the filtration. Let $I_n = (p, v_1, \dots, v_{n-1})$ denote the invariant prime ideals of V . The associated substacks correspond to formal groups of height at least n .

$$\mathcal{M}^{\geq n} = \mathrm{Spec}(V/I_n, V/I_n \otimes_V W \otimes_V V/I_n)$$

There is a filtration on \mathcal{M} by closed substacks

$$\mathcal{M} = \mathcal{M}^{\geq 0} \supseteq \mathcal{M}^{\geq 1} \supseteq \dots \supseteq \mathcal{M}^{\geq \infty}.$$

Let $U^n = \mathcal{M} - \mathcal{M}^{\geq n}$ ($0 \leq n \leq \infty$) be the open substack of \mathcal{M} complementary to $\mathcal{M}^{\leq n}$. There is an ascending chain on open immersions

$$\emptyset = U^0 \subseteq U^1 \subseteq \dots \subseteq U^\infty \subseteq \mathcal{M}.$$

Since for $0 \leq n < \infty$, I_n is finitely generated, the open immersion $U^n \subseteq \mathcal{M}$ is quasi-compact and U^n is an algebraic stack. The R -points of U^n consist of formal groups of height at most $n-1$ over $\mathrm{Spec} R$ along with trivializations of their corresponding cotangent bundles as before. The following corollary to [15, Theorem 26] gives an explicit atlas for U^{n+1} .

Proposition 2.1. *Let $E(n)$ be the n -th Johnson Wilson spectrum with $E(n)_* = \mathbb{Z}_{(p)}[v_1, \dots, v_n^{\pm 1}]$. Let*

$$(V_n, W_n) := (E(n)_*, E(n)_* \otimes_{BP_*} BP_*BP \otimes_{BP_*} E(n)_*)$$

be the Hopf algebroid induced from (BP_, BP_*BP) by the Landweber exact map $BP_* \rightarrow E(n)_*$. Then the Hopf algebroid (V_n, W_n) is flat and its associated algebraic stack is U^{n+1} . Equivalently,*

$$\mathrm{colim}_{\Delta^{op}} \mathrm{Spec} \pi_* E(n)^{\wedge \bullet+1} \simeq U^{n+1}.$$

Lemma 2.1 gives us the following representation of the pullback.

$$\operatorname{colim}_{\Delta^{\circ p}} \operatorname{Spec} \pi_*(E(n)^{\wedge \bullet+1} \wedge X) \simeq \mathcal{U}^{n+1} \times_{\mathcal{M}} \mathcal{M}_X$$

Let $\mathcal{M}^n = \mathcal{M}^{\geq n}[v_n^{-1}]$ denote the height n -layer. The stack \mathcal{M}^n is contained inside \mathcal{U}^{n+1} as a closed substack. Define the formal neighborhood of \mathcal{M}^n by taking the completion of \mathcal{U}^{n+1} at \mathcal{M}^n .

Definition 2.2. $\widehat{\mathcal{M}}^n = (\mathcal{U}^{n+1})_{I_n}^{\wedge}$

Let

$$\eta : \operatorname{Spec} \mathbb{F}_{p^n}[u^{\pm 1}] \rightarrow \mathcal{M}^n$$

be the Honda formal group of height n . This is a presentation for \mathcal{M}^n and a pro-étale $\mathbb{S}_n \rtimes \operatorname{Gal}(\mathbb{F}_{p^n}/\mathbb{F}_p)$ -torsor. Let $\operatorname{Def}(\Gamma_n, \mathbb{F}_{p^n})$ be the Lubin-Tate space of deformations of the Honda formal group Γ_n over \mathbb{F}_{p^n} . Let $\operatorname{Def}(\Gamma_n, \mathbb{F}_{p^n})[u^{\pm 1}] := \operatorname{Def}(\Gamma, \mathbb{F}_{p^n}) \times \operatorname{Spec} \mathbb{Z}[u^{\pm 1}]$. There is a map

$$\operatorname{Def}(\Gamma_n, \mathbb{F}_{p^n})[u^{\pm 1}] \rightarrow \widehat{\mathcal{M}}^n$$

which is a presentation and an pro-étale $\mathbb{S}_n \rtimes \operatorname{Gal}(\mathbb{F}_{p^n}/\mathbb{F}_p)$ -torsor. Furthermore there is a pullback of stacks,

$$\begin{array}{ccc} \operatorname{Spec} \mathbb{F}_{p^n}[u^{\pm 1}] & \longrightarrow & \operatorname{Def}(\Gamma_n, \mathbb{F}_{p^n})[u^{\pm 1}] \\ \downarrow & & \downarrow \\ \mathcal{M}^n & \longrightarrow & \widehat{\mathcal{M}}^n \end{array}$$

Let E_n denote the Hopkins-Miller spectrum associated to $\operatorname{Def}(\Gamma_n, \mathbb{F}_{p^n})$. There is an isomorphism $\operatorname{Def}(\Gamma_n, \mathbb{F}_{p^n})[u^{\pm 1}] \simeq \operatorname{Spf} \pi_* E_n$. The diagonal,

$$\begin{aligned} \operatorname{Def}(\Gamma_n, \mathbb{F}_{p^n}) \times_{\widehat{\mathcal{M}}^n} \operatorname{Def}(\Gamma_n, \mathbb{F}_{p^n})[u^{\pm 1}] &\simeq \operatorname{Spf} E_{n*} \times (\mathbb{S}_n \rtimes \operatorname{Gal}(\mathbb{F}_{p^n}/\mathbb{F}_p)) \\ &\simeq \operatorname{Spf} \pi_* L_{K(n)}(E_n \wedge E_n). \end{aligned}$$

This produces an atlas for $\widehat{\mathcal{M}}^n$.

Proposition 2.2. $\operatorname{colim}_{\Delta^{\circ p}} \operatorname{Spf} \pi_* L_{K(n)} E_n^{\wedge \bullet+1} \simeq \widehat{\mathcal{M}}^n$

Lemma 2.1 gives us the following representation of the pullback.

$$\operatorname{colim}_{\Delta^{\circ p}} \operatorname{Spf} \pi_* L_{K(n)}(E_n^{\wedge \bullet+1} \wedge X) \simeq \widehat{\mathcal{M}}^n \times_{\mathcal{M}} \mathcal{M}_X$$

For X that is $E(n)$ -local, the stack \mathcal{M}_X can be assembled from the pieces $\widehat{\mathcal{M}}^k \times_{\mathcal{M}} \mathcal{M}_X$ for $0 \leq k \leq n$. This follows from the observation that

$$\begin{array}{ccc} \widehat{\mathcal{M}}^n \times_{\mathcal{M}} \mathcal{U}^n & \longrightarrow & \mathcal{U}^n \\ \downarrow & & \downarrow \\ \widehat{\mathcal{M}}^n & \longrightarrow & \mathcal{U}^{n+1} \end{array}$$

is a pushout square of stacks (see [17, Theorem C] for reference on stack pushouts) and $\mathcal{M}_X \simeq \mathcal{U}^{n+1} \times_{\mathcal{M}} \mathcal{M}_X$ if X is $E(n)$ -local.

The remainder of this paper concerned with working out the structure of $\mathcal{M}_{ER(2)}$ near the height 2 point.

2.2. Elliptic curves. A Weierstrass curve over R is the closure in \mathbb{P}_R^2 of the affine curve

$$(3) \quad y^2 + a_1xy + a_3y = x^3 + a_2x^2 + a_4x + a_6$$

over R . The curve is smooth if and only if $\Delta = \Delta(a_1, \dots, a_6)$ is invertible in R . A strict isomorphism of Weierstrass curves is given by the change of coordinates

$$x' = x + r, \quad y' = y + sx + t.$$

The Weierstrass curves along with their coordinate changes form an algebraic stack $\mathcal{M}_{(A,\Gamma)}$ determined by the Hopf algebroid (A, Γ) where

$$A = \mathbb{Z}[a_1, a_2, a_3, a_4, a_6], \quad \Gamma = A[s, r, t]$$

The Hopf algebroid structure maps are implicit in the definitions. (see [6, section 3]).

We now explain how this stack $\mathcal{M}_{(A,\Gamma)}$ is associated to elliptic curves. Let Ell denote the moduli stack of elliptic curves over $\text{Spec } \mathbb{Z}$. A morphism $\text{Spec}(R) \rightarrow Ell$ classifies an elliptic curve $C \rightarrow R$, which is a smooth proper morphism whose geometric fibers are elliptic curves. Let \overline{Ell} denote the compactified moduli stack classifying generalized elliptic curves. There exists a line bundle $\omega \rightarrow \overline{Ell}$ associated to the cotangent space at the identity section of a generalized elliptic curve. Given a smooth elliptic curve $C : \text{Spec } R \rightarrow Ell$, the set of sections $\Gamma(\text{Spec } R, \omega(C))$ is the set of invariant 1-forms on C .

Let Ell be the \mathbb{G}_m -torsor over Ell whose R -points are given by pairs $(C/R, \alpha : \omega(C) \simeq R)$. Here C/R is an elliptic curve over R and α is a choice of trivialization of the R -sections of ω .

Any generalized elliptic curve $C \rightarrow S$ admits a presentation in the Weierstrass normal form locally over S in the flat topology. The identity element of the elliptic curve is identified with the unique point at infinity of the Weierstrass curve. This gives a map of stacks $\overline{\text{Ell}} \rightarrow \mathcal{M}_{(A,\Gamma)}$ which is an equivalence on the substack of smooth elliptic curves, $\text{Ell} \simeq \mathcal{M}_{(A[\Delta^{-1}], \Gamma[\Delta^{-1}])}$.

There is a natural map $Ell \rightarrow X_{1,fg}$ classifying the formal group associated to an elliptic curve. This map lifts canonically to a map of the \mathbb{G}_m -torsors.

$$\text{Ell} \rightarrow \mathcal{M}$$

Consider the substack $\overline{\text{Ell}}_p \rightarrow \overline{\text{Ell}}$ which is the p -completion of $\overline{\text{Ell}}$. Note that $\overline{\text{Ell}}_p$ is a *formal* Deligne-Mumford stack. For any p -complete ring R the map $\text{Spf}(R) \rightarrow \overline{\text{Ell}}$ classifies an ind-system $C_m/\text{Spec}(R/p^n)$ of generalized elliptic curves.

Define $(\overline{\text{Ell}})_{\mathbb{F}_p} = \overline{\text{Ell}} \times_{\mathbb{Z}} \text{Spec}(\mathbb{F}_p)$. Let $(\text{Ell}^{ord})_{\mathbb{F}_p} \subset (\overline{\text{Ell}})_{\mathbb{F}_p}$ denote the locus of ordinary generalized elliptic curves in characteristic p , and let

$$(\text{Ell}^{ss})_{\mathbb{F}_p} = (\overline{\text{Ell}})_{\mathbb{F}_p} - (\text{Ell}^{ord})_{\mathbb{F}_p}$$

denote the locus of supersingular elliptic curves in characteristic p .

Consider the substack

$$\text{Ell}_p^{ss} \subset \overline{\text{Ell}}_p$$

where Ell_p^{ss} is the completion of $\overline{\text{Ell}}$ at $(\text{Ell}^{ss})_{\mathbb{F}_p}$.

Define the Hopf algebroid,

$$A' = A[\Delta^{-1}]_{(p,a_1)}^\wedge, \quad \Gamma' = A' \otimes_A \Gamma \otimes_A A'.$$

Then,

$$\mathrm{Ell}_p^{ss} = \mathcal{M}_{(A', \Gamma')}.$$

The following is a special case of the Serre-Tate theorem for abelian schemes [9, Theorem 1.2.1].

Theorem 2.1. *(Serre, Tate) Let R be a Noetherian ring with p nilpotent and I a nilpotent ideal in R , $R_0 = R/I$. Let $\mathrm{Ell}_p^{ss}(R)$ denote the category of supersingular elliptic curves in characteristic p over R and $D(R, R_0)$ denote the category of triples*

$$(C_0, G, \epsilon)$$

consisting of a supersingular elliptic curve C_0 over R_0 , a formal group G over R and an isomorphism of formal groups over R_0 , $\epsilon : G_0 \simeq C_0^\wedge$, where G_0 is reduction modulo I of G . Then the functor

$$\mathrm{Ell}_p^{ss}(R) \rightarrow D(R, R_0)$$

$$C \mapsto (C_0, C^\wedge, \text{natural } \epsilon)$$

is an equivalence of categories.

The following is implied by the Serre-Tate theorem.

Proposition 2.3. *There is an equivalence of formal Deligne-Mumford stacks,*

$$\mathrm{Ell}_p^{ss} \simeq \widehat{\mathcal{M}}^2 \times_{\mathcal{M}} \overline{\mathrm{Ell}}.$$

Let E be a supersingular elliptic curve over a field k of characteristic p classifying a point $\eta : \mathrm{Spec} k \rightarrow (\mathrm{Ell})_{\mathbb{F}_p}$. Serre-Tate theory gives an isomorphism of the formal neighborhood of η with the universal deformation space for the formal group \widehat{E} . Therefore there is an equivalence of deformation spaces

$$\mathrm{Def}(E, k) \simeq \mathrm{Def}(\widehat{E}, k).$$

2.3. Level structures. In this section we review the modular curves. Let S be a scheme over $\mathbb{Z}[1/N]$ and C a smooth elliptic curve over S . Let $C[N]$ denote the N -torsion points of C . The group scheme $C[N]$ is étale locally over S isomorphic to the discrete group scheme $(\mathbb{Z}/N\mathbb{Z})^2$ over C . Let $\mathrm{Ell}(N)$ denote the moduli stack of pairs (C, ϕ) where C is a smooth elliptic curve and ϕ is a *full level- n structure*, a choice of isomorphism

$$\phi : (\mathbb{Z}/N\mathbb{Z})^2 \xrightarrow{\simeq} C[N].$$

Equivalently, the points of $\mathrm{Ell}(N)$ are triples (C, P, Q) where P and Q are a pair of sections $S \rightarrow C[N]$ that are, locally over S , linearly independent.

Let $\mathrm{Ell}_1(N)$ denote the moduli stack of pairs (C, P) where P is a primitive N -torsion point over C . Finally, let $\mathrm{Ell}_0(N)$ denote the moduli stack of pairs (C, H) where H is a choice of a subgroup scheme $H \subset C[N]$ isomorphic to $\mathbb{Z}/N\mathbb{Z}$.

In this paper we'll work with stacks $\mathrm{Ell}(N)$, $\mathrm{Ell}_1(N)$ and $\mathrm{Ell}_0(N)$ which are \mathbb{G}_m -torsors over the modular curves $\mathrm{Ell}(N)$, $\mathrm{Ell}_1(N)$ and $\mathrm{Ell}_0(N)$. The R -points of $\mathrm{Ell}(N)$ consists of triples $(C, \eta, (P, Q))$, where C is an elliptic curve over R , η is a choice of a nowhere vanishing invariant 1-form of C and (P, Q) is a full level- N structure described as before.

Theorem 2.2. *(Deligne, Rapoport) ([10]) The moduli stack $\mathrm{Ell}(N)$ is a smooth affine scheme over $\mathrm{Spec} \mathbb{Z}[1/N]$ for $N \geq 3$. For $N \geq 4$ the moduli stack $\mathrm{Ell}_1(N)$ is a smooth affine scheme over $\mathrm{Spec} \mathbb{Z}[1/N]$.*

The maps forgetting level structures induce a diagram

$$\begin{array}{ccc}
 & \text{Ell}(N) & \\
 & \Gamma_1(N) \downarrow & \\
 & \text{Ell}_1(N) & \Gamma_0(N) \\
 & (\mathbb{Z}/N\mathbb{Z})^\times \downarrow & \\
 & \text{Ell}_0(N) & \\
 & \downarrow & \\
 \text{Ell} \times \text{Spec } \mathbb{Z}[1/N] & &
 \end{array}$$

$GL(2, N)$ (curved arrow from $\text{Ell}(N)$ to $\text{Ell} \times \text{Spec } \mathbb{Z}[1/N]$)

where all the arrows are finite étale and the labeled ones are Galois.

The Galois groups are defined as follows:

$$\Gamma_1(N) = \left\{ \begin{pmatrix} 1 & * \\ 0 & * \end{pmatrix} \in GL(2, N) \right\}$$

$$\Gamma_0(N) = \left\{ \begin{pmatrix} * & * \\ 0 & * \end{pmatrix} \in GL(2, N) \right\}$$

Let $\text{Ell}(N)_p$, $\text{Ell}_1(N)_p$ and $\text{Ell}_0(N)_p$ denote the completions at p . Let $\text{Ell}(N)_p^{ss}$ denote the pullback,

$$\begin{array}{ccc}
 \text{Ell}(N)_p^{ss} & \longrightarrow & \text{Ell}(N)_p \\
 \downarrow & & \downarrow \\
 \text{Ell}_p^{ss} & \longrightarrow & \text{Ell}_p
 \end{array}$$

Since $\text{Ell}(N)_p$ is formal affine (assuming $N \geq 3$) and the right vertical arrow is an étale $GL(2, N)$ -torsor, Serre-Tate theory implies there is an equivalence

$$\text{Ell}(N)_p^{ss} \simeq \prod_i \text{Spf } W(k_i)[[u_1]][u^{\pm 1}]$$

for a finite set of fields k_i (depending on N).

2.4. Restatement of the main theorem. Given a formal Deligne-Mumford stack \mathcal{S} over \mathbb{Z}_p , define its maximal unramified cover \mathcal{S}^{nr} to be the pullback $\mathcal{S} \times_{\mathbb{Z}_p} \text{Spf } W(\overline{\mathbb{F}}_p)$. The map $\mathcal{S}^{nr} \rightarrow \mathcal{S}$ is a pro-étale cover with Galois group $\text{Gal}(\overline{\mathbb{F}}_p/\mathbb{F}_p) = \hat{\mathbb{Z}}$. Our main Theorem 1.1 can be restated in the following way.

Theorem 2.3. *There is a map of stacks over $\widehat{\mathcal{M}}^2$,*

$$\widehat{\mathcal{M}}^2 \times_{\mathcal{M}} \mathcal{M}_{ER(2)}^{nr} \rightarrow \text{Ell}_2^{ss}.$$

The map over $\widehat{\mathcal{M}}^2$ induced by forgetting level structure

$$(\text{Ell}_1(3)_2^{ss})^{nr} \rightarrow (\text{Ell}_0(3)_2^{ss})^{nr}$$

is equivalent to the map

$$\widehat{\mathcal{M}}^2 \times_{\mathcal{M}} \mathcal{M}_{E(2)}^{nr} \rightarrow \widehat{\mathcal{M}}^2 \times_{\mathcal{M}} \mathcal{M}_{ER(2)}^{nr}$$

over $\widehat{\mathcal{M}}^2$ induced by the inclusion of fixed points $ER(2) \rightarrow E(2)$.

3. REAL JOHNSON-WILSON THEORY FROM MODULAR CURVES

3.1. **Level 3-structures at the prime 2.** Consider the supersingular elliptic curve

$$(4) \quad C : x^3 + y^2 + y = 0 \in \mathbb{P}_{\mathbb{F}_4}^2.$$

The automorphism group $G_{24} = \text{Aut}_{\mathbb{F}_4}(C)$ is the group of units in the maximal order of a rational quaternion algebra $\mathbb{Q}\{i, j, k\}$. It's isomorphic to the binary tetrahedral group $\widehat{A}_4 = Q_8 \rtimes C_3$ of order 24, where C_3 acts on Q_8 by conjugation. It contains the quaternion group $Q_8 = \{\pm 1, \pm i, \pm j \pm k\}$ and 16 other elements $(\pm 1 \pm i \pm j \pm k)/2$.

Let C^\wedge be the completion of C at the identity section. C^\wedge is a formal group of height 2 over \mathbb{F}_4 . The automorphism group $\text{Aut}(C^\wedge) = \mathcal{O}_{D_{1/2}, \mathbb{Q}_2}^\times$ is the group of units in the maximal order of the 2-adic quaternion algebra

$$D_{\frac{1}{2}, \mathbb{Q}_2} = \mathbb{Q}_2\{i, j, k\}.$$

Abstractly this is the completion of the Hurwitz lattice $\mathbb{Z}(\pm 1, \pm i, \pm j, \pm k) \coprod (\pm 1 \pm i \pm j \pm k)/2$ at the ideal (2). Notice that G_{24} is the maximal finite subgroup.

Proposition 3.1. *The map $\text{Spec } \mathbb{F}_4 \rightarrow (\text{Ell}^{ss})_{\mathbb{F}_2}$ classifying C is a presentation and an étale $G_{24} \rtimes \text{Gal}(\mathbb{F}_4/\mathbb{F}_2)$ -torsor.*

It follows that the map $\text{Def}(C, \mathbb{F}_4) \rightarrow \text{Ell}_2^{ss}$ classifying the universal deformation of C is an étale $G_{24} \rtimes \text{Gal}(\mathbb{F}_4/\mathbb{F}_2)$ -torsor. By Serre-Tate theory there is an isomorphism of deformation spaces

$$\text{Def}(C, \mathbb{F}_4) \simeq \text{Spf } W(\mathbb{F}_4)[[a_1]].$$

The curve C lifts to $W(\mathbb{F}_4)[[a_1]]$ as $\widetilde{C} : y^2 + a_1xy + y = x^3$, which is the universal deformation curve over $\text{Def}(C, \mathbb{F}_4)$. \widetilde{C} lifts further to

$$(5) \quad y^2 + a_1uxy + u^3y = x^3$$

over $\text{Spf } W(\mathbb{F}_4)[[a_1]][[u^{\pm 1}]]$. We shall call this \widetilde{C} from now.

Proposition 3.2. [6, Theorem 3.1] *The map*

$$\text{Spf } W(\mathbb{F}_4)[[a_1]][[u^{\pm 1}]] \rightarrow \text{Ell}_2^{ss} \simeq \mathcal{M}_{(A', \Gamma')}$$

classifying the curve

$$\widetilde{C} : y^2 + a_1uxy + u^3y = x^3$$

is a presentation and an étale $G_{24} \rtimes \text{Gal}(\mathbb{F}_4/\mathbb{F}_2)$ -torsor.

This is a restatement of

$$L_{K(2)}\text{TMF} = \text{EO}_2.$$

The level 3-structures on elliptic curves and their associated moduli stacks are related by finite étale morphisms

$$\text{Ell}(3) \xrightarrow{6} \text{Ell}_1(3) \xrightarrow{2} \text{Ell}_0(3) \xrightarrow{4} \text{Ell} \times \text{Spec } \mathbb{Z}[1/3]$$

of degrees 6, 2 and 4. The related modular groups are as follows:

$$\Gamma_1(3) = \left\{ \begin{pmatrix} 1 & * \\ 0 & * \end{pmatrix} \in GL(2, 3) \right\} \simeq C_3 \rtimes C_2$$

$$\Gamma_0(3) = \left\{ \begin{pmatrix} * & * \\ 0 & * \end{pmatrix} \in GL(2, 3) \right\} \simeq C_6 \rtimes C_2$$

The point $(0, 0)$ on the Weierstrass normal form of a smooth elliptic curve is a point of order 3 if it is of the form

$$C(a_1, a_3) : y^2 + a_1xy + a_3y = x^3, \quad a_3 \neq 0.$$

The curve $C(a_1, a_3)$ comes with the invariant 1-form

$$\eta = \frac{dx}{2y + a_1x + a_3} = \frac{dy}{3x^2 - a_1y}.$$

Furthermore, if (C', P, η') is a smooth elliptic curve with a point P of order 3 and an invariant 1-form η' then there exists a *unique* isomorphism of the triple (see [14, Prop. 3.2])

$$(C(a_1, a_3), (0, 0), \eta) \simeq (C', P, \eta').$$

This gives us the following equivalence of stacks. Notation: $\Delta = a_3^3(a_1^3 - 27a_3)$.

$$\text{Ell}_1(3) = \text{Spec } \mathbb{Z}[1/3][a_1, a_3^{\pm 1}, \Delta^{-1}]$$

$$\text{Ell}_1(3)_2^{ss} = \text{Spf } \mathbb{Z}_2[[a_1]][a_3^{\pm 1}]$$

Proposition 3.3. *The map*

$$\text{Spf } W(\mathbb{F}_4)[[a_1]][u^{\pm 1}] \rightarrow \text{Ell}_1(3)_2^{ss}$$

classifying the point $(\tilde{C}, (0, 0), \frac{dx}{2y+a_1ux+u^3})$ is a presentation and an étale $C_3 \rtimes \text{Gal}(\mathbb{F}_4/\mathbb{F}_2)$ -torsor. The embedding $C_3 \subset \mathbb{S}_2$ is induced by lifting the 2-adic Teichmüller character $\mathbb{F}_4^\times \rightarrow W(\mathbb{F}_4)^\times$.

There is a C_2 -étale map forgetting level structure,

$$\text{Ell}_1(3) \rightarrow \text{Ell}_0(3)$$

$$(C, P, \eta) \mapsto (C, \langle P \rangle, \eta).$$

The group C_2 acts on $\text{Ell}_1(3)$ by $(C, P, \eta) \mapsto (C, -P, \eta)$.


Proposition 3.4. *The map*

$$\text{Spf } W(\mathbb{F}_4)[[a_1]][u^{\pm 1}] \rightarrow \text{Ell}_0(3)_2^{ss}$$

classifying the point $(\tilde{C}, \langle (0, 0) \rangle, \frac{dx}{2y+a_1ux+u^3})$ is a presentation and an étale $C_2 \times C_3 \rtimes \text{Gal}(\mathbb{F}_4/\mathbb{F}_2)$ -torsor. The group C_2 is the normal subgroup of $Q_8 \rtimes C_3$ generated by the formal inverse $[-1]_{C^\wedge}$.

The following diagram shows how the various modular curves at the supersingular locus in characteristic 2 are related. All the arrows are finite étale and the

labeled ones are Galois. Notation: $Gal = \text{Gal}(\mathbb{F}_4/\mathbb{F}_2)$.



As a consequence there is a diagram of $K(2)$ -local elliptic spectra.

$$\begin{array}{ccc}
 L_{K(2)}TMF_1(3) & \xrightarrow{\simeq} & E_2^{h(C_3 \times Gal)} \\
 \uparrow & & \uparrow \\
 L_{K(2)}TMF_0(3) & \xrightarrow{\simeq} & E_2^{h(C_2 \times C_3 \times Gal)} \\
 \uparrow & & \uparrow \\
 L_{K(2)}TMF & \xrightarrow{\simeq} & E_2^{h(\hat{A}_4 \times Gal)}
 \end{array}$$

Here E_2 is the Hopkins-Miller Morava E -theory spectrum associated to the deformation space $\text{Def}(C, \mathbb{F}_4)$.

3.2. The structure of $\mathcal{M}_{E(2)}$ near the height 2 point. Let the point $f : \text{Spec } \mathbb{F}_4 \rightarrow \mathcal{U}^3$ classify the formal group C^\wedge associated to the supersingular elliptic curve $C : y^2 + y + x^3 = 0$. Let \tilde{C} denote the universal deformation of the curve C over $\text{Spf } W(\mathbb{F}_4)[[a_1]][u^{\pm 1}]$ as in equation (5).

Since \mathcal{U}^3 denotes the moduli stack of formal groups of height at most 2 we know from Prop. 2.1 there is a faithfully flat presentation

$$\text{Spec } \mathbb{Z}_{(2)}[v_1, v_2^{\pm 1}] \rightarrow \mathcal{U}^3.$$

As a consequence there exists an extension K of \mathbb{F}_4 such that

$$f : \text{Spf } W(K)[[a_1]][u^{\pm 1}] \rightarrow \text{Spf } W(\mathbb{F}_4)[[a_1]][u^{\pm 1}]$$

is an fpqc cover and the pullback $f^*(\tilde{C}^\wedge)$ is isomorphic to a 2-typical formal group law $\psi : \text{Spf } W(K)[[a_1]][u^{\pm 1}] \rightarrow \text{Spec } \mathbb{Z}_{(2)}[v_1, v_2^{\pm 1}]$.

$$\alpha : f^*(\tilde{C}^\wedge) \simeq \psi$$

Consider the map

$$\text{Spf } W(K)[[a_1]][u^{\pm 1}] \xrightarrow{(f^*(\tilde{C}^\wedge), \alpha, \psi)} \widehat{\mathcal{M}}^2 \times_{\mathcal{M}} \mathcal{M}_{E(2)}$$

classifying the formal group $f^*(\tilde{C}^\wedge)$, ψ and the isomorphism α . This map is a surjection and an étale presentation.

The Teichmüller character map $\mathbb{F}_4^\times \rightarrow \mathbb{Z}_2^\times$ lifts to give an embedding $\mathbb{F}_4^\times \subset \text{Aut}(f^*(C^\wedge))$. For a $\omega \in \mathbb{F}_4^\times$ the linear formal power series $g(t) = \omega t$ gives an element of \mathbb{S}_2 . Since the action of $\text{Aut}(f^*(C^\wedge))$ extends to an action on the deformation space $\text{Spf } W(K)[[a_1]][u^{\pm 1}]$, we obtain an action of g on the ring $W(K)[[a_1]][u^{\pm 1}]$ by

$$\begin{aligned} g(u) &= \omega u \\ g(ua_1) &= \omega^2 ua_1 \end{aligned}$$

which leaves v_1 and v_2 invariant. Moreover these are the only elements of \mathbb{S}_2 which acts invariantly on v_1 and v_2 . Since $\widehat{\mathcal{M}}^2 \times_{\mathcal{M}} \text{Spec } \mathbb{Z}_{(2)}[v_1, v_2^{\pm 1}] \simeq \text{Spf } \mathbb{Z}_2[[v_1]][v_2^{\pm 1}]$ we can identify the pullback in the following diagram of stacks

$$\begin{array}{ccc} \text{Spf } W(K)[[a_1]][u^{\pm 1}] \times (C_3 \rtimes \text{Gal}(K/\mathbb{F}_2)) & \longrightarrow & \text{Spf } W(K)[[a_1]][u^{\pm 1}] \\ \downarrow & & \downarrow \\ \text{Spf } W(K)[[a_1]][u^{\pm 1}] & \longrightarrow & \widehat{\mathcal{M}}^2 \times_{\mathcal{M}} \mathcal{M}_{E(2)} \end{array}$$

Therefore, the map $\text{Spf } W(K)[[a_1]][u^{\pm 1}] \rightarrow \widehat{\mathcal{M}}^2 \times_{\mathcal{M}} \text{Spec } \mathbb{Z}_{(2)}[v_1, v_2^{\pm 1}]$ is an étale $C_3 \rtimes \text{Gal}(K/\mathbb{F}_2)$ -torsor. The conjugation action of $\text{Gal}(K/\mathbb{F}_2)$ on C_3 factors through the quotient $\text{Gal}(K/\mathbb{F}_2) \rightarrow \text{Gal}(\mathbb{F}_4/\mathbb{F}_2)$ since every automorphism of \widetilde{C}^\wedge is already defined over \mathbb{F}_4 . In terms of $K(2)$ -localization,

$$L_{K(2)}E(2) = E_2^{hH}$$

where $H = \mathbb{F}_4^\times \rtimes \text{Gal}(K/\mathbb{F}_2)$ and E_2 is the Hopkins-Miller spectrum associated to $\text{Def}(C^\wedge, K)$.

More generally, we can use the maximal unramified extension E_2^{nr} , which is the Hopkins-Miller spectrum associated to the deformation space $\text{Def}(C, \overline{\mathbb{F}}_2)$. The extended Morava stabilizer group in this case is $\mathbb{S}_2 \rtimes \text{Gal}(\overline{\mathbb{F}}_2/\mathbb{F}_2) = \mathbb{S}_2 \rtimes \widehat{\mathbb{Z}}$.

Proposition 3.5. *There is a map of stacks $\text{Def}(C, \overline{\mathbb{F}}_2) \rightarrow \widehat{\mathcal{M}}^2 \times_{\mathcal{M}} \mathcal{M}_{E(2)}$ which is a surjection and an étale $C_3 \rtimes \widehat{\mathbb{Z}}$ -torsor.*

The maximal unramified cover of the deformation space $\text{Def}(C, \mathbb{F}_4)$ can be identified with $\text{Def}(C, \overline{\mathbb{F}}_2)$. There is a natural map of stacks (we use the notation $//$ for quotient stacks)

$$(6) \quad \text{Def}(C, \overline{\mathbb{F}}_2) \rightarrow \text{Def}(C, \mathbb{F}_4) // \text{Gal}(\mathbb{F}_4/\mathbb{F}_2),$$

which realizes as a $K(n)$ -local $\widehat{\mathbb{Z}}$ -Galois extension of S -algebras

$$E_2^{Gal} \rightarrow E_2^{nr}.$$

The $\widehat{\mathbb{Z}}$ -extension is obtained by adjoining all the roots of unity of order prime to p to the p -complete spectrum E_2^{Gal} ([16, 5.4.6]).

The map (6) factors through the quotient

$$\text{Def}(C, \overline{\mathbb{F}}_2) // C_3 \rightarrow \text{Def}(C, \mathbb{F}_4) // \widehat{A}_4 \rtimes \text{Gal}(\mathbb{F}_4/\mathbb{F}_2).$$

In terms of $K(2)$ -localization there is a map

$$(7) \quad L_{K(2)}TMF \simeq (E_2)^{\widehat{A}_4 \rtimes Gal} \rightarrow (E_2^{nr})^{hC_3} \simeq L_{K(2)}E(2)(\mu_{\infty, 2}).$$

Combining Prop. 3.3 and Prop. 3.5 proves the following.

Proposition 3.6. *There are equivalences of stacks over $\widehat{\mathcal{M}}^2$.*

$$\mathrm{Def}(C, \overline{\mathbb{F}}_2) // C_3 \simeq \widehat{\mathcal{M}}^2 \times_{\mathcal{M}} \mathcal{M}_{E(2)}^{nr} \simeq (\mathrm{Ell}_1(3)_2^{ss})^{nr}.$$

3.3. The structure of $\mathcal{M}_{ER(2)}$ near the height 2 point.

Proposition 3.7. *The map of stacks*

$$\widehat{\mathcal{M}}^2 \times_{\mathcal{M}} \mathcal{M}_{E(2)} \rightarrow \widehat{\mathcal{M}}^2 \times_{\mathcal{M}} \mathcal{M}_{ER(2)}$$

is a surjection and an étale C_2 -torsor.

Proof. Combining Averett's formula (1) and [16, Theorem 5.4.4] we see that the map of ring spectra $L_{K(2)}ER(2) \rightarrow L_{K(2)}E(2)$ is a $K(2)$ -local C_2 -Galois extension of 2-complete commutative S -algebras. \square

Given the description of $\widehat{\mathcal{M}}^2 \times_{\mathcal{M}} \mathcal{M}_{E(2)}$ as the quotient stack $\mathrm{Def}(C, \overline{\mathbb{F}}_2) // C_3 \rtimes \hat{\mathbb{Z}}$, the action of C_2 must come from the inverse map $[-1]_C$ on the curve C .

The induced action on the deformation space is as follows:

$$\begin{aligned} [-1]_C : W(\mathbb{F}_4)[[a_1]][u^{\pm 1}] &\longrightarrow W(\mathbb{F}_4)[[a_1]][u^{\pm 1}] \\ [-1]_C(u) &= -u \\ [-1]_C(ua_1) &= ua_1 \end{aligned}$$

Proposition 3.8. *There is a map of stacks $\mathrm{Def}(C, \overline{\mathbb{F}}_2) \rightarrow \widehat{\mathcal{M}}^2 \times_{\mathcal{M}} \mathcal{M}_{ER(2)}$ which is an étale surjection and a $C_2 \times C_3 \rtimes \hat{\mathbb{Z}}$ -torsor. The C_2 -action on $\mathrm{Def}(C, \overline{\mathbb{F}}_2)$ comes from the inverse $[-1]_C$.*

This produces a map

$$\mathrm{Def}(C, \overline{\mathbb{F}}_2) // C_2 \times C_3 \rightarrow \mathrm{Def}(C, \mathbb{F}_4) // \hat{A}_4 \rtimes \mathrm{Gal}(\mathbb{F}_4/\mathbb{F}_2).$$

In terms of $K(2)$ -localization the map of spectra (7) factors through the real part.

$$(8) \quad L_{K(2)}TMF \rightarrow (E_2^{nr})^{h(C_2 \times C_3)} \simeq L_{K(2)}ER(2)(\mu_{\infty, 2}) \rightarrow L_{K(2)}E(2)(\mu_{\infty, 2}).$$

This proves the first part of Theorem 2.3. Combining Prop. 3.4 with Prop. 3.8 proves the following which together with Prop. 3.6 proves the rest of Theorem 2.3.

Proposition 3.9. *There are equivalences of stacks over $\widehat{\mathcal{M}}^2$,*

$$\mathrm{Def}(C, \overline{\mathbb{F}}_2) // C_2 \times C_3 \simeq \widehat{\mathcal{M}}^2 \times_{\mathcal{M}} \mathcal{M}_{ER(2)}^{nr} \simeq (\mathrm{Ell}_0(3)_2^{ss})^{nr}.$$

REFERENCES

- [1] Atiyah, M.F. *K-theory and Reality*, Quar. J. Math., 17(1966), 367-386.
- [2] Averett, M. *Completion of Real Johnson-Wilson theory $ER(n)$ gives fixed points of Morava E-theory*, PhD thesis, UCSD
- [3] Baker, A. I_n -local Johnson-Wilson spectra and their Hopf algebroids, Documenta Math., 5 (2000) 351-364.
- [4] Behrens, M; Hopkins, M.J. *Higher real K-theories and topological automorphic forms*, J. Topology 4 (2011), 39-72.
- [5] Deligne, P; Rapoport, M. *Les schémas de modules de courbes elliptiques*
- [6] Hopkins, M.J; Mahowald, M. *From elliptic curves to homotopy theory* Preprint, <http://hopf.math.purdue.edu/>.
- [7] Hu, P. *On Real-Oriented Johnson-Wilson cohomology*, Algebraic and Geometric Topology, 2(2002), 937-947.

- [8] Hu, P; Kriz, I. *Real-oriented Homotopy Theory and an analogue of the Adams-Novikov Spectral Sequence*, *Topology*, 40(2):317-399, 2001.
- [9] Katz, N. *Serre-Tate local moduli from: Algebraic surfaces (Orsay, 197678)*, *Lecture Notes in Math.* 868, Springer, Berlin (1981) 138202
- [10] Katz, N; Mazur, B. *Arithmetic Moduli of Elliptic Curves*, *Annals of Mathematics Studies.* AM-108
- [11] Kitchloo, N; Wilson, W.S. *On fibrations related to real spectra*. *Proceedings of the Nishida fest (Kinosaki 2003)*, volume 10, *Geometry and Topology Monographs*, pages 237-244, 2007.
- [12] Kitchloo, N; Wilson, W.S. *The second real Johnson-Wilson theory and non-immersions of RP^n* , *Homology, Homotopy Appl.*, 10(3), 2008, 223-268.
- [13] Laumon, G; Moret-Bailly, L. *Champs algébriques* volume 39 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*, Springer-Verlag, Berlin (2000)
- [14] Mahowald, M; Rezk, C. *Topological Modular Forms of Level 3* (arXiv:0812.2009)
- [15] Naumann, N. *Comodule categories and the geometry of the stack of formal groups* (arXiv:math/0503308)
- [16] Rognes, J. *Galois Extensions of Structured Ring Spectra: Stably Dualizable Groups* *Memoirs of the American Mathematical Society*, March 2008, Vol 192, Number 898, ISSN 0065-9266
- [17] Rydh, D. *Étale Dévissage, descent and pushouts of stacks* *J.Algebra* 331(1), 2011, p.194-223.

SCHOOL OF MATHEMATICS, TATA INSTITUTE OF FUNDAMENTAL RESEARCH, MUMBAI 400005
E-mail address: banerjee@math.tifr.res.in