ON ORNSTEIN-UHLENBECK DRIVEN BY ORNSTEIN-UHLENBECK PROCESSES

BERNARD BERCU, FRÉDÉRIC PROIA, AND NICOLAS SAVY

ABSTRACT. We investigate the asymptotic behavior of the maximum likelihood estimators of the unknown parameters of positive recurrent Ornstein-Uhlenbeck processes driven by Ornstein-Uhlenbeck processes.

1. INTRODUCTION AND MOTIVATION

Since the seminal work of Ornstein and Uhlenbeck [17], a wide literature is available on Ornstein-Uhlenbeck processes driven by Brownian or fractional Brownian motions [12], [14]. Many interesting papers are also available on Ornstein-Uhlenbeck processes driven by Lévy processes

(1.1)
$$dX_t = \theta X_t dt + dL_t$$

where $\theta < 0$ and (L_t) is a continuous-time stochastic process starting from zero with stationary and independent increments. We refer the reader to Barndorff-Nielsen and Shephard [2] for the mathematical foundation on Ornstein-Uhlenbeck processes driven by Lévy processes, and also to [1] for a recent extension to fractional Lévy processes. Parametric estimation results for Ornstein-Uhlenbeck driven by α -stable Lévy processes are established in [10] whereas nonparametric estimation results are given in [11]. Two interesting applications related to money exchange rates and stock prices may be found in [2] and [15], see also the references therein.

To the best of our knowledge, no results are available on Ornstein-Uhlenbeck driven by Ornstein-Uhlenbeck processes defined, over the time interval [0, T], by

(1.2)
$$\begin{cases} dX_t = \theta X_t dt + dV_t \\ dV_t = \rho V_t dt + dW_t \end{cases}$$

where $\theta < 0, \rho \leq 0$ and (W_t) is a standard Brownian motion. For the sake of simplicity and without loss of generality, we choose the initial values $X_0 = 0$ and $V_0 = 0$. Our motivation for studying (1.2) comes from two observations. On the one hand, the increments of Ornstein-Uhlenbeck processes are not independent which means that the weighted maximum likelihood estimation approach of [10] does not apply directly to our situation. On the other hand, Ornstein-Uhlenbeck driven by Ornstein-Uhlenbeck processes are clearly related with stochastic volatility models in financial mathematics [16]. Furthermore, (1.2) is the continuous-time version of

Key words and phrases. Ornstein-Uhlenbeck process, Maximum likelihood estimation, Continuous-time Durbin-Watson statistic, Almost sure convergence, Asymptotic normality.

the first-order stable autoregressive process driven by a first-order autoregressive process recently investigated in [3].

The paper organizes as follows. Section 2 is devoted to the maximum likelihood estimation for θ and ρ . We also introduce the continuous-time Durbin-Watson statistic which will allow us to propose a serial correlation test for Ornstein-Uhlenbeck driven by Ornstein-Uhlenbeck processes. In Section 3, we establish the almost sure convergence as well as the asymptotic normality of our estimates. One shall realize that there is a radically different behavior of the estimator of ρ in the two situations where $\rho < 0$ and $\rho = 0$. Our analysis relies on technical tools postponed to Section 4. Finally, in Section 5, we propose a statistical procedure based on the continuous-time Durbin-Watson statistic, in order to test whether or not $\rho = 0$.

2. MAXIMUM LIKELIHOOD ESTIMATION

The maximum likelihood estimator of θ is given by

(2.1)
$$\widehat{\theta}_T = \frac{\int_0^T X_t \, \mathrm{d}X_t}{\int_0^T X_t^2 \, \mathrm{d}t} = \frac{X_T^2 - T}{2\int_0^T X_t^2 \, \mathrm{d}t}$$

In the standard situation where $\rho = 0$, it is well-known that $\hat{\theta}_T$ converges to θ almost surely. Moreover, as $\theta < 0$, the process (X_T) is positive recurrent and we have the asymptotic normality

$$\sqrt{T}\left(\widehat{\theta}_T - \theta\right) \xrightarrow{\mathcal{L}} \mathcal{N}(0, -2\theta).$$

We shall see in Section 3 that the almost sure limiting value of $\hat{\theta}_T$ and its asymptotic variance will change as soon as $\rho < 0$. The estimation of ρ requires the evaluation of the residuals generated by the estimation of θ at stage T. For all $0 \le t \le T$, denote

(2.2)
$$\widehat{V}_t = X_t - \widehat{\theta}_T \Sigma_t$$

where

(2.3)
$$\Sigma_t = \int_0^t X_s \, \mathrm{d}s.$$

By analogy with (2.1) and on the basis of the residuals (2.2), we estimate ρ by

(2.4)
$$\widehat{\rho}_T = \frac{\widehat{V}_T^2 - T}{2\int_0^T \widehat{V}_t^2 \,\mathrm{d}t}$$

Therefore, we are in the position to define the continuous-time version of the discretetime Durbin-Watson statistic [3], [5], [6], [7],

(2.5)
$$\widehat{D}_T = \frac{2\int_0^T \widehat{V}_t^2 \,\mathrm{d}t - \widehat{V}_T^2 + T}{\int_0^T \widehat{V}_t^2 \,\mathrm{d}t},$$

which clearly means that $\widehat{D}_T = 2(1 - \widehat{\rho}_T)$. In Section 3, we shall make use of \widehat{D}_T to build a serial correlation statistical test for the Ornstein-Uhlenbeck driven noise, that is to test whether or not $\rho = 0$.

3. MAIN RESULTS

The almost sure convergences of our estimates are as follows.

Theorem 3.1. We have the almost sure convergences

(3.1)
$$\lim_{T \to \infty} \widehat{\theta}_T = \theta^*, \qquad \lim_{T \to \infty} \widehat{\rho}_T = \rho^* \qquad \text{a.s.}$$

where

(3.2)
$$\theta^* = \theta + \rho$$
 and $\rho^* = \frac{\theta \rho (\theta + \rho)}{(\theta + \rho)^2 + \theta \rho}$

Proof. We immediately deduce from (1.2) that

(3.3)
$$\int_0^T X_t \, \mathrm{d}X_t = \theta S_T + \rho P_T + M_T^X$$

where

(3.4)
$$S_T = \int_0^T X_t^2 dt, \qquad P_T = \int_0^T X_t V_t dt, \qquad M_T^X = \int_0^T X_t dW_t.$$

We shall see in Corollary 4.1 below that

(3.5)
$$\lim_{T \to \infty} \frac{1}{T} S_T = -\frac{1}{2(\theta + \rho)} \qquad \text{a.s.}$$

and in the proof of Corollary 4.2 that

(3.6)
$$\lim_{T \to \infty} \frac{1}{T} P_T = -\frac{1}{2(\theta + \rho)} \qquad \text{a.s.}$$

Moreover, if (\mathcal{F}_t) stands for the natural filtration of the standard Brownian motion (W_t) , then (M_t^X) is a continuous-time (\mathcal{F}_t) -martingale with quadratic variation S_t . Hence, it follows from the strong law of large numbers for continuous-time martingales given e.g. in [8] or [13], that $M_T^X = o(T)$ a.s. Consequently, we obtain from (3.3) that

(3.7)
$$\lim_{T \to \infty} \frac{1}{T} \int_0^T X_t \, \mathrm{d}X_t = -\frac{\theta}{2(\theta + \rho)} - \frac{\rho}{2(\theta + \rho)} = -\frac{1}{2} \qquad \text{a.s.}$$

which leads, via (2.1), to the first convergence in (3.1). The second convergence in (3.1) is more difficult to handle. We infer from (1.2) that

(3.8)
$$\int_0^T V_t \, \mathrm{d}V_t = \rho L_T + M_T^V$$

where

(3.9)
$$L_T = \int_0^T V_t^2 dt$$
 and $M_T^V = \int_0^T V_t dW_t.$

On the one hand, if $\rho < 0$, it is well-known, see e.g. [8] page 728, that

(3.10)
$$\lim_{T \to \infty} \frac{1}{T} L_T = -\frac{1}{2\rho} \qquad \text{a.s}$$

In addition, (M_t^V) is a continuous-time (\mathcal{F}_t) -martingale with quadratic variation L_t . Consequently, $M_T^V = o(T)$ a.s. and we find from (3.8) that

(3.11)
$$\lim_{T \to \infty} \frac{1}{T} \int_0^T V_t \, \mathrm{d}V_t = -\frac{1}{2} \qquad \text{a.s.}$$

However, we know from Itô's formula that

$$\frac{1}{T} \int_0^T X_t \, \mathrm{d}X_t = \frac{1}{2} \left(\frac{X_T^2}{T} - 1 \right) \quad \text{and} \quad \frac{1}{T} \int_0^T V_t \, \mathrm{d}V_t = \frac{1}{2} \left(\frac{V_T^2}{T} - 1 \right).$$

Then, we deduce from (3.7) and (3.11) that

(3.12)
$$\lim_{T \to \infty} \frac{X_T^2}{T} = 0 \quad \text{and} \quad \lim_{T \to \infty} \frac{V_T^2}{T} = 0 \quad \text{a.s.}$$

As $X_T = \theta \Sigma_T + V_T$, it clearly follows from (2.2) and (3.12) that

(3.13)
$$\lim_{T \to \infty} \frac{1}{2} \left(\frac{\widehat{V}_T^2}{T} - 1 \right) = -\frac{1}{2} \qquad \text{a.s.}$$

Hereafter, we have from (2.4) the decomposition

(3.14)
$$\widehat{\rho}_T = \frac{T}{2\widehat{L}_T} \left(\frac{\widehat{V}_T^2}{T} - 1 \right)$$

where

$$\widehat{L}_T = \int_0^T \widehat{V}_t^2 \,\mathrm{d}t.$$

We shall see in Corollary 4.2 below that

(3.15)
$$\lim_{T \to \infty} \frac{1}{T} \widehat{L}_T = -\frac{1}{2\rho^*} \qquad \text{a.s.}$$

Therefore, (3.14) together with (3.13) and (3.15) directly imply (3.1). On the other hand, if $\rho = 0$, it is clear from (1.2) that for all $t \ge 0$, $V_t = W_t$. Hence, we have from (2.2) and Itô's formula that

(3.16)
$$\widehat{V}_T^2 - T = 2M_T^W - 2W_T \Sigma_T (\widehat{\theta}_T - \theta) + \Sigma_T^2 (\widehat{\theta}_T - \theta)^2$$

and

(3.17)
$$\widehat{L}_T = L_T - 2(\widehat{\theta}_T - \theta) \int_0^T W_t \Sigma_t \, \mathrm{d}t + (\widehat{\theta}_T - \theta)^2 \int_0^T \Sigma_t^2 \, \mathrm{d}t$$

where

$$L_T = \int_0^T W_t^2 \, \mathrm{d}t \qquad \text{and} \qquad M_T^W = \int_0^T W_t \, \mathrm{d}W_t$$

It is now necessary to investigate the a.s. asymptotic behavior of L_T . We deduce from the self-similarity of the Brownian motion (W_t) that

(3.18)
$$L_T = \int_0^T W_t^2 \, \mathrm{d}t \stackrel{\mathcal{L}}{=} T \int_0^T W_{t/T}^2 \, \mathrm{d}t \stackrel{\mathcal{L}}{=} T^2 \int_0^1 W_s^2 \, \mathrm{d}s = T^2 L$$

5

Consequently, it clearly follows from (3.18) that for any power 0 < a < 2,

(3.19)
$$\lim_{T \to \infty} \frac{1}{T^a} L_T = +\infty \qquad \text{a.s}$$

As a matter of fact, since L is almost surely positive, it is enough to show that

(3.20)
$$\lim_{T \to \infty} \mathbb{E}\left[\exp\left(-\frac{1}{T^a}L_T\right)\right] = 0.$$

However, we have from standard Gaussian calculations, see e.g. [14] page 232, that

$$\mathbb{E}\left[\exp\left(-\frac{1}{T^a}L_T\right)\right] = \mathbb{E}\left[\exp\left(-\frac{T^2}{T^a}L\right)\right] = \frac{1}{\sqrt{\cosh(v_T(a))}}$$

where $v_T(a) = \sqrt{2T^{2-a}}$ goes to infinity, which clearly leads to (3.20). Furthermore, (M_t^W) is a continuous-time (\mathcal{F}_t) -martingale with quadratic variation L_t . We already saw that L_T goes to infinity a.s. which implies that $M_T^W = o(L_T)$ a.s. In addition, we obviously have $\Sigma_T^2 \leq TS_T$. One can observe that convergence (3.5) still holds when $\rho = 0$, which ensures that $\Sigma_T^2 \leq T^2$ a.s. Moreover, we deduce from the strong law of large numbers for continuous-time martingales that

$$(\widehat{\theta}_T - \theta)^2 = O\left(\frac{\log T}{T}\right)$$
 a.s.

which implies that $\Sigma_T^2(\widehat{\theta}_T - \theta)^2 = O(T \log T) = o(L_T)$ a.s. By the same token, as $X_T^2 = o(T)$ and $W_T^2 = o(T \log T)$ a.s., we find that

$$W_T \Sigma_T (\widehat{\theta}_T - \theta) = o(L_T)$$
 a.s.

Consequently, we obtain from (3.16) that

$$\widehat{V}_T^2 - T = o(L_T) \qquad \text{a.s.}$$

It remains to study the a.s. asymptotic behavior of \hat{L}_T . One can easily see that

$$\int_0^T \Sigma_t^2 \, \mathrm{d}t \le \frac{2}{\theta^2} (S_T + L_T).$$

However, it follows from (3.5) and (3.19) that $S_T = o(L_T)$ a.s. which ensures that

(3.22)
$$(\widehat{\theta}_T - \theta)^2 \int_0^T \Sigma_t^2 \, \mathrm{d}t = o(L_T)$$
 a.s.

Via the same arguments,

(3.23)
$$(\widehat{\theta}_T - \theta) \int_0^T W_t \Sigma_t \, \mathrm{d}t = o(L_T) \qquad \text{a.s.}$$

Then, we find from (3.17), (3.22) and (3.23) that

(3.24)
$$\widehat{L}_T = L_T(1+o(1))$$
 a.s

Finally, the second convergence in (3.1) follows from (3.21) and (3.24) which achieves the proof of Theorem 3.1.

Our second result deals with the asymptotic normality of our estimates

Theorem 3.2. If $\rho < 0$, we have the joint asymptotic normality

(3.25)
$$\sqrt{T} \begin{pmatrix} \widehat{\theta}_T - \theta^* \\ \widehat{\rho}_T - \rho^* \end{pmatrix} \xrightarrow{\mathcal{L}} \mathcal{N}(0, \Gamma)$$

where the asymptotic covariance matrix

(3.26)
$$\Gamma = \begin{pmatrix} \sigma_{\theta}^2 & \ell \\ \ell & \sigma_{\rho}^2 \end{pmatrix}$$

with
$$\sigma_{\theta}^{2} = -2\theta^{*}$$
, $\ell = \frac{2\rho^{*} ((\theta^{*})^{2} - \theta\rho)}{(\theta^{*})^{2} + \theta\rho}$ and
 $\sigma_{\rho}^{2} = -\frac{2\rho^{*} ((\theta^{*})^{6} + \theta\rho ((\theta^{*})^{4} - \theta\rho (2(\theta^{*})^{2} - \theta\rho)))}{((\theta^{*})^{2} + \theta\rho)^{3}}.$

In particular, we have

(3.27)
$$\sqrt{T}\left(\widehat{\theta}_T - \theta^*\right) \xrightarrow{\mathcal{L}} \mathcal{N}(0, \sigma_{\theta}^2),$$

and

(3.28)
$$\sqrt{T} \left(\widehat{\rho}_T - \rho^* \right) \xrightarrow{\mathcal{L}} \mathcal{N}(0, \sigma_{\rho}^2).$$

Proof. We obtain from (2.1) the decomposition

(3.29)
$$\widehat{\theta}_T - \theta^* = \frac{M_T^X}{S_T} + \frac{R_T^X}{S_T}$$

where

$$R_T^X = \rho \int_0^T X_t (V_t - X_t) \, \mathrm{d}t = -\theta \rho \int_0^T \Sigma_t \, \mathrm{d}\Sigma_t = -\frac{\theta \rho}{2} \Sigma_T^2.$$

We shall now establish a similar decomposition for $\hat{\rho}_T - \rho^*$. It follows from (2.2) that for all $0 \le t \le T$,

$$\widehat{V}_t = X_t - \widehat{\theta}_T \Sigma_t = V_t - (\widehat{\theta}_T - \theta) \Sigma_t = V_t - (\widehat{\theta}_T - \theta^*) \Sigma_t - \rho \Sigma_t, = V_t - \frac{\rho}{\theta} (X_t - V_t) - \frac{1}{\theta} (\widehat{\theta}_T - \theta^*) (X_t - V_t) = \frac{\theta^*}{\theta} V_t - \frac{\rho}{\theta} X_t - \frac{1}{\theta} (\widehat{\theta}_T - \theta^*) (X_t - V_t),$$

which leads to

(3.30)
$$\widehat{L}_T = I_T + (\widehat{\theta}_T - \theta^*) \left(J_T + (\widehat{\theta}_T - \theta^*) K_T \right),$$

where

$$I_{T} = \frac{1}{\theta^{2}} \Big(\rho^{2} S_{T} + (\theta^{*})^{2} L_{T} - 2\theta^{*} \rho P_{T} \Big),$$

$$J_{T} = \frac{1}{\theta^{2}} \Big(2\rho S_{T} + 2\theta^{*} L_{T} - 2(\theta + 2\rho) P_{T} \Big),$$

$$K_{T} = \frac{1}{\theta^{2}} \Big(S_{T} + L_{T} - 2P_{T} \Big).$$

7

Then, we deduce from (2.4) and (3.30) that

(3.31)
$$\widehat{L}_T\left(\widehat{\rho}_T - \rho^*\right) = \frac{I_T^V}{2} + \frac{1}{2}(\widehat{\theta}_T - \theta^*)\left(J_T^V + (\widehat{\theta}_T - \theta^*)K_T^V\right)$$

in which $I_T^V = \hat{V}_T^2 - T - 2\rho^* I_T$, $J_T^V = -2\rho^* J_T$, and $K_T^K = -2\rho^* K_T$. At this stage, in order to simplify the complicated expression (3.31), we make repeatedly use of Itô's formula. For all $0 \le t \le T$, we have

$$L_t = \frac{1}{2\rho} V_t^2 - \frac{1}{\rho} M_t^V - \frac{t}{2\rho},$$

$$P_t = \frac{1}{\theta^*} X_t V_t - \frac{1}{2\theta^*} V_t^2 - \frac{1}{\theta^*} M_t^X - \frac{t}{2\theta^*},$$

$$S_t = \frac{1}{2\theta} X_t^2 + \frac{\rho}{2\theta^*\theta} V_t^2 - \frac{\rho}{\theta^*\theta} X_t V_t - \frac{1}{\theta^*} M_t^X - \frac{t}{2\theta^*},$$

where the continuous-time martingales M_t^X and M_t^V were previously defined in (3.4) and (3.9). Therefore, it follows from tedious but straightforward calculations that

(3.32)
$$\widehat{L}_{T}(\widehat{\rho}_{T} - \rho^{*}) = C_{X}M_{T}^{X} + C_{V}M_{T}^{V} + \frac{J_{T}^{V}}{2}(\widehat{\theta}_{T} - \theta^{*}) + R_{T}^{V}$$

where

$$C_V = \frac{(\theta^*)^2 \rho^*}{\theta^2 \rho}$$
 and $C_X = -\frac{\rho(2\theta + \rho)\rho^*}{\theta^2 \theta^*}$

The remainder R_T^V is similar to R_T^X and they play a negligible role. The combination of (3.29) and (3.32) leads to the vectorial expression

(3.33)
$$\sqrt{T} \begin{pmatrix} \widehat{\theta}_T - \theta^* \\ \widehat{\rho}_T - \rho^* \end{pmatrix} = \frac{1}{\sqrt{T}} A_T Z_T + \sqrt{T} R_T$$

where

$$A_T = \begin{pmatrix} S_T^{-1}T & 0\\ B_T \widehat{L}_T^{-1}T & C_V \widehat{L}_T^{-1}T \end{pmatrix}, \qquad R_T = \begin{pmatrix} S_T^{-1} R_T^X\\ \widehat{L}_T^{-1} D_T \end{pmatrix}$$

with $B_T = C_X + J_T^V (2S_T)^{-1}$ and $D_T = R_T^V + J_T^V (2S_T)^{-1} R_T^X$. The leading term in (3.33) is the continuous-time vector (\mathcal{F}_t) -martingale (Z_t) with predictable quadratic variation $\langle Z \rangle_t$ given by

$$Z_t = \begin{pmatrix} M_t^X \\ M_t^V \end{pmatrix} \quad \text{and} \quad \langle Z \rangle_t = \begin{pmatrix} S_t & P_t \\ P_t & L_t \end{pmatrix}$$

We deduce from (3.5), (3.6) and (3.10) that

$$(3.34) \qquad \lim_{T \to \infty} A_T = A \qquad \text{a.s}$$

where A is the limiting matrix given by

$$A = \begin{pmatrix} -2\theta^* & 0\\ -2\rho^*(C_X - 2(\theta\rho)^{-1}\theta^*\rho^*) & -2\rho^*C_V \end{pmatrix}.$$

By the same token, we immediately have from (3.5), (3.6) and (3.10) that

(3.35)
$$\lim_{T \to \infty} \frac{\langle Z \rangle_T}{T} = \Lambda = -\frac{1}{2\theta^*} \begin{pmatrix} 1 & 1\\ 1 & \theta^* \rho^{-1} \end{pmatrix} \quad \text{a.s.}$$

Furthermore, it clearly follows from Corollary 4.3 below that

(3.36)
$$\frac{X_T^2}{\sqrt{T}} \xrightarrow{\mathcal{P}} 0$$
 and $\frac{V_T^2}{\sqrt{T}} \xrightarrow{\mathcal{P}} 0.$

Finally, as $\Gamma = A\Lambda A'$, the joint asymptotic normality (3.25) follows from the conjunction of (3.33), (3.34), (3.35), (3.36) together with Slutsky's lemma and the central limit theorem for continuous-time vector martingales given e.g. in [8], which achieves the proof of Theorem 3.2.

Theorem 3.3. If $\rho = 0$, we have the convergence in distribution

$$(3.37) T \,\widehat{\rho}_T \stackrel{\mathcal{L}}{\longrightarrow} \mathcal{W}$$

where the limiting distribution \mathcal{W} is given by

(3.38)
$$\mathcal{W} = \frac{\int_0^1 B_s \, \mathrm{d}B_s}{\int_0^1 B_s^2 \, \mathrm{d}s} = \frac{B_1^2 - 1}{2\int_0^1 B_s^2 \, \mathrm{d}s}$$

and (B_t) is a standard Brownian motion.

Proof. Via the same reasoning as in Section 2 of [9], it follows from the self-similarity of the Brownian motion (W_t) that

(3.39)
$$\left(\int_0^T W_t^2 \, \mathrm{d}t, \, \frac{1}{2} \left(W_T^2 - T \right) \right) \stackrel{\mathcal{L}}{=} \left(T \int_0^T W_{t/T}^2 \, \mathrm{d}t, \, \frac{T}{2} \left(W_1^2 - 1 \right) \right), \\ = \left(T^2 \int_0^1 W_s^2 \, \mathrm{d}s, \, \frac{T}{2} \left(W_1^2 - 1 \right) \right).$$

Moreover, we obtain from (3.30) that

$$(3.40) \qquad \qquad \widehat{L}_T = \alpha_T S_T + \beta_T P_T + \gamma_T L_T$$

where

$$\alpha_T = \frac{1}{\theta^2} (\widehat{\theta}_T - \theta)^2,$$

$$\beta_T = -\frac{2}{\theta} (\widehat{\theta}_T - \theta) - \frac{2}{\theta^2} (\widehat{\theta}_T - \theta)^2,$$

$$\gamma_T = 1 + \frac{2}{\theta} (\widehat{\theta}_T - \theta) + \frac{1}{\theta^2} (\widehat{\theta}_T - \theta)^2.$$

By Theorem 3.1, $\hat{\theta}_T$ converges a.s. to θ which implies that α_T , β_T , and γ_T converge a.s. to 0, 0 and 1. Hence, we deduce from (3.5), (3.6) and (3.40) that

(3.41)
$$\widehat{L}_T = L_T(1+o(1))$$
 a.s.

Furthermore, one can observe that \hat{V}_T^2/T shares the same asymptotic distribution as W_T^2/T . Finally, (3.37) follows from (3.39) and (3.41) together with the continuous mapping theorem.

Remark 3.1. The asymptotic behavior of $\hat{\rho}_T$ when $\rho < 0$ and $\rho = 0$ is closely related to the results previously established for the unstable discrete-time autoregressive process, see [4], [9], [18]. According to Corollary 3.1.3 of [4], we can express

$$\mathcal{W} = \frac{\mathcal{T}^2 - 1}{2\mathcal{S}}$$

where \mathcal{T} and \mathcal{S} are given by the Karhunen-Loeve expansions

$$\mathcal{T} = \sqrt{2} \sum_{n=1}^{\infty} \gamma_n Z_n$$
 and $\mathcal{S} = \sum_{n=1}^{\infty} \gamma_n^2 Z_n^2$

with $\gamma_n = 2(-1)^n/((2n-1)\pi)$ and (Z_n) is a sequence of independent random variables with $\mathcal{N}(0,1)$ distribution.

Remark 3.2. For all $0 \le t \le T$, the residuals \widehat{V}_t given by (2.2) depend on $\widehat{\theta}_T$. It would have been more natural to make use of the estimator of θ at stage t instead of stage T, in order to produce a recursive estimate. In this situation, Theorem 3.1 still holds but we have been unable to prove Theorem 3.2.

4. SOME TECHNICAL TOOLS

First of all, most of our results rely on the following keystone lemma.

Lemma 4.1. The process (X_t) is geometrically ergodic.

Proof. It follows from (1.2) that

(4.1)
$$dX_t = (\theta + \rho)X_t dt - \theta \rho \Sigma_t dt + dW_t$$

where we recall that

$$\Sigma_t = \int_0^t X_s \, \mathrm{d}s$$

Consequently, if

$$\Phi_t = \begin{pmatrix} X_t \\ \Sigma_t \end{pmatrix},$$

we clearly deduce from (4.1) that

$$\mathrm{d}\Phi_t = A\Phi_t \,\mathrm{d}t + \,\mathrm{d}B_t$$

where

$$A = \begin{pmatrix} \theta + \rho & -\theta\rho \\ 1 & 0 \end{pmatrix} \quad \text{and} \quad B_t = \begin{pmatrix} W_t \\ 0 \end{pmatrix}.$$

The geometric ergodicity of (Φ_t) only depends on the sign of $\lambda_{\max}(A)$, *i.e.* the largest eigenvalue of A, which has to be negative. An immediate calculation shows that

$$\lambda_{\max}(A) = \max(\theta, \rho)$$

which ensures that $\lambda_{\max}(A) < 0$ as soon as $\rho < 0$. Moreover, if $\rho = 0$, (X_t) is an ergodic Ornstein-Uhlenbeck process since $\theta < 0$, which completes the proof of Lemma 4.1.

Corollary 4.1. We have the almost sure convergence

(4.2)
$$\lim_{T \to \infty} \frac{1}{T} S_T = -\frac{1}{2(\theta + \rho)} \qquad \text{a.s.}$$

Proof. According to Lemma 4.1, it is only necessary to establish the asymptotic behavior of $\mathbb{E}[X_t^2]$. Denote $\alpha_t = \mathbb{E}[X_t^2]$, $\beta_t = \mathbb{E}[\Sigma_t^2]$ and $\gamma_t = \mathbb{E}[X_t\Sigma_t]$. One obtains from Itô's formula that

$$\frac{\partial U_t}{\partial t} = CU_t + I$$

where

$$U_t = \begin{pmatrix} \alpha_t \\ \beta_t \\ \gamma_t \end{pmatrix}, \quad C = \begin{pmatrix} 2(\theta + \rho) & 0 & -2\theta\rho \\ 0 & 0 & 2 \\ 1 & -\theta\rho & \theta + \rho \end{pmatrix}, \quad I = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}$$

It is not hard to see that $\lambda_{\max}(C) = \max(\theta + \rho, 2\theta, 2\rho)$. On the one hand, if $\rho < 0$, $\lambda_{\max}(C) < 0$ which implies that

$$\lim_{t \to \infty} U_t = -C^{-1}I.$$

It means that

$$\lim_{t \to \infty} \alpha_t = -\frac{1}{2(\theta + \rho)}, \quad \lim_{t \to \infty} \beta_t = -\frac{1}{2\theta\rho(\theta + \rho)}, \quad \lim_{t \to \infty} \gamma_t = 0.$$

Hence, (4.2) follows from Lemma 4.1 together with the ergodic theorem. On the other hand, if $\rho = 0$, (X_t) is a positive recurrent Ornstein-Uhlenbeck process and convergence (4.2) is well-known.

Corollary 4.2. If $\rho < 0$, we have the almost sure convergence

$$\lim_{T \to \infty} \frac{1}{T} \widehat{L}_T = -\frac{(\theta + \rho)^2 + \theta\rho}{2\theta\rho(\theta + \rho)} \qquad \text{a.s.}$$

Proof. If $\rho < 0$, (V_t) is a positive recurrent Ornstein-Uhlenbeck process and it is well-known that

$$\lim_{T \to \infty} \frac{1}{T} L_T = -\frac{1}{2\rho} \qquad \text{a.s}$$

In addition, as $X_t = \theta \Sigma_t + V_t$,

$$\int_0^T X_t \Sigma_t \, \mathrm{d}t = \frac{1}{\theta} (S_T - P_T).$$

However, we already saw in the proof of Corollary 4.1 that

$$\lim_{T \to \infty} \frac{1}{T} \int_0^T X_t \Sigma_t \, \mathrm{d}t = 0 \qquad \text{a.s.}$$

which leads, via (4.2), to the almost sure convergence

$$\lim_{T \to \infty} \frac{P_T}{T} = -\frac{1}{2(\theta + \rho)} \qquad \text{a.s.}$$

Consequently, we deduce from (3.1) together with (3.30) that

$$\lim_{T \to \infty} \frac{1}{T} \widehat{L}_T = \lim_{T \to \infty} \frac{1}{T} I_T = -\frac{(\theta + \rho)^2 + \theta \rho}{2\theta \rho (\theta + \rho)} \qquad \text{a.s}$$

which achieves the proof of Corollary 4.2.

Corollary 4.3. If $\rho < 0$, we have the asymptotic normalities

$$X_T \xrightarrow{\mathcal{L}} \mathcal{N}\left(0, -\frac{1}{2(\theta + \rho)}\right) \quad and \quad V_T \xrightarrow{\mathcal{L}} \mathcal{N}\left(0, -\frac{1}{2\rho}\right).$$

The asymptotic normality of X_T still holds in the particular case where $\rho = 0$.

Proof. This asymptotic normality is a well-known result for the Ornstein-Uhlenbeck process (V_t) with $\rho < 0$. In addition, one can observe that for all $t \ge 0$, $\mathbb{E}[X_t] = 0$. The end of the proof is a direct consequence of the Gaussianity of (X_t) together with Lemma 4.1 and Corollary 4.1.

5. A STATISTICAL TESTING PROCEDURE

Our purpose is now to propose a statistical procedure in order to test

$$\mathcal{H}_0$$
: " $\rho = 0$ " against \mathcal{H}_1 : " $\rho < 0$ ".

We shall make use of the Durbin-Watson statistic given by (2.5). Its asymptotic properties are as follows.

Theorem 5.1. We have the almost sure convergence

(5.1)
$$\lim_{T \to \infty} \widehat{D}_T = D^* \qquad \text{a.s.}$$

where $D^* = 2(1 - \rho^*)$. In addition, if $\rho < 0$, we have the asymptotic normality

(5.2)
$$\sqrt{T}\left(\widehat{D}_T - D^*\right) \xrightarrow{\mathcal{L}} \mathcal{N}(0, \sigma_D^2)$$

where

$$\sigma_D^2 = 4 \, \sigma_\rho^2 = -\frac{8\rho^* \left((\theta^*)^6 + \theta \rho \left((\theta^*)^4 - \theta \rho \left(2(\theta^*)^2 - \theta \rho \right) \right) \right)}{\left((\theta^*)^2 + \theta \rho \right)^3}.$$

while, if $\rho = 0$,

(5.3)
$$T\left(\widehat{D}_T-2\right) \xrightarrow{\mathcal{L}} -2\mathcal{W}$$

with W given by (3.38).

Proof. The proof of Theorem 5.1 is a straightforward application of (3.1), (3.28) and (3.37) since $\hat{D}_T = 2(1 - \hat{\rho}_T)$.

From now on, let us define the test statistic

$$\widehat{Z}_T = T^2 \left(\widehat{D}_T - 2\right)^2$$

It follows from Theorem 5.1 that under \mathcal{H}_0 ,

$$\widehat{Z}_T \xrightarrow{\mathcal{L}} 4\mathcal{W}^2$$

while, under \mathcal{H}_1 ,

$$\lim_{T \to \infty} \widehat{Z}_T = +\infty \qquad \text{a.s.}$$

From a practical point of view, for a significance level α where $0 < \alpha < 1$, the acceptance and rejection regions are given by $\mathcal{A} = [0, z_{\alpha}]$ and $\mathcal{R} =]z_{\alpha}, +\infty[$ where z_{α} stands for the $(1 - \alpha)$ -quantile of the distribution of $4\mathcal{W}^2$. The null hypothesis \mathcal{H}_0 will not be rejected if the empirical value

$$\widehat{Z}_T \le z_\alpha$$

and will be rejected otherwise. Assume to conclude that \mathcal{H}_0 is rejected, which means that we admit the existence of a serial correlation $\rho < 0$. Then, the best way to produce unbiased estimates is to study the process given by (4.1). As a matter of fact, for all $t \geq 0$,

$$X_t = (\theta + \rho)\Sigma_t - \theta\rho\Pi_t + W_t$$

where

$$\Sigma_t = \int_0^t X_s \, \mathrm{d}s$$
 and $\Pi_t = \int_0^t \Sigma_s \, \mathrm{d}s.$

The maximum likelihood estimator of the vector

$$\vartheta = \begin{pmatrix} \theta + \rho \\ -\theta \rho \end{pmatrix}$$

is given by

$$\widehat{\vartheta}_T = \left(\int_0^T \Phi_t \Phi_t' \,\mathrm{d}t\right)^{-1} \int_0^T \Phi_t \,\mathrm{d}X_t$$

where $\Phi_t = (X_t \Sigma_t)'$. We can show the almost sure convergence

$$\lim_{T \to \infty} \widehat{\vartheta}_T = \vartheta \qquad \text{a.s}$$

as well as the asymptotic normality

$$\sqrt{T}\left(\widehat{\vartheta}_T - \vartheta\right) \xrightarrow{\mathcal{L}} \mathcal{N}(0, \Delta)$$

where the asymptotic covariance matrix is given by

$$\Delta = \begin{pmatrix} -2\theta^* & 0\\ 0 & -2\theta\rho\,\theta^* \end{pmatrix}.$$

Accordingly, the maximum likelihood estimator $\widehat{\vartheta}_T$ is strongly consistent and one can see that its components are asymptotically independent.

References

- BARNDORFF-NIELSEN, O. E., AND BASSE-O'CONNOR, A. Quasi Ornstein-Uhlenbeck processes. *Bernoulli* 17, 3 (2011), 916–941.
- [2] BARNDORFF-NIELSEN, O. E., AND SHEPHARD, N. Non-Gaussian Ornstein-Uhlenbeck-based models and some of their uses in financial economics. J. R. Stat. Soc. Ser. B Stat. Methodol. 63, 2 (2001), 167–241.
- [3] BERCU, B., AND PROÏA, F. A sharp analysis on the asymptotic behavior of the Durbin-Watson statistic for the first-order autoregressive process. *ESAIM Probab. Stat.* 16 (2012).
- [4] CHAN, N. H., AND WEI, C. Limiting distributions of least squares estimates of unstable autoregressive processes. Ann. Statist. 16 (1) (1988), 367–401.
- [5] DURBIN, J., AND WATSON, G. S. Testing for serial correlation in least squares regression. I. Biometrika 37 (1950), 409–428.
- [6] DURBIN, J., AND WATSON, G. S. Testing for serial correlation in least squares regression. II. Biometrika 38 (1951), 159–178.
- [7] DURBIN, J., AND WATSON, G. S. Testing for serial correlation in least squares regession. III. Biometrika 58 (1971), 1–19.
- [8] FEIGIN, P. Maximum likelihood estimation for continuous-time stochastic processes. Advances in Appl. Probability 8, 4 (1976), 712–736.
- [9] FEIGIN, P. Some comments concerning a curious singularity. J. Appl. Probab. 16, 2 (1979), 440–444.
- [10] HU, Y., AND LONG, H. Parameter estimation for Ornstein-Uhlenbeck processes driven by α-stable Lévy motions. Commun. Stoch. Anal. 1, 2 (2007), 175–192.
- [11] JONGBLOED, G., VAN DER MEULEN, F. H., AND VAN DER VAART, A. W. Nonparametric inference for Lévy-driven Ornstein-Uhlenbeck processes. *Bernoulli* 11, 5 (2005), 759–791.
- [12] KUTOYANTS, Y. Statistical inference for ergodic diffusion processes. Springer Series in Statistics. Springer-Verlag London Ltd., London, 2004.
- [13] LÉPINGLE, D. Sur le comportement asymptotique des martingales locales. In Séminaire de Probabilités, XII, vol. 649 of Lecture Notes in Math. Springer, Berlin, 1978, pp. 148–161.
- [14] LIPTSER, R., AND SHIRYAEV, A. Statistics of random processes. II, vol. 6 of Applications of Mathematics (New York). Springer-Verlag, Berlin, 2001.
- [15] ONALAN, O. Financial modelling with ornstein-uhlenbeck processes driven by Lévy process. Proceedings of the world congress engineering 2 (2009), 1–6.
- [16] SCHOUTENS, W. Stochastic processes and orthogonal polynomials, vol. 146 of Lecture Notes in Statistics. Springer-Verlag, New York, 2000.
- [17] UHLENBECK, G. E., AND ORNSTEIN, L. S. On the theory of brownian motion. Phys. Rev. 36 (1930), 823–841.
- [18] WHITE, J. S. The limiting distribution of the serial correlation coefficient in the explosive case. Ann. Math. Statist. 29 (1958), 1188–1197.

E-mail address: Bernard.Bercu@math.u-bordeaux1.fr *E-mail address*: Frederic.Proia@inria.fr *E-mail address*: Nicolas.Savy@math.univ-toulouse.fr

UNIVERSITÉ BORDEAUX 1, INSTITUT DE MATHÉMATIQUES DE BORDEAUX, UMR 5251, AND INRIA BORDEAUX SUD-OUEST, TEAM ALEA, 351 COURS DE LA LIBÉRATION, 33405 TALENCE CEDEX, FRANCE.

UNIVERSITÉ PAUL SABATIER, INSTITUT DE MATHÉMATIQUES DE TOULOUSE, UMR C5583, 31062 TOULOUSE CEDEX 09, FRANCE.