Convolution type form of the Ito representation of the infinitesimal generator for Levy processes

Lev Sakhnovich

99 Cove ave., Milford, CT, 06461, USA E-mail: lsakhnovich@gmail.com

Mathematics Subject Classification (2010): Primary 60G51; Secondary 60J45; 45A05

Keywords. Semigroup, generator, Ito form, convolution form, potential, quasi-potential.

Abstract

In the present paper we show that the Ito representation of the infinitesimal generator L for Levy processes can be written in a convolution type form. Using the obtained convolution form and the theory of integral equations with difference kernels we study the properties of Levy processes.

1 Main notions

Let us introduce the notion of the Levy processes.

Definition 1.1 A stochastic process $\{X_t : t \geq 0\}$ is called Levy process , if the following conditions are fulfilled:

1. Almost surely $X_0 = 0$, i.e. $P(X_0 = 0) = 1$.

(One says that an event happens almost surely $(a.s.)$ if it happens with probability one.)

2. For any $0 \le t_1 < t_2 ... < t_n < \infty$ the random variables $X_{t_2}-X_{t_1}, X_{t_3}-X_{t_4},...,X_{t_n}-X_{t_{n-1}}$

are independent (independent increments).

(To call the increments of the process X_t independent means that increments $X_{t_2} - X_{t_1}, X_{t_3} - X_{t_4}, ..., X_{t_n} - X_{t_{n-1}}$ are mutually (not just pairwise) independent.)

3. For any $s < t$ the distributions of $X_t - X_s$ and X_{t-s} are equal (stationary increments).

4. Process X_t is almost surely right continuous with left limits. Then Levy-Khinchine formula gives (see [\[1\]](#page-7-0), [\[5\]](#page-7-1))

$$
\mu(z,t) = E\{\exp[izX_t]\} = \exp[-t\lambda(z)], \quad t \ge 0,
$$
\n(1.1)

where

$$
\lambda(z) = \frac{1}{2}Az^2 - i\gamma z - \int_{-\infty}^{\infty} (e^{ixz} - 1 - ixz1_{|x| < 1})\mu(dx). \tag{1.2}
$$

Here $A\geq 0$, $\gamma = \overline{\gamma}$, $z = \overline{z}$ and $\mu(dx)$ is a measure on the axis $(-\infty, \infty)$ satisfying the conditions

$$
\int_{-\infty}^{\infty} \frac{x^2}{1+x^2} \mu(dx) < \infty. \tag{1.3}
$$

The Levy-Khinchine formula is determined by the Levy-Khinchine triplet $(A, \gamma, \mu(dx)).$

By $P_t(x_0, \Delta)$ we denote the probability $P(X_t \in \Delta)$ when $P(X_0 = x_0) = 1$ and $\Delta \in R$. The transition operator P_t is defined by the formula

$$
P_t f(x) = \int_{-\infty}^{\infty} P_t(x, dy) f(y).
$$
 (1.4)

Let C_0 be the Banach space of continuous functions $f(x)$, satisfying the condition $\lim f(x) = 0$, $|x| \to \infty$ with the norm $||f|| = \sup_x |f(x)|$. We denote by C_0^n the set of $f(x) \in C_0$ such that $f^{(k)}(x) \in C_0$, $(1 \leq k \leq n)$. It is known that [\[5\]](#page-7-1)

$$
P_t f \in C_0,\tag{1.5}
$$

if $f(x) \in C_0^2$.

Now we formulate the following important result (see [\[5\]](#page-7-1)) .

Theorem 1.2 (Levy-Ito decomposition.) The family of the operators P_t (t≥0) defined by the Levy process X_t is a strongly continuous semigroup on C_0 with the norm $||P_t|| = 1$. Let L be its infinitesimal generator. Then

$$
Lf = \frac{1}{2}A\frac{d^2f}{dx^2} + \gamma\frac{df}{dx} + \int_{-\infty}^{\infty} (f(x+y) - f(x) - y\frac{df}{dx}1_{|y|<1})\mu(dy), \qquad (1.6)
$$

where $f \in C_0^2$.

2 Convolution type form of infinitesimal generator

1. In this section we prove that the infinitesimal generator L can be represented in the special convolution type form

$$
Lf = \frac{d}{dx} S \frac{d}{dx} f,\tag{2.1}
$$

where the operator S is defined by the relation

$$
Sf = \frac{1}{2}Af + \int_{-\infty}^{\infty} k(y - x)f(y)dy.
$$
 (2.2)

We note that for arbitrary $a (0 < a < \infty)$ the inequality

$$
\int_{-a}^{a} |k(t)| dt < \infty \tag{2.3}
$$

is true.

Formula [\(2.2\)](#page-2-0) was proved before in our works [\[4\]](#page-7-2) under some additional conditions. In the present paper we omit these additional conditions and prove the formula (2.2) for the general case. The representation of L in form (2.1) is convenient as the operator L is expressed with the help of the classic differential and convolution operators. Using the obtained convolution form of the generator L and the theory of integral equations with difference kernels [\[3\]](#page-7-3) we investigate the properties of a wide class of Levy processes.

By $C(a)$ we denote the set of functions $f(x) \in C_0$ which have the following property:

$$
f(x) = 0, \quad x \notin [-a, a] \tag{2.4}
$$

i.e. the function $f(x)$ is equal to zero in the neighborhood of $x = \infty$. We note, that parameter a can be different for different f .

We introduce the functions

$$
\mu_{-}(x) = \int_{-\infty}^{x} \mu(dx), \, x < 0,\tag{2.5}
$$

$$
\mu_{+}(x) = -\int_{x}^{\infty} \mu(dx), \, x > 0,\tag{2.6}
$$

where the functions $\mu_-(x)$ and $\mu_+(x)$ are monotonically increasing on the half-axis $(-\infty, 0]$ and $[0, \infty)$ respectively and

$$
\mu_+(x) \to 0, \ x \to +\infty; \ \mu_-(x) \to 0, \ x \to -\infty. \tag{2.7}
$$

We note that

$$
\mu_{-}(x) \ge 0, \ x < 0; \ \mu_{+}(x) \le 0, \ x > 0. \tag{2.8}
$$

In view of (1.3) the integrals in the right sides of (2.6) and (2.5) are convergent. Hence we have

$$
\int_{-\infty}^{\infty} f(x)\mu(dx) = \int_{-\infty}^{0} f(x)d\mu_{-}(x) + \int_{0}^{\infty} f(x)d\mu_{+}(x).
$$
 (2.9)

Theorem 2.1 The following relations

$$
\varepsilon^2 \mu_{\pm}(\pm \varepsilon) \to 0, \ \varepsilon \to + 0,\tag{2.10}
$$

$$
-\int_{-a}^{0} x\mu_{-}(x)dx < \infty, -\int_{0}^{a} x\mu_{+}(x)dx < \infty, 0 < a < \infty \tag{2.11}
$$

are true

Proof. According to [\(1.3\)](#page-1-0) we have

$$
0 \le \int_{-a}^{-\varepsilon} x^2 d\mu_-(x) \le M,\tag{2.12}
$$

where M does not depend from ε . Integrating by parts the integral of [\(2.12\)](#page-3-1) we obtain:

$$
\int_{-a}^{-\varepsilon} x^2 d\mu_-(x) = \varepsilon^2 \mu_-(-\varepsilon) - a^2 \mu_-(-a) - 2 \int_{-a}^{-\varepsilon} x \mu_-(x) dx \le M, \quad (2.13)
$$

The function $-\int_{-a}^{\infty} x\mu_-(x)dx$ of ε is monotonic increasing. According to [\(2.13\)](#page-3-2) this function is bounded. Hence we have

$$
\lim_{\varepsilon \to +0} \int_{-a}^{-\varepsilon} x \mu_-(x) dx = \int_{-a}^0 x \mu_-(x) dx \tag{2.14}
$$

It follows from (2.13) and (2.14) that

$$
\lim_{\varepsilon \to +0} \varepsilon^2 \mu_-(-\varepsilon) = 0. \tag{2.15}
$$

Thus, relations [\(2.10\)](#page-3-4) and [\(2.11\)](#page-3-5) are proved for $\mu_-(x)$. In the same way relations [\(2.10\)](#page-3-4) and [\(2.11\)](#page-3-5) can be proved for $\mu_+(x)$. \Box

2. Let us introduce the functions

$$
k_{-}(x) = \int_{-1}^{x} \mu_{-}(t)dt, \ -\infty \le x < 0,\tag{2.16}
$$

$$
k_{+}(x) = -\int_{x}^{1} \mu_{+}(t)dt, \ 0 < x \leq +\infty. \tag{2.17}
$$

In view of (2.11) the integrals in the right sides of (2.16) and (2.17) are absolutely convergent. From (2.16) and (2.17) we obtain the assertions:

Theorem 2.2 1. The function $k_-(x)$ is monotonically increasing on the $(-\infty, 0)$ and

$$
k_{-}(x) \ge 0, \ -1 \le x < 0. \tag{2.18}
$$

2. The function $k_{+}(x)$ is monotonically decreasing on the $(0, +\infty)$ and

$$
k_+(x) \ge 0, \ 0 < x \le 1. \tag{2.19}
$$

Further we need the following result:

Theorem 2.3 The following relations

$$
\varepsilon k_-(-\varepsilon) \to 0, \ \varepsilon \to + 0; \ \varepsilon k_+(\varepsilon) \to 0, \ \varepsilon \to + 0; \tag{2.20}
$$

$$
\int_{-1}^{0} k_{-}(x)dx < \infty; \int_{0}^{1} k_{+}(x)dx < \infty \tag{2.21}
$$

are valid.

Proof. According to [\(2.11\)](#page-3-5) we have

$$
0 \le -\int_{-1}^{-\varepsilon} x\mu_-(x)dx \le M,\tag{2.22}
$$

where M does not depend from ε . Integrating by parts the integral of [\(2.22\)](#page-4-2) we obtain:

$$
-\int_{-1}^{-\varepsilon} x\mu_-(x)dx = \varepsilon k_-(-\varepsilon) - k_-(-1) + \int_{-1}^{-\varepsilon} k_-(x)dx \le M,\tag{2.23}
$$

The function $\int_{-1}^{-\varepsilon} k_-(x)dx$ of ε is monotonic increasing. This function is bounded $(see (2.23))$ $(see (2.23))$ $(see (2.23))$. Hence we have

$$
\lim_{\varepsilon \to +0} \int_{-1}^{-\varepsilon} k_-(x) dx = \int_{-1}^0 k_-(x) dx \tag{2.24}
$$

It follows from [\(2.23\)](#page-5-0) and [\(2.24\)](#page-5-1) that

$$
\lim_{\varepsilon \to +0} \varepsilon k_-(-\varepsilon) = 0. \tag{2.25}
$$

Thus, relations [\(2.20\)](#page-4-3) and [\(2.21\)](#page-4-4) are proved for $k_-(x)$. In the same way relations [\(2.20\)](#page-4-3) and [\(2.21\)](#page-4-4) can be proved for $k_+(x)$. \Box

3. We use the following notation

$$
J(f) = J_1(f) + J_2(f), \tag{2.26}
$$

where

$$
J_1(f) = \frac{d}{dx} \int_{-\infty}^x f'(y)k_-(y-x)dy, \quad f(x) \in C(a), \tag{2.27}
$$

$$
J_2(f) = \frac{d}{dx} \int_x^{\infty} f'(y)k_+(y-x)dy, \quad f(x) \in C(a).
$$
 (2.28)

Lemma 2.4 The operator $J(f)$ defined by [\(2.26\)](#page-5-2) can be represented in the form

$$
J(f) = \int_{-\infty}^{\infty} [f(y+x) - f(x) - y \frac{df(x)}{dx} 1_{|y| \le 1}] \mu(dy) + \Gamma f'(x), \tag{2.29}
$$

where $\Gamma = \overline{\Gamma}$ and $f(x) \in C(a)$.

Proof. From [\(2.27\)](#page-5-3) we obtain the relation

$$
J_1(f) = -\int_{x-1}^x [f'(y) - f'(x)]k'_{-}(y-x)dy - \int_{-a}^{x-1} f'(y)k'_{-}(y-x)dy. \tag{2.30}
$$

By proving (2.30) we used relations $(2.16),(2.20)$ $(2.16),(2.20)$ and equality

$$
\int_{x-1}^{x} k_{-}(y-x) dy = \int_{-1}^{0} k_{-}(v) dv.
$$
 (2.31)

We introduce the notations

$$
P_1(x, y) = f(y) - f(x) - (y - x)f'(x), \quad P_2(x, y) = f(y) - f(x). \tag{2.32}
$$

Using notations (2.32) we represent (2.30) in the form

$$
J_1(f) = -\int_{-1}^{0} \frac{\partial}{\partial y} P_1(x, y+x)\mu_-(y) dy - \int_{-a-x}^{-1} \frac{\partial}{\partial y} P_2(x, y+x)\mu_-(y) dy, \tag{2.33}
$$

Integrating by parts the integrals of [\(2.33\)](#page-6-2) we deduce that

$$
J_1(f) = f'(x)\gamma_1 + \int_{-1}^0 P_1(x, y+x)d\mu_-(y) + \int_{-a-x}^{-1} P_2(x, y+x)d\mu_-(y) + P_2(x, -a)\mu_-(-a),
$$
\n(2.34)

where $\gamma_1 = k'_{-}(-1)$. It follows from [\(1.3\)](#page-1-0) that the integrals in [\(2.34\)](#page-6-3) are absolutely convergent. Passing to the limit in (2.34) , when $a \rightarrow +\infty$, and taking into account (2.30) , (2.31) we have

$$
J_1(f) = \int_{-\infty}^x [f(y+x) - f(x) - y \frac{df(x)}{dx} 1_{|y| \le 1}] d\mu_-(y) + \gamma_1 f'(x). \tag{2.35}
$$

In the same way it can be proved that

$$
J_2(f) = \int_x^{\infty} [f(y+x) - f(x) - y \frac{df(x)}{dx} 1_{|y| \le 1}] d\mu_+(y) + \gamma_2 f'(x), \qquad (2.36)
$$

where $\gamma_2 = k'_+(1)$. The relation [\(2.29\)](#page-5-4) follows directly from [\(2.35\)](#page-6-5) and [\(2.36\)](#page-6-6). Here $\Gamma = \gamma_1 + \gamma_2$. The lemma is proved. \Box

Remark 2.1 The operator $L_0 f = \frac{d}{dx} f$ can be represented in form [\(2.1\)](#page-2-1), (2.2) , where

$$
S_0 f = \int_{-\infty}^{\infty} p_0(x - y) f(y) dy,
$$
\n(2.37)

$$
p_0(x) = \frac{1}{2}\text{sign}(x).
$$
 (2.38)

From Lemmas [2.4,](#page-5-5) and Remark [2.1](#page-6-7) we deduce the following assertion.

Theorem 2.5 The infinitesimal generator L has a convolution type form $(2.1), (2.2).$ $(2.1), (2.2).$ $(2.1), (2.2).$ $(2.1), (2.2).$

References

- [1] J. Bertoin, Levy Processes. University Press, Cambridge, 1996.
- [2] K. Ito, On Stochastic Differential Equations. Memoirs Amer. Math. Soc. 4, 1951.
- [3] L. A. Sakhnovich, Integral Equations with Difference Kernels. Operator Theory 84, Birkhäuser, 1996.
- [4] L. A. Sakhnovich, Levy Processes, Integral Equations, Statistical Physics: Connections and Interactions. Operator Theory 225, Birkhäuser, 2012.
- [5] K.Sato, Levy Processes and Infinitely Divisible Distributions. University Press, Cambridge, 1999.
- [6] M. Thomas and O. Barndorff (ed.), Levy Processes: Theory and Applications. Birkhäuser, 2001.