Epistasis not needed to explain low dN/dS

David M. McCandlish¹, Etienne Rajon¹, Premal Shah¹, Yang Ding¹, and Joshua B. Plotkin^{1,*}

¹Department of Biology, University of Pennsylvania, Philadelphia, PA 19104

^{*}Corresponding author: jplotkin@sas.upenn.edu

An important question in molecular evolution is whether an amino acid that occurs at a given position makes an independent contribution to fitness, or whether its effect depends on the state of other loci in the organism's genome, a phenomenon known as epistasis¹⁻⁵. In a recent letter to Nature, Breen *et al.*⁶ argued that epistasis must be "pervasive throughout protein evolution" because the observed ratio between the per-site rates of non-synonymous and synonymous substitutions (dN/dS)⁷ is much lower than would be expected in the absence of epistasis. However, when calculating the expected dN/dS ratio in the absence of epistasis, Breen *et al.* assumed that all amino acids observed in a protein alignment at any particular position have equal fitness. Here, we relax this unrealistic assumption and show that any dN/dS value can in principle be achieved at a site, without epistasis. Furthermore, for all nuclear and chloroplast genes in the Breen *et al.* dataset, we show that the observed dN/dS values and the observed patterns of amino acid diversity at each site are jointly consistent with a non-epistatic model of protein evolution.

For a variety of proteins under long-term purifying selection, Breen *et al*. constructed alignments and recorded the amino acids observed at each position; these observed amino acids were deemed to be "acceptable" with respect to natural selection. Breen *et al*. then assumed that substitutions occur at neutral rates among these acceptable amino acids, and used these rates to calculate, for each protein, an expected value for dN/dS in the absence of epistasis. Because their empirical observations of dN/dS were much lower than these expected values, Breen *et al*. concluded that epistasis must be extremely prevalent.

The flaw in this reasoning is that Breen *et al.* considered only a single class of fitness assignments, such that all amino acids observed at a site were assumed equally fit. However, a more realistic alternative is simply that some amino acids observed at a site are more fit than others^{8,9}.

As a proof of principle, we considered a non-epistatic model in which, among the acceptable amino acids at a given site, one of these is preferable to the rest. We performed the following experiment: in a hypothetical protein of length 300 aa, for each position we randomly designated 8 amino acids as acceptable (the average number of acceptable amino acids reported by Breen *et al.*) but gave one of these a selective advantage over the rest. We then calculated the equilibrium dN/dS¹⁰ for this protein as a function of the selective advantage of the preferred amino acid at each site, 2Ns (Figure 1). While dN/dS is high for the case 2Ns=0, corresponding to the Breen *et al.* assumption, dN/dS is much lower for larger 2Ns. Thus, a large range of dN/dS values are consistent with non-epistatic models of protein evolution.




Figure 1: dN/dS and the strength of selection. dN/dS as a function of 2Ns for a hypothetical protein of length 300, in which 8 acceptable amino acids are chosen at random for each position and one of these amino acids at random is assigned a selective advantage of size 2Ns. The remaining 12 amino acids are lethal. The Breen *et al.* expectation for dN/dS in the absence of epistasis corresponds to 2Ns=0.

Although site-independent fitness assignments can in principle produce very low dN/dS values, are the Breen *et al.* data in fact consistent with a non-epistatic model? To answer this question, for each protein in the Breen *et al.* dataset, we assigned amino acids fitnesses at each site so that their equilibrium frequencies precisely matched the frequencies observed by Breen *et al.* for that site¹¹. For each protein, we then simulated the divergence between pairs of sequences and computed dN/dS for each pair. For the mitochondrial genes in the Breen *et al.* dataset, the average simulated dN/dS values, while on the whole substantially lower than the Breen *et al.* expectations, were still greater than the empirically observed values. However, for all nuclear and chloroplast genes in the Breen *et al.* dataset, the average dN/dS values under our non-epistatic model are comparable to or even lower than the empirical dN/dS values reported by Breen *et al.* Thus, the low dN/dS in these genes need not be attributed to epistasis (Table 1), but rather can be explained by the more parsimonious assumption that some amino acids are more fit than others, at each site.

Gene	Breen et al. Expected dN/dS	Our Average Simulated dN/dS	Breen et al. Empirical dN/dS
Mitochondrial			
ATP6	0.44	0.215	0.056
ATP8	0.56	0.624	0.224
COX1	0.28	0.078	0.015
COX2	0.43	0.140	0.025
COX3	0.32	0.144	0.036
CYTB	0.51	0.117	0.039
ND1	0.39	0.208	0.040
ND2	0.51	0.262	0.067
ND3	0.49	0.242	0.069
ND4	0.42	0.239	0.045
ND4L	0.49	0.369	0.076
ND5	0.32	0.211	0.057
ND6	0.42	0.397	0.073
Nuclear			
EEF1A1	0.11	0.031	0.020
H3.2	0.14	0.014	0.037
Chloroplast			_
rbcL	0.40	0.024	0.072

Table 1: Observed and expected dN/dS. Comparison of expected dN/dS values with the empirical values for each gene in the Breen *et al.* dataset. The Breen *et al.* expected dN/dS is based on the assumption that all amino acids observed at a given site are neutral relative to each other. Our average simulated dN/dS is based on the assumption that the amino acids observed at a given site have different fitnesses; these fitnesses are chosen so that the equilibrium amino acid frequencies at each site match the empirical frequencies at that site in the Breen *et al.* dataset.

It is important to note that the effects of natural selection and phylogeny are confounded in the observed amino acid frequencies. For small datasets, and with a known phylogeny, methods exist to distinguish these effects ^{12,13}. However, in the absence of a phylogeny, our fitness estimates are maximum-likelihood, and in equilibrium they are guaranteed to reproduce both the site-specific amino acid frequencies and the mean pairwise divergences between the empirical amino acid sequences. Our method makes the standard assumption (e.g. ref. 14, as used by Breen *et al.*) that molecular evolution can be modeled as an equilibrium Markov chain running on the branches of a phylogeny. Relaxing this assumption would make it even more difficult to reject the non-epistatic null hypothesis, because it would then be necessary to rule out the even larger class of non-equilibrium site-independent models before concluding that epistasis is present.

In summary, the data of Breen *et al*. do not support the conclusion that epistasis is the primary factor in molecular evolution. Breen *et al*. provide no direct evidence of epistasis whatsoever, but rather their conclusions were based on the rejection of an overly simplistic model of site-independent evolution. At best, the data support the conclusion that epistasis may be an important factor in the evolution of mitochondrial genes, but to a lesser extent than originally claimed.

Methods:

We assume that each codon evolves according to an independent Markov chain whose rate matrix is determined by the scaled selection coefficient assigned to each amino acid 15 . The equilibrium frequency of each amino acid is then proportional to $u_i e^{2Ns(i)}$ (ref. 15) where u_i is the number of codons that code for amino acid i and 2Ns(i) is its scaled selection coefficient. Simulations were conducted by running an independent Markov chain for each codon represented in an at least half the sequences of the Breen *et al*. alignment, until the divergence between each pair of sequences was dS=0.25, which is within the range of dS=0.05 to 0.5 used by Breen *et al*. dN/dS was then estimated using PAML again following the procedure described by Breen *et al*. The ancestral sequence at each site was drawn from the equilibrium distribution. All computer codes can be made available upon request.

References:

- Kondrashov A.S., Sunyaev, S. & Kondrashov F.A. Dobzhansky-Muller incompatibilities in protein evolution. Proc. Natl. Acad. Sci. USA 99, 14878-14883 (2002).
- 2. DePristo, M.A., Weinreich, D.M., & Hartl, D.L. Missense meanderings in sequence space: a biophysical view of protein evolution. Nature Rev. Genet. 6, 678-687 (2005).
- 3. Krazhimskiy, S., Dushoff, J., Brazykin, G.A. & Plotkin, J.B. Prevalence of epistasis in the evolution of Influenza A surface proteins. PLoS Gent 7, e1001301 (2011).
- 4. Salverda, M.L.M. *et al.* Initial mutations direct alternative pathways of protein evolution. PLoS Genet 7, e1001321 (2011).
- 5. Hansen, T.F. & Wagner, G.P. Modeling genetic architecture: a multilinear theory of gene interaction. Theor. Popul. Biol. 59, 61-86 (2001).
- 6. Breen, M.S., Kemena, C., Vlasov, P.K., Notredame, C. & Kondrashov, F.A. Epistasis as the primary factor in molecular evolution. Nature 490, 535-538 (2012).
- 7. Li, W. H. Molecular Evolution (Sinauer, 1997).
- 8. da Silva, J. Site-specific amino acid frequency, fitness and the mutational landscape model of adaptation in HIV-1. Genetics 174, 1689–1694 (2006).
- 9. Fowler, D.M. *et al.* High-resolution mapping of protein sequence-function relationships. Nature Methods 7, 741–746 (2010).
- 10. Yang, Z. & Nielsen, R. Synonymous and nonsynonymous rate variation in nuclear genes of mammals. J. Mol. Evol. 46, 409-418 (1998).
- 11. Choi, S.C., Redelings, B.D. & Thorne, J.L. Basing population genetic inferences and models of molecular evolution upon desired stationary distributions of DNA or protein sequences. Phil. Trans. R. Soc. B 363, 3931-3939 (2008).
- 12. Rodrigue, N., Phillippe, H. & Lartillot N. Mutation-selection models of coding sequence evolution with site-heterogeneous amino acid fitness profiles. Proc. Natl. Acad. Sci. USA 107, 4629-4634 (2010).
- 13. Tamuri, A. U., dos Reis, M. & Goldstein R. A. Estimating the distribution of selection coefficients from phylogenetic data using sitewise mutation-selection models. Genetics 190, 1101-1115 (2012).
- 14. Yang, Z. PAML 4: a program package for phylogenetic analysis by maximum likelihood. Mol. Biol. Evol. 24, 1586-1591 (2007).
- 15. Halpern, A.L. & Bruno, W.J. Evolutionary distances for protein-coding sequences: modeling site-specific residue frequencies. Mol. Biol. Evol. 15, 910-917 (1998).