COMPLETE MONOTONICITY OF FUNCTIONS INVOLVING THE q-TRIGAMMA AND q-TETRAGAMMA FUNCTIONS

FENG QI

ABSTRACT. Let $\psi_q(x)$, $\psi_q'(x)$, and $\psi_q''(x)$ for q>0 stand respectively for the q-digamma, q-trigamma, and q-tetragamma functions. In the paper, the author proves along two different approaches that the functions $[\psi_q'(x)]^2 + \psi_q''(x)$ for q>1 and $[\psi_q'(x)-\ln q]^2 + \psi_q''(x)$ for 0< q<1 are completely monotonic on $(0,\infty)$. Applying these results, the author derives monotonic properties of four functions involving the q-digamma function $\psi_q(x)$ and two double inequalities for bounding the q-digamma function $\psi_q(x)$.

1. Introduction

Recall from [14, Chapter XIII] that a function f is said to be completely monotonic on an interval I if f has derivatives of all orders on I and

$$0 \le (-1)^n f^{(n)}(x) < \infty \tag{1.1}$$

for $x \in I$ and $n \ge 0$. For more information about properties and applications of completely monotonic functions, please refer to [14, Chapter XIII] and closely related references therein.

It is common knowledge that the logarithmic derivative of the classical Euler's gamma function $\Gamma(x)$, denoted by $\psi(x) = \frac{\Gamma'(x)}{\Gamma(x)}$, the derivatives $\psi'(x)$ and $\psi''(x)$ are respectively called the digamma, trigamma and tetragamma functions. As a whole, the derivatives $\psi^{(i)}(x)$ for $i \in \mathbb{N}$ are respectively called polygamma functions.

Recall from [2, pp. 493–496] and [7, Section 1.10] that the q-gamma function, the q-analogue of the gamma function $\Gamma(x)$, is defined for x > 0 by

$$\Gamma_{q}(x) = \begin{cases}
(1-q)^{1-x} \prod_{i=0}^{\infty} \frac{1-q^{i+1}}{1-q^{i+x}}, & 0 < q < 1 \\
(q-1)^{1-x} q^{\binom{x}{2}} \prod_{i=0}^{\infty} \frac{1-q^{-(i+1)}}{1-q^{-(i+x)}}, & q > 1 \\
\Gamma(x), & q = 1
\end{cases}$$
(1.2)

and satisfies

$$\Gamma_q(x) = q^{\binom{x-1}{2}} \Gamma_{1/q}(x).$$
 (1.3)

 $^{2010\} Mathematics\ Subject\ Classification.$ Primary 33D05; Secondary 26A12, 26A48, 26D07, 33B15.

 $Key\ words\ and\ phrases.$ completely monotonic function; monotonicity; inequality; q-digamma function; q-trigamma function; q-tetragamma function.

This paper was typeset using $A_{\mathcal{M}}S$ -IAT_EX.

The q-digamma function $\psi_q(x)$, the q-analogue of the digamma function $\psi(x)$, is defined for q>0 and x>0 by

$$\psi_q(x) = \begin{cases} \frac{\Gamma'_q(x)}{\Gamma_q(x)}, & q \neq 1, \\ \psi(x), & q = 1. \end{cases}$$

$$(1.4)$$

The functions $\psi_q^{(k)}(x)$ for $k \in \mathbb{N}$, the q-analogues of polygamma functions $\psi_q^{(k)}(x)$, are called q-polygamma functions. In particular, the functions $\psi_q'(x)$ and $\psi_q''(x)$ are respectively called the q-trigamma and q-tetragamma functions. From (1.2), we obtain that

(1) when 0 < q < 1 and $x \in (0, \infty)$,

$$\psi_q(x) = -\ln(1-q) + (\ln q) \sum_{k=0}^{\infty} \frac{q^{k+x}}{1 - q^{k+x}}$$

$$= -\ln(1-q) + (\ln q) \sum_{k=1}^{\infty} \frac{q^{kx}}{1 - q^k},$$
(1.5)

(2) when q > 1 and $x \in (0, \infty)$,

$$\psi_q(x) = -\ln(q-1) + (\ln q) \left[x - \frac{1}{2} - \sum_{i=1}^{\infty} \frac{q^{-ix}}{1 - q^{-i}} \right]. \tag{1.6}$$

The inequality

$$\psi''(x) + [\psi'(x)]^2 > 0, \quad x > 0 \tag{1.7}$$

was deduced or recovered in [3, Theorem 2.1] and [5, Lemma 1.1], employed in [1, Theorem 4.3], [3, Theorem 2.1], and [5, Theorem 2.1], and generalized in [4, Lemma 1.2 and Remark 1.3]. In [8, 18, 19], among other things, the inequality (1.7) was generalized to complete monotonicity of the function $\psi''(x) + [\psi'(x)]^2$. Recently, some other results in this field were published in [17, 20] and closely related references therein.

In [1, pp. 80–81, Lemma 4.6], the inequality (1.7) was generalized as

$$[\psi_q'(x)]^2 + \psi_q''(x) > 0 \tag{1.8}$$

for q>1 and x>0. Applying this inequality, the following conclusion was established in [1, p. 82, Theorem 4.8]: for q>1 and $0< a< b\leq \infty$, the double inequality

$$\exp\left\{\alpha\left[e^{\psi_q(x)}\left(\psi_q(x)-1\right)+1\right]\right\} \le \frac{\Gamma_q(x)}{\Gamma_q(x_0)} \le \exp\left\{\beta\left[e^{\psi_q(x)}\left(\psi_q(x)-1\right)+1\right]\right\}$$

is valid on (a, b), with the best possible constant factors

$$\alpha = \begin{cases} Q_q(b), & \text{if } b < \infty \\ 0, & \text{if } b = \infty \end{cases}$$

and $\beta = Q_q(a)$, where $x_0 = x_0(q)$ is the unique positive zero of the function ψ_q for $0 < q \neq 1$, and

$$Q_q(x) = \begin{cases} \frac{\ln \Gamma_q(x) - \ln \Gamma_q(x_0)}{[\psi_q(x) - 1]e^{\psi_q(x)} + 1}, & x \neq x_0, \\ \frac{1}{\psi_q'(x_0)}, & x = x_0. \end{cases}$$

For more information on the subject of q-gamma and q-polygamma functions, please refer to [2, 9, 12] and closely related references therein.

The aim of this paper is to extend and generalize along two different approaches the inequality (1.8) to the following complete monotonicity.

Theorem 1.1. The functions

$$\mathcal{F}_q(x) = [\psi_q'(x) - \ln q]^2 + \psi_q''(x), \quad 0 < q < 1$$
(1.9)

and

$$\mathfrak{F}_q(x) = \left[\psi_q'(x)\right]^2 + \psi_q''(x), \quad q > 1$$
 (1.10)

are completely monotonic on $(0, \infty)$.

As consequences of Theorem 1.1, the following monotonicity and inequalities may be derived.

Corollary 1.1. The functions

$$\phi_q(x) = \psi_q(x) + \ln\left[\exp\frac{(\ln q)q^x}{q^x - 1} - 1\right], \quad q > 1,$$
(1.11)

$$\varphi_q(x) = \psi_q(x) + \ln \left(\exp \frac{\ln q}{q^x - 1} - 1 \right), \quad q > 1, \tag{1.12}$$

$$\Phi_q(x) = \psi_q(x) - (\ln q)x + \ln \left[\exp \frac{(\ln q)q^x}{q^x - 1} - 1 \right], \quad 0 < q < 1,$$
 (1.13)

$$\Theta_q(x) = \psi_q(x) - (\ln q)x + \ln\left(\exp\frac{\ln q}{q^x - 1} - 1\right), \quad 0 < q < 1$$
(1.14)

are increasing on $(0, \infty)$. Consequently, the double inequalities

$$\psi_{q}(1) + (\ln q)x - \ln\left[\exp\frac{(\ln q)q^{x}}{q^{x} - 1} - 1\right] < \psi_{q}(x)$$

$$< \ln\frac{\ln q}{q - 1} + (\ln q)x - \ln\left[\exp\frac{(\ln q)q^{x}}{q^{x} - 1} - 1\right] \quad (1.15)$$

for 0 < q < 1 and

$$\psi_{q}(1) - \ln q - \ln \left(\exp \frac{\ln q}{q^{x} - 1} - 1 \right) < \psi_{q}(x)$$

$$< \ln \frac{\ln q}{q - 1} - \frac{\ln q}{2} - \ln \left(\exp \frac{\ln q}{q^{x} - 1} - 1 \right) \quad (1.16)$$

for q > 1 hold on $(0, \infty)$, where the constants $\psi_q(1)$ and $\ln\left(\frac{\ln q}{q-1}\right)$ in (1.15) and the scalars $\psi_q(1) - \ln q$ and $\ln\frac{\ln q}{q-1} - \frac{\ln q}{2}$ in (1.16) are the best possible.

2. First proof of Theorem 1.1

For q > 1, using the expression (1.6) gives

$$\frac{\left[\psi_q'(x)\right]^2 + \psi_q''(x)}{(\ln q)^2} = 1 + \sum_{i=2}^{\infty} \sum_{j=1}^{i-1} c_j(q, x) c_{i-j}(q, x) - \sum_{j=1}^{\infty} (i-2)c_i(q, x),$$
(2.1)

where

$$c_i(q, x) = \frac{i \ln q}{1 - q^{-i}} q^{-ix} > 0.$$

In order to prove complete monotonicity of the function (1.10), by definition, it suffices to show

$$(-1)^k \frac{\mathrm{d}^k}{\mathrm{d}x^k} \left\{ \frac{\left[\psi_q'(x)\right]^2 + \psi_q''(x)}{(\ln q)^2} \right\} \ge 0, \quad k \ge 0.$$
 (2.2)

For k=0, the inequality (2.2) is equivalent to (1.8). For $k\geq 1$, it is equivalent to

$$(-1)^k \sum_{i=2}^{\infty} \sum_{j=1}^{i-1} \frac{\partial^k [c_j(q, x)c_{i-j}(q, x)]}{\partial x^k} \ge (-1)^k \sum_{i=1}^{\infty} (i-2) \frac{\partial^k [c_i(q, x)]}{\partial x^k}.$$
 (2.3)

In order to prove (2.3), it is sufficient to verify

$$(-1)^k \sum_{i=1}^{i-1} \frac{\partial^k [c_j(q,x)c_{i-j}(q,x)]}{\partial x^k} \ge (-1)^k (i-2) \frac{\partial^k [c_i(q,x)]}{\partial x^k}, \quad i \ge 3,$$

that is,

$$(-1)^{k} \sum_{j=1}^{i-1} \frac{(-1)^{k+1} i^{k} j(i-j) (\ln q)^{k+2} q^{i(1-x)+j}}{(q^{j}-1) (q^{j}-q^{i})}$$

$$\geq (-1)^{k} (i-2) \frac{(-1)^{k+1} i^{k+1} (\ln q)^{k+1} q^{i(1-x)}}{1-q^{i}}, \quad i \geq 3,$$

which can be further simplified as

$$\sum_{i=1}^{i-1} \frac{j(i-j)}{(q^j-1)(1-q^{i-j})} \le \frac{(i-2)i}{(\ln q)(1-q^i)}, \quad i \ge 3.$$
 (2.4)

Since

$$\sum_{i=1}^{i-1} \frac{j(i-j)(q^{i-j}-q^j)}{(q^j-1)(q^{i-j}-1)} = 0,$$

we have

$$\sum_{j=1}^{i-1} \frac{j(i-j)(q^i-1)}{(q^j-1)(q^{i-j}-1)} = \sum_{j=1}^{i-1} \frac{j(i-j)(q^j+1)}{q^j-1}.$$
 (2.5)

The inequality

$$e^{\tau} + 1 - \frac{2}{\tau}(e^{\tau} - 1) = \sum_{k=2}^{\infty} \frac{k-1}{(k+1)!} \tau^k > 0, \quad \tau = j \ln q > 0$$

implies that

$$\frac{j(q^j+1)}{q^j-1} > \frac{2}{\ln q}.$$

This leads to

$$\sum_{j=1}^{i-1} \frac{j(i-j)(q^j+1)}{q^j-1} > \frac{2}{\ln q} \sum_{j=1}^{i-1} (i-j) = \frac{i(i-1)}{\ln q} > \frac{i(i-2)}{\ln q}.$$

Substituting this inequality into (2.5) yields

$$\sum_{i=1}^{i-1} \frac{j(i-j)(q^i-1)}{(q^j-1)(q^{i-j}-1)} > \frac{i(i-2)}{\ln q}$$
 (2.6)

which is equivalent to the strict inequality in (2.4). Thus, the inequality (2.3) is proved. Hence, the function $\mathfrak{F}_q(x)$ is completely monotonic on $(0,\infty)$.

For 0 < q < 1, taking the logarithm and differentiating on both sides of (1.3) reveal

$$\psi_q(x) = (\ln q) \left(x - \frac{3}{2} \right) + \psi_{1/q}(x),$$
(2.7)

$$\psi_{q}'(x) = \ln q + \psi_{1/q}'(x), \tag{2.8}$$

$$\psi_q^{(k)}(x) = \psi_{1/q}^{(k)}(x), \quad k \ge 2. \tag{2.9}$$

Therefore, when 0 < q < 1, we have

$$\mathfrak{F}_{1/q}(x) = \left[\psi'_{1/q}(x)\right]^2 + \psi''_{1/q}(x) = \left[\psi'_{q}(x) - \ln q\right]^2 + \psi''_{q}(x) = \mathcal{F}_{q}(x) \tag{2.10}$$

on $(0, \infty)$. As a result, the function $\mathcal{F}_q(x)$ is completely monotonic on $(0, \infty)$. The first proof of Theorem 1.1 is complete.

3. Second proof of Theorem 1.1

The formula (1.11) in the paper [11] and its corrected preprint [12] reads that

$$\psi_q(x) = -\ln(1-q) - \int_0^\infty \frac{e^{-xt}}{1 - e^{-t}} d\gamma_q(t)$$
 (3.1)

for 0 < q < 1 and x > 0, where

$$\gamma_q(t) = \begin{cases} -\ln q \sum_{k=1}^{\infty} \delta(t + k \ln q), & 0 < q < 1 \\ t, & q = 1 \end{cases}$$
 (3.2)

and $\delta(t)$ represents the Dirac delta function, that is, $d\gamma_q(t)$ is a discrete measure with positive masses $|\ln q|$ at the positive points $k|\ln q|$ for $k\in\mathbb{N}$. Differentiating on both sides of the equation (3.1) acquires

$$\psi_q'(x) = \int_0^\infty \frac{te^{-xt}}{1 - e^{-t}} \,\mathrm{d}\,\gamma_q(t), \quad 0 < q < 1.$$
 (3.3)

By the definition of $\gamma_q(t)$, we obtain that

$$\int_0^\infty e^{-xt} \,\mathrm{d}\,\gamma_q(t) = -\frac{q^x \ln q}{1 - q^x} \tag{3.4}$$

and

$$\int_0^\infty t e^{-xt} \, \mathrm{d} \, \gamma_q(t) = \frac{q^x (\ln q)^2}{(1 - q^x)^2} \tag{3.5}$$

for 0 < q < 1 and x > 0. It was presented in [9, p. 1245, Theorem 4.4, (4.15)], [15, Lemma 2.3 and Remark 2.1], [16, Theorem 7.2, (7.5)], and [21, p. 152, Theorem 4.22, (4.20)] that, when 0 < q < 1, the identity

$$\psi_q^{(k-1)}(x) - \psi_q^{(k-1)}(x+1) = (\ln q) \frac{\mathrm{d}^{k-1}}{\mathrm{d} x^{k-1}} \left(\frac{q^x}{1 - q^x} \right)$$
 (3.6)

is valid for $x \in (0, \infty)$ and $k \in \mathbb{N}$. Accordingly, it follows that

$$\mathcal{F}_{q}(x) - \mathcal{F}_{q}(x+1) = \left[\psi_{q}''(x) - \psi_{q}''(x+1) \right] + \left[\psi_{q}'(x) - \psi_{q}'(x+1) \right] \left[\psi_{q}'(x) + \psi_{q}'(x+1) - 2 \ln q \right]$$

$$\begin{split} &= (\ln q) \frac{\mathrm{d}^2}{\mathrm{d}\,x^2} \bigg(\frac{q^x}{1-q^x} \bigg) + (\ln q) \frac{\mathrm{d}}{\mathrm{d}\,x} \bigg(\frac{q^x}{1-q^x} \bigg) \bigg[2\psi_q'(x) - (\ln q) \frac{\mathrm{d}}{\mathrm{d}\,x} \bigg(\frac{q^x}{1-q^x} \bigg) - 2\ln q \bigg] \\ &= \frac{q^x (\ln q)^2}{(q^x-1)^2} \bigg[2\psi_q'(x) - \frac{q^x (\ln q)^2}{(q^x-1)^2} - 2\ln q \bigg] - \frac{q^x (q^x+1) (\ln q)^3}{(q^x-1)^3} \\ &= \frac{q^x (\ln q)^2}{(q^x-1)^2} \bigg[2\psi_q'(x) - \frac{q^x (\ln q)^2}{(q^x-1)^2} - 2\ln q - \frac{(q^x+1) (\ln q)}{q^x-1} \bigg] \\ &= \frac{2q^x (\ln q)^2}{(q^x-1)^2} \bigg[\int_0^\infty \frac{te^{-xt}}{1-e^{-t}} \, \mathrm{d}\,\gamma_q(t) - \frac{1}{2} \int_0^\infty te^{-xt} \, \mathrm{d}\,\gamma_q(t) - \frac{1}{2} \ln q - \int_0^\infty e^{-xt} \, \mathrm{d}\,\gamma_q(t) \bigg] \\ &= \frac{2q^x (\ln q)^2}{(q^x-1)^2} \bigg[\int_0^\infty \bigg(\frac{t}{1-e^{-t}} - \frac{1}{2}t - 1 \bigg) e^{-xt} \, \mathrm{d}\,\gamma_q(t) - \frac{1}{2} \ln q \bigg] \\ &= 2 \bigg[\int_0^\infty \bigg(\frac{t}{1-e^{-t}} - \frac{1}{2}t - 1 \bigg) e^{-xt} \, \mathrm{d}\,\gamma_q(t) - \frac{1}{2} \ln q \bigg] \int_0^\infty te^{-xt} \, \mathrm{d}\,\gamma_q(t). \end{split}$$

By [22, p. 161, Theorem 12b] which reads that a function f(x) is completely monotonic on $(0, \infty)$ if and only if it can be represented in the form

$$f(x) = \int_0^\infty e^{-xt} \,\mathrm{d}\,\alpha(t),\tag{3.7}$$

where $\alpha(t)$ is non-decreasing and the integral converges for $0 < x < \infty$, and by the fact that the sum and product of finitely many completely monotonic functions are all still completely monotonic, we obtain that the difference $\mathcal{F}_q(x) - \mathcal{F}_q(x+1)$ is completely monotonic on $(0,\infty)$, that is,

$$0 \le (-1)^{k-1} [\mathcal{F}_q(x) - \mathcal{F}_q(x+1)]^{(k-1)}$$

= $(-1)^{k-1} \mathcal{F}_q^{(k-1)}(x) - (-1)^{k-1} \mathcal{F}_q^{(k-1)}(x+1)$

on $(0, \infty)$ for $k \in \mathbb{N}$. Consequently, by induction,

$$(-1)^{k-1}\mathcal{F}_q^{(k-1)}(x) \ge (-1)^{k-1}\mathcal{F}_q^{(k-1)}(x+1) \ge (-1)^{k-1}\mathcal{F}_q^{(k-1)}(x+2)$$

$$\ge (-1)^{k-1}\mathcal{F}_q^{(k-1)}(x+3) \ge \dots \ge (-1)^{k-1}\mathcal{F}_q^{(k-1)}(x+\ell) \to 0$$

as $\ell \to \infty$. This means that the function $\mathcal{F}_q(x)$ is completely monotonic on $(0, \infty)$. By (2.10), it is immediately deduced that the function $\mathfrak{F}_q(x)$ is completely monotonic on $(0, \infty)$. The second proof of Theorem 1.1 is complete.

4. Proof of Corollary 1.1

4.1. Substituting (2.8) and (2.9) into (3.6), we easily procure that

$$\psi_{1/q}^{(k-1)}(x) - \psi_{1/q}^{(k-1)}(x+1) = (\ln q) \frac{\mathrm{d}^{k-1}}{\mathrm{d} \, x^{k-1}} \left(\frac{q^x}{1 - q^x} \right)$$

for $0 < q < 1, k \ge 2$, and x > 0. Replacing q by $\frac{1}{q}$ in the above identity leads to

$$\psi_q^{(k-1)}(x+1) - \psi_q^{(k-1)}(x) = (\ln q) \frac{\mathrm{d}^{k-1}}{\mathrm{d} x^{k-1}} \left(\frac{1}{q^x - 1}\right) \tag{4.1}$$

for q > 1 and $k \ge 2$. Similarly, combining (2.7) with (3.6), we obtain

$$\psi_q(x+1) - \psi_q(x) = \frac{(\ln q)q^x}{q^x - 1} \tag{4.2}$$

for q > 1 and x > 0.

When $0 < q \neq 1$, we have

$$\exp[\phi_q(x)] = \exp[\psi_q(x)] \left[\exp\frac{(\ln q)q^x}{q^x - 1} - 1 \right]$$
$$= \exp\left[\psi_q(x) + \frac{(\ln q)q^x}{q^x - 1}\right] - \exp[\psi_q(x)]$$
$$= \exp[\psi_q(x+1)] - \exp[\psi_q(x)],$$

in which the last equality is obtained by (3.6) and (4.2). Further differentiating gives

$$\frac{\mathrm{d} \exp[\phi_q(x)]}{\mathrm{d} x} = \psi_q'(x+1) \exp[\psi_q(x+1)] - \psi_q'(x) \exp[\psi_q(x)],$$

$$\frac{\mathrm{d} \{\psi_q'(x) \exp[\psi_q(x)]\}}{\mathrm{d} x} = \{\psi_q''(x) + [\psi_q'(x)]^2\} \exp[\psi_q(x)].$$

The complete monotonicity of the function $\mathfrak{F}_q(x)$, obtained in Theorem 1.1, implies the positivity of $\mathfrak{F}_q(x)$ on $(0,\infty)$, that is, the inequality (1.8). Hence, the function $\psi_q'(x) \exp[\psi_q(x)]$ is increasing, and so $\frac{\mathrm{d}\exp[\phi_q(x)]}{\mathrm{d}x} > 0$ for q > 1 on $(0,\infty)$. As a result, the function $\exp[\phi_q(x)]$, and thus $\phi_q(x)$, is increasing for q > 1 on $(0,\infty)$.

For $0 < q \neq 1$, we also have

$$\exp[\Phi_q(x)] = \exp[\psi_q(x) - (\ln q)x] \left[\exp\frac{(\ln q)q^x}{q^x - 1} - 1 \right]$$

$$= \exp\left[\psi_q(x) + \frac{(\ln q)q^x}{q^x - 1} - (\ln q)x\right] - \exp[\psi_q(x) - (\ln q)x]$$

$$= \exp[\psi_q(x + 1) - (\ln q)x] - \exp[\psi_q(x) - (\ln q)x],$$

in which the last equality is deduced by using (3.6) and (4.2). A direct computation gives

$$\frac{\mathrm{d} \exp[\Phi_q(x)]}{\mathrm{d} x} = [\psi_q'(x+1) - \ln q] \exp[\psi_q(x+1) - (\ln q)x] - [\psi_q'(x) - \ln q] \exp[\psi_q(x) - (\ln q)x]$$

and

$$\frac{\mathrm{d}\{[\psi_q'(x) - \ln q] \exp[\psi_q(x) - (\ln q)x]\}}{\mathrm{d} x}$$

$$= \{\psi_q''(x) + [\psi_q'(x) - \ln q]^2\} \exp[\psi_q(x) - (\ln q)x].$$

The complete monotonicity of the function $\mathcal{F}_q(x)$, obtained in Theorem 1.1, implies the positivity of $\mathcal{F}_q(x)$ on $(0,\infty)$. Hence, the function

$$\{ [\psi_q'(x) - \ln q] \exp[\psi_q(x) - (\ln q)x] \}$$

is increasing, and so $\frac{\mathrm{d} \exp[\Phi_q(x)]}{\mathrm{d} x} > 0$ for 0 < q < 1 on $(0, \infty)$. As a result, the function $\exp[\Phi_q(x)]$, and thus $\Phi_q(x)$, is increasing for 0 < q < 1 on $(0, \infty)$.

For 0 < q < 1, utilizing (2.7) yields

$$\begin{split} \Phi_q(x) &= (\ln q) \left(x - \frac{3}{2} \right) + \psi_{1/q}(x) - (\ln q) x + \ln \left[\exp \frac{(\ln q) q^x}{q^x - 1} - 1 \right] \\ &= \psi_{1/q}(x) + \ln \left[\exp \frac{(\ln q) q^x}{q^x - 1} - 1 \right] - \frac{3}{2} \ln q \end{split}$$

$$= \psi_{1/q}(x) + \ln \left[\exp \frac{\ln(1/q)}{(1/q)^x - 1} - 1 \right] + \frac{3}{2} \ln \frac{1}{q}.$$

Replacing $\frac{1}{q}$ by q in the above equation leads to increasing monotonicity of the function $\varphi_q(x)$ for q > 1 on $(0, \infty)$.

For q > 1, employing (2.7) gives

$$\begin{split} \phi_q(x) &= \psi_{1/q}(x) - \left(\ln\frac{1}{q}\right) \left(x - \frac{3}{2}\right) + \ln\left[\exp\frac{(\ln q)q^x}{q^x - 1} - 1\right] \\ &= \psi_{1/q}(x) - \left(\ln\frac{1}{q}\right)x + \ln\left[\exp\frac{\ln(1/q)}{(1/q)^x - 1} - 1\right] + \frac{3}{2}\ln\frac{1}{q}. \end{split}$$

Substituting $\frac{1}{q}$ by q in the above equation reduces to increasing monotonicity of the function $\Theta_q(x)$ for 0 < q < 1 on $(0, \infty)$.

4.2. From (1.5), it is easy to see that

$$\lim_{x \to \infty} \psi_q(x) = -\ln(1 - q), \quad q \in (0, 1). \tag{4.3}$$

In addition, for $q \in (0,1)$, we have

$$\lim_{x \to \infty} \left\{ \ln \left[\exp \frac{(\ln q)q^x}{q^x - 1} - 1 \right] - (\ln q)x \right\} = \lim_{x \to \infty} \ln \left[\frac{\exp \frac{(\ln q)q^x}{q^x - 1} - 1}{q^x} \right] = \ln \left(\ln \frac{1}{q} \right).$$

Combining them with increasing monotonicity of $\Phi_q(x)$, we acquire

$$\psi_q(x) - (\ln q)x + \ln \left[\exp \frac{(\ln q)q^x}{q^x - 1} - 1\right] < \ln \left(\ln \frac{1}{q}\right) - \ln(1 - q) = \ln \left(\frac{\ln q}{q - 1}\right)$$

for $q \in (0,1)$ on $(0,\infty)$, which is equivalent to the right-hand side of (1.15). From (3.6) for k=1 and the limit

$$\lim_{x \to 0^+} \psi_q(x) = -\infty \tag{4.4}$$

for 0 < q < 1, it follows that

$$\psi_{q}(x) + \ln\left[\exp\frac{(\ln q)q^{x}}{q^{x} - 1} - 1\right] = \ln\exp[\psi_{q}(x)] + \ln\left[\exp\frac{(\ln q)q^{x}}{q^{x} - 1} - 1\right]$$

$$= \ln\left\{\left[\exp\frac{(\ln q)q^{x}}{q^{x} - 1} - 1\right] \exp[\psi_{q}(x)]\right\}$$

$$= \ln\left\{\exp\left[\frac{(\ln q)q^{x}}{q^{x} - 1} + \psi_{q}(x)\right] - \exp[\psi_{q}(x)]\right\}$$

$$= \ln\left\{\exp\left[\psi_{q}(x + 1)\right] - \exp[\psi_{q}(x)]\right\} \rightarrow \psi_{q}(1)$$

as $x \to 0^+$, so the limit $\lim_{x\to 0^+} \Phi_q(x) = \psi_q(1)$ for 0 < q < 1. Combining this with increasing monotonicity of $\Phi_q(x)$, we acquire $\Phi_q(x) > \psi_q(1)$ for 0 < q < 1 on $(0, \infty)$, that is,

$$\psi_q(x) - (\ln q)x + \ln \left[\exp \frac{(\ln q)q^x}{q^x - 1} - 1 \right] > \psi_q(1), \quad 0 < q < 1, \quad x \in (0, \infty).$$

The required inequality in the left-hand side of (1.15) is proved.

For 0 < q < 1, by using (3.6) for k = 1, we have

$$\Theta_{q}(x) = \psi_{q}(x) + \frac{(\ln q)q^{x}}{q^{x} - 1} - \frac{(\ln q)q^{x}}{q^{x} - 1} - (\ln q)x + \ln\left(\exp\frac{\ln q}{q^{x} - 1} - 1\right)
= \psi_{q}(x+1) + \ln\frac{\exp[\ln q/(q^{x} - 1)] - 1}{q^{x}\exp[(\ln q)q^{x}/(q^{x} - 1)]}
\rightarrow \psi_{q}(1) - \ln q$$
(4.6)

as $x \to 0^+$. Since $\Theta_q(x)$ is increasing on $(0, \infty)$, we obtain that

$$\psi_q(x) - (\ln q)x + \ln\left(\exp\frac{\ln q}{q^x - 1} - 1\right) > \psi_q(1) - \ln q$$

for 0 < q < 1 on $(0, \infty)$, which is included in the left-hand side of (1.15). For q > 1, by a similar argument as in (4.6), we have

$$\varphi_q(x) = \psi_q(x+1) + \ln \frac{\exp[\ln q/(q^x-1)] - 1}{\exp[(\ln q)q^x/(q^x-1)]} \to \psi_q(1) - \ln q$$

as $x \to 0^+$. Furthermore, from (2.7) and (4.3), it follows that

$$\varphi_q(x) = \psi_{1/q}(x) + (\ln q) \left(x - \frac{3}{2} \right) + \ln \left(\exp \frac{\ln q}{q^x - 1} - 1 \right)$$

$$= \psi_{1/q}(x) - \frac{3 \ln q}{2} + \ln \left[q^x \left(\exp \frac{\ln q}{q^x - 1} - 1 \right) \right]$$

$$\to -\ln \left(1 - \frac{1}{q} \right) - \frac{3 \ln q}{2} + \ln(\ln q)$$

$$= \ln \frac{\ln q}{q - 1} - \frac{\ln q}{2}$$

as $x \to \infty$. Hence, we have

$$\psi_q(1) - \ln q < \varphi_q(x) < \ln \frac{\ln q}{q - 1} - \frac{\ln q}{2}$$

for q > 1 on $(0, \infty)$. The required double inequality (1.16) is proved.

For q>1, the limits (4.4) and (4.5) are still valid, then $\lim_{x\to 0^+}\phi_q(x)=\psi_q(1)$ for q>1, and so $\phi_q(x)>\psi_q(1)$ which is contained in the left-hand side of (1.16). The proof of Corollary 1.1 is complete.

5. Remarks

Finally we remark some closely related concerns.

Remark 5.1. It is easy to see that

$$\lim_{q \to 1} \phi_q(x) = \lim_{q \to 1} \varphi_q(x) = \lim_{q \to 1} \Phi_q(x) = \lim_{q \to 1} \Theta_q(x) = \psi(x) + \ln(e^{1/x} - 1).$$
 (5.1)

It is known [10, 13] that the function (5.1) is increasing and concave on $(0, \infty)$ and that

$$a - \ln(e^{1/x} - 1) < \psi(x) < b - \ln(e^{1/x} - 1)$$
(5.2)

if and only if $a \leq -\gamma$ and $b \geq 0$, where $\gamma = 0.577...$ stands for Euler-Mascheroni's constant. The inequality (5.2) may also be derived from taking the limit $q \to 1$ in (1.15) or (1.16). We conjecture that the functions $\phi_q(x)$, $\varphi_q(x)$, $\Phi_q(x)$, and $\Theta_q(x)$ should be concave on $(0, \infty)$.

Remark 5.2. From the proof of Corollary 1.1, it may be obtained that the functions $\psi_q'(x)e^{\psi_q(x)}$ for $q\geq 1$ and $[\psi_q'(x)-\ln q]e^{\psi_q(x)-(\ln q)x}$ for $0< q\leq 1$ are increasing on $(0,\infty)$. In [5, Lemma 1.2] and [6, pp. 241–242], it was obtained that the function $\psi'(x)e^{\psi(x)}$ is increasing and less than 1 on $(0,\infty)$. Can one find bounds of the functions $\psi_q'(x)e^{\psi_q(x)}$ for $q\geq 1$ and $[\psi_q'(x)-\ln q]e^{\psi_q(x)-(\ln q)x}$ for $0< q\leq 1$?

Acknowledgements. The author thanks anonymous referees for their comments on and helpful suggestions to the original version of this paper.

References

- H. Alzer and A. Z. Grinshpan, Inequalities for the gamma and q-gamma functions, J. Approx. Theory 144 (2007), 67–83; Available online at http://dx.doi.org/10.1016/j.jat.2006.04. 008. 2
- [2] G. E. Andrews, R. Askey, and R. Roy, *Special Functions*, Encyclopedia of Mathematics and its Applications **71**, Cambridge University Press, Cambridge, 1999. 1, 3
- [3] N. Batir, An interesting double inequality for Euler's gamma function, J. Inequal. Pure Appl. Math. 5 (2004), no. 4, Art. 97; Available online at http://www.emis.de/journals/ JIPAM/article452.html. 2
- [4] N. Batir, On some properties of digamma and polygamma functions, J. Math. Anal. Appl. 328 (2007), no. 1, 452-465; Available online at http://dx.doi.org/10.1016/j.jmaa.2006. 05.065. 2
- [5] N. Batir, Some new inequalities for gamma and polygamma functions, J. Inequal. Pure Appl. Math. 6 (2005), no. 4, Art. 103; Available online at http://www.emis.de/journals/JIPAM/article577.html. 2, 10
- [6] N. Elezović, C. Giordano, and J. Pečarić, The best bounds in Gautschi's inequality, Math. Inequal. Appl. 3 (2000), 239–252; Available online at http://dx.doi.org/10.7153/mia-03-26.
- [7] G. Gasper and M. Rahman, Basic hypergeometric series, 2nd ed., Encyclopedia of Mathematics and its Applications 96, Cambridge University Press, Cambridge, 2004. 1
- [8] B.-N. Guo and F. Qi, A class of completely monotonic functions involving divided differences of the psi and tri-gamma functions and some applications, J. Korean Math. Soc. 48 (2011), no. 3, 655-667; Available online at http://dx.doi.org/10.4134/JKMS.2011.48.3.655.
- [9] B.-N. Guo and F. Qi, Properties and applications of a function involving exponential functions, Commun. Pure Appl. Anal. 8 (2009), no. 4, 1231–1249; Available online at http://dx.doi.org/10.3934/cpaa.2009.8.1231. 3, 5
- [10] B.-N. Guo and F. Qi, Some properties of the psi and polygamma functions, Hacet. J. Math. Stat. 39 (2010), no. 2, 219–231. 9
- [11] M. E. H. Ismail and M. E. Muldoon, Inequalities and monotonicity properties for gamma and q-gamma functions, in: R.V.M. Zahar (Ed.), Approximation and Computation: A Festschrift in Honour of Walter Gautschi, ISNM, Vol. 119, BirkhRauser, Basel, 1994, 309–323; Available online at http://dx.doi.org/10.1007/978-1-4684-7415-2_19.
- [12] M. E. H. Ismail and M. E. Muldoon, Inequalities and monotonicity properties for gamma and q-qamma functions, available online at http://arxiv.org/abs/1301.1749. 3, 5
- [13] W.-H. Li, F. Qi, and B.-N. Guo, On proofs for monotonicity of a function involving the psi and exponential functions, Analysis (Munich) 33 (2013), no. 1, 45–50; Available online at http://dx.doi.org/10.1524/anly.2013.1175. 9
- [14] D. S. Mitrinović, J. E. Pečarić, and A. M. Fink, Classical and New Inequalities in Analysis, Kluwer, Dordrecht, 1993. 1
- [15] F. Qi, A completely monotonic function related to the q-trigamma function, Politehn. Univ. Bucharest Sci. Bull. Ser. A Appl. Math. Phys. 76 (2014), no. 1, 107–114. 5
- [16] F. Qi, Bounds for the ratio of two gamma functions, J. Inequal. Appl. 2010 (2010), Article ID 493058, 84 pages; Available online at http://dx.doi.org/10.1155/2010/493058.
- [17] F. Qi, P. Cerone, and S. S. Dragomir, Complete monotonicity of a function involving the divided difference of psi functions, Bull. Aust. Math. Soc. 88 (2013), no. 2, 309-319; Available online at http://dx.doi.org/10.1017/S0004972712001025.

- [18] F. Qi and B.-N. Guo, Completely monotonic functions involving divided differences of the di- and tri-gamma functions and some applications, Commun. Pure Appl. Anal. 8 (2009), no. 6, 1975–1989; Available online at http://dx.doi.org/10.3934/cpaa.2009.8.1975.
- [19] F. Qi and B.-N. Guo, Necessary and sufficient conditions for functions involving the triand tetra-gamma functions to be completely monotonic, Adv. Appl. Math. 44 (2010), no. 1, 71-83; Available onine at http://dx.doi.org/10.1016/j.aam.2009.03.003. 2
- [20] F. Qi, Q.-M. Luo, and B.-N. Guo, Complete monotonicity of a function involving the divided difference of digamma functions, Sci. China Math. 56 (2013), no. 11, 2315–2325; Available online at http://dx.doi.org/10.1007/s11425-012-4562-0. 2
- [21] F. Qi and Q.-M. Luo, Bounds for the ratio of two gamma functions—From Wendel's and related inequalities to logarithmically completely monotonic functions, Banach J. Math. Anal. 6 (2012), no. 2, 132–158. 5
- [22] D. V. Widder, The Laplace Transform, Princeton University Press, Princeton, 1946. 6

Institute of Mathematics, Henan Polytechnic University, Jiaozuo City, Henan Province, $454010,\,\mathrm{China}$

 $E{-}mail~address: \verb| qifeng618@gmail.com|, qifeng618@hotmail.com|, qifeng618@qq.com| \\ URL: \verb| http://qifeng618.wordpress.com| \\$