

# Pairs of Fan-type heavy subgraphs for pancyclicity of 2-connected graphs

Bo Ning\*

Department of Applied Mathematics, School of Science,  
Northwestern Polytechnical University, Xi'an, Shaanxi 710129, P.R. China

## Abstract

A graph  $G$  on  $n$  vertices is Hamiltonian if it contains a spanning cycle, and pancyclic if it contains cycles of all lengths from 3 to  $n$ . In 1984, Fan presented a degree condition involving every pair of vertices at distance two for a 2-connected graph to be Hamiltonian. Motivated by Fan's result, we say that an induced subgraph  $H$  of  $G$  is  $f_1$ -heavy if for every pair of vertices  $u, v \in V(H)$ ,  $d_H(u, v) = 2$  implies  $\max\{d(u), d(v)\} \geq (n + 1)/2$ . For a given graph  $R$ ,  $G$  is called  $R$ - $f_1$ -heavy if every induced subgraph of  $G$  isomorphic to  $R$  is  $f_1$ -heavy. In this paper we show that for a connected graph  $S$  with  $S \neq P_3$  and a 2-connected claw- $f_1$ -heavy graph  $G$  which is not a cycle,  $G$  being  $S$ - $f_1$ -heavy implies  $G$  is pancyclic if  $S = P_4, Z_1$  or  $Z_2$ , where claw is  $K_{1,3}$  and  $Z_i$  is the path  $a_1a_2a_3 \dots a_{i+2}a_{i+3}$  plus the edge  $a_1a_3$ . Our result partially improves a previous theorem due to Bedrossian on pancyclicity of 2-connected graphs.

**Keywords:** Fan-type heavy subgraph; Hamilton cycle; Cycle; Pancyclicity

**AMS Subject Classification (2010):** 05C38, 05C45

## 1 Introduction

We use Bondy and Murty [5] for terminology and notation not defined here and consider simple graphs only.

Let  $G$  be a graph and  $H$  be a subgraph. Let  $x, y$  be two vertices of  $V(H)$ . An  $(x, y)$ -path in  $H$  is a path  $P$  connecting  $x$  and  $y$  in  $H$ . The *distance* between  $x$  and  $y$  in  $H$ , denoted by  $d_H(x, y)$ , is the length of a shortest  $(x, y)$ -path in  $H$ . When there is no danger of ambiguity, we use  $d(x, y)$  instead of  $d_G(x, y)$ .

Let  $G$  be a graph on  $n$  vertices. For a given graph  $R$ ,  $G$  is called  $R$ -free if  $G$  contains no induced subgraph isomorphic to  $R$ , and  $R$ - $f_i$ -heavy if for every induced subgraph  $H$

---

\*Email-address: ningbo\_math84@mail.nwpu.edu.cn (B. Ning).




Figure 1: Graphs  $P_i, C_3, Z_i, B, N$  and  $W$

of  $G$  isomorphic to  $R$  and every pair of vertices  $u, v \in V(H)$ ,  $d_H(u, v) = 2$  implies that  $\max\{d(u), d(v)\} \geq (n + i)/2$ . For a family  $\mathcal{R}$  of graphs,  $G$  is called  $\mathcal{R}$ -free ( $\mathcal{R}$ - $f_i$ -heavy) if  $G$  is  $R$ -free ( $R$ - $f_i$ -heavy) for each  $R \in \mathcal{R}$ . In particular, similar as in [9], we use  $R$ - $f$ -heavy ( $\mathcal{R}$ - $f$ -heavy) instead of  $R$ - $f_0$ -heavy ( $\mathcal{R}$ - $f_0$ -heavy). Note that every  $\mathcal{R}$ -free graph is also  $\mathcal{R}$ - $f_1$ -heavy ( $\mathcal{R}$ - $f$ -heavy).

The bipartite graph  $K_{1,3}$  is called the *claw*. We say that its (only) vertex of degree 3 is the *center* and the other vertices are its *end vertices*. In this paper, we use the terminology *claw- $f_1$ -heavy* instead of  $K_{1,3}$ - $f_1$ -heavy.

A graph  $G$  on  $n$  vertices is said to be *Hamiltonian* if it contains a *Hamilton cycle*, i.e., a cycle containing all vertices of  $G$ , and *pancyclic* if  $G$  contains cycles of all lengths from 3 to  $n$ . Bedrossian [1] completely characterized all the pairs of forbidden subgraphs for a 2-connected graph to be Hamiltonian and to be pancyclic.

**Theorem 1** (Bedrossian [1]). *Let  $R$  and  $S$  be connected graphs with  $R, S \neq P_3$  and let  $G$  be a 2-connected graph. Then  $G$  being  $\{R, S\}$ -free implies  $G$  is Hamiltonian if and only if (up to symmetry)  $R = K_{1,3}$  and  $S = P_4, P_5, P_6, C_3, Z_1, Z_2, B, N$  or  $W$  (see Figure 1).*

**Theorem 2** (Bedrossian [1]). *Let  $R$  and  $S$  be connected graphs with  $R, S \neq P_3$  and let  $G$  be a 2-connected graph which is not a cycle. Then  $G$  being  $\{R, S\}$ -free implies  $G$  is pancyclic if and only if (up to symmetry)  $R = K_{1,3}$  and  $S = P_4, P_5, Z_1$  or  $Z_2$ .*

In 1984, Fan [6] presented a degree condition (so-called Fan's condition) involving every pair of vertices at distance two for a 2-connected graph to be Hamiltonian.

**Theorem 3** (Fan [6]). *Let  $G$  be a 2-connected graph on  $n$  vertices. If  $\max\{d(u), d(v)\} \geq n/2$  for every pair of vertices  $u, v$  such that  $d(u, v) = 2$ , then  $G$  is Hamiltonian.*




Figure 2: The Graph  $F_{4r}$

Obviously, Fan's condition is equivalent to every 2-connected  $P_3$ - $f$ -heavy graph is Hamiltonian. By restricting Fan's condition to some induced subgraphs of 2-connected graphs, Ning and Zhang [9] extended Theorem 1 as follows.

**Theorem 4** (Ning and Zhang [9]). *Let  $R$  and  $S$  be connected graphs with  $R, S \neq P_3$  and let  $G$  be a 2-connected graph. Then  $G$  being  $\{R, S\}$ - $f$ -heavy implies  $G$  is Hamiltonian if and only if (up to symmetry)  $R = K_{1,3}$  and  $S = P_4, P_5, P_6, Z_1, Z_2, B, N$  or  $W$ .*

In this paper, our aim is to find the corresponding Fan-type heavy subgraph conditions for a 2-connected graph to be pancyclic. First, from a well known result, we can deduce that every 2-connected  $P_3$ - $f_1$ -heavy graph is pancyclic.

**Theorem 5** (Benhocine and Wojda [3]). *Let  $G$  be a 2-connected graph on  $n \geq 3$  vertices. If  $G$  is  $P_3$ - $f$ -heavy, then  $G$  is pancyclic unless  $n = 4r$ ,  $r > 2$ , and  $G = F_{4r}$  (see Figure 2), or  $n$  is even and  $G = K_{n/2, n/2}$  or else  $n \geq 6$  is even and  $G = K_{n/2, n/2} - e$ .*

Furthermore, we can see that  $P_3$  is the only connected graph  $S$  such that every 2-connected  $S$ - $f_1$ -heavy graph is pancyclic. For details, see [7, Theorem 13]. So we can pose the following problem naturally.

**Problem 1.** Which two connected graphs  $R$  and  $S$  other than  $P_3$  imply that every 2-connected  $\{R, S\}$ - $f_1$ -heavy graph is pancyclic?

By Theorem 2, we know that  $R = K_{1,3}$  (up to symmetry) and  $S$  must be one of  $Z_1, Z_2, P_4$  and  $P_5$ .

In this paper, we mainly prove the following result.

**Theorem 6.** *Let  $G$  be a 2-connected graph which is not a cycle. If  $G$  is  $\{K_{1,3}, Z_2\}$ - $f_1$ -heavy, then  $G$  is pancyclic.*

As a corollary of Theorem 6, we have

**Theorem 7.** *Let  $G$  be a 2-connected graph which is not a cycle. If  $G$  is  $\{K_{1,3}, P_4\}$ - $f_1$ -heavy, then  $G$  is pancyclic.*

In [2], Bedrossian et al. proved a theorem as follows.

**Theorem 8** (Bedrossian, Chen and Schelp [2]). *Let  $G$  be a 2-connected graph on  $n$  vertices. If  $G$  is  $\{K_{1,3}, Z_1\}$ - $f$ -heavy, then  $G$  is pancyclic unless  $G = F_{4r}$  or  $G = K_{n/2, n/2}$  or  $G = K_{n/2, n/2} - e$  or else  $G$  is a cycle.*

By Theorem 8, we have

**Theorem 9.** *Let  $G$  be a 2-connected graph which is not a cycle. If  $G$  is  $\{K_{1,3}, Z_1\}$ - $f_1$ -heavy, then  $G$  is pancyclic.*

Combining with Theorems 6, 7 and 9, we obtain Theorem 10, which partially answers Problem 1.

**Theorem 10.** *Let  $S$  be a connected graph with  $S \neq P_3$  and let  $G$  be a 2-connected claw- $f_1$ -heavy graph which is not a cycle. Then  $G$  being  $S$ - $f_1$ -heavy implies  $G$  is pancyclic if  $S = P_4, Z_1$  or  $Z_2$ .*

The rest of this paper is organized as follows. In Section 2, we will give additional terminology and list some useful lemmas. The proof of Theorem 6 will be postponed to Section 3.

## 2 Preliminaries

In this section, we first introduce some additional terminology and notation and then present four lemmas which will be used in our proof of Theorem 6.

Let  $G$  be a graph and  $S$  be a subset of  $V(G)$ . We use  $G[S]$  to denote the subgraph of  $G$  induced by  $S$  and  $G - S$  to denote  $G[V(G) \setminus S]$ . In particular, if  $S = \{u\}$ , then we use  $G - u$  instead of  $G - \{u\}$ . If  $S = \{x_i : 1 \leq i \leq 5\}$  and  $G[S]$  is isomorphic to  $Z_2$ , then we say that  $\{x_1, x_2, x_3, x_4, x_5\}$  induces a  $Z_2$ , where  $x_1x_2x_3x_4$  is a triangle and  $x_1$  is the vertex of degree 3 in  $G[S]$ . If  $S = \{x_i : 1 \leq i \leq 4\}$  and  $G[S]$  is isomorphic to  $K_{1,3}$ , then we say that  $\{x_1, x_2, x_3, x_4\}$  induces a claw, where  $x_1$  is the center, and  $x_2, x_3, x_4$  are the end vertices.

Let  $k, l$  ( $k < l$ ) be two integers. We say that  $G$  contains a  $k$ -cycle if  $G$  contains a cycle of length  $k$ , and  $G$  contains  $[k, l]$ -cycles if  $G$  contains cycles of all lengths from  $k$  to  $l$ . In

particular, for a vertex  $u \in V(G)$ , we say that  $G$  contains a  $u$ -triangle if  $G$  contains the cycle  $uxyu$ , where  $x, y \in V(G)$ .

A vertex  $v$  of a graph  $G$  on  $n$  vertices is called *heavy* if  $d(v) \geq n/2$ , and *super-heavy* if  $d(v) \geq (n+1)/2$ . For two vertices  $u, v$  of  $G$ ,  $\{u, v\}$  is called a *heavy-pair* if  $d(u) + d(v) \geq n$  and a *super-heavy pair* if  $d(u) + d(v) \geq n + 1$ .

**Lemma 1** (Benhocine and Wojda [3]). *Let  $G$  be a graph on  $n \geq 4$  vertices and  $C$  be a cycle of length  $n - 1$  in  $G$ . If  $d(x) \geq n/2$  for the vertex  $x \in V(G) \setminus V(C)$ , then  $G$  is pancyclic.*

**Lemma 2** (Bondy [4]). *Let  $G$  be a graph on  $n$  vertices with a Hamilton cycle  $C$ . If there exist two vertices  $x, y \in V(G)$  such that  $d_C(x, y) = 1$  and  $d(x) + d(y) \geq n + 1$ , then  $G$  is pancyclic.*

**Lemma 3** (Hakimi and Schmeichel [10]). *Let  $G$  be a graph on  $n$  vertices with a Hamilton cycle  $C$ . If there exist two vertices  $x, y \in V(G)$  such that  $d_C(x, y) = 1$  and  $d(x) + d(y) \geq n$ , then  $G$  is pancyclic unless  $G$  is bipartite or else  $G$  is missing only an  $(n - 1)$ -cycle.*

**Lemma 4** (Ferrara, Jacobson and Harris [8]). *Let  $G$  be a graph on  $n$  vertices with a Hamilton cycle  $C$ . If there exist two vertices  $x, y \in V(G)$  such that  $d_C(x, y) = 2$  and  $d(x) + d(y) \geq n + 1$ , then  $G$  is pancyclic.*

### 3 Proof of Theorem 6

We prove Theorem 6 by contradiction. Suppose that  $G$  satisfies the condition of Theorem 6 but is not pancyclic. Since the result is easy to verify for  $3 \leq n \leq 5$ , we assume that  $n \geq 6$ .

If  $G$  is  $\{K_{1,3}, Z_2\}$ -free, then by Theorem 2,  $G$  is pancyclic. Thus we assume that  $G$  contains an induced claw or an induced  $Z_2$ . Therefore, there is a super-heavy vertex, say  $u \in V(G)$ . Set  $G' = G - u$ . Since  $G$  is  $\{K_{1,3}, Z_2\}$ - $f_1$ -heavy,  $G'$  is  $\{K_{1,3}, Z_2\}$ - $f$ -heavy. If  $G'$  is 2-connected, then by Theorem 4,  $G'$  is Hamiltonian. Hence  $G$  is pancyclic by Lemma 1. Now, it will be assumed that  $G'$  is not 2-connected. Then there exists a vertex  $v \in V(G)$  ( $v \neq u$ ) such that  $G - \{u, v\}$  is not connected. By Theorem 4,  $G$  is Hamiltonian. Hence  $G - \{u, v\}$  consists of two components  $H_1$  and  $H_2$ . Without loss of generality, we assume that  $|V(H_1)| \leq |V(H_2)|$ , where  $V(H_1) = \{x_1, x_2, \dots, x_{h_1}\}$  and  $V(H_2) = \{y_1, y_2, \dots, y_{h_2}\}$ . Let  $C = uy_1 \cdots y_{h_2} vx_{h_1} \cdots x_1 u$  be a Hamilton cycle with the given orientation. In the following, for any two vertices  $w_1, w_2 \in V(C)$ , we use  $C[w_1, w_2]$  to denote the segment of  $C$  from  $w_1$  to  $w_2$  along the orientation. Set  $G_1 = G[V(H_1) \cup \{u\}]$  and  $G_2 = G[V(H_2) \cup \{u\}]$ .

**Claim 1.** There are no super-heavy vertices in  $H_1$ .

*Proof.* For any vertex  $x \in V(H_1)$ ,  $x$  is adjacent to at most  $u, v$  and all the vertices in  $H_1$  except for itself. Therefore,  $d(x) \leq d_{H_1}(x) + 2 \leq h_1 - 1 + 2 \leq n/2$ . Hence  $H_1$  contains no super-heavy vertices.  $\square$

**Claim 2.**  $N_{G_2}(u) \setminus \{y_1\} \subseteq N(y_1)$ .

*Proof.* If there exists a vertex  $y_i \in N_{G_2}(u) \setminus \{y_1\}$  such that  $y_i y_1 \notin E(G)$ , then  $\{u; x_1, y_1, y_i\}$  induces a claw. By Claim 1,  $x_1$  is not super-heavy. Since  $G$  is claw- $f_1$ -heavy,  $y_1$  is super-heavy. Hence  $\{u, y_1\}$  is a super-heavy pair such that  $d_C(u, y_1) = 1$ . By Lemma 2,  $G$  is pancyclic.  $\square$

**Claim 3.** There are no super-heavy pairs with distance one or two along the orientation of a Hamilton cycle in  $G$ .

*Proof.* Suppose not. By Lemma 2 or 4,  $G$  is pancyclic.  $\square$

**Case 1.**  $h_1 = 1$ .

**Subcase 1.1.**  $uv \in E(G)$ .

Note that  $G$  cannot be bipartite or missing an  $(n - 1)$  cycle, so if Lemma 3 applies to  $G$  then  $G$  is pancyclic. If  $u$  is adjacent to every vertex in  $C$ , then  $G$  is pancyclic. Now we can choose a vertex  $y_i \in N_{G_2}(u)$  such that  $u y_{i+1} \notin E(G)$ . Let  $y_j$  be the first vertex on  $C[y_i, y_{h_2}]$  such that  $u y_{j+1} \in E(G)$ , where assume that  $y_{h_2+1} = v$ . Obviously,  $j \geq i + 1$ .

**Claim 4.**  $i \geq 2$ .

*Proof.* Assume there exists  $y \in V(H_2)$  such that  $y_1 y \in E(G)$  and  $u y \notin E(G)$ . By Claim 2, we have  $N_{G_2}(u) \setminus \{y_1\} \subset N(y_1)$ . Since  $d(u) \geq (n + 1)/2$  and  $u, y \in N(y_1) \setminus N(u)$ ,  $d(y_1) \geq d(u) - 3 + 2 \geq (n - 1)/2$ . Therefore,  $\{u, y_1\}$  is a heavy-pair such that  $d_C(u, y_1) = 1$ . By Lemma 3,  $G$  is pancyclic. Also, since  $y_1 y_2 \in E(G)$ , then  $u y_2 \in E(G)$  and  $i \geq 2$ .  $\square$

Next we assume that  $i \leq h_2 - 2$ . Note that  $y_i, y_{i+1}, y_{i+2} \in C[y_2, y_{h_2}]$ .

**Claim 5.**  $j \geq i + 2$ .

*Proof.* Assume that  $j = i + 1$ . First, we have  $u y_i, u y_{i+2} \in E(G)$  and  $u y_{i+1} \notin E(G)$ .

If  $y_i y_{i+2} \notin E(G)$ , then  $\{u; x_1, y_i, y_{i+2}\}$  induces a claw. Since  $d(x_1) = 2 < (n + 1)/2$  and  $G$  is claw- $f_1$ -heavy,  $\{y_i, y_{i+2}\}$  is a super-heavy pair such that  $d_C(y_i, y_{i+2}) = 2$ , which contradicts to Claim 3.

Now assume that  $y_i y_{i+2} \in E(G)$ . If  $y_1 y_{i+1} \in E(G)$ , then it follows  $d(y_1) \geq (n-1)/2$  from Claim 2. Hence  $\{u, y_1\}$  is a heavy pair with  $d_C(u, y_1) = 1$ , and  $G$  is pancyclic by Lemma 3. Therefore,  $y_1 y_{i+1} \notin E(G)$ . We set  $G' = G - y_i$ . Clearly,  $C' = C[y_{i+2}, y_i] y_i y_{i+2}$  is a Hamilton cycle in  $G'$ . Moreover,  $u, y_1$  satisfy that  $d_{G'}(u) + d_{G'}(y_1) = d(u) + d(y_1) \geq (n+1)/2 + (n-3)/2 = n-1$  and  $d_{G'}(u, y_1) = 1$ . By Lemma 3,  $G'$  is pancyclic. Together with the cycle  $C$ ,  $G$  is pancyclic.  $\square$

By Claim 5, we obtain  $u y_{i+2} \notin E(G)$ .

**Claim 6.**  $v y_{i+1} \in E(G)$ .

*Proof.* Assume that  $v y_{i+1} \notin E(G)$ .

**Claim 6.1.**  $v y_{i+2} \notin E(G)$ .

*Proof.* Assume that  $v y_{i+2} \in E(G)$ . Then  $\{v, x_1, u; y_{i+2}, y_{i+1}\}$  induces a  $Z_2$ . If  $v$  is a super-heavy vertex, then  $\{u, v\}$  is a super-heavy pair such that  $d_C(u, v) = 2$ , contradicting to Claim 3. Now assume that  $v$  is not super-heavy. Note that  $x_1$  is not super-heavy. Since  $G$  is  $Z_2$ - $f_1$ -heavy,  $\{y_{i+1}, y_{i+2}\}$  is a super-heavy pair such that  $d_C(y_i, y_{i+1}) = 1$ , contradicting to Claim 3.  $\square$

**Claim 6.2.**  $v y_i \notin E(G)$ .

*Proof.* Assume that  $v y_i \in E(G)$ . By Claim 6.1, we have  $v y_{i+2} \notin E(G)$ . Note that  $v y_{i+1} \notin E(G)$  by the initial hypothesis. If  $y_i y_{i+2} \notin E(G)$ , then  $\{y_i, u, v; y_{i+1}, y_{i+2}\}$  induces a  $Z_2$ . Since  $v$  is not super-heavy,  $y_{i+1}$  is super-heavy. Hence either  $\{y_{i+1}, y_{i+2}\}$  or  $\{y_{i+1}, y_i\}$  is a super-heavy pair, a contradiction by Claim 3. If  $y_i y_{i+2} \in E(G)$ , then  $\{y_i, y_{i+1}, y_{i+2}; v, x_1\}$  induces a  $Z_2$ . Since  $v$  is not super-heavy,  $\{y_{i+1}, y_{i+2}\}$  is a super-heavy pair such that  $d_C(y_i, y_{i+1}) = 1$ , a contradiction by Claim 3.  $\square$

**Claim 6.3.**  $y_i$  is super-heavy.

*Proof.* By Claims 6.2 and the initial hypothesis, we have  $v y_i \notin E(G)$  and  $v y_{i+1} \notin E(G)$ . Since  $\{u, v, x_1; y_i, y_{i+1}\}$  induces a  $Z_2$  and  $x_1$  is not super-heavy,  $y_i$  is super-heavy.  $\square$

By Claim 4, we have  $i \geq 2$ , and this implies  $y_{i-1}$  is well-defined.

**Claim 6.4.**  $y_{i-1} y_{i+1} \notin E(G)$ ,  $u y_{i-1} \in E(G)$ ,  $y_i y_{i+2} \notin E(G)$  and  $y_{i-1} y_{i+2} \notin E(G)$ .

*Proof.* By Claim 6.3,  $y_i$  is super-heavy. If  $y_{i-1} y_{i+1} \in E(G)$ , then  $G$  is pancyclic by Lemma 1.

If  $uy_{i-1} \notin E(G)$ , then  $\{y_i; y_{i-1}, y_{i+1}, u\}$  induces a claw. Hence either  $y_{i-1}$  or  $y_{i+1}$  is super-heavy. Therefore, either  $\{y_{i-1}, y_i\}$  or  $\{y_i, y_{i+1}\}$  is a super-heavy pair such that  $d_C(y_{i-1}, y_i) = d_C(y_i, y_{i+1}) = 1$ , a contradiction by Claim 3.

By Claim 2 and Lemma 3,  $y_1y_{i+1} \notin E(G)$ . If  $y_iy_{i+2} \in E(G)$ , then set  $G' = G - y_{i+1}$ . Now  $C' = vx_1uy_1 \dots y_iy_{i+2} \dots y_{h_2}v$  is a Hamilton cycle in  $G'$ , and  $d_{G'}(u) + d_{G'}(y_1) \geq n - 1 = |G'|$  by Claim 2. By Lemma 3,  $G'$  is either pancyclic, bipartite, or missing only an  $(n - 2)$ -cycle. Since  $C' = vx_1uy_1 \dots y_iy_{i+2} \dots y_{h_2}v$  is an  $(n - 1)$ -cycle and  $C'' = vuy_1 \dots y_iy_{i+2} \dots y_{h_2}v$  is an  $(n - 2)$ -cycle in  $G'$ ,  $G'$  is pancyclic. Therefore,  $G$  is pancyclic.

If  $y_{i-1}y_{i+2} \in E(G)$ , then set  $G' = G - y_{i+1}$ . Now  $C' = uy_1 \dots y_{i-1}y_{i+2} \dots y_{h_2}vx_1u$  is a Hamilton cycle in  $G'' = G' - y_i$  and  $d_{G'}(y_i) \geq (n - 1)/2 = |G'|/2$ . By Lemma 1,  $G'$  is pancyclic. Together with the cycle  $C$ ,  $G$  is pancyclic.  $\square$

By Claim 6.4,  $\{y_i, u, y_{i-1}; y_{i+1}, y_{i+2}\}$  induces a  $Z_2$ . Since  $G$  is  $Z_2$ - $f_1$ -heavy, either  $y_{i-1}$  or  $y_{i+1}$  is super-heavy. Then either  $\{y_{i-1}, y_i\}$  or  $\{y_{i+1}, y_i\}$  is a super-heavy pair such that  $d_C(y_{i-1}, y_i) = d_C(y_{i+1}, y_i) = 1$ . By Claim 3, a contradiction.  $\square$

**Claim 7.** For any  $k \in \{i + 1, \dots, j\}$ ,  $vy_k \in E(G)$ .

*Proof.* By Claim 6, we have  $vy_{i+1} \in E(G)$ . Now we show that  $vy_k \in E(G)$  for any  $k \in \{i + 2, \dots, j\}$ . Otherwise, assume that  $y_t$  is the first vertex on  $C[y_{i+2}, y_j]$  such that  $vy_t \notin E(G)$ . Note that for any  $k \in \{i + 1, \dots, j\}$ ,  $uy_k \notin E(G)$ . We have  $uy_{t-1}, uy_t \notin E(G)$ . Then  $\{v, x_1, u; y_{t-1}, y_t\}$  induces a  $Z_2$ . Since  $x_1, v$  are not super-heavy,  $\{y_{t-1}, y_t\}$  is a super-heavy pair such that  $d_C(y_{t-1}, y_t) = 1$ . By Claim 3, a contradiction, hence  $vy_k \in E(G)$ .  $\square$

Note that since  $j \geq i + 2$  and  $i$  could be selected to be  $\leq h_2 - 2$ , then if  $(j + 1) \leq h_2 - 2$ , let  $i = j + 1$  and repeat the previous arguments to conclude that for any vertex  $y \in \{y_2, y_3, \dots, y_{h_2-2}\}$  such that  $uy \notin E(G)$ , we have  $vy \in E(G)$ . Hence  $d_{C[y_1, y_{h_2-2}]}(u) + d_{C[y_1, y_{h_2-2}]}(v) \geq h_2 - 2$ . If  $uy_{h_2-1} \in E(G)$  or  $vy_{h_2-1} \in E(G)$  or  $uy_{h_2} \in E(G)$ , then  $d_{G_2}(u) + d_{G_2}(v) \geq h_2 = n - 3$ . This implies that  $d(u) + d(v) \geq n + 1$ . By Claim 3, a contradiction. Otherwise, assume that  $uy_{h_2-1}, uy_{h_2} \notin E(G)$  and  $vy_{h_2-1} \notin E(G)$ . Then  $\{v, x_1, u; y_{h_2}, y_{h_2-1}\}$  induces a  $Z_2$ . It follows that  $\{y_{h_2}, y_{h_2-1}\}$  is a super-heavy pair such that  $d_C(y_{h_2-1}, y_{h_2}) = 1$ , contradicting to Claim 3.

**Subcase 1.2.**  $uv \notin E(G)$ .

By Claim 2,  $N_{G_2}(u) \setminus \{y_1\} \subseteq N(y_1)$ . If  $uy_2 \notin E(G)$ , then since  $u$  is super-heavy and  $u, y_2 \in N(y_1) \setminus N(u)$ ,  $y_1$  is super-heavy. Hence  $\{u, y_1\}$  is a super-heavy pair such that  $d_C(u, y_1) = 1$ , a contradiction by Claim 3. If  $uy_2 \in E(G)$ , then we have  $d(y_1) \geq (n - 1)/2$  and  $\{u, y_1\}$  is a heavy-pair such that  $d_C(u, y_1) = 1$ . By Lemma 3,  $G$  is either pancyclic,



bipartite, or missing only an  $(n - 1)$ -cycle. The cycle  $uy_1y_2u$  (a triangle) is odd, so  $G$  is not bipartite. Since  $C' = ux_1vy_{h_2}, \dots, y_2u$  is an  $(n - 1)$ -cycle,  $G$  is pancyclic.

**Case 2.**  $h_1 \geq 2$ .

**Subcase 2.1.**  $G_1$  contains a  $u$ -triangle.

Without loss of generality, we denote a  $u$ -triangle in  $G_1$  by  $ux_kx_{k'}u$  where  $k < k'$ .

**Subsubcase 2.1.1.**  $u$  is not adjacent to every vertex of  $H_2$ .

Let  $y_i \in V(H_2)$  be the vertex such that  $uy_i \notin E(G)$  and  $i$  is as small as possible. Note that  $\{u, x_k, x_{k'}; y_{i-1}, y_i\}$  induces a  $Z_2$ . By Claim 1,  $y_{i-1}$  is super heavy. So if  $i = 2$  then  $\{u, y_1\}$  is a super-heavy pair such that  $d_C(u, y_1) = 1$ , a contradiction by Claim 3. Therefore  $i \geq 3$  and  $uy_2 \in E(G)$ .

If there exists  $t \in \{1, 2, \dots, h_1 - 1\}$  such that  $ux_t \in E(G)$  and  $ux_{t+1} \notin E(G)$ , then  $\{u, y_1, y_2; x_t, x_{t+1}\}$  induces a  $Z_2$ . Note that  $x_t$  is not super-heavy. Since  $G$  is  $Z_2$ - $f_1$ -heavy,  $y_1$  is super-heavy. Hence  $\{u, y_1\}$  is a super-heavy pair such that  $d_C(u, y_1) = 1$ , contradicting to Claim 3. Therefore,  $u$  is adjacent to every vertex of  $H_1$ . Note that  $C' = ux_1 \cdots x_i u$  is an  $(i + 1)$ -cycle, where  $2 \leq i \leq h_1$ , and  $G$  contains  $[3, h_1 + 1]$ -cycles. If  $i = h_2$ , then  $u$  is adjacent to every vertex of  $H_2$  other than  $y_{h_2}$ . It follows  $G$  contains  $[h_1 + 4, n]$ -cycles. Furthermore,  $C' = ux_2 \cdots x_{h_1}vy_{h_2}y_{h_2-1}u$  is an  $(h_1 + 3)$ -cycle. If  $h_1 \geq 3$ , then  $C' = ux_3 \cdots x_{h_1}vy_{h_2}y_{h_2-1}u$  is an  $(h_1 + 2)$ -cycle, and  $G$  is pancyclic. If  $h_1 = 2$  and  $h_2 \geq 4$ , then we can easily find a 4-cycle in  $G$ , and  $G$  is pancyclic. If  $h_1 = 2$  and  $h_2 = 2$  or 3, then  $n = 6$  or 7. In these two cases, the result is easy to verify.

Now we suppose that  $3 \leq i \leq h_2 - 1$  and try to get a contradiction. If there exists  $y_k \in N_{G_2}(u)$  such that  $y_k y_{i-2} \notin E(G)$  and  $y_k \neq y_{i-2}$ , then  $\{u; x_1, y_k, y_{i-2}\}$  induces a claw. Since  $G$  is claw- $f_1$ -heavy and  $x_1$  is not super-heavy,  $y_{i-2}$  is super-heavy. Therefore,  $\{y_{i-2}, y_{i-1}\}$  is a super-heavy pair such that  $d_C(y_{i-2}, y_{i-1}) = 1$ , a contradiction by Claim 3. So,  $N_{G_2}(u) \setminus \{y_{i-2}\} \subseteq N(y_{i-2})$ .

If  $uv \in E(G)$ , then we set  $G' = G - V(H_1)$ . Since  $N(u) \cup \{u\} \setminus (V(H_1) \cup \{v, y_{i-2}\}) \subseteq N(y_{i-2})$ , we have  $d(y_{i-2}) \geq d(u) + 1 - h_1 - 2 \geq (n + 1)/2 - h_1 - 1$ . Furthermore, we obtain  $d_{G'}(y_{i-2}) + d_{G'}(y_{i-1}) = d(y_{i-2}) + d(y_{i-1}) \geq n - h_1 = |G'|$ . Let  $C' = uv y_{h_2} \cdots y_1 u$ . Then  $C'$  is a Hamilton cycle in  $G'$  and  $d_{C'}(y_{i-2}, y_{i-1}) = 1$ . By Lemma 3,  $G'$  is either pancyclic, bipartite, or missing only a  $(|G'| - 1)$ -cycle. But  $G'$  contains the triangle  $uy_1y_2u$ , hence it is not bipartite. Note that  $G$  contains the cycle  $C'' = uv y_{h_2} \cdots y_2 u$  of length  $|G'| - 1$ . Hence  $G'$  is pancyclic, and this implies that  $G$  contains  $[3, |G'|]$ -cycles. Since  $u$  is adjacent to every vertex of  $H_1$ ,  $G$  contains  $[|G'| + 1, n]$ -cycles. Hence  $G$  is pancyclic.

If  $uv \notin E(G)$ , then we set  $G' = G - (V(H_1) \setminus \{x_{h_1}\})$ . Now we have  $d(y_{i-2}) \geq d(u) - h_1 - 1 + 1 \geq (n+1)/2 - h_1$ . And we obtain  $d_{G'}(y_{i-2}) + d_{G'}(y_{i-1}) \geq d(y_{i-2}) + d(y_{i-1}) \geq n+1 - h_1 = |G'|$ . Similarly, we can prove that  $G$  is pancyclic.

**Subsubcase 2.1.2.**  $u$  is adjacent to every vertex of  $H_2$ .

Note that  $uy_1y_2u$  is a  $u$ -triangle. If there exists a vertex  $x_t \in V(H_1)$  such that  $ux_t \in E(G)$  and  $ux_{t+1} \notin E(G)$ , then  $\{u, y_1, y_2; x_t, x_{t+1}\}$  induces a  $Z_2$ . This implies that  $y_1$  is super-heavy. Hence  $\{u, y_1\}$  is a super-heavy pair such that  $d_C(u, y_1) = 1$ , a contradiction by Claim 3. If  $u$  is adjacent to every vertex in  $H_1$ , then  $u$  is adjacent to all vertices of  $V(G) \setminus \{u, v\}$ . This implies that  $d(u) \geq n-2$ , and  $d(u) + d(y_1) \geq n$ . By Lemma 3,  $G$  is either pancyclic, bipartite, or missing only an  $(n-1)$ -cycle. Since  $u$  is adjacent to every vertex in  $H_2$ ,  $G$  is neither bipartite nor missing  $(n-1)$ -cycles. It follows that  $G$  is pancyclic.

**Subcase 2.2.**  $G_1$  contains no  $u$ -triangles.

We first show that  $N_{G_1}(u) = \{x_1\}$ . Suppose not. If there is a vertex  $x \in N_{G_1}(u)$  such that  $x \neq x_1$ , then since  $G_1$  contains no  $u$ -triangles, we have  $xx_1 \notin E(G)$ . Now  $\{u; x, x_1, y_1\}$  induces a claw. It follows that either  $x$  or  $x_1$  is super-heavy, which contradicts to Claim 1.

If there exist two consecutive vertices, say  $y_i, y_{i+1} \in V(H_2)$ , such that  $uy_i, uy_{i+1} \in E(G)$ , then  $\{u, y_i, y_{i+1}; x_1, x_2\}$  induce a  $Z_2$ . Hence  $\{y_i, y_{i+1}\}$  is a super-heavy pair such that  $d_C(y_i, y_{i+1}) = 1$ , a contradiction by Claim 3.

Therefore for any  $y_i \in V(H_2) \setminus \{y_{h_2}\}$ ,  $|\{uy_i, uy_{i+1}\} \cap E(G)| \leq 1$ . This implies that  $u$  is adjacent to only one vertex  $x_1$  in  $H_1$  and at most  $(h_1+1)/2$  vertices in  $H_2$  and maybe adjacent to  $v$  or not. Hence we have  $(n+1)/2 \leq d(u) \leq 1 + 1 + (h_2+1)/2$ . This implies that  $h_2 \geq n-4$ . Noting that  $h_2 = n-2-h_1 \leq n-2-2 = n-4$ , we have  $h_2 = n-4, h_1 = 2, uv \in E(G)$  and  $N_{G_2}(u) = \{y_{2k+1} : k = 0, 1, \dots, (n-5)/2\}$ , where  $n$  is odd.

If  $y_1y_3 \notin E(G)$ , then  $\{u; x_1, y_1, y_3\}$  induces a claw. Since  $G$  is claw- $f_1$ -heavy,  $\{y_1, y_3\}$  is a super-heavy pair such that  $d_C(y_1, y_3) = 2$ . By Claim 3, a contradiction.

If  $y_1y_3 \in E(G)$ , then  $\{u, y_1, y_3; x_1, x_2\}$  induces a  $Z_2$ . Since  $G$  is  $Z_2$ - $f_1$ -heavy,  $\{y_1, y_3\}$  is a super-heavy pair such that  $d_C(y_1, y_3) = 2$ . By Claim 3, also a contradiction.

The proof is complete.  $\square$

## Acknowledgements

This work was supported by NSFC (No. 11271300) and the Doctorate Foundation of Northwestern Polytechnical University (cx201326). The author is very indebted to Pro-

fessor E.F. Schmeichel for his kindness and helpful documents. The author would like to thank Dr. Jun Ge and Lu Qiao for many helpful discussions, and would like to express deep gratitude to editors and reviewers for their carefully reading the earlier version of this work and so many suggestions.

## References

- [1] P. Bedrossian, *Forbidden Subgraph and Minimum Degree Conditions for Hamiltonicity*, Ph.D. Thesis, Memphis State University, USA (1991).
- [2] P. Bedrossian, G. Chen and R.H. Schelp, A generalization of Fan's condition for Hamiltonicity, pancyclicity, and Hamiltonian connectedness, *Discrete Math.* **115** (1993), 39-50.
- [3] A. Benhocine and A.P. Wojda, The Geng-Hua Fan conditions for pancyclic or Hamilton-connected graphs, *J. Combin. Theory Ser. B* **42** (1987), 167-180.
- [4] J.A. Bondy, Pancyclic graphs I, *J. Combin. Theory Ser. B* **11** (1971), 80-84.
- [5] J.A. Bondy and U.S.R. Murty, *Graph Theory with Applications*, Macmillan London and Elsevier, New York (1976).
- [6] G. Fan, New sufficient conditions for cycles in graphs, *J. Combin. Theory Ser. B* **37** (1984), 221-227.
- [7] R.J. Faudree and R.J. Gould, Characterizing forbidden pairs for hamiltonian properties, *Discrete Math.* **173** (1997), 45-60.
- [8] M. Ferrara, M.S. Jacobson and A. Harris, Cycle lengths in Hamiltonian graphs with a pair of vertices having large degree sum, *Graphs and Combin.* **26** (2010), 215-223.
- [9] B. Ning and S. Zhang, Ore- and Fan-type heavy subgraphs for Hamiltonicity of 2-connected graphs, *Discrete Math.* **313** (2013), 1715-1725.
- [10] E.F. Schmeichel and S.L. Hakimi, A cycle structure theorem for hamiltonian graphs, *J. Combin. Theorey Ser. B* **45** (1988), 99-107.